

Cal/Ecotox Exposure Factors for Brown Pelican (*Pelecanus occidentalis*)*

Page 1

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Age at Fledging, Metamorphosis, Weaning			74-77	days	NR	Fledgling	FL	a	1
Age at Sexual Maturity	3			yr	NR	Adult	LA	b	2
Body Weight - Mean			3,300-3,500	g	NR	Adult	FL	c	1
Body Weight - Mean	27.6	1.2 SE		g	B	Hatchling	CANADA	d	3
Body Weight - Mean	1698.4	35.2 SE		g	B	Nestling	CANADA	e	3
Body Weight - Mean			45-80	g	NR	Nestling	FL	f	1
Clutch or Litter Size	2.85		2-4	eggs	F	Adult	SC	g	4
Clutch or Litter Size	3.2	0.2 SE		eggs	F	Adult	CANADA	h	3
Clutch or Litter Size	2.16		1-3	eggs	F	Juvenile	SC	i	4
Dietary Composition	northern anchovy, <i>Engraulis mordax</i> (80%)				NR	Nestling	CA	j	5
Fledging or Weaning Rate	2.3	0.2		fledglings/nest	B	Fledgling	CANADA	k	3
Fledging or Weaning Rate			0.18-0.88		NR	Fledgling	CA	l	6
Fledging or Weaning Rate			0.23-1.20		NR	Fledgling	CA	m	6
Food Ingestion Rate			16-18%		NR	Both Adult and Juv.	FL	n	1
Foraging Distance			10.2-49.4	km/day	NR	Juvenile	CA	o	7
Growth Rate	90.5	1.8 SE		g/day	B	Nestling	CANADA	p	3
Growth Rate	see figure				NR	Nestling	FL	q	1
Hatching Success	60.2%				F	Adult	SC	r	4
Hatching Success	2.8	0.2 SE		eggs/nest	B	Hatchling	CANADA	s	3
Hatching Success	57.1%				F	Juvenile	SC	t	4
Metabolic Rate	75		75-76	cal/kg/d	NR	Adult	Lab	u	8
Population Density	0.23	0.23 SE		birds/km ²	B	Both Adult and Juv.	CA	v	9
Survival/ Mortality	94%				NR	Adult	CA	w	10
Survival/ Mortality	69%				NR	Adult	CA	x	10
Time of Migration or Dispersal	early to mid-summer				B	Both Adult and Juv.	CA	y	9

Notes

- a range of mean fledging ages over four years; N=7-12 birds/year; March-July; Tarpon Key
- b age of first nesting; N=13 nests; Age=3 yr; March; Barataria Bay
- c body weight of captive adult; N=1 bird; March-July; Tarpon Key
- d hatch weight; N=11 chicks; St. Lawrence estuary
- e asymptotic weight (40 days); N=65 chicks; St. Lawrence estuary
- f body weight at hatching; N=47 birds; March-July; Tarpon Key; asymptotic weight reached at about 50 days
- g mean clutch size; N=89 clutches; Cape Romain National Wildlife Refuge
- h mean clutch size among successful nests; N=32 nests; St. Lawrence estuary
- i mean clutch size; N=37 clutches; Cape Romain National Wildlife Refuge
- j percent of estimated prey biomass, from pre-fledgling regurgitations; N=3 colonies; Southern California Bight
- k fledging rate among successful nests; N=32 nests; St. Lawrence estuary
- l range of mean fledging rates (number of fledglings per built nest) over a 6 year period
; N=76-1258 nests/year; West Anacapa Island; fledging success strongly related to prey biomass and density near the breeding colony
- m range of mean fledging rates (number of fledglings per built nest) over a 6 year period
; N=76-1258 nests/year; Isla Coronado Norte; fledging success strongly related to prey biomass and density near the breeding colony
- n percent of body weight consumed daily during maintenance of adult weight; N=4 birds; March-July; Tarpon Key

Exposure Factors for Brown Pelican (*Pelecanus occidentalis*)*

Page 2

o distance travelled by radiotagged bird; N=1 bird; October; south Monterey Bay
p growth rate during maximum growth phase; N=65 chicks; St. Lawrence estuary
q daily growth rates of wild nestlings; N=varied; March-July; Tarpon Key
r percent of eggs producing downy young in successful nests (nests producing at least one young); N=133 nests; Cape Romain National Wildlife Refuge
s proportion of eggs hatched per nest among successful nests; N=32 nests; St. Lawrence estuary
t percent of eggs producing downy young in successful nests (nests producing at least one young); N=7 nests; Cape Romain National Wildlife Refuge
u average basal metabolism (heat production); N=1 bird (3 measurements); Condition=fasted
v mean density at sea off central and northern coast; N=3 years; central and northern coast
w estimated survival rate of radiotagged control birds at 150-180 days post-release; N=17; Southern California Bight
x estimated survival rate of radiotagged rehabilitated oiled birds at 150-180 days post-release; N=16; Condition=rehabilitated oiled birds; Southern California Bight
y time of northward movement from southern breeding areas; N=3 years; northern and central coast

References

- 1 Schreiber, Ralph W. 1976. Growth and development of nestling brown pelicans. *Bird-Band*. 47(1):19-39.
- 2 Williams, Lovett E., Jr. and Ted Joanen. 1974. Age of first nesting in the brown pelican. *Wilson Bull.* 86(3):279-280.
- 3 Desgranges, Jean-Luc. 1982. Weight growth of young double-crested cormorants in the St. Lawrence Estuary, Quebec. *Colon. Waterbirds*. 5:79-86.
- 4 Blus, Lawrence J. and Julie A. Keahey. 1978. Variation in reproductivity with age in the brown pelican. *Auk*. 95:128-134.
- 5 Kelly, P.R., Franklin Gress and D.W. Anderson. 1981. Dietary composition of brown pelicans breeding in the southern California bight. *Pac. Seabird Group Bull.* 8(2):105.
- 6 Anderson, Daniel W., Franklin Gress, and Kenneth F. Mais. 1982. Brown pelicans: influence of food supply on reproduction. *Oikos*. 39:23-31.
- 7 Croll, Donald A., Lisa T. Ballance, Bernd G. Wursig and Wm. Breck Tyler. 1986. Movements and daily activity patterns of a brown pelican in Central California. *Condor*. 88:258-260.
- 8 Benedict, Francis G., and Edward L. Fox. 1927. The gaseous metabolism of large wild birds under aviary life, In: *American Philosophical Society. The Record of the Celebration of the Two Hundredth Anniversary of the Founding of the American Philosophical Society*; April 27-30, 1927, Philadelphia. Philadelphia, PA: American Philosophical Society. p 511-534.
- 9 Briggs, Kenneth T., Wm. Breck Tyler, David B. Lewis, Paul R. Kelly and Donald A. Croll. 1983. Brown pelicans in central and northern California. *J. Field Ornithol.* 54(4):353-373.
- 10 Anderson, Daniel W., Franklin Gress, and D. Michael Fry. 1996. Survival and dispersal of oiled brown pelicans after rehabilitation and release. *Mar. Pollut. Bull.* 32(10):711-718.

*Cal/EPA, OEHHA and the University of California Regents are not responsible for damages of any kind resulting from the use of or reliance on information in this report. Users are encouraged to consult the original data. Updated: February 1999.