

DRAFT

**Public Health Goal for
1,2,3-Trichloropropane
in Drinking Water**

Prepared by

**Pesticide and Environmental Toxicology Branch
Office of Environmental Health Hazard Assessment
California Environmental Protection Agency**

September 2007

LIST OF CONTRIBUTORS

PHG PROJECT MANAGEMENT

REPORT PREPARATION

SUPPORT

Project Director

Anna Fan, Ph.D.

Authors

Richard Sedman, Ph.D.

Administrative Support

Hermelinda Jimenez

Michael Baez

Sharon Davis

PHG Program Leader

Robert A. Howd, Ph.D.

Primary Reviewers

Dan Sultana, M.S.

Sara Hoover, M.S.

Library Support

Charleen Kubota, M.L.S.

Comment Coordinator

Thomas Parker, M.S.

Final Reviewers

Anna Fan, Ph.D.

George Alexeeff, Ph.D.

Robert Howd, Ph.D.

Web site Posting

Laurie Monserrat

PREFACE

**Drinking Water Public Health Goals
Pesticide and Environmental Toxicology Branch
Office of Environmental Health Hazard Assessment
California Environmental Protection Agency**

This Public Health Goal (PHG) technical support document provides information on health effects from contaminants in drinking water. PHGs are developed for chemical contaminants based on the best available toxicological data in the scientific literature. These documents and the analyses contained in them provide estimates of the levels of contaminants in drinking water that would pose no significant health risk to individuals consuming the water on a daily basis over a lifetime.

The California Safe Drinking Water Act of 1996 (Health and Safety Code, Section 116365) requires the Office of Environmental Health Hazard Assessment (OEHHA) to perform risk assessments and adopt PHGs for contaminants in drinking water based exclusively on public health considerations. The Act requires that PHGs be set in accordance with the following criteria:

1. PHGs for acutely toxic substances shall be set at levels at which no known or anticipated adverse effects on health will occur, with an adequate margin of safety.
2. PHGs for carcinogens or other substances that may cause chronic disease shall be based solely on health effects and shall be set at levels that OEHHA has determined do not pose any significant risk to health.
3. To the extent the information is available, OEHHA shall consider possible synergistic effects resulting from exposure to two or more contaminants.
4. OEHHA shall consider potential adverse effects on members of subgroups that comprise a meaningful proportion of the population, including but not limited to infants, children, pregnant women, the elderly, and individuals with a history of serious illness.
5. OEHHA shall consider the contaminant exposure and body burden levels that alter physiological function or structure in a manner that may significantly increase the risk of illness.
6. OEHHA shall consider additive effects of exposure to contaminants in media other than drinking water, including food and air, and the resulting body burden.
7. In risk assessments that involve infants and children, OEHHA shall specifically assess exposure patterns, special susceptibility, multiple contaminants with toxic mechanisms in common, and the interactions of such contaminants.

8. In cases of insufficient data for OEHHA to determine a level that creates no significant risk, OEHHA shall set the PHG at a level that is protective of public health with an adequate margin of safety.
9. In cases where scientific evidence demonstrates that a safe dose response threshold for a contaminant exists, then the PHG should be set at that threshold.
10. The PHG may be set at zero if necessary to satisfy the requirements listed above in items seven and eight.
11. PHGs adopted by OEHHA shall be reviewed at least once every five years and revised as necessary based on the availability of new scientific data.

PHGs adopted by OEHHA are for use by the California Department of Health Services (DHS) in establishing primary drinking water standards (State Maximum Contaminant Levels, or MCLs). Whereas PHGs are to be based solely on scientific and public health considerations without regard to economic cost considerations or technical feasibility, drinking water standards adopted by DHS are to consider economic factors and technical feasibility. Each primary drinking water standard adopted by DHS shall be set at a level that is as close as feasible to the corresponding PHG, placing emphasis on the protection of public health. PHGs established by OEHHA are not regulatory in nature and represent only non-mandatory goals. By state and federal law, MCLs established by DHS must be at least as stringent as the federal MCL, if one exists.

PHG documents are used to provide technical assistance to DHS, and they are also informative reference materials for federal, state and local public health officials and the public. While the PHGs are calculated for single chemicals only, they may, if the information is available, address hazards associated with the interactions of contaminants in mixtures. Further, PHGs are derived for drinking water only and are not intended to be utilized as target levels for the contamination of other environmental media.

Additional information on PHGs can be obtained at the OEHHA web site at www.oehha.ca.gov.

TABLE OF CONTENTS

LIST OF CONTRIBUTORS	II
PREFACE	III
TABLE OF CONTENTS	V
PUBLIC HEALTH GOAL FOR 1,2,3-TRICHLOROPROPANE IN DRINKING WATER	1
SUMMARY	1
INTRODUCTION	2
Chemical Identity.....	2
Production and Uses	3
ENVIRONMENTAL OCCURRENCE AND HUMAN EXPOSURE	3
Air	3
Soil.....	4
Water.....	4
Food.....	5
Exposure	5
METABOLISM AND PHARMACOKINETICS	6
Absorption/Distribution/Elimination.....	6
Metabolism	7
TOXICOLOGY	10
Toxicological Effects in Animals	10
Acute Toxicity	10
Subchronic Toxicity.....	10
Genetic Toxicity	12
Developmental and Reproductive Toxicity	13
Immunotoxicity.....	14
Neurotoxicity	14
Chronic Toxicity/Carcinogenicity	15
Toxicological Effects in Humans	19

DRAFT

Acute Toxicity	19
Subchronic Toxicity.....	19
Neurotoxicity	20
Immunotoxicity.....	20
Developmental/Reproductive Toxicity.....	20
Chronic Toxicity	20
Evidence for Carcinogenic Activity of 1,2,3-Trichloropropane.....	20
Animal Studies.....	20
Human Studies.....	20
Genotoxicity.	20
Mechanism.....	20
Conclusion	21
DOSE-RESPONSE ASSESSMENT.....	21
Noncarcinogenic Effects.....	21
Carcinogenic Effects.....	21
CALCULATION OF PHG	24
Noncarcinogenic Effects.....	24
Carcinogenic Effects.....	26
RISK CHARACTERIZATION	27
OTHER REGULATORY STANDARDS.....	28
REFERENCES	30

PUBLIC HEALTH GOAL FOR 1,2,3-TRICHLOROPROPANE IN DRINKING WATER

SUMMARY

The Office of Environmental Health Hazard Assessment (OEHHA) has developed a proposed Public Health Goal (PHG) of 7×10^{-7} mg/L (0.0007 μ g/L, 0.0007 ppb or 0.7 ppt) for 1,2,3-trichloropropane (1,2,3-TCP). The proposed PHG is based on carcinogenic effects observed in animals (NTP, 1993). Significant increases in tumors have been observed in a number of tissues in male and female mice and rats administered 1,2,3-TCP by the oral route. 1,2,3-TCP was genotoxic in studies *in vivo* and *in vitro*. DNA adducts have been observed in animals following exposure to 1,2,3-TCP. Studies of 1,2,3-TCP metabolism, while incomplete, link 1,2,3-TCP genotoxic and carcinogenic activity with metabolism to a reactive form. Evidence that links exposure to 1,2,3-TCP to an increased incidence of cancer in humans is lacking.

A time-to-tumor model was employed to obtain a lower-bound estimate of the dose associated with a 10 percent increased incidence of tumors at various tumor sites in male and female rats and mice. Tumors in the female mouse forestomach yielded the lowest lower-bound estimate of the dose associated with a 10 percent increased incidence of tumors, and therefore was used to estimate cancer potency. A cancer potency for 1,2,3-TCP of $(24 \text{ mg/kg-day})^{-1}$ was derived using the lower bound dose associated with a 10 percent increased incidence of forestomach tumors and an assumed linear relationship between dose and response at low doses. The proposed PHG was derived employing this cancer potency and exposure to 4 L_{eq} /day of water.

A health protective level was also derived based on non-carcinogenic effects. A no observed adverse effect level (NOAEL) of 8 mg/kg-day at 8 and 17 weeks was identified from a 17 week subchronic gavage study, based on decreased erythrocyte mass (lower mean hematocrit, hemoglobin and erythrocyte counts) at doses of 16 mg/kg-day or greater in female rats. A health protective level of 0.08 mg/L or 80 ppb was derived employing an uncertainty factor of 1,000, and the ingestion of 4 L_{eq} /day of water.

Equivalent total exposure to 1,2,3-TCP (L_{eq} /day) was calculated using the CalTOX model to estimate the relative exposure to 1,2,3-TCP by the oral, dermal and inhalation routes. Because 1,2,3-TCP is a volatile organic compound, both dermal and inhalation exposure could be a significant contributor to the overall daily dose. While exposure via the dermal route was minimal (predicted to be approximately 2 percent of the overall exposure), the model predicted that inhalation exposure was equivalent to that from the oral route (2 liters/day). Therefore, the PHG was based on daily exposure to 4 L_{eq} of water.

There is no California or federal Maximum Contaminant Level (MCL) for 1,2,3-TCP. The California Notification Level for 1,2,3-TCP is 0.005 μ g/L, which is also the detection limit for the purposes of reporting (DLR). Notification levels are health-based advisory levels established by California Department of Public Health (DPH, formerly

the Department of Health Services [DHS]) for chemicals in drinking water that lack MCLs. DPH advises “If a chemical is over its notification level in drinking water that is provided to consumers, [DPH] recommends that the utility inform its customers and consumers about the presence of the contaminant, and about the health concerns associated with exposure to it” (<http://www.dhs.ca.gov/ps/ddwem/chemicals/al/PDFs/notificationoverview.pdf>).

INTRODUCTION


The purpose of this document is to develop a proposed PHG for 1,2,3-TCP. PHGs are based on a comprehensive analysis of information on the toxicology of the compounds. PHGs are based solely on protection of public health without regard to cost impacts or other factors. PHGs for carcinogens are set at a *de minimis* risk level of one in a million (10^{-6}), assuming a lifetime of exposure to the chemical in the drinking water. PHGs for non-carcinogens are based on levels estimated to be without risk of any adverse effects for exposures up to a lifetime, to the general population as well as any significant identifiable sensitive subpopulations.

A PHG for 1,2,3-TCP is being developed by OEHHA at the request of the California DPH, because of repeated detections of this chemical in California drinking water sources. The major source of the 1,2,3-TCP is presumably leaching from hazardous waste sites (DHS, 2005). This chemical is listed on the priority list of hazardous substances maintained by the federal Agency for Toxic Substances and Disease Registry (ATSDR, 2005). 1,2,3-TCP has been reported to cause cancer in laboratory animals (U.S. EPA, 1997), which is the basis for the DPH notification level of 0.005 $\mu\text{g/L}$ (DHS, 2005). It was also listed in 1992 as known to the State of California to cause cancer [Title 22, California Code of Regulations, Section 12000] and has been identified as reasonably anticipated to be a human carcinogen by the National Toxicology Program (NTP, 2005).

Available scientific information on 1,2,3-TCP was identified using U.S. EPA’s Integrated Risk Information System (IRIS), PubMed, and other relevant reference databases. Two reviews were identified and were particularly useful in identifying and evaluating relevant toxicity studies (WHO, 2003; ATSDR, 1992). There is no California or federal MCL for 1,2,3-TCP.

Chemical Identity

1,2,3-Trichloropropane is also known as allyl trichloride, trichlorohydrin, and glycerol trichlorohydrin. The structure of 1,2,3-trichloropropane shown below is from WHO (2003).



The physical and chemical properties of 1,2,3-trichloropropane are summarized below in Table 1.

Table 1. Physical and Chemical Properties of 1,2,3-Trichloropropane¹

Property	Value
CAS No.	96-18-4
Physical state	Liquid
Molecular weight	147.43
Density	1.3888 g/cm ³ at 20° C
Solubility in water	1759 mg/L at 20° C
Solubility in organic solvents	Soluble in ethyl alcohol, chloroform, ethyl ether, benzene
Vapor Pressure	3.1 mm Hg at 25° C
Henry's Law constant	3.17 x 10 ⁻⁴ atm-m ³ /mol at 25° C
Absorption to organic carbon (K _{oc})	77 to 95
Octanol-water partition coefficient (Log K _{ow})	2.27
Conversion factor:	1 ppm = 6.03 mg/m ³

¹from ATSDR, 1992; WHO, 2003; HSDB, 2005.

Production and Uses

1,2,3-Trichloropropane has been used as a solvent and degreasing agent and in the synthesis of other compounds such as epichlorohydrin and certain polymers (WHO, 2003; NTP, 2005). The continued use of 1,2,3-TCP as a solvent is unclear. The TSCA Inventory Update lists usage in 2002 as >1 to 10 million pounds, which is a decrease in usage from earlier reporting periods (U.S. EPA, 2007a). 1,2,3-trichloropropane also occurs as a byproduct in pesticides and the dichloropropene soil fumigants.

ENVIRONMENTAL OCCURRENCE AND HUMAN EXPOSURE

Air

Because of the relatively high volatility of 1,2,3-TCP, it will be readily transferred into air. Low levels of 1,2,3-TCP (<1 µg/m³) have been detected in ambient air in some studies but not in others (reviewed by WHO, 2003). Much higher levels were detected in ambient air in Kuwait (mean 491 µg/m³ from 19 samples) than in other areas sampled. 1,2,3-Trichloropropane will react with hydroxyl radicals in the atmosphere, with an estimated half-life of 46 days (HSDB, 2005). Because of its high water solubility, 1,2,3-TCP can also be removed from the atmosphere by rain. The combination of

volatilization, washout, and resistance to degradation may result in a recycling of 1,2,3-trichloropropane among the environmental compartments (HSDB, 2005).

Soil

Low levels of 1,2,3-TCP have been detected in soil at some hazardous waste sites (ATSDR, 1992). Its detection in groundwater near hazardous waste sites implies limited adsorption to soils (high mobility) and a low rate of biodegradation in soil and groundwater (HSDB, 2005). An INCHEM review cites limited reports of detection of 1,2,3-TCP in groundwater after its use as a nematocide (INCHEM, 2007).

Water

As discussed by HSDB (2005), 1,2,3-TCP should readily volatilize from surface water because of its high Henry's Law constant. Its half-life was estimated to be 6.7 hours in a river and 5.7 days in a lake (HSDB, 2005). Due to its low adsorptivity, it should not be substantially adsorbed to sediment or particulates in the water column. 1,2,3-Trichloropropane is resistant to biodegradation and hydrolysis, so breakdown is not expected to be an important fate process in the aquatic environment. Also, according to HSDB (2005), it would not be expected to bioconcentrate in aquatic organisms.

1,2,3-Trichloropropane has been detected (two or more times) in 83 water systems in 16 counties in California as of 2005. The number of detections and range of detections as reported by the California DPH are shown in Tables 2 and 3.

Table 2. 1,2,3-TCP Detections in California Groundwater Sources ^a

Reported 1,2,3-TCP Detections		
County	No. of Sources	No. of Systems
Kern	95	17
Los Angeles	35	15
Fresno	28	8
Tulare	23	4
San Bernardino	21	6
Merced	19	11
San Joaquin	8	2
Riverside	7	5
San Mateo	7	2
San Diego	5	2

Stanislaus	6	5
Monterey	2	2
Kings	1	1
Madera	1	1
Sacramento	1	1
Solano	1	1
TOTAL	260	83

^aFrom DHS, 2005

Table 3. Distribution of Peak Concentrations of 1,2,3-TCP Detected in California Drinking Water Sources ^a

Peak concentrations of 1,2,3-TCP		
Peak (µg/L)	No. of sources	% of detects
>50	1	-
5.1 – 50	4	2
0.51 - 5.0	17	7
0.051 – 0.5	91	35
0.0051 – 0.05	146	56
<0.0051	1	-
TOTAL	260	100

^aFrom DHS, 2005

Food

No data were located on food analyses or content of 1,2,3-TCP.

Exposure

Studies of human exposure to volatile chemicals in drinking water, including radon and trihalomethanes indicate that in addition to ingestion, inhalation exposure (from chemical volatilized from showering and other indoor activities) and dermal exposure can significantly contribute to the overall daily intake (Jo *et al.* 1990a,b; McKone, 1987; Nieuwenhuijsen *et al.*, 2000; McKone and Layton, 1986). U.S. EPA (1991) also suggested this in a guidance memorandum from its Risk Assessment Forum: “Exposure

to volatile organic compounds in tap water during showering has been found to be approximately equivalent, within an order of magnitude (i.e., plus or minus a factor of three), to exposure from ingesting two liters per day of the same water.”

The CalTOX multimedia exposure model (DTSC, 1994) was employed to determine if inhalation and dermal exposures to 1,2,3-trichloropropane, mainly during showering, would be expected to substantially add to the daily exposure to 1,2,3-trichloropropane due to ingestion. CalTOX is a seven-compartment regional and dynamic fugacity model in a Microsoft Excel format. The chemical parameters for 1,2,3-trichloropropane were identified and the Kp (dermal transfer coefficient) of 0.01 cm/hr was estimated based on Kow and molecular weight (U.S. EPA, 1992; Potts and Guy, 1992).

Human parameters were set for a 70 kg average human consuming 2.0 L/day of tap water (0.028 L/kg-day). The vadose zone soil compartment was loaded with various concentrations of 1,2,3-trichloropropane and the model was run to determine average in-house exposures by different exposure routes. Since the model predicts exposure rather than absorbed dose, the inhalation pathway values were adjusted to assume 50 percent absorption of 1,2,3-trichloropropane at low environmental concentrations while ingested 1,2,3-trichloropropane was assumed to be 100 percent absorbed at low doses.

Dermal exposure was estimated to contribute less than 2 percent of the overall exposure. CalTOX predicted that inhalation exposure was essentially equivalent to exposure due to the ingestion of drinking water. Therefore, the calculation of the PHG will be based on exposure to 4 L_{eq} of water per day.

METABOLISM AND PHARMACOKINETICS

Absorption/Distribution/Elimination

Radiolabeled 1,2,3-trichloropropane (3.6 mg/kg) was administered to F-344 rats (three rats per time point) by injection into the tail vein (Volp *et al.* 1984). Following sacrifice, excreta and samples of tissue were collected at various times. In a separate study, rats were anesthetized and the bile duct was cannulated to determine the amount of 1,2,3-TCP excreted into the bile.

TCP was rapidly distributed to adipose tissue skin and muscle with 70 percent of the dose accounted for in these tissues after 15 minutes. After 4 hours, the amount of 1,2,3-TCP in the liver was larger than in the aforementioned tissues. The amount in the small intestine increased to over nine percent of the administered dose after one hour.

Radioactivity was excreted by three routes, in the urine, feces and expired air. By 24 hours, 40 percent of the dose was accounted for in the urine, 18 percent in the feces and 30 percent in expired air. More than 99 percent of the dose was recovered by day six with 47 percent of the dose recovered in the urine. Five percent of the dose in expired air was recovered unchanged while 25 percent of the administered dose was recovered as carbon dioxide. Little of the dose was recovered unchanged in the urine and feces.

Overall, approximately 30 percent of the dose was excreted in the bile while only 18 percent was recovered in the feces, suggesting some reabsorption of biliary metabolites from the intestine. The maximum rate of biliary excretion occurred at 1 to 1.5 hours post-administration. Administration of glycidol, which decreased hepatic glutathione levels, reduced biliary excretion to only 16 to 30 percent of the rate in control animals. These findings suggested that conjugation with glutathione is an important element in 1,2,3-TCP metabolism.

Radiolabeled 1,2,3-trichloropropane was administered in corn oil to male and female rats and male mice by gavage. Rats were administered 30 mg/kg, while mice were administered either 30 or 60 mg/kg (Mahmood *et al.* 1991). These doses were selected because they were employed in the NTP cancer bioassay (NTP, 1993).

Similar to the study of Volp *et al.* (1984), 1,2,3-TCP was initially distributed in the rat (6 hours post-administration) to the adipose tissue, liver and kidney. Levels of 1,2,3-TCP in adipose tissues were markedly lower by 24 and 60 hours post administration while levels in the liver and kidney remained elevated. Less than 50 percent of the radioactivity remaining in the forestomach, liver and kidney was extractable in solvents at 24 and 60 hours post-administration, suggesting that it was covalently bound to macromolecules. There were little differences in the levels in male and female rats. Radioactivity levels in the rat tissues were substantially higher than in the mouse tissues. The radioactivity levels in mouse tissue were below those in the rat even when twice the dose was administered to the mice.

Metabolism

The metabolism of 1,2,3-trichloropropane is not fully understood. Various metabolic pathways have been proposed (Figure 1) based on recovered metabolites and what is known about the metabolism of similar molecules such as dibromochloropropane (Mahmood *et al.*, 1991; Weber and Sipes, 1992). Metabolism appears to involve cytochrome P-450 and glutathione.

1,2,3-TCP was rapidly metabolized in both the mouse and rat. More than half of the dose was recovered in excreta within 24 hours. Metabolism, measured by the amount of CO₂ collected in expired air, was more rapid in the male mouse than in the male rat during the first 24 hours. Doubling of the dose in the mouse did not alter the pattern of excretion in the mouse, indicating that metabolism did not appear to be saturated. Over 50 percent of the radioactivity was recovered in the urine (50 to 57 percent in the rat and 65 percent in the mouse) with 15 to 20 percent of the administered dose recovered in expired air and feces. Expired volatiles represented approximately 2 percent of the administered dose.

One metabolite, N-acetyl-S-(3-chloro-2-hydroxypropyl)-L-cysteine was identified as the major metabolite in the urine of rats but was a minor constituent in mouse urine, representing 40 percent of urinary radioactivity in the rat and only 3 percent in the mouse at six hours. S-(3-chloro-2-hydroxypropyl)cysteine was the other major constituent in rat urine. These metabolites indicate the involvement of glutathione and also suggest the formation of reactive intermediates that are consistent with the observed unextracted (covalently bound) radioactivity in the tissues.

DNA, RNA and protein were extracted from the liver of male F-344 rats at various times after radiolabeled 1,2,3-TCP was administered i.p. in vegetable oil (30 mg/kg) (Weber and Sipes, 1990). Radioactivity was covalently bound to the protein, DNA and RNA fractions. Inducers of drug metabolism, phenobarbital or beta-naphthoflavone, reduced (phenobarbital) or had no effect (beta-naphthoflavone) on the labeling of protein or DNA. Administration of SKF 525-A, an inhibitor of cytochrome P-450 metabolism, produced an increase in labeling compared to the untreated control group. Administration of an agent that reduces the level of glutathione in the liver (L-buthionine-(S,R)-sulfoximine) resulted in a marked increase of binding of radioactivity to proteins and a marked reduction of binding to DNA.

Given that activation of 1,2,3-TCP appears necessary for its mutagenic activity, agents that stimulate drug metabolism might be expected to increase the binding of 1,2,3-TCP metabolites to macromolecules. However, induction of cytochrome P450 reduced the amount of binding to DNA, and blocking metabolism by cytochrome P450 with SKF 525-A increased DNA binding. The authors suggest that different forms of cytochrome P450 may shunt TCP into metabolic pathways that produce metabolites that do not bind to DNA. A reduction of cellular glutathione levels that resulted in reduced binding of 1,2,3-TCP to DNA suggests that glutathione is involved in the activation of 1,2,3-TCP.

1,2,3-Trichloropropane metabolism was investigated *in vitro* in microsomal and cytosol preparations obtained from human and male F-344 rat livers (Weber and Sipes, 1992). In the rat, 1,2,3-TCP was metabolized to 1,3-dichloroacetone in the presence of microsomal protein and a NADPH generating system. The investigators noted that 1,3-dichloroacetone (DCA) is a direct-acting mutagen and may be responsible for the mutagenic and carcinogenic activity associated with 1,2,3-TCP. The *in vitro* rate of DCA formation using rat liver microsomes was approximately 10 times that using microsomes prepared from human livers.

Pretreatment of rats with phenobarbital and dexamethasone, inducers of cytochrome P450, substantially increased the rate of metabolism, while adding a cytochrome P450 inhibitor; SKF 525-A, to the incubations markedly reduced the rate of metabolism. Incubation of 1,2,3-TCP with rat microsomes resulted in the formation of reactive species that were covalently bound to the microsomal proteins. As expected, the pretreatment of rats with inducers of cytochrome P450 markedly increased binding to microsomal protein. Interestingly, induction of cytochrome P450 resulted in increased binding to microsomal protein *in vitro* in this study, while binding to DNA *in vivo* was decreased by the inducer of cytochrome P450 (Weber and Sipes, 1990). Thus it is unclear whether the lower rate of metabolism in the human liver observed *in vitro* would be reflected in an increase or decreased binding of 1,2,3-TCP metabolite(s) to DNA in human liver relative to the rat liver.

Little binding occurred when a NADPH generating system was omitted from the *in vitro* system. Addition of 1,2,3-TCP to a cytosol fraction did not yield reactive species that covalently bound to cytosol proteins. Addition of glutathione to the microsomes inhibited the binding, even in the absence of glutathione-S-transferase. These findings

indicate that reactive species of 1,2,3-TCP produced by cytochrome P450 are reacting with glutathione.

The addition of alcohol dehydrogenase and NADH to the microsomes yielded mostly 1,3-dichloro-2-propanol, and some 2,3-dichloropropanol. This finding suggests the formation of a reactive intermediate *in vivo* (converted to alcohols by the addition of alcohol dehydrogenase in this *in vitro* system) that is normally difficult to detect because it rapidly binds to large macromolecules such as microsomal proteins (Weber and Sipes, 1990, 1992).

TOXICOLOGY

Toxicological Effects in Animals

Acute Toxicity

Oral LD₅₀s ranging from 150 mg/kg to 500 mg/kg have been reported in the rat (reviewed by ATSDR, 1992; WHO, 2003). Exposure of mice to 1,2,3-TCP at 1 to 130 parts per million (ppm) by inhalation for less than two weeks resulted in irritation of the eye and nose, decreased thickness of olfactory epithelium, and increased liver weight (reviewed by ATSDR, 1992; WHO, 2003).

Subchronic Toxicity

Male and female Sprague-Dawley rats (10 animals/group) were administered 1,2,3-TCP at 0, 10, 100, or 1,000 ppm in drinking water for 90 days (Villeneuve *et al.* 1985). The authors reported only the amount of chemical ingested (based on measured water intake and weight gain) for high dose males (113 mg/kg-day) and the two highest female dose groups (17.6 and 149 mg/kg-day, respectively). Both body weights and water intakes were reduced in male and female rats in the high-dose group.

Liver and kidney weights relative to body weight were elevated in high-dose males and the two highest-dose groups in females. Brain weight relative to body weight also was elevated in the high-dose group. Some of these findings appear to reflect a loss in body weight, as the average organ weights were essentially unchanged in the treatment groups. Indicators of hepatic effects such as serum cholesterol levels, serum hepatic aniline hydroxylase activity and/or aminopyrine demethylase active were elevated, particularly in males and in the high-dose groups. Certain hematological changes were noted but the investigators reported that the levels were considered to be in the normal range. Histological changes were observed in the liver, kidney, and thyroid. Changes were mild and were most evident in the highest-dose groups. The authors concluded that 1,2,3-TCP did not cause overt toxicity in this study.

1,2,3-Trichloropropane was administered in corn oil by gavage to male and female Sprague-Dawley rats at 0, 1.5, 7.4, 29.5, 118 mg/kg-day for 10 days or 0, 1.5, 7.4, 14.7, 59 mg/kg-day for 90 days (Merrick *et al.*, 1991). Final body weights were significantly

decreased in male and female rats in the high dose groups after 10 or 90 days of exposure. The weights of the brain, testis, kidney, and liver were elevated relative to body weight in the high-dose groups, which probably reflected a lower body weight compared to control. The most notable toxicity in these animals was a diffuse inflammation-related necrosis of the heart. The inflammation was more pronounced at higher doses and after 90 days of exposure. Males appeared to be more sensitive than females. Effects of 1,2,3-TCP on other tissues appeared to be minimal, transient (thymus), or were also observed in the control group (liver, male rats).

Male and female CD rats were exposed to 1,2,3-TCP by inhalation, 6 hours per day five days/week (Johannsen *et al.*, 1988). Animals were exposed to 0, 100, 300, 600 and 900 ppm in a four-week pilot study and 0, 5, 15 and 50 ppm or 0, 0.5, 1.5 ppm in a followup 13-week study. In the four-week study, nine out of ten rats died with a single exposure to 900 ppm, while three out of 10 rats died that received 600 ppm and 1 out of ten rats died at 300 ppm. A dose dependent reduction in body weight and an increase in liver/body weight ratios in all treatment groups were noted. Effects on ovary, spleen, kidney and testis weights were also observed.

In the 13-week study, no treatment-related deaths occurred. The body weights of female rats exposed to 15 and 50 ppm of 1,2,3-TCP were significantly lower than control after two weeks of exposure. No biologically significant effects on hematology or clinical chemistry were observed in male or female rats. Effects on absolute or relative liver weights were observed in rats exposed to 5, 15 or 50 ppm. Mild histopathology was reported in the lungs (male and female), liver (male) and spleen (female; mild to marked) in rats exposed to 5, 15 or 50 ppm. Midzonal hepatocellular hypertrophy was observed in the livers of male rats exposed to 15 or 50 ppm. No treatment-related histopathology was observed at lower concentrations.

In a study designed to select doses for a two-year cancer bioassay, 1,2,3-TCP was administered (0, 8, 16, 32, 63, 125, 250 mg/kg-day) in corn oil by gavage, 5 days per week to male and female F-344 rats and B6C3F₁ mice for 17 weeks with an 8-week interim evaluation (NTP, 1993; WHO, 2003). Rats appeared to be more sensitive to 1,2,3-TCP than mice. All male and female rats that received 250 mg/kg-day died and four of 10 females and one male rat that received 125 mg/kg-day died during the study. Mean body weight of male rats receiving 63 or 125 mg/kg-day and female rats receiving 125 mg/kg-day were significantly decreased.

Absolute and relative liver weights were significantly greater than control in male rats receiving 32 mg/kg-day or greater and in female rats receiving 16 mg/kg-day or greater. Hepatocellular necrosis was evident in female rats receiving 125 mg/kg-day. Absolute and relative kidney weights of male rats receiving 32 mg/kg-day or more or female rats receiving 63 mg/kg-day or more were significantly increased over control. Regenerative hyperplasia in the kidney was observed in the kidney of male and female rats receiving 64 mg/kg-day at 8 weeks. Decreased erythrocyte mass (lower mean hematocrit, hemoglobin and erythrocyte counts) were observed at doses of 16 mg/kg-day or greater at 8 and 17 weeks in female rats and 16 mg/kg-day or greater at 8 weeks and 63 mg/kg-day or greater at 17 weeks in male rats.

Clinical chemistry findings were consistent with the aforementioned changes in the liver, with female rats that received 125 mg/kg-day exhibiting prominent increases in serum alanine aminotransferase, aspartate aminotransferase and sorbitol dehydrogenase activity. Bilirubin levels were elevated in males and females receiving 63 and 125 mg/kg-day. Effects on erythrocytes in mice were observed at 63 mg/kg-day in male mice and 16 mg/kg-day in female mice (significant decreases in hematocrit and erythrocytes levels at 8 weeks in male mice and 16 weeks in female mice). Effects on the liver in mice were also observed but at higher doses than in rats. Significant increases in serum enzymes that are indicative of liver toxicity were not observed in mice.

Genetic Toxicity

In *in vitro* bioassays, 1,2,3-TCP was generally positive for genotoxic activity in the presence of metabolic activation and negative without metabolic activation. Results of NTP genotoxicity studies where 1,2,3-TCP was tested in various strain of *S. typhimurium* were generally positive (NTP, 1993). 1,2,3-Trichloropropane was also positive with metabolic activation, in inducing gene mutations in mouse lymphoma cells, sister chromatid exchange in CHO and V79 cells, chromosomal aberrations in CHO cells, and micronuclei in human lymphoblast cell lines (WHO, 2003). 1,2,3-TCP did not induce unscheduled DNA synthesis in primary rat hepatocytes.

1,2,3-Trichloropropane was negative in a micronucleus assay with or without added S9 mix in isolated human lymphocytes (Tafazoli and Kirsch-Volders, 1996). In isolated human lymphocytes, the toxicant was positive in an alkaline single cell gel electrophoresis test (comet assay) indicating 1,2,3-TCP ability to cause DNA breaks with and without added S9 mix (Tafazoli and Kirsch-Volders, 1996).

In vivo genotoxicity studies with 1,2,3-trichloropropane revealed DNA single strand breaks in rat liver (Weber and Sipes, 1991). Other tests described below demonstrated the formation of DNA adducts following 1,2,3-TCP administration. DNA adduct formation was investigated in male F-344 rats and male B6C3F₁ mice administered 1,2,3-TCP in corn oil by gavage at the highest doses and one-tenth the doses administered in the NTP bioassay (La *et al.*, 1995). DNA was isolated from various tissues and adducts were isolated by HPLC. In a separate study, 1,2,3-TCP was administered by intravenous injection to increase adduct formation in the liver so that adducts could be identified.

One DNA adduct predominated in the liver, S-[1-(hydroxymethyl)-2-(N7-guanyl)ethyl]glutathione. DNA adducts were detected in a number of tissues including tissues where tumors were observed in the NTP study. Adducts were also detected in tissues where tumors were not observed, sometimes at very high levels, e.g., in glandular stomach in rats and mice. The lack of concordance between adduct formation and tumors suggests that other factors are involved in tumorigenesis.

The effect of vehicle and route of administration on DNA adduct formation was investigated in male B6C3F₁ mice administered 6 mg/kg-day radiolabeled 1,2,3-TCP for five days either by gavage or in drinking water (La *et al.*, 1996). DNA from the forestomach and liver (target tissues in NTP cancer bioassay) and glandular stomach and the kidney (non-target tissues) was isolated and adducts were characterized. The S-[1-

(hydroxymethyl)-2-(N7-guanyl)ethyl]glutathione DNA adduct appeared to predominate in all tissues.

Statistically significant increased adduct levels were detected in kidney and liver but not the glandular stomach or the forestomach when 1,2,3-TCP was administered in corn oil compared to a drinking water vehicle. Statistically significant increases in cell proliferation were not observed in any of the four tissues when 1,2,3-TCP was administered in drinking water. In contrast, statistically significant increased cell proliferation was observed in all four tissues when 1,2,3-TCP was administered in corn oil. The changes in adduct formation and cell proliferation occurred in tissues with a tumorigenic response in the NTP bioassay (liver and forestomach) as well as tissues where no response was detected (glandular stomach and kidney). In this study, neither adduct formation nor cell proliferation appeared to be predictive of the tumorigenic response observed in the NTP bioassay.

Ito and coworkers investigated the relevance of DNA adduct (S-[1-(hydroxymethyl)-2-(N7-guanyl)ethyl]glutathione) on ras gene mutations in forestomach tumors from mice administered 1,2,3-TCP (Ito *et al.*, 1996). Tumors in ten of sixteen mice had mutations in the H-*ras* or K-*ras* gene. The observed AT to TA or GC to CG transversions were not consistent with miscoding properties of the (S-[1-(hydroxymethyl)-2-(N7-guanyl)ethyl]glutathione) adduct.

The available studies (summarized above) on 1,2,3-TCP genotoxic activity *in vivo* and *in vitro* are more extensively described and reviewed by IARC (1995) and WHO (2003).

Developmental and Reproductive Toxicity

Male and female CD rats were exposed to 1,2,3-TCP five days/week for 6 hour/day by inhalation at 0, 5 or 15 ppm for 10 weeks prior to mating and during the mating period (Johannsen *et al.*, 1988). The female mice that were successfully mated were then exposed daily during the first 14 days of gestation. For females where no mating occurred, exposure was continued while they were housed with other male(s) for additional 10 day periods until mating occurred. They were then exposed during the first 14 days of gestation. Additional mating periods were restricted to a maximum of four or forty days of exposure. In a follow up study, the mice were exposed to 0, 0.5 or 1.5 ppm 1,2,3-TCP due to low mating indices in the initial study.

In the initial study, the body weights of both sexes in the high-dose group (15 ppm) were lower than control, and body weight gain was significantly lower than control during the pre-mating period. Body weight in the high-dose females also remained below that of the control group during the gestation and lactation period. Severe head tilt was observed in one low-dose female and six high-dose animals (the authors suggest that this effect is consistent with mycoplasma infection).

No difference was observed in mating performance (successful mating/total) and fertility index (pregnancy/successful mating) between the treated versus control groups in the initial or followup study. There were no differences between litter size, gestation length, mean pup body weight and number of surviving pups in the treated and control groups in

the initial and followup study. No histopathology was observed in the testes, epididymes or ovaries in the treated animals.

1,2,3-Trichloropropane was evaluated for reproductive and developmental toxicology using a continuous breeding protocol in which toxicant exposure of males and females continues as animals breed, producing up to five litters per breeding pair (Chapin *et al.*, 1997; Gulati *et al.*, 1991; WHO, 2003). Beginning seven days prior to breeding, 30, 60 or 120 mg/kg was administered by gavage in corn oil daily to male and female CD-1 mice for 14 weeks (Task 2). Fewer breeding pairs delivered litters by the third, fourth and fifth litters than in control animals. In the fifth litter, 87, 78, 68 and 42 percent of the pairs exposed to 0, 30, 60, and 120 mg/kg-day, respectively, delivered a litter. The decrease was statistically significant (42 percent) compared to control (87 percent) in the fifth litters of the high dose group. Overall (aggregating the results across all five litters), there was a 16 percent decrease in litters per breeding pair exposed to the high dose of 1,2,3-TCP and there were 47 percent fewer pups per litter.

Pups in the final litters (litter five) were reared by the dam, weaned and then exposed to TCP through mating (Task 4). Neither pup weights nor viability were noticeably affected following the exposure of the F₁ generation to 1,2,3-TCP. Because of the reduced number of litters in the high dose in Task 2, only nine breeding pairs were available in Task 4. Only three of nine breeding pairs in the high-dose group delivered a litter with any pups.

A crossover study in which high-dose females or males were bred with an untreated partner (Task 3) revealed fewer live pups when females were exposed and no noticeable reproductive toxicity when only males were exposed to 1,2,3-TCP. After the crossover breeding study, the adults were sacrificed. Increased liver and kidney weight was observed in the high-dose males, and increased liver weight was observed in the high-dose females. Four of ten females had microscopic ovarian amyloidosis compared to zero females in the control group.

Five doses of 1,2,3-trichloropropane (80 mg/kg) were administered to male rats by gavage (Saito-Suzuki *et al.*, 1982). Following treatment, each rat was mated with an untreated female rat once a week for eight successive weeks. No significant changes from control levels were observed for the following toxic endpoints: number of females with implants, numbers of corpora lutea per female, number of implants per female, number of live embryos per female, and mean number of dead implants.

Immunotoxicity

No data were located.

Neurotoxicity

No data were located.


Chronic Toxicity/Carcinogenicity

Groups of 60 male or female rats (F-344) and mice (B6C3F₁) were administered 1,2,3-TCP by gavage in corn oil in a two-year cancer bioassay (NTP, 1993). Rats were administered 0, 5, 10 or 30 mg/kg-day of 1,2,3-TCP, 5 days per week while mice were administered 0, 10, 30 or 60 mg/kg-day, 5 days per week. The doses employed in this study were based on findings in a 17-week study in which higher doses of 1,2,3-TCP in rats and mice resulted in significant toxicity. It was reported that “up to 10” animals in each group were used for an interim sacrifice at 15 months.

The survival of mice was substantially reduced in the two highest dose groups administered 1,2,3-TCP (Figure 2). In mice, early deaths in the 60 mg/kg-day group were reported to result from chemically-related neoplasms, primarily forestomach tumors (NTP, 1993). Of the surviving mice in the 60 mg/kg-day group, the remaining males were sacrificed in week 79 and females in week 73 because of high mortality. Survival in the 20 mg/kg-day group was also reduced due to chemically-related tumors. Remaining mice in these dose groups were sacrificed in week 89 because of high mortality. The early death of mice in the two highest dose groups likely affected the incidence of tumors that would have been detected at certain sites (absent the early mortality), particularly tumors that were late developing.


Analogous to what was observed in mice, the survival of rats in the two highest dose groups was markedly reduced (Figure 3). Early deaths were attributed to the occurrence of chemically-related neoplasms, but the site(s) were not specified. In the 30 mg/kg-day dose group, surviving female rats were sacrificed at week 66 and males at week 76 because of high mortality.

Figure 2. Survival of mice administered various doses of 1,2,3-trichloropropane¹



¹from NTP (1993)

Figure 3. Survival of rats administered various doses of 1,2,3-trichloropropane¹



¹from NTP (1993)

The administration of 1,2,3-TCP resulted in statistically significant increases in tumors at a number of sites in both male and female mice and rats (Tables 4 and 5). In the male mouse, statistically significant or notable increases in neoplasms were observed in the forestomach, liver and Harderian gland. Statistically significant or notable increases in tumors were observed in the oral cavity, forestomach, liver, Harderian gland and uterus of female mice.

Table 4. Summary of Treatment-Related Tumors in the 2-Year 1,2,3-Trichloropropane Bioassay in B6C3F₁ Mice¹

Male Mice	Female Mice
Forestomach: squamous cell papilloma (3/52, 28/51, 22/54, 33/56) ² squamous cell carcinoma (0/52, 40/51, 50/54, 51/56)	Forestomach: squamous cell papilloma (0/50, 23/50, 18/51, 29/55) squamous cell carcinoma (0/50, 46/50, 49/51, 49/55)
Liver: hepatocellular adenoma (11/52, 18/51, 21/54, 29/56) hepatocellular carcinoma (4/52, 11/51, 5/54, 3/56)	Liver: hepatocellular adenoma (6/50, 9/50, 8/51, 31/55) hepatocellular carcinoma (1/50, 3/50, 0/51, 2/55)
Harderian gland: adenoma (1/52, 2/51, 10/54, 11/56)	Oral cavity: squamous cell carcinoma (0/50, 0/50, 1/51, 5/55)
	Uterus: adenoma (0/50, 1/50, 0/51, 3/54) adenocarcinoma (0/50, 4/50, 3/51, 6/54) stromal polyp (0/50, 2/50, 1/51, 6/54)

¹Adapted from summary table in NTP, 1993.

²Number of tumor-bearing mice/number of mice necropsied in each dose group.

In the male rat, statistically significant or notable increases in neoplasms were observed in the oral cavity (tongue and pharynx), forestomach, pancreas, kidney, preputial gland and Zymbal gland. Statistically significant or notable increases in tumors were observed in the oral cavity, forestomach, clitoral gland, mammary gland and Zymbal gland of female rat.

Table 5. Summary of Treatment-Related Tumors in the 2-Year 1,2,3-Trichloropropane Cancer Bioassay in the F-344 Rat¹

Male Rats	Female Rats
Oral cavity: squamous cell papilloma (0/50, 4/50, 9/49, 19/52) ² squamous cell carcinoma (1/50, 0/50, 11/49, 25/52)	Oral cavity: squamous cell papilloma (1/50, 5/49, 10/52, 18/52) squamous cell carcinoma (0/50, 1/49, 21/52, 21/52)
Forestomach: squamous cell papilloma (0/50, 29/50, 33/49, 38/52) squamous cell carcinoma (0/50, 9/50, 27/49, 13/52)	Forestomach: squamous cell papilloma (0/50, 13/49, 32/51, 17/52) squamous cell carcinoma (0/50, 3/49, 9/51, 4/52)
Pancreas: acinar adenoma (5/50, 21/50, 36/49, 29/52)	Clitoral gland: adenoma (5/46, 10/46, 13/50, 10/51) carcinoma (0/46, 0/46, 4/50, 6/51)
Kidney: renal tubule adenoma (0/50, 2/50, 20/49, 21/52)	Mammary gland: adenocarcinoma (1/50, 6/49, 12/52, 21/52)
Preputial gland: adenoma (5/49, 3/47, 5/49, 11/50) carcinoma (0/49, 3/47, 3/49, 5/50)	Zymbal's gland: carcinoma (0/50, 1/49, 0/52, 3/52)
Zymbal's gland: carcinoma (0/50, 0/50, 0/49, 3/52)	

¹Adapted from summary table in NTP, 1993).

²Number of tumor bearing rats/number of rats necropsied in each dose group.

Toxicological Effects in Humans

Acute Toxicity

Acute exposure to 1,2,3-TCP resulted in eye and throat irritation (reviewed by ATSDR, 1992; WHO, 2003). Acute CNS effects have also been reported (HSDB, 2005).

Subchronic Toxicity

No data were located.

Neurotoxicity

No data were located.

Immunotoxicity

No data were located.

Developmental/Reproductive Toxicity

No data were located.

Chronic Toxicity

No data were located.

Evidence for Carcinogenic Activity of 1,2,3-Trichloropropane

Animal Studies

The administration of 1,2,3-TCP in corn oil by gavage to rats and mice resulted in a statistically significant increase in tumors in multiple organs in both male and female mice and rats. Tumors occurred in tissues that were in direct contact with the administered compound (oral cavity, forestomach) and also in tissues distal to the site of administration (liver, Harderian gland, uterus). The increase in tumors also appeared to be dose-dependent.

Human Studies

No studies were identified that evaluated carcinogenic effects of 1,2,3-TCP in humans.

Genotoxicity

1,2,3-Trichloropropane was positive in both *in vivo* and *in vitro* genotoxicity studies. The compound was positive in *in vitro* assays in the presence of metabolic activation and negative without metabolic activation. *In vivo* studies revealed single strand chromosome breaks in rat liver and kidney. Studies also revealed the formation of DNA adducts following the administration of 1,2,3-TCP to rats or mice.

Mechanism

Studies have indicated cytochrome P450 has a role in metabolizing 1,2,3-TCP to an active form. Activation may also involve reaction with glutathione. Glutathione conjugation also appears to inactivate the toxicant. Genotoxicity studies have

demonstrated the formation of DNA adducts following the administration of TCP. Agents that alter TCP metabolism by inducing cytochrome P450 or inhibiting metabolism by cytochrome P-450 or agents that deplete glutathione in the cell altered the amount of 1,2,3-TCP metabolite that bound to protein or DNA.

Conclusion

1,2,3-Trichloropropane was a potent carcinogen by gavage in the available cancer bioassays conducted in rats and mice. 1,2,3-TCP also tested positive in a number of *in vivo* and *in vitro* genotoxicity studies. Therefore it is prudent to assume that 1,2,3-TCP represents a significant carcinogenic risk when it occurs in drinking water.

DOSE-RESPONSE ASSESSMENT

Noncarcinogenic Effects

1,2,3-Trichloropropane was administered in corn oil by gavage to rats and mice five days a week for up to 90 days in a study aimed at identifying doses for a two-year cancer bioassay (NTP, 1993). Rats appeared to be more sensitive than mice to the adverse effects. Effects on the liver (increased absolute and relative liver weight) were observed at doses as low as 32 mg/kg-day (males) and 16 mg/kg-day (females). Histopathology in the liver was observed at 125 or 250 mg/kg. Effects on absolute and relative kidney weights were observed in males treated with 32 mg/kg-day and females receiving 63 mg/kg-day. Regenerative hyperplasia was observed in the kidney of male and female rats receiving doses as low as 64 mg/kg-day after eight weeks of treatment.

Total bilirubin values were increased at 63 and 125 mg/day in male and female rats. Increases in serum enzymes associated with liver toxicity were observed at dose as low as 63 to 125 mg/kg-day. Effects on erythrocytes (decreased erythrocyte mass, lower mean hematocrit, hemoglobin and erythrocyte counts) were observed in male and female rats at doses as low as 16 mg/kg-day. No toxic effects that were not related to the occurrence of tumors were identified in the subsequent two-year cancer bioassay.

Based on effects on erythrocytes at 16 mg/kg-day, 8 mg/kg-day (administered dose) is identified as the no observed adverse effects level (NOAEL). This dose, administered 5 days/week, is corrected to a 7 days/week equivalent as follows:

$$8 \text{ mg/kg} * (5 \text{ days/week} / 7 \text{ days/week}) = 5.7 \text{ mg/kg-day}$$

Carcinogenic Effects

The administration of 1,2,3-TCP to male or female rats and mice resulted in statistically significant increases in tumors at a number of sites (NTP, 1993). At some sites (forestomach and clitoral gland in female rats), incidence of tumors at the high dose was less than at the lower dose(s). This may have been due to shorter survival of the animals, which may have prevented the occurrence or detection of tumors at the site (although

shortened survival would not alter the incidence of a tumor at the site that caused the death, or for tumors that were detected prior to or at death). The incidence of late occurring tumors would be expected to be influenced by changes in survival. At some sites (e.g., the forestomach), tumors occurred in nearly 100 percent of the animals that received 1,2,3-TCP. This finding complicates dose-response calculations. However, the tumors tended to occur earlier in animals that received higher doses.

Both the notable shortening of lifespan and its probable effect on incidence of tumors at many of the sites, and the high incidence of tumors in a number of dose groups suggest that at certain sites, a time-to-tumor model is most appropriate to estimate 1,2,3-TCP cancer potency. The time-to-tumor model takes into consideration both how many tumors occurred and when they occurred. Reporting in the NTP (1993) study of when and what types of tumors were observed in each animal allows for a time-to-tumor analysis. Unfortunately the cause of death of individual animals was not reported.

A time-to-tumor model that is multistage in dose and Weibull in time (Tox_Risk v. 5.2) was employed to develop a dose-response relationship for various tumor sites in male and female rats and mice (Crump *et al.*, 2000). The dose in animals was adjusted to reflect daily exposure to 1,2,3-TCP (multiplied by 5/7) and scaled to humans based on the ratio of human and animal body weights to the $\frac{3}{4}$ power. The model default body weights of 350 g for rats and 30 g for mice were used. While a number of tumor sites could have been selected, the modeling focused on sites with a high incidence of tumors.

NTP reported that mortality was related to the occurrence of tumors. However, because tumors occurred at multiple sites in many of the animals, it is unclear which tumor was the cause of death in a given animal (although the NTP did report that early deaths in mice from the high dose groups (60 mg/kg-day) were due to chemical-related neoplasms, primarily in the forestomach (NTP, 1993). In a separate publication, the investigators reported: "Neoplasms of the forestomach in rats and mice, of the oral mucosa in rats and mammary gland in female rats, were the principal cause of death of most animals dying or killed moribund before the end of the study" (Irwin *et al.*, 1995).

Given the multiplicity of tumor sites, an analysis based on the assumption that tumors at each of the sites were responsible for the animal's death appeared to be inappropriate except perhaps for forestomach tumors in mice, based on the NTP (1993) comment that early deaths in mice from the high dose groups (60 mg/kg-day) were due to chemical-related neoplasms, primarily in the forestomach. Assuming that all tumors were incidental at the time of death (not the cause of it) also appears to be inappropriate given the animals were reported to be dying from chemically-related tumors. Therefore, a middle position was employed in the selection of data from the time-to-tumor modeling, where carcinomas were judged to be fatal while papillomas and adenomas were assumed to be incidental to the cause of death for the modeling. In addition, separate analyses of the findings in mice were performed for tumors of the forestomach: 1.) where all forestomach tumors were considered to be incidental to the cause of death of the animal, and 2.) where forestomach tumors were presumed to be the cause of death of the animal. These separate analyses were undertaken to determine the impact of the assumption regarding the cause of death on the dose response relationship based on a time-to-tumor model.

NTP did not report the findings of individual animals in the interval sacrifice groups in its technical report but did provide the specific findings on its internet site (NTP 2007a, NTP 2007b). A separate analysis was performed that included the results from animals in the interval (66 weeks) sacrifice groups. For this analysis, animals were censored (removed from the analysis) if they did not display a tumor, because they did not live until the end of the animal's natural lifetime.

The results of the time-to-tumor analysis with different assumption about cause of death are shown in Table 6. The lower-bound estimate of the dose associated with a 10 percent cancer risk (LED₁₀) is based on lifetime (70 years) exposure.

Table 6. 1,2,3-Trichloropropane Potency Estimates using a Time-to-Tumor Model

Species	Sex	Tumor Site	Human q ₁ * (mg/kg-day) ⁻¹	LED ₁₀ (μg/kg-day)
Rat	Male	Forestomach ¹	3.0	36
	Female	Forestomach ¹	0.9	110
	Male	Oral Cavity ¹	0.2	460
	Female	Oral Cavity ¹	0.3	350
	Male	Preputial ¹	0.2	590
	Female	Mammary ¹	1.0	110
	Male	Pancreas	1.4	77
	Mouse	Male	Forestomach ¹	16
Female		Forestomach ¹	17	6.1
Male		Forestomach ^{1,2}	17	6.2
Female		Forestomach ^{1,2}	25	4.2
Male		Forestomach ³	22	4.8
Female		Forestomach ³	1900	0.07
Male		Forestomach ^{2,3}	22	4.8
Female		Forestomach ^{2,3}	180	0.6
Male		Forestomach ⁴	4	23.7
Female		Forestomach ⁴	11	9.4
Male		Liver ¹	1.8	59
Female		Liver ¹	0.9	110

¹Where carcinomas are considered to be fatal tumors and adenomas are considered to be incidental (not the cause of death in the animal).

²Includes interval sacrifice groups.

³Where carcinomas and papillomas were considered to be incidental (incidental to the cause of death in the animal).

⁴Where carcinomas and papillomas were considered to be fatal (the cause of death in the animal).

The assumption that all tumors were incidental to the cause of death of the animals had a marked effect on the cancer potency estimate, q_1^* , while the assumption that all tumors were the cause of death in the animals had little effect on q_1^* . Including interval sacrifice groups in the model had little effect on the q_1^* . Again, a middle position was employed in the selection of data from the time-to-tumor modeling, where carcinomas were judged to be fatal while papillomas and adenomas were assumed to be incidental to the cause of death for the modeling. With this strategy, the female mouse forestomach yielded the highest estimate of potency q_1^* , $25 \text{ (mg/kg-day)}^{-1}$ and the lowest estimate of the lower bound on the dose causing a 10 percent tumor incidence (LED_{10}), $4.2 \text{ } \mu\text{g/kg-day}$. The linear slope associated with the dose response relationship is calculated from the LED_{10} for forestomach tumors in the female mouse as:

$$\frac{0.1}{\text{LED}_{10}} = \frac{0.1}{0.0042 \text{ mg/kg-day}} = 24 \text{ mg/kg-day}^{-1}$$

For comparative purposes, dose response relationships for tumors at each site were also determined using a linearized multistage (LMS) model although the goodness of fit for tumors in the forestomach in rats and mice was very poor ($p < 10^{-6}$). As expected, the time to tumor model yielded higher estimates of potency than the LMS model because it accounted for the early death in the animals that prevented the observation of late occurring tumors. The lower-bound dose associated with 10 percent risk (LED_{10}) for tumors in the forestomach in the male rat using a Linearized Multistage Model is 0.093 mg/kg-day . Using a linear model, the slope of the dose-response relationship associated with forestomach tumors in male rats (the highest rat potency) is $0.1 / 0.093 \text{ mg/kg-day} = 1.1 \text{ (mg/kg-day)}^{-1}$. The LED_{10} for tumors in the forestomach in the female mice using a Linearized Multistage Model is 0.047 mg/kg-day . Using a linear model, the slope of the dose-response relationship associated with forestomach tumors in female mice is $0.1 / 0.047 \text{ mg/kg-day} = 2.1 \text{ (mg/kg-day)}^{-1}$.

CALCULATION OF PHG

Noncarcinogenic Effects

For estimation of a health-protective concentration of a chemical in drinking water, an acceptable daily dose (ADD) of the chemical from all sources will first be calculated. This involves incorporation of appropriate estimates of uncertainty in the extrapolation of the critical toxic dose from human or animal studies to the estimation of a lifetime daily dose for that is unlikely to result in any toxic effects. For this purpose, the following equation will be used:

$$\text{ADD} = \frac{\text{NOAEL/LOAEL in mg/kg-day}}{\text{UF}}$$

where,

ADD = acceptable daily dose, an estimate of the maximum daily dose which can be consumed by humans for an entire lifetime without toxic effects;

NOAEL/LOAEL = no-observed-adverse-effect level or lowest-observed-adverse-effect level in the critical study;

UF = uncertainty factor.

Calculation of a public health-protective concentration (C, in mg/L) for a chemical in drinking water uses the following equation for noncarcinogenic endpoints:

$$C = \frac{\text{ADD mg/kg-day} \times \text{BW kg/day} \times \text{RSC}}{\text{L/day}}$$

where,

BW = body weight (a default of 70 kg for adult males, 60 kg for adult females, or 25 kg for a child);

RSC = relative source contribution, usually 20 to 80 percent (0.20 to 0.80);

L/day = volume of water consumed daily for an adult or child (defaults of 2 L/day and 1 L/day, respectively, which may be corrected for exposures to the chemical from other household uses of tap water).

The most robust data available for calculation of the health protective level based on noncarcinogenic effects were from a 17-week subchronic study conducted in F-344 rats (NTP, 1993). A NOAEL of 8 mg/kg-day was identified based on effects on erythrocytes at the next higher dose, 16 mg/kg-day. A dose of 8 mg/kg administered 5 days/week corresponds to 5.7 mg/kg-day when adjusted to reflect daily dosing. Use of an uncertainty factor of 1,000 was considered appropriate to account for the use of subchronic toxicity data (10), for interspecies differences (10), and for human variability (10). Thus, the ADD is calculated as:

$$\text{ADD} = \frac{5.7 \text{ mg/kg-day}}{1,000} = 5.7 \text{ } \mu\text{g/kg-day}$$

Based on data for water consumption by the general population in the Western Region of the U.S. (OEHHA, 2000), a water ingestion amount of 2.0 L/day was assumed. The contributions from inhalation and dermal exposure to 1,2,3-trichloropropane in tap water in the home were 2 L_{eq}/day and 0 L_{eq}/day, respectively (see Exposure section). Adding

up exposure by the different routes leads to a total drinking water consumption of 4 L_{eq}/day for 1,2,3-trichloropropane.

Using the adjusted NOAEL value, 70 kg body weight, 80 percent relative source contribution (no other substantive sources of environmental exposure were identified), and a combined 4 L_{eq}/day drinking water consumption rate, the public health protective concentration for noncarcinogenic effects is calculated as follows:

$$C = \frac{5.7 \mu\text{g/kg-day} \times 70 \text{ kg} \times 0.80}{4.0 \text{ L}_{\text{eq}}/\text{day}} = 0.08 \text{ mg/L} = 80 \text{ ppb}$$

The health-protective concentration for 1,2,3-trichloropropane based on a noncancer endpoint for the general population is therefore 0.08 mg/L or 80 ppb.

Carcinogenic Effects

For carcinogens, the following general equation can be used to calculate the public health-protective concentration (C) of a chemical in drinking water (in mg/L):

$$C = \frac{R \times BW}{CSF \times WC} = \text{mg/L (ppm)}$$

where,

R = *de minimis* lifetime excess individual cancer risk (a default of 10⁻⁶);

BW = adult body weight (a default of 70 kg);

CSF = cancer slope factor, a potency derived from the lower 95% confidence limit on the 10 percent tumor dose (LED₁₀);

WC = volume of water consumed daily (L/d), which may include a multiroute exposure estimate for volatile organic compounds (i.e., L_{eq}/d).

The most sensitive cancer endpoint for 1,2,3-trichloropropane was an increase in tumors in the forestomach of the female mouse (NTP, 1993). The concentration of 1,2,3-TCP in water equivalent to a lifetime risk of 10⁻⁶ was calculated using a time-to-tumor model to estimate the low bound of the dose associated with a risk of 10⁻¹. A linear dose response model was then employed to obtain an oral slope factor of 24 (mg/kg-day)⁻¹. The health protective concentration was calculated using oral data only, as no data on the carcinogenicity of 1,2,3-trichloropropane via the inhalation route were available.

The default adult body weight of 70 kg was used in the calculation. Based on data for water consumption by the general population in the Western Region of the U.S. (OEHHA, 2000), a water ingestion amount of 2.0 L/day was assumed. The contributions from inhalation and dermal exposure to 1,2,3-trichloropropane in tap water in the home were 2 L_{eq}/day and 0 L_{eq}/day, respectively (see Exposure section). Adding up exposure

by the different routes leads to a total drinking water consumption of 4 L_{eq}/day for 1,2,3-trichloropropane.

Using the cancer slope factor of 24 (mg/kg-day)⁻¹, the estimated health-protective value is therefore:

$$C = \frac{10^{-6} \times 70 \text{ kg}}{24 (\text{mg/kg-day})^{-1} \times 4 \text{ L}_{\text{eq}}/\text{day}} = 0.0007 \text{ } \mu\text{g/L} = 0.0007 \text{ ppb}$$

It is proposed to derive the PHG for 1,2,3-trichloropropane using a linear cancer extrapolation model, and a *de minimis* 10⁻⁶ lifetime theoretical cancer risk, resulting in a proposed PHG of 0.0007 μg/L or 0.0007 ppb. The corresponding values at 10⁻⁵ and 10⁻⁴ risk are 0.007 and 0.07 μg/L (ppb), respectively. The proposed PHG based on cancer effects would also be protective against noncancer effects.

RISK CHARACTERIZATION

The proposed PHG for 1,2,3-trichloropropane of 0.0007 ppb is based on risk associated with ingestion of drinking water and inhalation of 1,2,3-TCP that volatilizes from the tap water during showering and other household water uses. Dermal exposures were estimated to be insignificant relative to direct ingestion and inhalation. Various sources of uncertainty regarding the development of health-protective criteria for the oral and inhalation routes are discussed.

Hazard Identification - There is considerable evidence that exposure to 1,2,3-TCP results in increased incidences of cancer in animals, based on the results of the 1993 NTP cancer bioassays. No other comparable cancer studies were identified. Marked increases in tumors were observed in both male and female rats and mice at multiple sites, and the increases appeared to be dose-related. 1,2,3-Trichloropropane was positive in genotoxicity assays and DNA adducts have been detected in rats and mice and identified following exposure to 1,2,3-trichloropropane.

Dose Response – A time-to tumor model was employed to develop a dose-response relationship for various tumors in male and female rats and mice. 1,2,3-TCP was a potent tumorigenic agent at many sites. In both rats and mice, the animals died early and the deaths were judged to be due to the tumors. The early deaths would also affect the incidence of tumors at other sites. The time to tumor model used information regarding how many tumors occurred and when they occurred to develop a dose response relationship. The NTP (1993) study did not provide details on the cause of death of individual animals but did indicate that neoplasms of the forestomach were the principal cause of most of the early deaths. Given the uncertainty of the cause of death in a given animal, carcinomas were assumed to be lethal while adenomas and other benign tumors were considered to be incidental to the cause of death. Alternatively, the modeling could have been based on all tumors being fatal and all tumors being incidental. Given the multiplicity of tumors in most animals, it is imprudent to assume that tumors at all sites caused the death of each animal, or, given the early mortality of the animals, that the

tumors were incidental to the early death of the animals (especially since NTP concluded that the deaths were related to tumors).

Exposure Assessment – The default upper-bound estimate of drinking water ingestion (2 L/day) was employed to assess oral exposure to 1,2,3-trichloropropane. The contribution of other exposure routes was estimated using CalTOX model (DTSC, 1994). While exposure via the dermal route was determined to contribute a negligible portion of the overall dose, inhalation exposure was estimated to contribute approximately 50 percent of the daily dose (2 L_{eq}/day). The default adult body weight of 70 kg was used in the calculation. The non-cancer health-based criterion also reflects a relative source contribution of 80 percent of the total exposure coming from drinking water, because no other significant sources of exposure were identified. The RSC is not used in the cancer calculation because the cancer risk is calculated as “extra” risk, which would be in addition to other sources of exposure to the chemical.

Risk Characterization - The various sources of uncertainty attendant in the hazard identification, dose response, and exposure assessment are reflected in the estimates of a health-protective concentration of 1,2,3-TCP in drinking water. As better studies of the toxicity of 1,2,3-TCP, better methods to characterize the dose response relationship, and better methods to characterize exposure become available, the uncertainties associated with the risk assessment can be reduced. With the presently available information, the risk associated with exposure to 1,2,3-TCP may have been under- or overestimated. To address this uncertainty, the cancer risk assessment utilized an upper bound estimate of potency in the development of the health-based criterion, to ensure that risk is not markedly underestimated.

No sensitive populations were identified. It is judged that pregnant women and their fetuses, infants, the elderly, and other potentially sensitive populations are adequately protected by the proposed PHG.

OTHER REGULATORY STANDARDS

There are no State of California or federal MCLs for 1,2,3-TCP. U.S. EPA published an oral slope factor of 7 (mg/kg-day)⁻¹ in its HEAST summary table (U.S. EPA, 1997). DPH has developed a California Notification Level of 0.005 ppb based on the HEAST oral slope factor, which is also the detection limit for reporting (DHS, 2005).

The U.S. EPA developed an RfD of 0.006 mg/kg-day based on a non-cancer endpoint, alterations in clinical chemistry and reduction in red cell mass (U.S. EPA, 2007; file last updated 08/01/1990). The Permissible Exposure Limit (PEL) developed by the federal Occupational Safety and Health Administration (OSHA) for 1,2,3-TCP is 50 ppm or 300 mg/m³. The concentration in air rated as an Immediate Danger to Life and Health (IDLH) is 100 ppm (NIOSH, 2004). The National Institute for Occupational Safety and Health (NIOSH) recommended exposure limit (REL) for 1,2,3-TCP is 10 ppm and NIOSH considers 1,2,3-TCP as a potential occupational carcinogen (NIOSH, 2004).

The overall evaluation of the International Agency for Research on Cancer (IARC) is that 1,2,3-TCP is probably carcinogenic to humans (Group2A) (IARC, 1995). IARC

DRAFT

concluded that there is inadequate evidence in humans and sufficient evidence of carcinogenicity in experimental animals. NTP concluded that 1,2,3-TCP is reasonably anticipated to be a human carcinogen based on sufficient evidence of malignant tumor formation at multiple sites in multiple species of experimental animals (NTP, 2005).

REFERENCES

- ATSDR (1992). Toxicological profile for 1,2,3-Trichloropropane. Agency for Toxic Substances and Disease Registry, U.S. Department of Health and Human Services, Atlanta, Georgia, Sept 1992.
- ATSDR (2003). 2003 CERCLA priority list of hazardous substances. Agency for Toxic Substances and Disease Registry, U.S. Department of Health and Human Services, Atlanta, Georgia. Accessed July 2005 at: <http://www.atsdr.cdc.gov/clist.html>.
- Chapin R, Gulati D, Mounce R, Russell S, Poonacha K (1997). Trichloropropane. *Environ Health Perspect* 105(S1)1-2. NTIS #PB91129676. Accessed at: <http://ehp.niehs.nih.gov/members/1997/Suppl-1/dfa184.html>.
- Crump KS, Howe RB, van Landingham C, Fuller WG (2000). TOX_RISK, A Toxicology Risk Assessment Program. Version 5.2. KS Crump Division, Clement International Corp., Ruston, Louisiana, October.
- DHS (2005). 1,2,3-Trichloropropane: Notification Level & Monitoring Results. California Department of Health Services. Accessed May 5, 2005 at: <http://www.dhs.ca.gov/ps/ddwem/chemicals/123tcp/notificationlevel.htm>.
- DTSC (1994). CalTOX, A Multimedia Total Exposure Model for Hazardous Waste Sites. Spreadsheet User's Guide. Version 1.5. Prepared by the University of California, Davis, in cooperation with Lawrence Livermore National Laboratory for the Department of Toxic Substances Control (DTSC), California Environmental Protection Agency, Sacramento, California.
- Gulati DK, Hope E, Teague JL, Chapin RE (1991). Reproductive toxicity assessment by Continuous Breeding in Sprague-Dawley rats: a comparison of two study designs. *Fundam Appl Toxicol* 17:270-279.
- HSDB (2005). 1,2,3-Trichloropropane. Hazardous Substances Data Bank. Accessed at: <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB>.
- IARC (1995). 1,2,3-Trichloropropane. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans, Vol 63, International Agency for Research on Cancer, Lyon, France, pp. 223–244.
- INCHEM (2003). 1,2,3-Trichloropropane. Concise International Chemical Assessment Doc. 56. International Programme on Chemical Safety, World Health Organization, Geneva, Switzerland. Accessed at: <http://www.inchem.org/documents/cicads/cicads/cicad56.htm#5.1>.
- Irwin RD, Haseman JK, Eustis SL (1995). 1,2,3-Trichloropropane: a multisite carcinogen in rats and mice. *Fundam Appl Toxicol* 25(2):241-252.
- Ito N, La D, Holt S, Craft TR, Sills RC, Swenberg JA (1996). Analysis of ras mutations in forestomach tumors from B6C3F1 mice exposed to 1,2,3-trichloropropane. 87th Annual Meeting of the American Assoc. for Cancer Research, Washington, DC, USA, April 20-24, 1996. *Proc. American Assoc. for Cancer Research* 37(0):137.

- Jo WK, Weisel CP, Liroy PJ (1990a). Chloroform exposure and the health risk associated with multiple uses of chlorinated tap water. *Risk Anal* 10(4):581-585.
- Jo WK, Weisel CP, Liroy PJ (1990b). Routes of chloroform exposure and body burden from showering with chlorinated tap water. *Risk Anal* 10(4):575-580.
- Johannsen FR, Levinskas GJ, Rusch GM, Terrill JB, Schroeder RE (1988). Evaluation of the subchronic and reproductive effects of a series of chlorinated propanes in the rat. I. Toxicity of 1,2,3-trichloropropane. *J Toxicol Environ Health* 25(3):299-315.
- La DK, Lilly PD, Anderegg RJ, Swenberg JA (1995). DNA adduct formation in B6C3F1 mice and Fischer-344 rats exposed to 1,2,3-trichloropropane. *Carcinogenesis* 16(6):1419-1424.
- La DK, Schoonhoven R, Ito N, Swenberg JA (1996). The effects of exposure route on DNA adduct formation and cellular proliferation by 1,2,3-trichloropropane. *Toxicol Appl Pharmacol* 140(1):108-114.
- Mahmood NA, Overstreet D, Burka LT (1991). Comparative disposition and metabolism of 1,2,3-trichloropropane in rats and mice. *Drug Metab Dispos* 19(2):411-418.
- McKone TE (1987). Human exposure to volatile organic compounds in household tap water: the indoor inhalation pathway. *Environ Sci Technol* 21:1194-201.
- McKone TE, Layton DW (1986). Screening the potential risks of toxic substances using a multimedia compartment model: estimation of human exposure. *Regul Toxicol Pharmacol* 6(4):359-380.
- Merrick BA, Robinson M, Condie LW (1991). Cardiopathic effect of 1,2,3-trichloropropane after subacute and subchronic exposure in rats. *J Appl Toxicol* 11(3):179-187.
- Nieuwenhuijsen MJ, Toledano MB, Elliott P (2000). Uptake of chlorination disinfection by-products; a review and a discussion of its implications for exposure assessment in epidemiological studies. *J Expo Anal Environ Epidemiol* 10(6 Pt 1):586-599.
- NIOSH (2004). 1,2,3-Trichloropropane. Pocket Guide to Chemical Hazards. National Institute of Occupational Safety and Health (NIOSH), Centers for Disease Control and Prevention, U.S. Department of Health and Human Services, Cincinnati, Ohio. Accessed May 12, 2005 at: <http://www.cdc.gov/niosh/npg/npg.html>.
- NTP (1993). Toxicology and carcinogenesis of 1,2,3-trichloropropane (CAS No. 96-18-4) in F344/N Rats and B6C3F1 Mice (gavage studies). Technical Report Series No 384. NIH publication No. 94-2839. Research Triangle Park, NC: National Toxicology Program. 348 pp.
- NTP (2005). 1,2,3-Trichloropropane, Report on Carcinogens, Eleventh Ed.; U.S. Department of Health and Human Services, Public Health Service, National Toxicology Program, January 2005
- NTP (2007a). Results for long-term bioassay for Trichloropropane, All Animals with Neoplastic Diagnosis Stomach, Forestomach Squamous Cell Carcinoma. (Study TR-384) Accessed at: <http://ntp->

apps.niehs.nih.gov/ntp_tox/index.cfm?fuseaction=longtermbioassaydata.allanimalswithneoplasticdiagnosisstomach¤t_organ_id=25465¤t_lesion_id=551&study_no=C60220B&study_length=2%20Years.

NTP(2007b). Results for long-term bioassay for Trichloropropane, All Animals with Neoplastic Diagnosis Stomach, Forestomach Squamous Cell Papilloma. (Study TR-384) Accessed at: http://ntp-apps.niehs.nih.gov/ntp_tox/index.cfm?fuseaction=longtermbioassaydata.allanimalswithneoplasticdiagnosisstomach¤t_organ_id=25465¤t_lesion_id=330&study_no=C60220B&study_length=2%20Years.

OEHHA (2000). The Air Toxics Hot Spots Risk Assessment Guidelines, Part IV, Exposure Assessment and Stochastic Analysis Technical Support Document. Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Oakland, California. September.

Potts RO, Guy RH (1992). Predicting skin permeability. *Pharm Res* 9(5):663-669.

Saito-Suzuki R, Teramoto S, Shirasu Y (1982). Dominant lethal studies in rats with 1,2-dibromo-3-chloropropane and its structurally related compounds. *Mutat Res* 101(4):321-327.

Tafazoli M, Kirsch-Volders M (1996). In vitro mutagenicity and genotoxicity study of 1,2-dichloroethylene, 1,1,2-trichloroethane, 1,3-dichloropropane, 1,2,3-trichloropropane and 1,2,3-trichloropropene using the micronucleus test and the alkaline single cell gel electrophoresis technique (comet assay) in human lymphocytes. *Mut Res* 371:185-202.

U.S. EPA (1991). Guidance on Estimating Exposure to VOCs During Showering. Memorandum from Patton DE, Risk Assessment Forum to Habicht FH, Risk Assessment Council, Office of Research and Development, U.S. Environmental Protection Agency, Washington, D.C. July 10.

U.S. EPA (1997). Health Effects Advisory Summary Tables (HEAST), FY 1997 Update, U.S. Environmental Protection Agency, Solid Waste and Emergency Response, 9200.6-303 (97-1), EPA-540-R-97-036, July 1997.

U.S. EPA (2007a). Inventory Update Reporting: Non-confidential IUR Production Volume Information (1986-2002). U.S. Environmental Protection Agency, Washington, D.C. Accessed at: <http://www.epa.gov/opptintr/iur/tools/data/2002-vol.htm>

U.S. EPA (2007b). 1,2,3-Trichloropropane (last revised 1990). Integrated Risk Information System (IRIS), Office of Research and Development, National Center for Environmental Assessment, U.S. Environmental Protection Agency, Washington, D.C. Accessed at: <http://www.epa.gov/iris/subst/0200.htm>.

Villeneuve DC, Chu I, Secours VE, Cote MG, Plaa GL, Valli VE (1985). Results of a 90-day toxicity study on 1,2,3- and 1,1,2-trichloropropane administered via the drinking water. *Sci Total Environ* 47:421-426.

Volp RF, Sipes IG, Falcoz C, Carter DE, Gross JF (1984). Disposition of 1,2,3-trichloropropane in the Fischer 344 rat: conventional and physiological pharmacokinetics. *Toxicol Appl Pharmacol* 75(1):8-17.

Weber GL, Sipes IG (1990). Covalent interactions of 1,2,3-trichloropropane with hepatic macromolecules: studies in the male F-344 rat. *Toxicol Appl Pharmacol* 104(3):395-402.

Weber GL, Sipes IG (1991). Rat hepatic DNA damage induced by 1,2,3-trichloropropane. *Adv Exp Med Biol* 283:853-855.

Weber GL, Sipes IG (1992). In vitro metabolism and bioactivation of 1,2,3-trichloropropane. *Toxicol Appl Pharmacol* 113(1):152-158.

WHO (2003). 1,2,3-Trichloropropane. Concise International Chemical Assessment Document 56, Geneva Switzerland, Accessed May 16, 2005 at: <http://www.who.int/ipcs/publications/cicad/en/cicad56.pdf>.