

DRAFT
For Review Only

Public Health Goal for
Methyl Tertiary Butyl Ether
(MTBE)
in Drinking Water

Prepared by
Pesticide and Environmental Toxicology Section
Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

June 1998

DRAFT

LIST OF CONTRIBUTORS

PREFACE

Drinking Water Public Health Goal of the Office of Environmental Health Hazard Assessment

This Public Health Goal (PHG) technical support document provides information on health effects from contaminants in drinking water. The PHG describes concentrations of contaminants at which adverse health effects would not be expected to occur, even over a lifetime of exposure. PHGs are developed for chemical contaminants based on the best available toxicological data in the scientific literature. These documents and the analyses contained in them provide estimates of the levels of contaminants in drinking water that would pose no significant health risk to individuals consuming the water on a daily basis over a lifetime.

The California Safe Drinking Water Act of 1996 (amended Health and Safety Code, Section 116365) requires the Office of Environmental Health Hazard Assessment (OEHHA) to adopt PHGs for contaminants in drinking water based exclusively on public health considerations. The Act requires OEHHA to adopt PHGs that meet the following criteria:

1. PHGs for acutely toxic substances shall be set at levels at which scientific evidence indicates that no known or anticipated adverse effects on health will occur, plus an adequate margin-of-safety.
2. PHGs for carcinogens or other substances which can cause chronic disease shall be based solely on health effects without regard to cost impacts and shall be set at levels which OEHHA has determined do not pose any significant risk to health.
3. To the extent the information is available, OEHHA shall consider possible synergistic effects resulting from exposure to two or more contaminants.
4. OEHHA shall consider the existence of groups in the population that are more susceptible to adverse effects of the contaminants than a normal healthy adult.
5. OEHHA shall consider the contaminant exposure and body burden levels that alter physiological function or structure in a manner that may significantly increase the risk of illness.
6. In cases of scientific ambiguity, OEHHA shall use criteria most protective of public health and shall incorporate uncertainty factors of noncarcinogenic substances for which scientific research indicates a safe dose-response threshold.
7. In cases where scientific evidence demonstrates that a safe dose-response threshold for a contaminant exists, then the PHG should be set at that threshold.
8. The PHG may be set at zero if necessary to satisfy the requirements listed above.
9. OEHHA shall consider exposure to contaminants in media other than drinking water, including food and air and the resulting body burden.
10. PHGs adopted by OEHHA shall be reviewed periodically and revised as necessary based on the availability of new scientific data.

PHGs adopted by OEHHA are for use by the California Department of Health Services (DHS) in establishing primary drinking water standards (State Maximum Contaminant Levels, or MCLs). Whereas PHGs are to be based solely on scientific and public health considerations without regard to economic cost considerations, drinking water standards adopted by DHS are to consider economic factors and technical feasibility. For this reason PHGs are only one part of the information used by DHS for establishing drinking water standards. PHGs established by OEHHA are not regulatory in nature and represent only non-mandatory goals. By federal law, MCLs established by DHS must be at least as stringent as the federal MCL if one exists.

PHG documents are developed for technical assistance to DHS, but may also benefit federal, state and local public health officials. While the PHGs are calculated for single chemicals only, they may, if the information is available, address hazards associated with the interactions of contaminants in mixtures. Further, PHGs are derived for drinking water only and are not to be utilized as target levels for the

DRAFT

contamination of environmental waters where additional concerns of bioaccumulation in fish and shellfish may pertain. Often environmental water contaminant criteria are more stringent than drinking water PHGs, to account for human exposures to a single chemical in multiple environmental media and from bioconcentration by plants and animals in the food chain.

OEHHA specifically requests comments on the following areas:

1. Considering the mandates of the California Health and Safety Code specifically Section 116300 (a), (d), (e), and (f) and Section 116365 (c), are the methods, assumptions and results of our analyses used in developing PHGs consistent with the intent of the statute? OEHHA solicits comments on toxicity study selection, no observed adverse effect levels (NOAELs), lowest observed adverse effect levels (LOAELs), uncertainty factors, severity of effect modifications, relative source contributions, multi-route exposure assumptions for volatile chemicals and other relevant assumptions and analyses for each chemical. Are the assumptions employed sufficiently health-protective in view of the statutory definition of the PHG (above)?
2. In proposing PHGs for carcinogens, OEHHA employed new methodology proposed by U.S. EPA in their 1996 Guidelines for Carcinogen Risk Assessment. These methods were applied to low-dose extrapolation and inter-species scaling and generally resulted in approximately two-fold lower estimated carcinogen potencies or slope factors as compared to earlier methods. We invite your comments on our methodology.
3. In developing the PHGs, OEHHA considered different levels of risk. Previously, when OEHHA developed Recommended Public Health Goals (RPHGs), the recommended levels were based on a 10^{-6} level of risk, a level that has been considered negligible or de minimis. This level corresponds to a theoretical extra lifetime cancer risk of 1×10^{-6} , or one fatal cancer per million exposed population over 70 years. This risk level has been identified by federal and state agencies as a level at or below which there are no public health concerns. Higher risk levels of 1×10^{-5} and 1×10^{-4} were also considered and are provided in the supporting documentation for the aid of risk managers. State law allows PHGs to be set at zero. U.S. EPA policy employs zero as a numerical goal (Maximum Contaminant Level goal, or MCLG) for drinking water standards for selected carcinogens. OEHHA welcomes comments on the various options that were considered in identifying the proposed PHGs presented in these technical support documents.

OEHHA expects the following process to pertain to these PHG documents:

- The Draft documents will be released for external peer review and public comment including a public workshop.
- Public comments will be received and reviewed by OEHHA and the documents revised as may be appropriate.
- In accordance with the Health and Safety Code Section 57003, the revised document drafts will be circulated for a period of 30 days following the public workshop.
- Following this 30-day comment period the documents will be finalized and the PHGs adopted by OEHHA.

LIST OF ABBREVIATIONS

AB	Assembly Bill
AL	Action Level
ACGIH	American Conference of Governmental Industrial Hygienists
API	American Petroleum Institute
ARB	California Air Resources Board
ATSDR	Agency for Toxic Substances and Disease Registry
AUC	area under the concentration-time curve
BAAQMD	Bay Area Air Quality Management District, San Francisco, California
BIBRA	British Industrial Biological Research Association
BTEX	benzene, toluene, ethylbenzene, and xylenes
BUN	blood urea nitrogen
BW	body weight
CAAA	1990 U.S. Clean Air Act Amendments
Cal/EPA	California Environmental Protection Agency
CAS	Chemical Abstracts Service
CCR	California Codes of Register
CDC	Centers for Disease Control and Prevention
CFS	chronic fatigue syndrome
CENR	Committee on Environment and Natural resources
CHRIS	Chemical Hazard Response Information System
CNS	central nervous system
CO	carbon monoxide
CSF	cancer slope factor, a cancer potency derived from the lower 95% confidence bound on the dose associated with a 10% (0.1) increased risk of cancer (LED ₁₀) calculated by the LMS model. CSF = 0.1/LED ₁₀ .
CPF	cancer potency factor, cancer potency, carcinogenic potency, or carcinogenic potency factor

DRAFT

DHS	California Department of Health Services
DOT	Department of Transportation
DOT/UN/NA/IMCO	Department of Transportation/United Nations/North America/ International Maritime Dangerous Goods Code
DLR	detection limit for purposes of reporting
DWC	daily water consumption
DWEL	Drinking Water Equivalent Level
EBMUD	East Bay Municipal Utility District
EHS	Extremely Hazardous Substances, SARA Title III
EOHSI	Environmental and Occupational Health Sciences Institute, New Jersey
ETBE	ethyl tertiary butyl ether
GAC	granulated activated charcoal
gd	gestation day
g/L	grams per liter
HA	Health Advisory
HAP	Hazardous Air Pollutant
HCHO	formaldehyde
HEI	Health Effects Institute
HSDB	Hazardous Substances Data Bank
IARC	International Agency for Research on Cancer
i.p.	intraperitoneal
IRIS	Integrated Risk Information Systems
i.v.	intravenous
kg	kilograms
L	liter
LC ₅₀	lethal concentrations with 50% kill
LD ₅₀	lethal doses with 50% kill
LED ₁₀	lower 95% confidence bound on the dose associated with a 10% increased risk of cancer
Leq/day	liter equivalent per day
LMS	linearized multistage
LOAEL	lowest observed adverse effect level

DRAFT

MCCHD	Missoula City-County Health Department, Montana
MCL	Maximum Contaminant Level
MCLG	Maximum Contaminant Level Goal
mg/L	milligrams per liter
µg/L	micrograms per liter
MCS	multiple chemical sensitivities
mL	milliliter
MOE	margin of exposure
MORS	Office of Research and Standards, Department of Environmental Protection, the Commonwealth of Massachusetts
MRL	minimal risk levels
MTBE	methyl tertiary butyl ether
MTD	maximum tolerated dose
MWDSC	Metropolitan Water District of Southern California
NAERG	North American Emergency Response Guidebook Documents
NAS	National Academy of Sciences
NAWQA	National Water-Quality Assessment
NCDEHNR	North Carolina Department of Environment, Health, and Natural Resources
NCEH	National Center for Environmental Health
NCI	National Cancer Institute
ng	nanograms
NIOSH	National Institute for Occupational Safety and Health
NJHSFS	New Jersey Hazardous Substance Fact Sheets
NJDWQI	New Jersey Drinking Water Quality Institute
NOAEL	no observed adverse effect levels
NOEL	no observed effect levels
NRC	National Research Council
NSTC	National Science and Technology Council
NTP	U.S. National Toxicology Program

DRAFT

OEHHA	Office of Environmental Health Hazard Assessment
OEL	Occupational Exposure Limit
OHM/TADS	Oil and Hazardous Materials/Technical Assistance Data System
OSTP	Office of Science and Technology Policy
O ₃	ozone
oxyfuel	oxygenated gasoline
PBPK	Physiologically-Based Pharmacokinetic
PHG	Public Health Goal
pnd	postnatal day
POTW	publicly owned treatment works
ppb	parts per billion
ppbv	ppb by volume
ppm	parts per million
ppt	parts per trillion
pptv	ppt by volume
Proposition 65	California Safe Drinking Water and Toxic Enforcement Act of 1986
q ₁ *	a cancer potency that is the upper 95% confidence limit of the low dose extrapolation on cancer potency slope calculated by the LMS model
RfC	Reference Concentration
RfD	Reference Dose
RFG	reformulated gasoline
RSC	relative source contribution
RTECS	Registry of Toxic Effects of Chemical Substances
SARA	Superfund (CERCLA) Amendments and Reauthorization Act of 1986
SB	Senate Bill
SCVWD	Santa Clara Valley Water District
SGOT	serum glutamic-oxaloacetic transaminase
SS	statistically significant
STEL	Short-Term Occupational Exposure Limit
Superfund	Comprehensive Environmental Response, Compensation and Liability Act of 1980, a.k.a. CERCLA
SWRCB	State Water Resources Control Board

DRAFT

TAC	toxic air contaminant
TAME	tertiary amyl methyl ether
TBA	tertiary butyl alcohol
TBF	tertiary butyl formate
TERIS	Teratogen Information System
TOMES	Toxicology and Occupational Medicine System
TRI	Toxics Release Inventory
TSCA	Toxic Substances Control Act
TWA	Time-Weighted Average
t_e	experimental duration
t_l	lifetime of the animal used in the experiment
$t_{1/2}$	plasma elimination half-life
UF	uncertainty factors
U.S.	United States
USCG	United States Coast Guard
U.S. EPA	United States Environmental Protection Agency
USGS	United States Geological Survey
UST	underground storage tanks
VOC	volatile organic compound
VRG	vessel rich group
WDOH	Wisconsin Division of Health, Department of Natural Resources
WHO	World Health Organization

TABLE OF CONTENTS

LIST OF CONTRIBUTORS.....	ii
PREFACE.....	iii
LIST OF ABBREVIATIONS.....	v
TABLE OF CONTENTS	x
SUMMARY	1
INTRODUCTION.....	1
CHEMICAL PROFILE	4
CHEMICAL IDENTITY	4
PHYSICAL AND CHEMICAL PROPERTIES	5
ORGANOLEPTIC PROPERTIES.....	5
PRODUCTION AND USES	10
ENVIRONMENTAL OCCURRENCE AND HUMAN EXPOSURE.....	11
AIR, SOIL, FOOD, AND OTHER SOURCES	12
WATER	13
METABOLISM AND PHARMACOKINETICS.....	16
ABSORPTION	17
DISTRIBUTION	17
METABOLISM.....	17
EXCRETION	18
PHARMACOKINETICS	19
PHYSIOLOGICALLY-BASED PHARMACOKINETIC (PBPK) MODELS	20
TOXICOLOGY.....	20
TOXICOLOGICAL EFFECTS IN ANIMALS.....	21
Acute Toxicity	27
Subacute Toxicity	28
Subchronic Toxicity.....	28
Genetic Toxicity	29
Developmental and Reproductive Toxicity	30
Immunotoxicity.....	44
Neurotoxicity	45
Chronic Toxicity.....	45
Carcinogenicity.....	46

DRAFT

Ecotoxicity.....	57
TOXICOLOGICAL EFFECTS IN HUMANS.....	58
Acute Toxicity	58
Immunotoxicity.....	62
Neurotoxicity	62
DOSE-RESPONSE ASSESSMENT	63
INTERNAL DOSE ESTIMATION.....	63
NONCARCINOGENIC EFFECTS	68
CARCINOGENIC EFFECTS	68
Possible Modes of Action	68
Estimation of Carcinogenic Potency.....	68
CALCULATION OF PHG	74
NONCARCINOGENIC EFFECTS	74
EXPOSURE FACTORS	75
CARCINOGENIC EFFECTS	79
RISK CHARACTERIZATION.....	80
ACUTE HEALTH EFFECTS	80
CARCINOGENIC EFFECTS	80
OTHER REGULATORY STANDARDS.....	83
REFERENCES	86

Summary

A Public Health Goal (PHG) of 0.014 mg/L (14 µg/L or 14 ppb) is proposed for methyl tertiary butyl ether (MTBE) in drinking water. The PHG is based on carcinogenic effects observed in experimental animals. Carcinogenicity has been observed in both sexes of the rat in a lifetime gavage study (Belpoggi et al. 1995, 1997), in male rats of a different strain in a 24-month inhalation study (Chun et al. 1992, Bird et al. 1997), and in male and female mice in an 18-month inhalation study (Burleigh-Flayer et al. 1992, Bird et al. 1997). In Sprague-Dawley rats receiving MTBE by gavage, statistically significant increases in Leydig interstitial cell tumors of the testes were observed in males, and statistically significant increases in lymphomas and leukemias (combined) were observed in females. In Fischer 344 rats exposed to MTBE by inhalation, statistically significant increases in the incidences of Leydig interstitial cell tumors of the testes were also observed in males, as well as renal tubular tumors. In CD-1 mice exposed to MTBE by inhalation, statistically significant increases in the incidences of liver tumors were observed in females (hepatocellular adenomas; hepatocellular adenomas and carcinomas combined) and males (hepatocellular carcinomas; hepatocellular adenomas and carcinomas combined). The two inhalation studies (Burleigh-Flayer et al. 1992, Chun et al. 1992, Bird et al. 1997) and a gavage study (Belpoggi et al. 1995, 1997) cited in this document for the development of the PHG provided evidence for the carcinogenicity of MTBE at multiple sites in both sexes of the rat and mouse; MTBE is a carcinogen in two species, both sexes and multi-sites.

For the calculation of the PHG, cancer potency estimates were made, based on the recommended practices of the 1996 United States Environmental Protection Agency (U.S. EPA) proposed draft guidelines for carcinogenic risk assessment (U.S. EPA 1996f), in which the linearized multistage (LMS) model is fit to the experimental data in order to establish the lower 95% confidence bound on the dose associated with a 10% increased risk of cancer (LED₁₀). It is plausible that the true value of the human cancer potency has a lower bound of zero based on statistical and biological uncertainties. The PHG was calculated assuming a de minimis theoretical excess individual cancer risk level of 10⁻⁶ from exposure to MTBE. Based on these considerations, OEHHA proposes a PHG of 0.014 mg/L (14 µg/L or 14 ppb) for MTBE in drinking water. The range of possible values based either on different individual tumor sites or on different multiroute exposure estimates and the average cancer potency of the three sites was 2.5 to 18 ppb. The proposed PHG is considered to contain an adequate margin of safety for the potential noncarcinogenic effects including adverse effects on the renal, neurological and reproductive systems.

In addition to the 14 ppb value based on carcinogenicity, a value of 0.047 mg/L (47 ppb) was calculated based on non-cancer effects of increased relative kidney weights in the Robinson et al. (1990) 90-day gavage study in rats. This value incorporates four 10-fold uncertainty factors for a less than lifetime study, interspecies and interindividual variation and possible carcinogenicity. While the lower value of 14 ppb is proposed as the PHG the difference in the two approaches is only three-fold.

INTRODUCTION

The purpose of this document is to establish a PHG for the gasoline additive MTBE in drinking water. MTBE is a synthetic solvent used primarily as an oxygenate in unleaded gasoline to boost

DRAFT

octane and improve combustion efficacy by oxygenation. Reformulated fuel with MTBE has been used in 32 regions in 18 states in the United States (U.S.) to meet the 1990 federal Clean Air Act Amendments (CAAA) requirements for reducing carbon monoxide (CO) and ozone (O₃) levels because the added oxygenate promotes more complete burning of gasoline. MTBE is currently used (11% by volume) in California's cleaner-burning reformulated gasoline (California RFG) to improve air quality (Denton and Masur 1996). California is the third largest consumer of gasoline in the world. It is surpassed only by the rest of the U.S. and the former Soviet Union. Californians use more than 13.7 billion gallons of gasoline a year and another one billion gallons of diesel fuel. MTBE and other oxygenates such as ethyl tertiary butyl ether (ETBE) and ethanol are currently being studied to determine the extent of their presence in drinking water and what, if any, potential health implications could result from exposure to them (Freed 1997, Scheible 1997).

MTBE was the second most-produced chemical in the U.S. in 1997, whereas previously it was ranked the twelfth in 1995 and eighteenth in 1994 (Cal/EPA 1998, Kirschner 1996, Reisch 1994). In 1994 and 1995, it was estimated that about 70 million Americans were exposed to oxygenated gasoline (oxyfuel) and approximately 57 million were exposed to reformulated gasoline (RFG) (ATSDR 1996, HEI 1996, NRC 1996, NSTC 1996, 1997). About 40% of the U.S. population live in areas where MTBE is used in oxyfuel or RFG (USGS 1996) and most people find its distinctive terpene-like odor disagreeable (CDC 1993a, 1993b, 1993c, Kneiss 1995, Medlin 1995, U.S. EPA 1997a). MTBE is now being found in the environment in many areas of the U.S. because of its increased use over the last several years. Recently it has become a drinking water contaminant due to its high water solubility and persistence. When gasoline with 10% MTBE by weight comes in contact with water, about 5 grams per liter (g/L) can dissolve (Squillace et al. 1996, 1997a). MTBE has been detected in groundwater as a result of leaking underground storage tanks (USTs) or pipelines and in surface water reservoirs via recreational boating activities. MTBE does not appear to adsorb to soil particles or readily degrade in the subsurface environment. It is more expensive to remove MTBE-added gasoline than gasoline without MTBE from contaminated water (Cal/EPA 1998, U.S. EPA 1987a, 1992c, 1996a, 1997a).

MTBE is not regulated currently under the federal and the California drinking water regulations.

However, an interim non-enforceable Action Level (AL) of 0.035 mg/L (35 µg/L or 35 ppb) in drinking water was established by the California Department of Health Services (DHS) in 1991 to protect against adverse health effects. The Office of Environmental Health Hazard Assessment (OEHHA) recommended this level (OEHHA 1991) based on noncarcinogenic effects of MTBE in laboratory animals (Greenough et al. 1980). OEHHA applied large uncertainty factors to provide a substantial margin of safety for drinking water. Since February 13, 1997, DHS (1997) regulations (22 CCR Section 64450) have included MTBE as an unregulated chemical for which monitoring is required. Pursuant to this requirement, data on the occurrence of MTBE in groundwater and surface water sources are being collected from drinking water systems in order to document the extent of MTBE contamination in drinking water supplies.

In California, the Local Drinking Water Protection Act of 1997 (SB 1189, Hayden, and AB 592, Kuehl) requires DHS to develop a two-part drinking water standard for MTBE. The first part is a secondary maximum contaminant level (MCL) that addresses aesthetic qualities including taste and odor, to be established by July 1, 1998. The second part is a primary MCL that addresses health concerns, to be established by July 1, 1999. DHS is proceeding to establish drinking water standards for MTBE with a request to OEHHA to conduct a risk assessment by July 1998.

in order to meet the mandated schedule to set this regulation by July 1999. This Act also requires the State's qualified experts to evaluate MTBE for listing under the Safe Drinking Water and Toxic Enforcement Act of 1986 (Proposition 65) as a chemical known to the state to cause cancer or reproductive and developmental toxicity. OEHHA is responsible for these hazard identification and listing activities.

The U.S. EPA has not established primary or secondary MCLs or a Maximum Contaminant Level Goal (MCLG) for MTBE but included MTBE on the Contaminant Candidate List published in the Federal Register on March 2, 1998 (U.S. EPA 1998, 1997b, 1997d). An advisory released in December 1997 recommended that MTBE concentrations in the range of 20 to 40 ppb or below would assure both consumer acceptance of the water and a large margin of safety from any toxic effects (U.S. EPA 1997a, Du et al. 1998). U.S. EPA (1994a, 1994c) proposed to classify MTBE as a Group C possible human carcinogen in 1994 based upon animal inhalation studies published in 1992. The U.S. EPA noted that a Group B2 probable human carcinogen designation may be appropriate if oral MTBE exposure studies in animals (published later in 1995) result in treatment-related tumors. In 1987, MTBE was identified by the U.S. EPA (1987a) under Section Four of the Toxic Substances Control Act (TSCA) for priority testing because of its large production volume, potential widespread exposure, and limited data on long-term health effects (Duffy et al. 1992). The results of the testing have been published in a peer-reviewed journal (Bevan et al. 1997a, 1997b, Bird et al. 1997, Daughtrey et al. 1997, Lington et al. 1997, McKee et al. 1997, Miller et al. 1997, Stern and Kneiss 1997).

The American Conference of Governmental Industrial Hygienists (ACGIH) lists MTBE as an A3 Animal Carcinogen (ACGIH 1996). The U.S. National Toxicology Program (NTP) has listed MTBE as a candidate to be considered for testing and listing as "reasonably anticipated to be a human carcinogen based on positive carcinogenicity findings in laboratory animal studies". The U.S. EPA nominated MTBE for review in 1998 to NTP as an addition to their Ninth Edition Report on Carcinogens (NTP 1998). ACGIH (1996) indicated that the International Agency for Research on Cancer (IARC) classified MTBE as a Group 2B carcinogen, possibly carcinogenic to humans, however, no such classification has been published by IARC to date, based on a thorough search of IARC publications including a search of the IARC website. IARC has neither evaluated nor classified MTBE as to its carcinogenicity (IARC 1987, 1995) and is planning to perform the evaluation in the near future. MTBE has been reviewed by the Environmental Epidemiology Section of the North Carolina Department of Environment, Health, and Natural Resources (NCDEHNR) and it was determined that there was limited evidence for carcinogenicity in experimental animals and that the compound should be classified as a Group B2 probable human carcinogen (Rudo 1995). The North Carolina Scientific Advisory Board on Toxic Air Contaminants (TAC) considered MTBE to be eligible as a Group C possible human carcinogen (Lucier et al. 1995). In addition to the U.S. EPA advisory document and the TSCA testing program report mentioned above, five recent documents on MTBE have received nationwide attention. In February 1996 the Office of Science and Technology Policy (OSTP) through the Committee on Environment and Natural Resources (CENR) of the White House National Science and Technology Council (NSTC) released a draft report titled "Interagency Assessment of Potential Health Risks Associated with Oxygenated Gasoline" (NSTC 1996). This report focused primarily on inhalation exposure to MTBE and its principal metabolite, tertiary butyl alcohol (TBA). In March 1996 NSTC released the draft document "Interagency Oxygenated Fuels Assessment" which addressed issues related to public health, air and water quality, fuel economy, and engine performance associated with MTBE in gasoline relative to conventional gasoline. This draft was

finalized in 1997 (NSTC 1997). These two documents were followed by the April 1996 release of the Health Effects Institute (HEI) document "The Potential Health Effects of Oxygenates Added to Gasoline, A Special Report of the Institute's Oxygenates Evaluation Committee" (HEI 1996). HEI (1996) concluded "the possibility that ambient levels may pose some risk of carcinogenic effects in human populations cannot be excluded".

In August 1996 the Agency for Toxic Substances and Disease Registry (ATSDR) released the final report "Toxicological Profile for MTBE" which evaluated the toxic effects of MTBE in detail (ATSDR 1996). The latter 1996 NSTC draft document was peer reviewed by the National Academy of Sciences (NAS) under guidance from the National Research Council (NRC) which then published its findings and recommendations in the document "Toxicological and Performance Aspects of Oxygenated Motor Vehicle Fuels" (NRC 1996). The limited review on the potential health effects of MTBE in the NRC report (1996) considered the animal carcinogenicity to be positive. The NRC findings were used to revise the NSTC document and the final report was released in June of 1997. The NSTC (1997) concluded "there is sufficient evidence that MTBE is an animal carcinogen".

California Environmental Protection Agency (Cal/EPA) has reported some background information and ongoing activities on MTBE in California's "cleaner-burning fuel program" in a briefing paper (Cal/EPA 1998). California Senate Office of Research also released a position paper on MTBE (Wiley 1998). U.S. EPA (1996d, 1996e) published fact sheets on MTBE in water in addition to several advisory documents. While concerns have been raised about its potential health impacts, based on hazard evaluation of the available data, MTBE is substantially less hazardous than benzene (a Group A human carcinogen) and 1,3-butadiene (a Group B2 probable human carcinogen), two carcinogenic chemicals it displaces in California's new gasoline formulations (Spitzer 1997). Potential health benefits from ambient O₃ reduction related to the use of MTBE in RFG were evaluated (Erdal et al. 1997). Whether the addition of MTBE in gasoline represents a net increase in cancer hazard is beyond the scope of this document. However, NSTC (1997) concluded: "... the weight of evidence supports regarding MTBE as having a carcinogenic hazard potential for humans." U.S. EPA (1997a) agreed with the NSTC and concluded: "The weight of evidence indicates that MTBE is an animal carcinogen, and the chemical poses a carcinogenic potential to humans."

In this document, the available data on the toxicity of MTBE primarily by the oral route based on the five above-mentioned reports are evaluated, and information available since the previous assessment by NSTC (1997) and U.S. EPA (1997a) is included. To determine a public health-protective level of MTBE in drinking water, relevant studies were identified, reviewed and evaluated, and sensitive groups and exposure scenarios are considered.

chemical profile

CHEMICAL IDENTITY

MTBE [(CH₃)₃C(OCH₃), CAS Registry Number 1634-04-4] is a synthetic chemical without known natural sources. The chemical structure, synonyms, and identification numbers are listed in Table 1 and are adapted from the Merck Index (1989), Hazardous Substances Data Bank (HSDB) of the National Library of Medicine (1997), Integrated Risk Information Systems (IRIS) of U.S. EPA (1997c), TOMES PLUS® (Hall and Rumack 1998) computerized database, and the

ATSDR (1996), Cal/EPA (1998), HEI (1996), NRC (1996), NSTC (1996, 1997), and U.S. EPA (1997a) documents.

TOMES (Toxicology and Occupational Medicine System) PLUS® is a computerized database which includes the data systems of Hazard Management®, Medical Management®, INFOTEXT®, HAZARDTEXT®, MEDITEXT®, REPROTEXT®, SERATEXT®, HSDB, IRIS, Registry of Toxic Effects of Chemical Substances (RTECS®) of National Institute for Occupational Safety and Health (NIOSH), Chemical Hazard Response Information System (CHRIS) of U.S. Coast Guard, Oil and Hazardous Materials/Technical Assistance Data System (OHM/TADS) of U.S. EPA, Department of Transportation (DOT) Emergency Response Guide, New Jersey Hazardous Substance Fact Sheets (NJHSFS), North American Emergency Response Guidebook Documents (NAERG) of U.S. DOT, Transport Canada and the Secretariat of Communications and Transportation of Mexico, REPROTOX® System of the Georgetown University, Shepard's Catalog of Teratogenic Agents of the Johns Hopkins University, Teratogen Information System (TERIS) of the University of Washington, and NIOSH Pocket Guide(TM). For MTBE, TOMES PLUS® (Hall and Rumack 1998) contains entries in HAZARDTEXT®, MEDITEXT®, REPROTEXT®, REPROTOX®, HSDB, IRIS, RTECS®, NAERG and NJHSFS.

PHYSICAL AND CHEMICAL PROPERTIES

Important physical and chemical properties of MTBE are given in Table 2 and are adapted from Merck Index (1989), HSDB (1997), TOMES PLUS® (Hall and Rumack 1998), and the ATSDR (1996), Cal/EPA (1998), HEI (1996), NRC (1996), NSTC (1996, 1997), and U.S. EPA (1997a) documents.

MTBE, an aliphatic ether, is a volatile organic compound (VOC) with a characteristic odor. It is a colorless liquid at room temperature. It is highly flammable and combustible when exposed to heat or flame or spark, and is a moderate fire risk. Vapors may form explosive mixtures with air. It is unstable in acid solutions. Fire may produce irritating, corrosive or toxic gases. Runoff from fire control may contain MTBE and its combustion products (HSDB 1997).

MTBE is miscible in gasoline and soluble in water, alcohol, and other ethers. It has a molecular weight of 88.15 daltons, a vapor pressure of about 245 mmHg at 25 °C, an octane number of 110, and solubility in water of about 50 g/L at 25 °C. It disperses evenly in gasoline and water and stays suspended without requiring physical mixing. It does not increase volatility of other gasoline components when it is mixed in the gasoline. MTBE released to the environment via surface spills or subsurface leaks was found to initially partition between water and air (Jeffrey 1997). The log of the octanol-water partition coefficient ($\log K_{ow}$) is reported to range from 0.94 to 1.24 which indicates that there is 10 times more partitioning of MTBE in the lipophilic phase than in the aqueous phase of solvents. The molecular size and $\log K_{ow}$ of MTBE are characteristic of molecules which are able to penetrate across biological membranes of the skin, lungs and gastrointestinal tracts (Mackay et al. 1993, Nihlen et al. 1995). The octanol-water partition coefficient is reported to be 16 by Nihlen et al. (1997). Fujiwara et al. (1984) reported laboratory-derived octanol-water partition coefficients ranging from 17.2 to 17.5 with a $\log K_{ow}$ of 1.2. The blood-air, urine-air, saline-air, fat-air and oil-air partition coefficients (λ) are reported to be 20, 15.6, 15.3, 142 and 138, respectively (Imbriani et al. 1997). One part per million (1 ppm) of MTBE, volume to volume in air, is approximately 3.6 mg/m³ of air at 20 °C (ATSDR 1996).

ORGANOLEPTIC PROPERTIES

Taste or odor characteristics, often referred to as organoleptic properties, are not used by U.S. EPA or DHS for developing primary drinking water standards, but are used for developing secondary standards. The estimated thresholds for these properties of MTBE reported in the literature are given in Table 3 and are adapted from the ATSDR (1996), Cal/EPA (1998), HEI (1996), HSDB (1997), NSTC (1996, 1997), and U.S. EPA (1997a) documents. Taste and odor may alert consumers to the fact that the water is contaminated with MTBE (Angle 1991) and many people object to the taste and odor of MTBE in drinking water (Killian 1998, Reynolds 1998). However, not all individuals respond equally to taste and odor because of differences in individual sensitivity. It is not possible to identify point threshold values for the taste and odor of MTBE in drinking water, as the concentration will vary for different individuals, for the same individuals at different times, for different populations, and for different water matrices, temperatures, and many other variables. The odor threshold ranges from about 0.32 to 0.47 mg/m³ (about 90 to 130 ppb) in air and can be as low as five ppb (about 0.02 mg/m³) for some sensitive people. In gasoline containing 97% pure MTBE at mixture concentrations of three percent, 11% and 15% MTBE, the threshold for detecting MTBE odor in air was estimated to be 50 ppb (about 0.18 mg/m³), 280 ppb (about 1 mg/m³), and 260 ppb (about 0.9 mg/m³), respectively (ACGIH 1996). A range of five ppb to 53 ppb (about 0.19 mg/m³) odor threshold in the air was reported in an American Petroleum Institute (API) document (API 1994).

The individual taste and odor responses reported for MTBE in water are on average in the 15 to 180 ppb (µg/L) range for odor and the 24 to 135 ppb range for taste (API 1994, Prah et al. 1994, Young et al. 1996, Dale et al. 1997b, Shen et al. 1997, NSTC 1997). The ranges are indicative of the average variability in individual response. U.S. EPA (1997a) has analyzed these studies in detail and recommended a range of 20 to 40 ppb as an approximate threshold for organoleptic responses. The study (Dale et al. 1997b) by the Metropolitan Water District of Southern California (MWDSC) found people more sensitive to the taste than odor. This result is consistent with API's (1994) findings for MTBE taste and odor thresholds. But in the study by Young et al. (1996), test subjects were more sensitive to odor than taste.

The subjects described the taste of MTBE in water as "nasty", "bitter", "nauseating", and "similar to rubbing alcohol" (API 1994). It is noted that chlorination and temperature of the water would likely affect the taste and odor of MTBE in water. Thresholds for the taste and odor of MTBE in chlorinated water would be higher than those of MTBE in nonchlorinated water. Thresholds for the taste and odor of MTBE in water at higher temperatures (e.g., for showering) would likely be lower than those of MTBE in water at lower temperatures. There were undoubtedly individuals who could only detect the odor of MTBE at even higher concentrations than 180 ppb (Prah et al. 1994). Odor thresholds as high as 680 ppb have been reported (Gilbert and Calabrese 1992). On the other hand, some subjects in these studies were able to detect the odor of MTBE in water at much lower concentrations, i.e. 2.5 ppb (Shen et al. 1997), five ppb (McKinnon and Dyksen 1984), or 15 ppb (Young et al. 1996). Some sensitive subjects in the taste studies were able to detect MTBE in water at concentrations as low as two ppb (Dale et al. 1997b), 10 ppb (Barker et al. 1990), 21 ppb (Dale et al. 1997b), or 39 ppb (Young et al. 1996). Thus, in a general population, some unknown percentage of people will be likely to detect the taste and odor of MTBE in drinking water at concentrations below the U.S. EPA (1997a) 20 to 40 ppb advisory level. DHS (1997) has recently proposed five ppb as the secondary MCL for MTBE. The lowest olfaction threshold is likely to be at or about 2.5 ppb (Shen et al. 1997). The lowest taste threshold is likely to be at or about two ppb (Dale et al. 1997b).

Table 1. Chemical Identity of Methyl Tertiary Butyl Ether (MTBE)

Characteristic	Information	Reference
Chemical Name	Methyl tertiary butyl ether	Merck 1989
Synonyms	Methyl tertiary-butyl ether; methyl tert-butyl ether; tert-butyl methyl ether; tertiary-butyl methyl ether; methyl-1,1-dimethylethyl ether; 2-methoxy-2-methylpropane; 2-methyl-2-methoxypropane; methyl t-butyl ether; MtBE; MTBE	Merck 1989
Registered trade names	No data	
Chemical formula	C ₅ H ₁₂ O or (CH ₃) ₃ C(OCH ₃)	Merck 1989
Chemical structure	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{CH}_3 - \text{C} - \text{O} - \text{CH}_3 \\ \\ \text{CH}_3 \end{array} $	
Identification numbers:		
Chemical Abstracts Service (CAS) Registry number	1634-04-4	Merck 1989
National Institute for Occupational Safety and Health (NIOSH) Registry of Toxic Effects of Chemical Substances (RTECS) number	KN5250000	HSDB 1997
Department of Transportation/United Nations/North America/International Maritime Dangerous Goods Code (DOT/UN/NA/IMCO) Shipping number	UN 2398, IMO 3.2	HSDB 1997
Hazardous Substances Data Bank (HSDB) number	5847	HSDB 1997
North American Emergency Response Guidebook Documents (NAERG) number	127	HSDB 1997
National Cancer Institute (NCI) number	No data	
U.S. Environmental Protection Agency (U.S. EPA) Hazardous Waste number	No data	
U.S. EPA Oil and Hazardous Materials/ Technical Assistance Data System (OHM/TADS) number	No data	

Table 2. Chemical and Physical Properties of MTBE

Property	Value or information	Reference
Molecular weight	88.15 g/mole	Merck 1989
Color	colorless	Merck 1989
Physical state	liquid	Merck 1989
Melting point	-109 °C	HSDB 1997
Boiling point	53.6 - 55.2 °C	Mackay et al. 1993
Density at 20 °C	0.7404 - 0.7578 g/mL	Squillace et al. 1997a
Solubility		
in water	4.8 g/100 g water	Merck 1989
in water	23.2 - 54.4 g/L water	Garrett et al. 1986, Mackay et al. 1993
in water	43 - 54.3 g/L water	Squillace et al. 1997a
in water, 20 °C	4 - 5%	Gilbert and Calabrese 1992
in water, 25 °C	51 g/L water	HSDB 1997
Partition coefficients		
octanol-water	16	Nihlen et al. 1997
	17.2 - 17.5	Fujiwara et al. 1984
Log K _{ow}	0.94 - 1.16	Mackay et al. 1993
	1.2	Fujiwara et al. 1984
	1.24	U.S. EPA 1997a
Log K _{oc}	1.05 (estimated)	Squillace et al. 1997a
	2.89 (calculated)	U.S. EPA 1995b
Vapor pressure		
at 25 °C	245 - 251 mm Hg	Mackay et al. 1993
at 100 °F	7.8 psi (Reid Vapor Pressure)	ARCO 1995a
Henry's law constant	0.00058 - 0.003 atm-m ³ /mole	Mackay et al. 1993
at 25 °C	5.87 × 10 ⁻⁴ atm-m ³ /mole	ATSDR 1996
at 15 °C	0.011 (dimensionless)	Robbins et al. 1993
Ignition temperature	224 °C	Merck 1989
Flash point	-28 °C	Merck 1989
	28 °C (closed cup)	Gilbert and Calabrese 1992
Explosion limits	1.65 to 8.4% in air	Gilbert and Calabrese 1992
Heat of combustion	101,000 Btu/gal at 25 °C	HSDB 1997
Heat of vaporization	145 Btu/lb at 55 °C	HSDB 1997
Stability	MTBE is unstable in acidic solution	Merck 1989
Conversion factors		
ppm (v/v) to mg/m ³ in air at 25 °C	1 ppm = 3.61 mg/m ³	ACGIH 1996
mg/m ³ to ppm (v/v) in air at 25 °C	1 mg/m ³ = 0.28 ppm	ACGIH 1996

Table 3. Organoleptic Properties of MTBE

Property	Value or information	Reference
Odor	terpene-like at 25 °C	Gilbert and Calabrese 1992
Threshold in air	300 ppb	Smith and Duffy 1995
	0.32 - 0.47 mg/m ³	ACGIH 1996
	(~90 - 130 ppb)	
	5 - 53 ppb (detection)	API 1994
99% pure MTBE	8 ppb (recognition)	API 1994
97% pure MTBE	125 ppb (recognition)	API 1994
97% pure MTBE in gasoline		
15% MTBE	260 ppb	ACGIH 1996
11% MTBE	280 ppb	ACGIH 1996
3% MTBE	50 ppb	ACGIH 1996
Threshold in water	680 ppb	Gilbert and Calabrese 1992
	180 ppb	Prah et al. 1994
	95 ppb	ARCO 1995a
	55 ppb (recognition)	API 1994
	45 ppb (detection)	API 1994
	15 - 95 ppb (mean 34 ppb)	Young et al. 1996
	15 - 180 ppb	U.S. EPA 1997a
	13.5 - 45.4 ppb	Shen et al. 1997
	5 - 15 ppb	McKinnon and Dyksen 1984
	2.5 ppb	Shen et al. 1997
Taste	solvent-like at 25 °C	U.S. EPA 1997a
Threshold in water	21 - 190 ppb	Dale et al. 1997b
	24 - 135 ppb	U.S. EPA 1997a
	39 - 134 ppb (mean 48 ppb)	Young et al. 1996
	39 - 134 ppb	API 1994
	10 - 100 ppb	Barker et al. 1990
	2 ppb (one subject)	Dale et al. 1997b

PRODUCTION AND USES

MTBE is manufactured from isobutene, also known as isobutylene or 2-methylpropene (Merck 1989), which is a product of petroleum refining. It is made mainly by combining methanol with isobutene, or derived from combining methanol and TBA. It is used primarily as an oxygenate in unleaded gasoline, in the manufacture of isobutene, and as a chromatographic eluent especially in high pressure liquid chromatography (ATSDR 1996, HSDB 1997). MTBE also has had a limited use as a therapeutic drug for dissolving cholesterol gallbladder stones (Leuschner et al. 1994).

MTBE is the primary oxygenate used in gasoline because it is the least expensive and in greatest supply. It is promoted as a gasoline blending component due to its high octane rating, low cost of production, ability to readily mix with other gasoline components, ease in distribution through existing pipelines, distillation temperature depression, and beneficial dilution effect on undesirable components of aromatics, sulfur, olefin and benzene. In addition, the relatively low co-solvent volatility of MTBE does not result in a more volatile gasoline that could be hazardous in terms of flammability and explosivity. The use of MTBE has helped offset the octane specification loss due to the discontinued use of higher toxicity high octane aromatics and has reduced emissions of benzene, a known human carcinogen, and 1,3-butadiene, an animal carcinogen (Cal/EPA 1998, Spitzer 1997).

MTBE has been commercially used in Europe since 1973 as an octane enhancer to replace lead in gasoline and was approved as a blending component in 1979 by U.S. EPA. Since the early 1990s, it has been used in reformulated fuel in 18 states in the U.S. Under Section 211 of the 1990 CAAA, the federal oxyfuel program began requiring gasoline to contain 2.7% oxygen by weight which is equivalent to roughly 15% by volume of MTBE be used during the four winter months in regions not meeting CO reduction standards in November 1992. In January 1995, the federal RFG containing 2% oxygen by weight or roughly 11% of MTBE by volume was required year-round to reduce O₃ levels. Oxygenates are added to more than 30% of the gasoline used in the U.S. and this proportion is expected to rise (Squillace et al. 1997a).

In California, federal law required the use of Phase I RFG in the worst polluted areas including Los Angeles and San Diego as of January 1, 1995, and in the entire state as of January 1, 1996. By June 1, 1996, state law required that all gasoline sold be California Phase 2 RFG and federal Phase II RFG will be required by the year 2000 (Cornitius 1996). MTBE promotes more complete burning of gasoline, thereby reducing CO and O₃ levels in localities which do not meet the National Ambient Air Quality Standards (ATSDR 1996, USGS 1996). Almost all of the MTBE produced is used as a gasoline additive; small amounts are used by laboratory scientists (ATSDR 1996). When used as a gasoline additive, MTBE may constitute up to 15% volume to volume of the gasoline mixture. Currently, MTBE is added to virtually all of the gasoline consumed in California (Cal/EPA 1998).

The amount of MTBE used in the U.S. has increased from about 0.5 million gallons per day in 1980 to over 10 million gallons per day in early 1997. Of the total amount of MTBE used in the U.S., approximately 70% is produced domestically, about 29% is imported from other countries, and about 1% is existing stocks. Over 4.1 billion gallons of MTBE are consumed in the U.S. annually, including 1.49 billion gallons -- more than 36% of the national figure -- in California (Wiley 1998). California uses about 4.2 million gallons per day of MTBE, about 85% of which is imported into the state, primarily by ocean tankers from the Middle East (Cal/EPA 1998). California also imports MTBE from Texas and other major MTBE-producing states in the U.S.

MTBE production in the U.S. began in 1979 and increased rapidly after 1983. It was the second most-produced chemical, in terms of amount, in the U.S. in 1997, whereas previously it was ranked the twelfth in 1995 and eighteenth in 1994 (Cal/EPA 1998, Kirschner 1996, Reisch 1994). The production was 13.61 million pounds in 1994 and 17.62 million pounds in 1995 (Kirschner 1996). MTBE production was estimated at about 2.9 billion gallons in the U.S. and about 181 million gallons in California in 1997 (Wiley 1998). MTBE is manufactured at more than 40 facilities by about 27 producers primarily concentrated along the Houston Ship Channel in Texas and the Louisiana Gulf Coast. Texas supplies about 80% of the MTBE produced in the U.S. with about 10% produced in Louisiana and about five

percent in California (Cal/EPA 1998). The major portion of MTBE produced utilizes, as a co-reactant, isobutylene which is a waste product of the refining process (Wiley 1998).

ENVIRONMENTAL OCCURRENCE AND HUMAN EXPOSURE

The recent NSTC (1997) report provides extensive occurrence data for MTBE and other fuel oxygenates, as well as information on applicable treatment technologies. For additional information concerning MTBE in the environment, this report can be accessed through the NSTC Home Page via a link from the OSTP. The U.S. Geological Survey (USGS) has been compiling data sets for national assessment of MTBE and other VOCs in ground and surface water as part of the National Water-Quality Assessment (NAWQA) Program (Buxton et al. 1997, Lapham et al. 1997, Squillace et al. 1997a, 1997b, Zogorski et al. 1996, 1997). Information on analytical methods for determining MTBE in environmental media is compiled in the ATSDR (1996) Toxicological Profile document.

The U.S. EPA (1993, 1995a) estimated that about 1.7 million kilograms (kgs) MTBE were released from 141 facilities reporting in the Toxics Release Inventory (TRI) per year, 97.3% to air, 2.44% to surface water, 0.25% to underground injection, and 0.01% to land. Cohen (1998) reported that an estimated 27,000 kgs or 30 tons per day were emitted from 9,000 tons of MTBE consumed in California per day. The California Air Resources Board (ARB) estimated that the exhaust and evaporative emission was about 39,000 kgs or 43 tons per day in California in 1996 (Cal/EPA 1998).

A multimedia assessment of refinery emissions in the Yorktown region (Cohen et al. 1991) indicated that the MTBE mass distribution was over 73% in water, about 25% in air, less than two percent in soil, about 0.02% in sediment, about 10^{-6} % in suspended solids, and 10^{-7} % in biota. A recent laboratory study on liquid-gas partitioning (Rousch and Sommerfeld 1998) suggests that dissolved MTBE concentrations can vary substantially from nominal. The main route of exposure for occupational and non-occupational groups is via inhalation, ingestion is considered as secondary, and dermal contact is also possible. The persistence half-life of MTBE (Jeffrey 1997) is about four weeks to six months in soil, about four weeks to six months in surface water, and about eight weeks to 12 months in groundwater based on estimated anaerobic biodegradation, and about 20.7 hours to 11 days in air based on measured photooxidation rate constants (Howard et al. 1991, Howard 1993). Church et al. (1997) described an analytical method for detecting MTBE and other major oxygenates and their degradation products in water at sub-ppb concentrations. MTBE appears to be biodegraded under anaerobic conditions (Borden et al. 1997, Daniel 1995, Jensen and Arvin 1990, Mormile et al. 1994, Steffan et al. 1997). Brown et al. (1997) and Davidson and Parsons (1996) reviewed state-of-the-art remediation technologies for treatment of MTBE in water. The removal of MTBE from groundwater through aeration plus granulated activated charcoal (GAC) was

described by McKinnon and Dyksen (1984). Koenigsberg (1997) described a newly developed bioremediation technology for MTBE cleanup in groundwater.

AIR, SOIL, FOOD, AND OTHER SOURCES

The presence of MTBE in ambient air is documented and likely to be the principal source of human exposure. MTBE is released into the atmosphere during the manufacture and distribution of oxyfuel and RFG, in the vehicle refueling process, and from evaporative and tailpipe emissions from motor vehicles. The general public can be exposed to MTBE through inhalation while fueling motor vehicles or igniting fuel under cold start-up conditions (Lindstrom and Pleil 1996). The level of inhaled MTBE at the range relevant to human exposures appears to be directly proportional to the MTBE concentrations in air (Bio/dynamics, Inc. 1981, 1984c, Nihlen et al. 1994). In air, MTBE may represent five to 10% of the VOCs that are emitted from gasoline-burning vehicles, particularly in areas where MTBE is added to fuels as part of an oxygenated fuel program (ARCO 1995a). MTBE has an atmospheric lifetime of approximately four days and its primary byproducts are tert-butyl formate (TBF), formaldehyde (HCHO), acetic acid, acetone, and TBA.

MTBE was found in urban air in the U.S. (Zogorski et al. 1996, 1997) and the median concentrations ranged from 0.13 to 4.6 parts per billion by volume (ppbv). Grosjean et al. (1998) reported ambient concentrations of ethanol and MTBE at a downtown location in Porto Alegre, Brazil where about 74% of about 600,000 vehicles use gasoline with 15% MTBE, from March 20, 1996 to April 16, 1997. Ambient concentrations of MTBE ranged from 0.2 to 17.1 ppbv with an average of 6.6 ± 4.3 ppbv. This article also cited unpublished data including Cape Cod (four samples, July to August 1995): 39 to 201 parts per trillion by volume (pptv or 1/1,000 ppbv), Shenandoah National Park (14 samples, July to August 1995): ≤ 7 pptv, Brookhaven (16 samples, July to August 1995): 33 to 416 pptv, Wisconsin (62 samples, August 1994 to December 1996, with all but five samples yielding no detectable MTBE with a detection limit of 12 pptv): ≤ 177 pptv, and downtown Los Angeles, California (one sample, collected in 1993 prior to the introduction of California RFG with MTBE): 0.8 ppbv.

Ambient levels of MTBE in California are similar or slightly higher than the limited data suggest for other states. The results of two recent (from 1995 to 1996) monitoring surveys (Poore et al. 1997, Zielinska et al. 1997) indicate that ambient levels of MTBE averaged 0.6 to 7.2 ppbv with sampling for three hours at four southern California locations, and 1.3 to 4.8 ppbv with sampling for 24 hours at seven California locations. The Bay Area Air Quality Management District (BAAQMD) has an 18-station network and has been monitoring for MTBE since 1995. The average concentration of MTBE in the San Francisco Bay area is approximately one ppbv (Cal/EPA 1998). There are a few other areas in the country which have reported MTBE concentrations in ambient air. Fairbanks, Alaska reported concentrations ranging from two to six ppbv when the gasoline contained 15% MTBE (CDC 1993a).

The ARB established a 20-station TAC air-monitoring network in 1985, and began analyzing ambient air for MTBE in 1996 (ARB 1996). Preliminary data suggest a statewide average of approximately two ppbv with higher concentrations in the South Coast of about four ppbv. The limit of detection is 0.2 ppbv. The Desert Research Institute, under contract to ARB as a part of the 1997 Southern California Ozone Study (Fujita et al. 1997), monitored for MTBE in July through September of 1995 and 1996 in Southern California, at the Asuza, Burbank, and North Main monitoring sites. The monitoring was designed to determine peak morning rush hour concentrations (six to nine a.m.) and was part of a comprehensive study to analyze reactive

organics in the South Coast Air Basin. The results showed a mean of approximately four ppbv with a range of one to 11 ppbv. These concentrations are similar to the ARB findings. Although ARB sampled for 24 hours, the highest concentrations are seen in the morning rush hour traffic because MTBE is a tailpipe pollutant.

Industrial hygiene monitoring data for a MTBE operating unit shows an average eight-hour exposure of 1.42 ppm. Average exposure for dockworkers was determined to be 1.23 ppm. Occupational exposure to gasoline containing two to eight percent MTBE is estimated at one to 1.4 ppm per day (ARCO 1995a, 1995b). In a New Jersey study, MTBE concentrations as high as 2.6 ppm were reported in the breathing zone of individuals using self-service gasoline stations without vapor recovery equipment, and the average MTBE exposure among service station attendants was estimated to be below one ppm when at least 12% MTBE was used in fuels (Hartle 1993). The highest Canadian predicted airborne concentration of 75 ng/m³ is 3.9×10^7 times lower than the lowest reported effect level of 2,915 mg/m³ in a subchronic inhalation study in rats (Environmental Canada 1992, 1993, Long et al. 1994).

In a Finnish study based on inhalation exposure (Hakkola and Saarinen 1996), oil company road tanker drivers were exposed to MTBE during loading and delivery at concentrations between 13 and 91 mg/m³ (about 3.6 to 25 ppm) and the authors suggested some improvement techniques to reduce the occupational exposure. A recent Finnish study, Saarinen et al. (1998) investigated the exposure and uptake of 11 drivers to gasoline vapors during road-tanker loading and unloading. On average the drivers were exposed to vapors for 21 ± 14 minutes, three times during a work shift. The mean concentration of MTBE was 8.1 ± 8.4 mg/m³ (about 2.3 ppm).

Unlike most gasoline components which are lipophilic, the small, water-soluble MTBE molecule has low affinity for soil particles and moves quickly to reach groundwater. In estuaries, MTBE is not expected to stay in sediment soil but can accumulate at least on a seasonal basis in sediment interstitial water (ATSDR 1996). There are no reliable data on MTBE levels in food, but food is not suspected as a significant source of exposure to MTBE. There is little information on the presence of MTBE in plants or food chains. The bioconcentration potential for MTBE in fish is rated as insignificant based on the studies with Japanese carp by Fujiwara et al. (1984) generating bioconcentration factors for MTBE ranging from 0.8 to 1.5. Limited data suggest that MTBE will not bioaccumulate in fish or food chains (ATSDR 1996). Based on fugacity modeling and limited information on concentrations in shellfish, it is estimated that the average daily intake of MTBE for the age group of the Canadian population most exposed on a body weight basis, i.e., five to 11-year-olds, is 0.67 ng/kg/day (Environmental Canada 1992, 1993, Long et al. 1994).

WATER

MTBE, being a water-soluble molecule, binds poorly to soils and readily enters surface and underground water. MTBE appears to be resistant to chemical and microbial degradation in water (ATSDR 1996). When it does degrade, the primary product is TBA. The level of ingested MTBE from drinking water at the range relevant to human exposures appears to be directly proportional to the MTBE concentrations in water (Bio/dynamics, Inc. 1981, 1984c, Nihlen et al. 1994). The concentrations of MTBE in Canadian surface water predicted under a worst-case scenario is six ppt (or six ng/L), which is 1.12×10^8 times lower than the 96-hour LC₅₀ for fathead minnow of 672 ppm (or 672 mg/L) (Environmental Canada 1992, 1993).

MTBE can be a water contaminant around major production sites, pipelines, large tank batteries, transfer terminals, and active or abandoned waste disposal sites. It tends to be the most frequently detected VOC in shallow groundwater (Bruce and McMahon 1996). The primary release of MTBE into groundwater is from leaking USTs. Gasoline leaks, spills or exhaust, and recharge from stormwater runoff contribute to MTBE in groundwater. In small quantities, MTBE in air dissolves in water such as deposition in rain (Pankow et al. 1997). Recreational gasoline-powered boating and personal watercraft is thought to be the primary source of MTBE in surface water. MTBE has been detected in public drinking water systems based on limited monitoring data (Zogorski et al. 1997). Surveillance of public drinking water systems in Maine, begun in February 1997, has detected MTBE at levels ranging from one to 16 ppb in 7% of 570 tested systems with a median concentration of three ppb (Smith and Kemp 1998).

MTBE is detected in groundwater following a reformulated fuel spill (Garrett et al. 1986, Shaffer and Uchrin 1997). MTBE in water can be volatilized to air, especially at higher temperature or if the water is subjected to turbulence. However, it is less easily removed from groundwater than other VOCs such as benzene, toluene, ethylbenzene, and xylenes (BTEX) that are commonly associated with gasoline spills. MTBE and BTEX are the most water-soluble fractions in gasoline and therefore the most mobile in an aquifer system. Based on equilibrium fugacity models and especially during warm seasons, the high vapor pressure of MTBE leads to partitioning to air and half-lives in moving water are estimated around 4.1 hours (Davidson 1995, Hubbard et al. 1994). In shallow urban groundwater, MTBE was not found with BTEX. MTBE may be fairly persistent since it is refractory to most types of biodegradation (Borden et al. 1997, Daniel 1995, Jensen and Arvin 1990). Adsorption is expected to have little effect and dissolved MTBE will move at the same rate as the groundwater. MTBE may be volatilized into air or into soil gas from groundwater and these mechanisms may account for the removal of MTBE from groundwater.

MTBE has been detected in water, mainly by the USGS, in Colorado (Livo 1995, Bruce and McMahon 1996), Connecticut (Grady 1997), Georgia, Indiana (Fenelon and Moore 1996), Maine (Smith and Kemp 1998), Maryland (Daly and Lindsey 1996), Massachusetts (Grady 1997), Minnesota, Nevada, New Hampshire (Grady 1997), New Jersey (Terracciano and O'Brien 1997), New Mexico, New York (Stackelberg et al. 1997), North Carolina (Rudo 1995), Pennsylvania (Daly and Lindsey 1996), South Carolina (Baehr et al. 1997), Texas, Vermont (Grady 1997), Wisconsin and other states. USGS has published the results of the NAWQA Program (Squillace et al. 1995, 1996, 1997a, 1997b) of monitoring wells, which are not drinking water wells. This program analyzed concentrations of 60 VOCs from 198 shallow wells and 12 springs in eight urban areas (none in California) and 549 shallow wells in 21 agricultural areas (including the San Joaquin Valley). MTBE was detected in 27% of the urban wells and springs and 1.3% of the agricultural wells. The average concentration found in shallow groundwater was 0.6 ppb. MTBE was the second most frequently detected VOC (behind chloroform) in urban wells (Anonymous 1995). No MTBE was detected in 100 agricultural wells in the San Joaquin Valley. MTBE was detected in municipal stormwater in seven percent of the 592 samples from 16 U.S. cities during 1991 to 1995 with a range of 0.2 to 8.7 ppb and a median of 1.5 ppb (Delzer et al. 1997). MTBE was found to be the seventh-most frequently detected VOCs in municipal stormwater. Surveys by the U.S. EPA found that 51 public water suppliers in seven responding states had detected MTBE. There are ongoing regional studies of MTBE occurrence in California, New England, Long Island, New Jersey and Pennsylvania (Wiley 1998). MTBE was detected in aquifers (Landmeyer et al. 1997, Lindsey 1997).

DRAFT

Cal/EPA and other state agencies have taken a proactive approach toward investigating MTBE in water in California. MTBE has recently been detected in shallow groundwater at over 75% of about 300 leaking UST sites in the Santa Clara Valley Water District (SCVWD), at 90 out of 131 fuel leak sites under jurisdiction of the San Francisco Regional Water Quality Control Board and at over 200 leaking sites in the Orange County Water District. According to the Santa Ana Regional Water Quality Control Board, MTBE has been found at concentrations higher than 200 ppb at 68% of the leaking UST sites in its jurisdiction and at concentrations above 10,000 ppb at 24% of the leaking sites. In Solano County, concentrations of MTBE as high as 550,000 ppb have been reported in groundwater at sites with leaking USTs. However, these wells are not sources for drinking water (SCDEM 1997). At sites of gasoline leakage, MTBE concentrations as high as 200,000 ppb have been measured in groundwater (Davidson 1995, Garrett et al. 1986).

In 1994, Senate Bill 1764 established an independent advisory committee to the State Water Resources Control Board (SWRCB) to review the cleanup of USTs including requesting companies to monitor MTBE. State and federal statutes require that all USTs be removed, replaced or upgraded to meet current standards by December 22, 1998. In June, 1996, the SWRCB asked local regulatory agencies to require analysis at all leaking UST sites with affected groundwater. MTBE has been detected at a majority of the sites. Concentrations of MTBE in shallow groundwater near the source of the fuel release can exceed 10,000 ppb (or 10 ppm) (Cal/EPA 1998).

In 1995, ARB requested DHS' Division of Drinking Water and Environmental Management to test for MTBE in the state's drinking water. In February 1996, DHS sent an advisory letter to water suppliers it regulates, requesting voluntary testing for MTBE while a monitoring regulation was being developed. The regulation was adopted on February 13, 1997, and requires monitoring of MTBE as an unregulated chemical by the water suppliers from a drinking water well or a surface water intake at least once every three years. DHS routinely updates the reported detection of MTBE in groundwater and surface water sources on its website. DHS uses a detection limit for purposes of reporting (DLR) for MTBE of five ppb based on consideration of the State's commercial laboratories' use of MTBE in other common analyses and the potential for sample contamination and the reporting of false positives. Laboratories are only required to report MTBE analytical results at or above the five ppb DLR, but some laboratories are reporting lower concentrations.

According to the DHS report, from February 13 to June 13, 1997, MTBE had been detected in 14 of the 388 drinking water systems that had been monitored. As of December 22, 1997, 18 of the 516 systems monitored had reported MTBE detection. These are drinking water wells tapping deep aquifers and some aquifers at depths of 200 feet or greater. In addition, approximately 2,500 public drinking water sources had been sampled and reported. Only 33 sources including 19 groundwater sources and 14 surface water sources, nine of which are reservoirs, had reported detectable concentrations of MTBE. Three groundwater sources including City of Santa Monica (up to 300 ppb in February 1996), City of Marysville (up to 115 ppb in January 1997), and Presidio of San Francisco (up to 500 ppb in July 1990 from an abandoned well) had reported concentrations above the U.S. EPA 1997 advisory level of 20 to 40 ppb. Otherwise, the range of reported values was < 1 to 34.1 ppb in groundwater sources and < 1 to 15 ppb in surface water sources (DHS 1997).

The City of Santa Monica has shut down two well fields, Charnock and Arcadia, due to MTBE contamination. These well fields used to supply 80% of the drinking water to the city residents. Concentrations as high as 610 ppb were observed in the Charnock aquifer and the seven wells in the field have been closed. In the Arcadia well field, two wells have been closed due to MTBE

contamination from an UST at a nearby gasoline station (Cal/EPA 1998, Cooney 1997). DHS (1997) reported MTBE concentrations up to 130 ppb in a Charnock well and 300 ppb in another Charnock well in February 1996, and up to 72.4 ppb in an Arcadia well in August 1996. In Santa Clara county, the Great Oaks Water Company has closed a drinking water well in South San Jose due to trace MTBE contamination.

MTBE has also been found in many surface water lakes and reservoirs (DHS 1997). The reservoirs allowing gasoline powerboat activities have been detected with MTBE at higher concentrations than those reservoirs prohibiting boating activities. DHS reported MTBE in Lake Tahoe, Lake Shasta, Whiskeytown Lake in the City of Redding, San Pablo Reservoir in East Bay Municipal Utility District (EBMUD) in the San Francisco Bay area, Lobos Creek in Presidio of San Francisco, Del Valle and Patterson Pass of Zone Seven Water Agency serving east Alameda County, Clear Lake in Konocti County Water District, Canyon Lake in the Elsinore Valley Municipal Water District, Lake Perris in the MWDSC in the Los Angeles area, and Alvarado, Miramar, and Otay Plant influent in City of San Diego. MTBE concentrations ranged from < 1 to 15 ppb. Donner Lake, Lake Merced, Cherry and New Don Pedro Reservoirs in EBMUD, Anderson and Coyote Reservoirs in the SCVWD, Modesto Reservoir in the Stanislaus Water District, and Castaic Reservoir in MWDSC also had detectable levels of MTBE.

The City of Shasta Lake domestic water supply intake raw water was reported with 0.57 ppb MTBE in September 1996 although Lake Shasta had 88 ppb in a surface water sample next to a houseboat at a marina dock. BTEX were found in lower concentrations than MTBE. Water was analyzed for hydrocarbons before and after organized jet ski events held in the summer and fall of 1996 in Orange County and Lake Havasu (Dale et al. 1997a). MTBE was measured in the water at the small holding basin in Orange County at concentrations of up to 40 ppb a few days after the event while there was only negligible BTEX. At the larger Lake Havasu, the MTBE concentrations increased from below the level of detection to 13 ppb. A recent report to the SCVWD described the detection of an average concentration of three ppb MTBE in Anderson, Calero, and Coyote Reservoirs which are drinking water sources where powerboating is allowed.

The Carson publicly owned treatment works (POTW) in Carson, California has also reported MTBE in its wastewater. The Carson POTW processes the largest volume of refinery wastewater in the nation (13 refineries sporadically discharge wastewater to the POTW). Refineries in California perform their own pretreatment prior to discharging to sewers. The refineries' discharges contain average levels from one to seven ppm with concentrations occasionally as high as 40 ppm. California refineries are situated mainly along the coast and discharge directly or indirectly to marine waters. No California refineries discharge their wastewater to sources of drinking water.

Metabolism and Pharmacokinetics

MTBE can be absorbed into the body after inhalation, ingestion or skin contact. It is metabolized and eliminated from the body within hours. MTBE caused lipid peroxidation in the liver and induction of hepatic microsomal cytochrome P₄₅₀ content in mice (Katoh et al. 1993). The major metabolic pathway of MTBE in both animals and humans is oxidative demethylation leading to the production of TBA (Cederbaum and Cohen 1980, Li et al. 1991, Poet et al. 1997c). In animals, HCHO is also a major metabolite (Hutcheon et al. 1996). This reaction is catalyzed by cytochrome P₄₅₀ enzymes (Brady et al. 1990, Hong et al. 1997b).

MTBE and TBA have been detected in blood, urine, and breath of humans exposed to MTBE via inhalation for 12 hours. Nihlen et al. (1998b) in a human chamber study suggests that TBA in blood or urine is a more appropriate biological exposure marker for MTBE than the parent ether itself. Bonin et al. (1995) and Lee and Weisel (1998) described analytical methods for detecting MTBE and TBA in human blood and urine at concentrations below one ppb. A recent Finnish study, Saarinen et al. (1998) investigated the uptake of 11 drivers to gasoline vapors during road-tanker loading and unloading. The total MTBE uptake during the shift was calculated to be an average of 106 ± 65 μ mole. The mean concentrations of MTBE and TBA detected in the first urine after the work shift were 113 ± 76 and 461 ± 337 nanomole/L, and those found 16 hours later in the next morning were 18 ± 12 and 322 ± 213 nanomole/L, respectively.

ABSORPTION

In its liquid or gaseous state, MTBE is expected to be absorbed into the blood stream (Nihlen et al. 1995). MTBE is absorbed into the circulation of rats following oral, intraperitoneal (i.p.), intravenous (i.v.), or inhalation exposures (Bioresearch Laboratories 1990a, 1990b, 1990c, 1990d, Miller et al. 1997, NSTC 1997). Dermal absorption of MTBE is limited, as compared with other routes. MTBE is lipophilic which will facilitate its absorption across the lipid matrix of cell membranes (Nihlen et al. 1997). The concentration-time course of MTBE in blood plasma of male rats administered 40 mg/kg/day by oral, dermal, or i.v. routes was followed (Miller et al. 1997). Peak blood concentrations of MTBE (C_{max}) were obtained within five to 10 minutes. Higher levels of MTBE were seen after oral versus i.v. exposure indicating elimination of the latter via the lungs. Comparison of the area under the concentration-time curve (AUC) for MTBE following i.v. and oral administrations indicated that MTBE was absorbed from the gastrointestinal tract. Plasma levels of MTBE following dermal exposure were limited; peak concentrations were achieved two to four hours after dosing. Absorption ranged from 16 to 34% of applied doses of 40 mg/kg/day and 400 mg/kg/day respectively. After inhalation exposure, plasma concentrations of MTBE reached apparent steady state within two hours at both low (400 ppm) and high (8,000 ppm) doses. Peak MTBE concentrations were reached at four to six hours and were 14 and 493 ppb, respectively.

DISTRIBUTION

Once absorbed, MTBE is either exhaled as the parent compound or metabolized. Oxidative demethylation by cytochrome P_{450} -dependent enzymes is the first step in the metabolism that yields HCHO and TBA. TBA is detected in blood and urine and appears to have a longer half-life in blood than MTBE (Poet et al. 1996, Prah et al. 1994, Prescott-Mathews et al. 1996, Savolainen et al. 1985). Once in the blood, MTBE is distributed to all major tissues in the rat. Due to its hydrophilic properties, neither MTBE nor its metabolites would be expected to accumulate in body tissues. TBA appears to remain longer, and chronic exposure could result in accumulation to some steady-state level, but this needs further study.

METABOLISM

The metabolism of absorbed MTBE proceeds regardless of route of exposure. MTBE undergoes oxidative demethylation in the liver via the cytochrome P_{450} -dependent enzymes (P_{450} IIE1,

P₄₅₀IIB1, and P₄₅₀ IIA6 are thought to be involved) to give TBA and HCHO (Brady et al. 1990, Hong et al. 1997b). Rat olfactory mucosa displays a high activity in metabolizing MTBE via the cytochrome P₄₅₀-dependent enzymes (Hong et al. 1997a). In vitro studies of MTBE in human (Poet and Borghoff 1998) and rat (Poet and Borghoff 1997b) liver microsomes confirm that MTBE is metabolized by P₄₅₀-dependent enzymes and suggest that the metabolism of MTBE will be highly variable in humans. TBA may be eliminated unchanged in expired air or may undergo secondary metabolism forming 2-methyl-1,2-propanediol and α -hydroxyisobutyric acid. Both of these latter metabolites are excreted in the urine and account for about 14% and 70% respectively of urine radioactivity for ¹⁴C-MTBE dosed rats (Miller et al. 1997). Two unidentified minor metabolites are also excreted in urine. In vitro evidence suggests that TBA may also undergo oxidative demethylation to produce HCHO and acetone (Cederbaum and Cohen 1980). Identification of ¹⁴CO₂ in expired air of ¹⁴C-MTBE treated rats suggests some complete oxidation of MTBE or metabolites occurs, probably via HCHO. Studies in humans are more limited but TBA has been observed as a blood metabolite of MTBE. The participation of hepatic cytochrome P₄₅₀-dependent enzymes also indicates a potential role of co-exposure to other environmental chemicals in affecting MTBE metabolism and toxicity (Hong et al. 1997b, NSTC 1997).

EXCRETION

Elimination of MTBE and its metabolites by Fischer 344 rats is primarily via the lungs (expired air) and the kidneys (urine). In expired air, MTBE and TBA are the predominant forms. After i.v. administration of ¹⁴C-MTBE to male rats most of the radioactivity was excreted in the exhaled air (60%) and urine (34.9%) with only two percent in the feces and 0.4% remaining in the tissues/carcass. Most of the administered dose was eliminated as MTBE during the first three hours following administration. About 70% of the dose recovered in the urine was eliminated in the first 24 hours and 90% in 48 hours. After dermal exposure to MTBE for six hours, 70 to 77% of the applied radioactivity was unabsorbed while 7.6 to 18.9% was excreted in expired air, 6.3 to 16.2% in urine, and 0.25 to 0.39% in feces at 40 and 400 mg/kg/day respectively. A negligible amount (< 0.2%) was found in tissues/carcass. The composition of ¹⁴C-radiolabel in expired air was 96.7% MTBE and 3.3% TBA at the high dose. After inhalation exposures most of the ¹⁴C was eliminated in the urine with 64.7% after single and 71.6% after repeated low doses. At the high dose, a larger fraction was eliminated in exhaled air: 53.6% compared to 17% for single or 21% for repeated low doses. Less than 1% of the dose was recovered in the feces and < 3.5% in the tissues/carcass. The composition of ¹⁴C-radiolabel in exhaled breath in the first six hours following administration of MTBE was 66 to 69% MTBE and 21 to 34% TBA. By 24 hours post-dose 85 to 88% of the urine radioactivity was eliminated in rats from all exposure groups (Miller et al. 1997). Pulmonary elimination of MTBE after intraperitoneal injection in mice (Yoshikawa et al. 1994) at three treated doses (50, 100 and 500 mg/kg) indicated an initial rapid decrease of the elimination ratio followed by a slow decrease at the doses of 100 and 500 mg/kg. The calculated half-lives of the two elimination curves obtained by the least squares method were approximately 45 minutes and 80 minutes. The pulmonary elimination ratios at the three different doses were from 23.2% to 69%. Most of the excreted MTBE was eliminated within three hours.

In a human chamber study (Buckley et al. 1997), two subjects were exposed to 1.39 ppm MTBE, which is comparable to low levels which might be found in the environment for one hour, followed by clean air for seven hours. The results showed that urine accounted for less than one

percent of the total MTBE elimination. The concentrations of MTBE and TBA in urine were similar to that of the blood ranging from 0.37 to 15 µg/L and two to 15 µg/L, respectively. Human breath samples of end-expiration volume were collected from two individuals during motor vehicle refueling, one person pumping the fuel and a nearby observer, immediately before and for 64 minutes after the vehicle was refueled with premium grade gasoline (Lindstrom and Pleil 1996). Low levels of MTBE were detected in both subjects' breaths before refueling and levels were increased by a factor of 35 to 100 after the exposure. Breath elimination indicated that the half-life of MTBE in the first physiological compartment was between 1.3 and 2.9 minutes. The breath elimination of MTBE during the 64-minute monitoring period was about four-fold for the refueling subject comparing to the observer subject.

Johanson et al. (1995) and Nihlen et al. (1998a, 1998b) reported toxicokinetics and acute effects of inhalation exposure of 10 male subjects to MTBE vapor at five, 25, and 50 ppm for two hours during light physical exercise. MTBE and TBA were monitored in exhaled air, blood, and urine. The elimination of MTBE from blood was multi-phasic with no significant differences between exposure levels. The elimination phases had half-lives of one minute, 10 minutes, 1.5 hours, and 19 hours respectively. Elimination of MTBE in urine occurred in two phases with average half-lives of 20 minutes and three hours. Excretion of MTBE appeared to be nearly complete within 10 hours. For TBA excretion the average post-exposure half-lives in blood and urine were 10 and 8.2 hours respectively. Some exposure dependence was noted for the urinary half-life with shorter values seen at the highest exposure level (50 ppm x 2 hour). A low renal clearance for TBA (0.6 to 0.7 mL/hour/kg) may indicate extensive blood protein binding or renal tubular reabsorption of TBA.

PHARMACOKINETICS

The plasma elimination half-life ($t_{1/2}$) of MTBE in male rats was about 0.45 to 0.57 hour after i.v., oral (low dose), and inhalation exposures. A significantly longer $t_{1/2}$ of 0.79 hour was observed with the high oral dose of 400 mg/kg/day. For dermal exposure the initial MTBE elimination $t_{1/2}$ was 1.8 to 2.3 hours. TBA elimination $t_{1/2}$ values were 0.92 hour for i.v., 0.95 to 1.6 hours for oral, 1.9 to 2.1 hours for dermal, and 1.8 to 3.4 hours for inhalation exposures. The apparent volume of distribution for MTBE ranged from 0.25 to 0.41 L after i.v., oral, and inhalation dosing and from 1.4 to 3.9 liters (L) after dermal exposures. The total plasma clearance of MTBE, corrected for relative bioavailability, ranged from 358 to 413 mL/hour in i.v., oral, and dermal administrations. Inhalation values ranged from 531 mL/hour for low single dose to 298 mL/hour for high single dose. For oral administration of 40 or 400 mg/kg/day MTBE the AUC values were 17 and 230 (µg/mL)hour for MTBE and 39 and 304 (µg/mL)hour for TBA (Miller et al. 1997).

The disposition and pharmacokinetics observed in these studies are similar to those observed in human volunteers following inhalation and dermal exposures (U.S. EPA 1993). For inhalation exposure to five mg/m³ for one hour the $t_{1/2}$ value for MTBE was 36 minutes. Blood TBA levels rose during exposure and remained steady for up to seven hours post-exposure suggesting a longer $t_{1/2}$ for TBA in humans compared to rats. Other more recent data (cited in NSTC 1997) indicate a multiexponential character to MTBE elimination from human blood with $t_{1/2}$ values of two to five minutes, 15 to 60 minutes and greater than 190 minutes. These results possibly indicate a more complex distribution or binding of MTBE in humans than observed in rats. Such differences probably are related to larger fat compartments in humans compared to rats.

Overall, these studies show that following i.v., oral, or inhalation exposures MTBE is absorbed, distributed, and eliminated from the body with a half-life of about 0.5 hour. Dermal absorption is limited. The extent of metabolism to TBA (and HCHO) the major metabolite is somewhat dependent on route and dose. TBA is eliminated from the body with a half-life of one to three hours or longer in humans. Virtually all MTBE is cleared from the body 48 hours post-exposure.

PHYSIOLOGICALLY-BASED PHARMACOKINETIC (PBPK) MODELS

Computer-based PBPK models have been developed for rats (Borghoff et al. 1996a, Rao and Ginsberg 1997). These models vary in complexity, metabolic parameters, and one chemical specific parameter. The Borghoff et al. (1996a) model uses five compartments for MTBE and either five or two for TBA. While model predictions of MTBE blood concentrations and clearance following inhalation or oral exposures were generally good, the model underpredicted MTBE blood levels at 8,000 ppm by a factor of two. Accurate model predictions of TBA blood levels and clearance were more elusive with the two compartment model giving more accurate predictions at lower oral and inhalation doses than at higher doses or than the five compartment model. The Rao and Ginsberg (1997) model is more complex using eight compartments for MTBE and eight for TBA. While both models assume two Michaelis-Menten processes (V_{maxc}/K_m) from MTBE to TBA namely high capacity to low affinity (V_{maxc_2}/K_{m_2}), and low capacity to high affinity (V_{maxc_1}/K_{m_1}), the Rao and Ginsberg (1997) model uses different parameters than Borghoff et al. (1996a) with a lower V_{maxc_1}/K_{m_1} . Rao and Ginsberg (1997) use a lower tissue/blood partition coefficient for TBA in the slowly perfused compartment (e.g., muscle) of 0.4 versus 1. Predictions of blood levels and clearance rates for MTBE and TBA with MTBE inhalation exposures appear to be more accurate with this model. Similar validation is claimed for the oral and i.v. routes for MTBE exposure and for i.p. exposure to TBA although these data have not been seen in detail. Rao and Ginsberg (1997) used their model to evaluate some key uncertainties of acute inhalation exposures to MTBE during bathing and showering and concluded that the acute central nervous system (CNS) toxicity is likely due to MTBE rather than to its TBA metabolite. The simulated brain TBA concentration for CNS effects was in the 500 to 600 mg/L range. In contrast, the simulated brain concentration for MTBE's CNS effects was considerably lower (89 to 146 mg/L). By comparing TBA only versus MTBE exposure studies the authors concluded that under conditions where MTBE dosing produced acute CNS toxicity, the simulated TBA brain concentrations were too low to be effective.

Despite the lack of human data on tissue/blood partition coefficients and other key parameters, both models have been adjusted to human anatomical and physiological values and estimated metabolic and chemical parameters and compared with limited human blood data. Although the Borghoff et al. (1996a) model was able to predict MTBE levels seen in Cain et al. (1996) during inhalation exposure, it underpredicted MTBE blood concentrations after exposure, resulting in a faster clearance than seen experimentally. The Rao and Ginsberg (1997) model more closely simulated the data (1.7 ppm MTBE for one hour) of Cain et al. (1996) but underpredicted the peak and postexposure concentrations at higher inhalation exposures of five and 50 ppm MTBE for two hours (Johanson et al. 1995). It is clear that while human MTBE PBPK models may be improved considerably, they may prove useful in their present state to assess risks associated with some environmental exposures to MTBE (e.g., exposures when taking a shower).

Toxicology

The toxicology profile of MTBE has been summarized in the U.S. (Von Burg 1992, ATSDR 1996) and in Great Britain (BIBRA 1990). Zhang et al. (1997) used computer modeling to predict metabolism and toxicological profile of gasoline oxygenates including MTBE based on structure activity relationships. Health risk assessment of MTBE has been performed (Gilbert and Calabrese 1992, Hartly and Englande 1992, Hiremath and Parker 1994, Stern and Tardiff 1997, Tardiff and Stern 1997) and the general toxicity of MTBE is not considered as "highly hazardous" in a hazard ranking system for organic contaminants in refinery effluents (Siljeholm 1997). A substantial amount of health-related research has been conducted or initiated on MTBE in recent years (ATSDR 1996, U.S. EPA 1997a). A recent literature review (Borak et al. 1998) summarizes the exposure to MTBE and acute human health effects including nine epidemiological studies, ten industrial hygiene studies, and 12 clinical studies. However, most of the studies and reviews focus on the inhalation route of exposure in human health effects and laboratory animal toxicities. No studies were located regarding toxic effects in humans after oral exposure to MTBE alone. Because this document is mainly concerned with the effects of MTBE in drinking water, it focuses on oral toxicity studies in animals. There is limited information on dermal exposure effects in humans and animals. Very little is known about the toxic effects of MTBE in plants and ecosystems.

TOXICOLOGICAL EFFECTS IN ANIMALS

Table 4 summarizes the lowest concentrations resulting in toxicity in laboratory animals via inhalation or oral exposure as reported in the ATSDR (1996) document and the latest U.S. EPA (1997c) advisory. Clary (1997) reviewed the systemic toxicity of MTBE including 12 inhalation and four oral studies. Stelljes (1997) summarized similar information based on only the ATSDR (1996) document. The various noncancer health effects via oral route of exposure in all tested species and the duration of exposure are summarized in Table 5. The highest no observed adverse effect levels (NOAELs) and all the lowest observed adverse effect level (LOAELs) are also included in Table 5. Details of each of the studies listed in Table 5 are described in the following sections on acute, subacute, subchronic and chronic toxicity. The cancer effects observed in animals are discussed in a separate section on carcinogenicity in this chapter. There were no studies located regarding cancer in humans after oral, or any other exposure to MTBE.

In animal studies, oral exposure to MTBE for acute, subacute, subchronic, or chronic duration appears to be without effects on the cardiovascular, musculoskeletal, dermal, ocular, or reproductive systems. In acute and subacute oral exposure studies, limited effects on the respiratory, gastrointestinal, hematological, hepatic, renal, or neurological systems and some minor systemic toxicities have been observed. In subchronic oral exposure, limited effects on gastrointestinal, hematological, hepatic, or renal systems and some minor systemic toxicities have been observed. In chronic oral exposure, the main observation is cancer and preneoplastic effects (ATSDR 1996). In this document, all the potential toxic effects of MTBE have been reviewed with an emphasis on the oral exposure, particularly the potential reproductive, developmental and carcinogenic effects have been extensively reviewed by OEHHA staff.

DRAFT

Some acute, intermediate or chronic duration minimal risk levels (MRLs) have been derived by the ATSDR for inhalation or oral exposure to MTBE (ATSDR 1996). U.S. EPA (1997c) lists in IRIS a Reference Concentration (RfC) for inhalation that is similar to the ATSDR's inhalation MRL. However, the IRIS (U.S. EPA 1997c) does not list a Reference Dose (RfD) for ingestion (U.S. EPA 1987b) that is similar to the ATSDR's ingestion MRL. In addition to the key documents from governmental agencies and literature search articles mentioned above, toxicology information in the TOMES PLUS® database (Hall and Rumack 1998) also has been used in the following summary of toxic effects of MTBE.

**Table 4. Summary of Selected Data on MTBE:
Noncancer Toxic Effects in Animals***

Dose level	Inhalation (mg/m ³)			Oral (mg/kg/day)	
	ACUTE	SUBACUTE/ SUBCHRONIC	CHRONIC	ACUTE	SUBACUTE/ SUBCHRONIC
NOAEL	1,440	1,440	1,440	40	100
LOAEL	3,600	2,880	10,800	90	300
Lethal Dose	649,000	NA	NA	3,866	NA

*Values represent the lowest reported in ATSDR (1996) and U.S. EPA (1997a)

Table 5. Significant Noncancer Health Effects and Levels of Oral Exposure to MTBE in Animals*

Species/ (Strain)	Exposure/ Duration/ Frequency (Specific route)	System	NOAEL (mg/kg/day)	LOAEL (mg/kg/day)	Reference
ACUTE EXPOSURE					
Death					
Rat	once (gavage)			3,866 (LD ₅₀)	ARCO 1980
Mouse	once (gavage)			4,000 (LD ₅₀)	Little et al. 1979
Systemic Toxicity					
Rat	once (gavage)	Respiratory Neurological		4,080 (labored respiration) 1,900 (slight to marked CNS depression) 2,450 (ataxia)	ARCO 1980
Rat (Sprague- Dawley)	once (gavage in oil)	Gastrointestinal Neurological	900	100 (diarrhea) 1,200 (profound but transient anesthesia)	Robinson et al. 1990
Rat (Fischer 344)	once (gavage in water)	Neurological	40	400(drowsiness)	Bioresearch Labs. 1990b
Rat (Sprague- Dawley)	once (gavage)	Neurological		90 (salivation) 440 (Male) (hypoactivity, ataxia) 1,750 (Female)	Johnson and Boyne 1992, Klan et al. 1992

DRAFT

Species/ (Strain)	Exposure/ Duration/ Frequency (Specific route)	System	NOAEL (mg/kg/day)	LOAEL (mg/kg/day)	Reference
SUBACUTE EXPOSURE					
Systemic Toxicity					
Rat (Sprague- Dawley)	14 days 7 days/week once/day (gavage in oil)	Respiratory	1,428		Robinson et al. 1990
		Cardiovascular	1,428		
		Gastrointestinal		357 (diarrhea)	
		Hematological	1,428	357 (Male)	
			(Female)	(decreased monocytes)	
		Hepatic	714 (Male)	1,071 (Male)	
				[increased serum glutamic-oxaloacetic transaminase (SGOT) and lactic dehydrogenase] 1,428 (Female)	
				[decreased blood urea nitrogen (BUN) values]	
		Renal	1,071 (Male)	1,428 (Male)	
			1,428 (Female)	(increased hyaline droplets)	
		Endocrine Body weight	1,428		
			714 (Female)	1,071 (Female)	
				(unspecified reduced weight gain)	
Immunological/ Lymphoreticular	357 (Male)				
	1,428				
Neurological	1,071	1,428 (profound but transient anesthesia, hypoactivity, ataxia)			
Reproductive Other	1,428				
	1,071 (Male)	1,428 (Male)			
	357 (Female)	714 (Female)			
		(elevated cholesterol)			
Mouse (CD-1)	3 weeks, 5 days/week (gavage in oil)	Body weight	1,000		Ward et al. 1994, 1995
		Reproductive	1,000		

DRAFT

Species/ (Strain)	Exposure/ Duration/ Frequency (Specific route)	System	NOAEL (mg/kg/day)	LOAEL (mg/kg/day)	Reference
SUBCHRONIC EXPOSURE					
Death					
Rat (Sprague- Dawley)	16 weeks 4 days/week once/day (gavage in oil)			250 (Female) (increased mortality)	Belpoggi et al. 1995
Systemic Toxicity					
Rat (Sprague- Dawley)	4 weeks 5 days/week once/day (gavage)	Respiratory	1,750		Johnson and Boyne 1992, Klan et al. 1992
		Cardiovascular	1,750		
		Gastrointestinal	440	1,750 (inflammation, submucosal edema, epithelial hyperplasia, stomach ulcers)	
		Hematological	1,750		
		Muscle/skeleton	1,750		
		Hepatic	440	1,750 (increased relative liver weights)	
		Renal	1,750 (Female)	440 (Male) (increased hyaline droplets in proximal convoluted tubules and increased relative kidney weights)	
		Endocrine	1,750		
		Dermal	1,750		
		Ocular	1,750		
		Body weight	1,750		
		Immunological/ Lymphoreticular	1,750		
		Neurological		440 (hypoactivity, ataxia)	
		Reproductive	1,750		
		Other	440	1,750 (increased serum cholesterol)	
SUBCHRONIC EXPOSURE (Continued)					
Systemic Toxicity					
Species/ (Strain)	Exposure/ Duration/ Frequency (Specific route)	System	NOAEL (mg/kg/day)	LOAEL (mg/kg/day)	Reference

DRAFT

Rat (Sprague-Dawley)	90 days 7 days/week once/day (gavage in oil)	Respiratory	1,200		Robinson et al. 1990
		Cardiovascular	1,200		
		Gastrointestinal		all treated doses (diarrhea)	
		Hematological	900	1,200 (increased monocytes, decreased mean corpuscular volume in males, increased red blood cell, hemoglobin, hematocrit and decreased white blood cells in females)	
		Hepatic		all treated doses (decreased BUN values)	
		Renal	900 (Male) 1,200 (Female) 100	1,200 (Male) (hyaline droplets, granular casts) 300 (alterations in kidney weights)	
		Endocrine	1,200		
		Body weight	1,200		
		Immunological/ Lymphoreticular	1,200		
		Reproductive	1,200		
		Other	300 (Male)	900 (Male) 100 (Female) (elevated cholesterol)	

Species/ (Strain)	Exposure/ Duration/ Frequency (Specific route)	System	NOAEL (mg/kg/day)	LOAEL (mg/kg/day)	Reference
CHRONIC EXPOSURE					
Systemic Toxicity					
Rat (Sprague- Dawley)	104 weeks 4 days/week once/day (gavage in oil)	Respiratory	1,000		Belpoggi et al. 1995
		Cardiovascular	1,000		
		Gastrointestinal	1,000		
		Muscle/skeleton	1,000		
		Hepatic	1,000		
		Renal	1,000		
		Endocrine	1,000		
		Dermal	1,000		
		Body weight	1,000		
		Immunological/ Lymphoreticular	1,000 (Male)	250 (Female) (dysplastic proliferation of lymphoreticular tissues, possibly preneoplastic)	
	Reproductive	1,000			

*adapted from ATSDR (1996) and U.S. EPA (1997c)

Acute Toxicity

Studies of the systemic effects of MTBE have been conducted in animals, but the majority involve inhalation exposure (Clary 1997). Inhalation or contact with MTBE may irritate or burn skin and eyes. Vapors may cause dizziness or suffocation. Acute toxicity studies in animals demonstrate the extremely low toxicity of MTBE (ARCO 1980, Little et al. 1979, Reese and Kimbrough 1993). The oral LD₅₀s (lethal doses with 50% kill) are approximately 3,866 mg/kg or four mL/kg in rats, and 4,000 mg/kg or 5.96 mL/kg in mice. The inhalation four-hour LC₅₀s (lethal concentrations with 50% kill) in rats have been calculated to be approximately 39,395 ppm for 96.2% MTBE, 33,370 ppm for 99.1% MTBE and 23,576 ppm for MTBE. The inhalation 10-minute LC₅₀ in mice is 180,000 ppm and the inhalation 15-minute LC₅₀ in mice is 141 g/m³. The inhalation LT₅₀ (time at which death occurs in 50% of the exposed animals) in mice exposed to 209,300 ppm MTBE is 5.6 minutes (ATSDR 1996). The dermal LD₅₀ is estimated to be greater than 10 mL/kg in New Zealand rabbits (HSDB 1997). The i.p. LD₅₀ is 1.7 mL/kg in mice and greater than 148 mg/kg in rats (Arashidani et al. 1993, RTECS 1997). Zakko et al. (1997) reported cytotoxicity of MTBE to intestinal mucosa of rats via i.p. injection similar to the effects of MTBE treatment for gallstone dissolution in humans.

At lethal doses, ocular and mucous membrane irritation, ataxia, labored breathing, CNS depression, and general anesthetic effects precede death. An inhalation study also demonstrated inflammation in the nasal mucosa of rats at a dose of 3,000 ppm for six hours per day for nine days (HSDB 1997). Mice that inhaled up to approximately 8,400 ppm MTBE for one hour had approximately a 52% decrease in breathing frequency (Tepper et al. 1994). The decrease occurred immediately, reached a maximum by 10 minutes and returned to baseline 15 minutes after exposure. High oral doses of greater than 4,080 mg/kg of MTBE caused labored respiration in rats (ARCO 1980). A four-hour direct exposure to MTBE vapor at concentrations greater than 18,829 ppm in an inhalation study resulted in ocular discharges in rats (ARCO 1980). A six-hour inhalation study produced signs of reversible CNS depression following exposure to 8,000 ppm and, to a lesser extent, to 4,000 ppm vapor with a NOAEL of 800 ppm (Dodd and Kintigh 1989, Daughtrey et al. 1997). As indicated in Tables 4 and 5, a NOAEL of 40 mg/kg/day and a LOAEL of 90 mg/kg/day are established by these acute oral exposure experiments based on the neurological effects (Bioresearch Laboratories 1990b, Johnson and Boyne 1992, Klan et al. 1992).

Subacute Toxicity

In a consecutive 14-day study, Sprague-Dawley rats (10/sex/dose) were administered MTBE in corn oil by gavage at 0, 357, 714, 1,071 or 1,428 mg/kg/day. MTBE appears to be irritating to the gastrointestinal tract of rats as evidenced by diarrhea and histological lesions at all levels of MTBE by the third day of dosing throughout the 14-day study. Decreased lung weight was observed in female rats at all MTBE doses and at 714 mg/kg/day in male rats. Decreased levels of monocytes in blood were observed in male rats at all MTBE doses. Increased liver enzymes in males at 1,071 mg/kg/day and decreased blood urea nitrogen (BUN) values in females at 1,428 mg/kg/day were observed. At the highest dose, anesthesia was immediate, but recovery was complete within two hours. Ataxia and hyperactivity, an increase in the weight of kidneys, adrenal glands, and livers in both genders at 1,428 mg/kg/day, and an increase in hyaline droplet formation in kidneys of male rats at 1,428 mg/kg/day were observed. Increases in relative kidney weights were noted in the males at 1,071 and at 1,428 mg/kg/day and in females at the 1,428 mg/kg/day dose. Although there was a dose-related decrease in body weight gain, it was significant only in females at the highest treatment regimen. At 1,428 mg/kg/day in males and at 714 mg/kg/day in females, elevated cholesterol was observed. There were no gross lesions seen at any treatment level. Based on the increases in relative kidney weight, a NOAEL of 714 mg/kg/day and a LOAEL of 1,071 mg/kg/day are established by these experiments (Robinson et al. 1990). These studies indicate that the male kidney is the primary target of short-term toxicity at relatively high doses. Subchronic toxicity studies of TBA indicated that, in rodents, the urinary tract is a target system and males are more sensitive to TBA toxicity than females (NTP 1995).

Subchronic Toxicity

In a 104-week gavage cancer study, increased mortality was observed in female Sprague-Dawley rats at 250 mg/kg/day beginning at 16 weeks from the start of the study (Belpoggi et al. 1995). Daily oral administration in rats for four weeks resulted in increased hyaline droplets and kidney weight in males at 440 mg/kg/day and higher doses, and stomach ulcers, increased liver weights and serum cholesterol at 1,750 mg/kg/day (Johnson and Boyne 1992, Klan et al. 1992).

Sprague-Dawley rats (10/sex/dose) were treated orally with MTBE in corn oil for 90 days at 0, 100, 300, 900, or 1,200 mg/kg/day. Anesthesia was evident at the highest dose, but as in the 14-day study, full recovery occurred in two hours. There was a significant decrease in final body weight of females only at the highest level of treatment. The diarrhea seen in the treated animals was considered to be the consequence of the bolus dosing regime. In female rats, there were significantly increased heart weights at 900 mg/kg/day and increases in relative kidney weights at 300, 900, and 1,200 mg/kg/day. In male rats, increases were noted only at the two highest treatment levels. BUN levels were significantly reduced in both males and females at all MTBE doses. Reductions in serum calcium and creatinine were observed in males and a reduction in cholesterol in females was reported, but there were no clear dose-dependent results. Based on the alterations in kidney weights, a NOAEL and LOAEL of 100 and 300 mg/kg/day, respectively, are identified from this study (Robinson et al. 1990).

The subchronic data from the study by Robinson et al. (1990) were used by U.S. EPA (1996a) to develop a RfD and a Drinking Water Equivalent Level (DWEL) for kidney effects from MTBE. The increase in kidney weights at doses of 300 mg/kg/day and higher was considered to be an adverse effect, since increases in organ weights are a marker for adverse organ effects (Weil 1970). The diarrhea observed was considered to be a gastrointestinal complication of the gavage dosing. Based on the NOAEL of 100 mg/kg/day, a DWEL for kidney effects of 3,500 ppb can be derived for a 70 kg male adult with two liters (L) of daily water consumption (DWC), using an uncertainty factor of 1,000. The uncertainty factor reflects a 10 for the less-than-lifetime duration of the study, a 10 for interspecies variability, and a 10 for intraspecies variability. Using an additional uncertainty factor of 10 for potential carcinogenicity and a 20% default Relative Source Contribution (RSC), U.S.EPA (1996a) drafted a lifetime Health Advisory (HA) of 70 ppb or 70 µg/L. Details of the equation and calculation of the HA are described later in the chapter on the calculation of the PHG.

Genetic Toxicity

The results of genetic toxicity studies for MTBE were generally negative, but contained one well performed positive study which included information on mechanisms of action. As detailed later in this section, MTBE was not mutagenic in bacteria and tissue culture gene mutation assays, a sister chromatid exchange assay, a *Drosophila* sex-linked recessive lethal test, in vitro and in vivo chromosomal aberration assays, in vivo and in vitro unscheduled DNA synthesis assays, an in vivo DNA repair assay, an in vivo cytotoxicity assay, and in vitro and in vivo micronucleus assays. The only positive genotoxicity test was for forward mutations in the mouse lymphoma assay with exogenous activation (ARCO 1980, Mackerer et al. 1996). ATSDR (1996) indicated that MTBE has little or no genotoxic activity. However, the positive result in the Mackerer et al. (1996) study indicates that the genetic toxicity of MTBE needs to be investigated further.

MTBE was negative in the Ames in vitro assay for reverse mutation in *Salmonella typhimurium* strains TA1535, TA1537, TA1538, TA98, and TA100 in the absence or presence of metabolic activation (ARCO 1980, Cinelli et al. 1992, Life Science Research Roma Toxicology Centre S.P.A. 1989a). Since MTBE is volatile, a closed system was used in a recent microsuspension assay (Kado et al. 1998), and negative results were observed even though some elevated revertant values were seen with TA100 and TA104. MTBE produced no evidence of a dose-related increase for sister chromatid exchange (ARCO 1980), for gene mutation in Chinese hamster V79 cells (Life Science Research Roma Toxicology Centre S.P.A. 1989b) and for in vitro unscheduled DNA synthesis in primary rat hepatocytes (Life Science Research Roma

Toxicology Centre S.P.A. 1989c, Vergnes and Chun 1994). It was negative for micronuclei formation in erythrocytes (Vergnes and Kintigh 1993).

MTBE was assessed for its *in vivo* mutagenic potential (McKee et al. 1997). It was negative in the sex-linked recessive lethal assay in *Drosophila melanogaster* (Sernau 1989). It was negative for chromosomal aberrations in Fischer 344 rats exposed via inhalation (Vergnes and Morabit 1989), in Sprague-Dawley rats (ARCO 1980) and CD-1 mice (Ward et al. 1994) exposed orally. It was negative for hypoxanthine-guanine phosphoribosyl transferase (hprt) mutant frequency increase in spleen lymphocytes of CD-1 mice exposed orally for six weeks (Ward et al. 1994, 1995), for micronuclei formation in bone marrow in mice exposed via inhalation (Vergnes and Kintigh 1993) or via *i.p.* injection (Kado et al. 1998), for *in vivo* DNA repair increase in cultured primary hepatocytes of CD-1 mice exposed via inhalation (Vergnes and Chun 1994) and for an *in vivo* cytotoxicity assay in rats exposed via inhalation (Vergnes and Morabit 1989).

The only test system in which MTBE has tested positive is the activated mouse lymphoma forward mutation assay (ARCO 1980, Mackerer et al. 1996). TBA, one of MTBE's major metabolites, was negative in this assay (McGregor et al. 1988). MTBE was positive for forward mutations in mouse lymphoma L5178Y tk⁺/tk⁻ cells in the presence, but not the absence, of metabolic activation (ARCO 1980, Stoneybrook Labs. Inc. 1993). HCHO, another one of MTBE's major metabolites, is genotoxic causing both gene mutations and chromosomal damage with the exogenous metabolic activation system. HCHO is also a known carcinogen causing nasal tumors in rodents when inhaled at high concentrations, and may also cause nasopharyngeal tumors in humans via inhalation. Work by Mackerer et al. (1996) suggested that HCHO was the MTBE metabolite responsible for mutagenic activity in the activated mouse lymphoma forward mutation assay. The HCHO was produced from *in vitro* metabolism.

MTBE is volatile and water-soluble. Given the technical difficulties associated with testing volatile chemicals in bacterial and cultured cell systems, it is possible that careful delivery to genetic materials may have yielded data on reasons for the relative lack of genotoxic activity of MTBE *in vitro* (Mackerer et al. 1996, Kado et al. 1998). Additionally, the *in vivo* test systems used to test MTBE were primarily chromosomal damage assays, with two exceptions being the spleen lymphocyte hprt mutation assay (Ward et al. 1994) and the *in vivo-in vitro* mouse hepatocyte unscheduled DNA synthesis assay (Vergnes and Chun 1994). These assay systems would probably not detect gene mutations. They are also relatively insensitive in detecting genotoxic chemicals with known false negatives. *In vivo* genotoxicity and metabolism data is not available for a number of the organ systems such as rat kidney, testis, and spleen/bone marrow, which developed tumors in carcinogenicity bioassays.

Developmental and Reproductive Toxicity

No human studies relevant to MTBE reproductive and developmental toxicity were located. There are a limited number of animal developmental and reproductive toxicity studies, all using the inhalation route of exposure, as listed below:

- one developmental toxicity study in rats exposed to 250 to 2,500 ppm for six hours per day on gestation days (gd) six to 15 (Conaway et al. 1985, Bio/dynamics, Inc. 1984a),
- two developmental toxicity studies in mice exposed to 250 to 2,500 ppm for six hours per day on gestation days six to 15 (Conaway et al. 1985, Bio/dynamics, Inc. 1984b), or to 1,000 to 8,000 ppm for six hours per day on gestation days six to 15 (Bevan et al. 1997b, Tyl and Neepser-Bradley 1989),

- one developmental toxicity study in rabbits exposed to 1,000 to 8,000 ppm for six hours per day on gestation days six to 18 (Bevan et al. 1997b, Tyl 1989),
- one single generation reproductive toxicity study in rats exposed to 300 to 3,400 ppm (Biles et al. 1987),
- one two generation reproductive toxicity study in rats exposed to 400 to 8,000 ppm (Bevan et al. 1997a, Neeper-Bradley 1991).

Study designs and results are outlined in Table 6. Some information on reproductive organs can also be obtained from subchronic and chronic toxicity studies (also outlined in Table 6), and there are a few recent studies of possible endocrine effects.

While no effects on fertility endpoints were reported, these studies provide evidence for adverse effects of MTBE on development. Reduced fetal weight and increased frequency of fetal skeletal variations were reported in mice after MTBE exposure during organogenesis, with a NOAEL of 1,000 ppm (Bevan et al. 1997b, Tyl and Neeper-Bradley 1989). Also, in the rat two generation study, increased postnatal death and decreased postnatal weights were found; the NOAEL was 400 ppm MTBE (Bevan et al. 1997a). A provisional RfC of 173 ppm (48 mg/m³) has been derived using U.S. EPA risk assessment methodology (Sonawane 1994) on the basis of developmental toxicity that occurred in the two-generation rat study (Bevan et al. 1997a, Neeper-Bradley 1991). Additionally, a projected no-effect-concentration in drinking water for humans of 2.3 to 9.2 mg/L has been derived by U.S. EPA (1997a) based on a range of NOAELs (250 to 1,000 ppm) in the two developmental toxicity studies in mice. The NSTC (1997) report stated that "MTBE is not expected to pose a reproductive or developmental hazard under the intermittent, low-level exposure experienced by humans".

The developmental and reproductive toxicity studies were of good quality, and generally conformed to U.S. EPA testing guidelines. The highest inhalation concentration used (8,000 ppm) produced hypoactivity, ataxia, and reduced auditory responsiveness in adult males and females during exposure, reflecting the anesthetic properties of MTBE. Prostration, labored respiration, lacrimation, and periocular encrustation were among the clinical signs reported. There was no increase in adult male and female mortality or organ pathology at any inhalation concentration, but lower food intake and weight gain were sometimes seen at the 8,000 ppm concentration. The developmental toxicity study (Conaway et al. 1985) and single generation study (Biles et al. 1987) in rats, and one of the developmental toxicity studies in mice (Conaway et al. 1985) did not include a dose that was minimally toxic to adult males and females. Little developmental or reproductive toxicity was reported in these studies, but it is difficult to interpret this lack of findings because the concentrations were not high enough to induce adult maternal and paternal toxicity.

Developmental Toxicity

Animal Developmental Toxicity Studies

Dose-dependent effects on fetal weight and fetal skeletal variations were reported in mice; no fetal effects were reported in the rats and rabbits. Notably, the rat developmental toxicity study (Conaway et al. 1985, Bio/dynamics, Inc. 1984a) was conducted in a lower concentration range. In rabbits, maternal toxicity was reported at the highest concentration (8,000 ppm) as reduced maternal food intake, maternal weight loss, hypoactivity, and ataxia during treatment and increased relative liver weights at term. However, no fetal effects of treatment were reported in rabbits (Tyl 1989).

In mice (Bevan et al. 1997b, Tyl and Neeper-Bradley 1989), an 8,000 ppm concentration produced statistically significant lower pregnancy weight gain (approximately 30% lower compared to controls) as well as reduced corrected pregnancy weight gain. Food consumption of dams was lower during the exposure period only. Clinical signs of toxicity, statistically greater in incidence in the 8,000 ppm group on gestation day six to 15, were hypoactivity, ataxia, prostration, labored respiration, lacrimation and periocular encrustation. Group observations during daily exposures included hypoactivity, ataxia and forced respiration. Fetal toxicity endpoints at the 8,000 ppm concentration included: increased postimplantation loss, fewer live fetuses per litter, higher percent of litters with external and visceral malformations, increased incidence of cleft palate and partial atelectasis (absence of fetal lung inflation),

reduced fetal body weight (21%), and increase in the frequency of a number of skeletal variations reflecting delayed ossification.

At the 4,000 ppm exposure, two of these fetal effects (reduced fetal body weight and delayed ossification) were also statistically significant and no maternal toxicity in the form of body weights or clinical signs of toxicity occurred. Group observations at the 4,000 ppm concentrations included hypoactivity and ataxia. The fetal body weight effects and delayed ossification were generally concentration-related at 4,000 and 8,000 ppm, with no indication of treatment related effects at 1,000 ppm, the NOAEL. The mouse developmental toxicity study (Conaway et al. 1985) reported a nonsignificant but apparently concentration-related pattern of increased fetal skeletal malformations in mice exposed to 0, 250, 1,000, or 2,500 ppm (seven, 11, 16, and 22% affected litters), including fused ribs and sternebrae. Conaway et al. (1985) also evaluated skeletal ossification variations (Bio/dynamics, Inc. 1984b), but data were not provided or discussed.

Animal Reproductive Toxicity Studies

As noted above, the two rat reproductive toxicity studies used longer exposures than the developmental toxicity studies, beginning prior to mating and continuing through pregnancy and lactation in the dams. Developmental toxicity in the two generation rat study included reduced pup viability and body weights in the postnatal period for both generations (Bevan et al. 1997a, Neeper-Bradley 1991). Viability, as indexed by the number of dead pups on postnatal day four, was lower than controls in the 8,000 ppm group of both the F₁ and F₂ generations; survival indices were not affected. Group difference in pup body weights were not significant on lactation day one; group differences in body weight appeared later in lactation. Pup weights were consistently lower than controls in the 8,000 ppm group after postnatal day 14 in the F₁ generation and after postnatal day seven in the F₂ generation, and in the 3,000 ppm group after postnatal day 14 in the F₂ generation.

The finding of reduced pup weight gain during lactation in the absence of reduced maternal weight gain is a distinctive finding of the study. Pups were not directly exposed to MTBE during the lactation period but may have been indirectly exposed via dam's milk or MTBE condensation on the dam's fur. The postnatal effects could also have been the result of MTBE effects on maternal behavior or lactation. The findings on postnatal effects are partially supported by the earlier rat single generation study (Biles et al. 1987), which described reduced pup survival and reduced postnatal weights at exposure concentrations of 250 to 2,500 ppm. The statistical significance and dose-related characteristics of these effects varied in the single generation study (see Table 6).

Reproductive Toxicity

Fertility and general toxicity

The two rat reproductive toxicity studies used exposures beginning prior to mating and continuing through pregnancy and lactation in the dams. No indication of reduced fertility was reported in either study. No evaluation of ovarian cyclicity or sperm parameters were included in either study.

As mentioned above, a concentration toxic to the adult breeders was not reached in the single generation study (Biles et al. 1987), but was included in the two generation study (Bevan et al. 1997a, Neeper-Bradley 1991). Increased absolute liver weights (8,000 ppm males and females) and increased relative liver weights (3,000 and 8,000 ppm males and 8,000 ppm females) were reported in the F₁ generation. Liver weights of the F₁ generation were the only organ weights reported.

An unexplained effect was greater lactational body weight gain in the 3,000 ppm dams (F₁) and 8,000 ppm dams (F₀ and F₁) relative to controls. This was due to less maternal weight loss at the end of the lactation period, postnatal days 14 to 28. Lactational weight gain through postnatal day 14 did not differ from controls. Maternal body weight had not been reduced during gestation or at term. However, pups in the 3,000 and 8,000 ppm groups were smaller than controls at some postnatal ages (see section on developmental toxicity above) and this may have resulted in lower energy requirements for lactation.

Reproductive organs

Information on reproductive organs of rats from single and multigeneration studies is varied and incomplete. No effects on reproductive organ weights (testes, epididymides, seminal vesicles, prostate, ovaries) or pathology (testes, epididymides, ovaries) were reported in the rat single generation study (Biles et al. 1987). Reproductive organ weights were not obtained in the rat multigeneration study; no exposure related histopathology of reproductive organs (vagina, uterus, ovaries, epididymides, seminal vesicles, testes, prostate) was reported when 25 rats per sex per generation in the control and 8,000 ppm group were examined (Bevan et al. 1997a, Neeper-Bradley 1991).

Reproductive organ weights and pathology were sometimes reported in subchronic and chronic toxicity and oncogenicity studies in rats. No effects on weight or histopathology of gonads (ovaries and testes) were noted in 14 and 90-day gavage studies in rats (n = 10/sex/group) (Robinson et al. 1990). No effects on histopathology (testes, ovaries, prostate, uterus) were reported in a lifetime (eight weeks to natural death) gavage study in rats (n = 60/sex/group) (Belpoggi et al. 1995). Organ weights were not reported in this oncogenicity study.

Endocrine effects

Moser et al. (1996b, 1998) have conducted studies in mice of potential antiestrogenic effects of MTBE. Endocrine modulating effects of MTBE were suggested by the rodent tumor profile of endocrine sensitive organs in oncogenicity studies. An additional suggestive finding was reduced incidence of uterine endometrial hyperplasia in the mouse inhalation cancer bioassays (Burleigh-Flayer et al. 1991), which implies reduced estrogen action on the endometrium

throughout the lifetime. Moser et al. (1996b, 1998) demonstrated a number of adverse effects of MTBE on the reproductive system of mice:

- lower relative uterine and ovarian weights compared to controls
- increase in overall length of estrous cycle, as well as estrus and nonestrus stages
- lower rate of cell proliferation in the uterine, cervical and vaginal epithelium
- changes in histology of the uterus, cervix and vagina indicative of decreased estrogen action

Body weight gain was also lower in MTBE exposed mice than in controls.

In investigating the potential mechanism of MTBE-induced reduction in estrogen action, Moser et al. (1996b) found that estrogen metabolism was increased twofold in hepatocytes isolated from mice exposed to 1,800 mg MTBE/kg/day by gavage for three days. This change was associated with greater liver weight and P₄₅₀ content. This series of experiments suggested that MTBE might lower circulating estrogen concentrations by increasing estrogen metabolism. However, later studies failed to confirm effects on serum estrogen when female mice were exposed to 8,000 ppm MTBE for four or eight months (Moser et al., 1998). A further series of experiments (Moser et al. 1998) failed to find evidence that MTBE endocrine effects were mediated by the estrogen receptor by studying binding of MTBE and its metabolites to the estrogen receptor, changes in expression of estrogen receptor in MTBE exposed mice, and alterations of estrogen receptor activation and translocation in a transfection assay. The authors suggest that MTBE may exert an antiestrogenic action by a mechanism which does not involve a change in circulating estrogen or estrogen receptor binding.

The consequences of reduced estrogen action induced by MTBE in mice are not known; no fertility studies have been conducted in mice. It is also not clear whether similar effects occur in other species, at other doses, or with other exposure durations, since parallel studies have not been done. The specificity of the effect also needs to be determined. Unleaded gasoline has been found to have some antiestrogenic effects similar to MTBE (MacGregor et al. 1993, Moser et al. 1996b, Standeven et al. 1994). Also, an in vivo study, reported recently in abstract form (Okahara et al. 1998) described mild estrogenic and antiestrogenic effects in pubertal mice (21 to 25 days old) gavaged with 600 or 1,500 mg MTBE/kg body weight for five days.

Other Relevant Data

As discussed in the section on metabolism and pharmacokinetics, MTBE is distributed to all major tissues studied in the rat. MTBE is metabolized in the liver to TBA. TBA appears to be widely distributed (Aarstad et al. 1985, Borghoff et al. 1996a, Savolainen et al. 1985). No studies specifically examining distribution of MTBE or TBA to male or female reproductive organs, or the placenta, embryo, or fetus were located in the general published literature. In view of the general widespread distribution, it is plausible that MTBE and TBA distribute to these tissues.

Several studies have examined the developmental toxicity of TBA in mice (oral) and rats (inhalation and oral). No reproductive studies of TBA were located. NTP conducted subchronic and carcinogenesis studies in mice and rats by drinking water which examined some reproductive endpoints. There is also an in vitro study of TBA and mouse sperm.

The specific studies located were:

- one developmental toxicity study in mice, oral (liquid food), 0, 0.5, 0.75, or 1% weight to volume, gestation days six to 20 (Daniel and Evans 1982),

DRAFT

- one developmental toxicity study in mice, oral (gavage), 0 or 780 mg/kg, twice per day, gestation days six to 18 (Faulkner et al. 1989),
- one developmental toxicity study in rats, inhalation, 0, 2,000, 3,500, or 5,000 ppm, seven hours per day, gestation days one to 19 (Nelson et al. 1989a),
- one developmental toxicity study in rats, inhalation, 0, 6,000, 12,000 mg/m³ (0, 1,660, or 3,330 ppm), seven hours per day, gestation days one to 19 (abstract only) (Nelson et al. 1989b),
- one developmental toxicity study in rats, oral (liquid food), 0, 0.65, 1.3, or 10.9% volume to volume, gestation days eight to 22 (abstract only) (Abel and Bilitzke 1992),
- one developmental toxicity study in rats, gastric cannula, 0, or 0.6 to 2.7 g/kg/day, postnatal day four to seven (Grant and Samson 1982),
- subchronic (13 weeks) and carcinogenesis (two years) studies in rats and mice (both sexes), oral (water), various concentrations (NTP 1995),
- one in vitro study of mouse sperm fertilization capacity (Anderson et al. 1982).

With the exception of Nelson et al. (1989a), reporting of the data in the developmental studies was incomplete. Developmentally toxic effects were observed in mice and rats orally administered TBA, including prenatal and postnatal death (Abel and Bilitzke 1992, Faulkner et al. 1989, Daniel and Evans 1982) and postnatal developmental retardation (Daniel and Evans 1982). Malformations were not observed (Faulkner et al. 1989). The inhalation study in rats by Nelson et al. (1989a) found developmental retardation, as manifested in lower fetal weights, at concentrations of 2,000, 3,500 and 5,000 ppm TBA, and a higher percent of skeletal variations compared to controls at 3,500 and 5,000 ppm. No increases in resorptions or malformations were observed. Lower maternal weight was reported at 5,000 ppm. Maternal neurobehavioral effects associated with the exposures (narcosis at 5,000 ppm, unsteady gait at 3,500 and 5,000 ppm, unsteady at 2,000 ppm) were also observed in the Nelson et al. (1989a) study.

The NTP subchronic and carcinogenesis studies in mice and rats by drinking water used various concentrations of TBA. In these studies, systemic toxicity was observed at the high concentration, usually including death, reduced weight gain, and altered kidney weight. The studies found little indication of potential reproductive toxicity. Specifically, no effects on testes weight or sperm were observed. Minor and inconsistent effects on testes histopathology and estrous cyclicity were observed at the high concentrations. The in vitro study found no effect of TBA on mouse sperm fertilization capacity.

**Table 6. MTBE: Developmental and Reproductive Toxic Effects
(studies in alphabetical order by author)**

Study design⁽¹⁾	Reported effects⁽²⁾	Reference
Rat (Sprague-Dawley) oral (gavage) Male and female 104 weeks, 4 days/week 0, 250, 1,000 mg/kg/day	Male: No increased male death, reduced male weight gain, reduced food consumption, or testicular histopathological effects. Female: No reduced female weight gain, reduced food consumption, or ovarian histopathological effects. 250, 1,000 mg/kg/day: Increased female death (dose-responsive, SS not addressed).	Belpoggi et al. 1995

Study design⁽¹⁾	Reported effects⁽²⁾	Reference
Mouse (CD-1) inhalation gd 6-15, 6 hours/day Target concentrations: 0, 1,000, 4,000, 8,000 ppm Analytical concentrations: 0, 1,035, 4,076, 8,153 ppm	No maternal death, altered liver weight. 8,000 ppm: Reduced maternal weight (SS), reduced body weight gain (SS), reduced food consumption during treatment period (SS). Clinical signs (individual observations): maternal, hypoactivity (SS), ataxia (SS), prostration (SS), labored respiration (SS), lacrimation (SS), periocular encrustation (SS). Clinical signs (group observations during daily exposure periods): maternal hypoactivity, ataxia, labored breathing. 4,000 ppm: Clinical signs (group observations during daily exposure periods): maternal hypoactivity, ataxia. No increased pre-implant loss, early resorptions. 8,000 ppm: Increased post-implant loss (late resorptions and dead fetuses) (SS), reduced live litter size (SS), altered sex ratio (less males) (SS), increased cleft palate (SS), increased soft tissue malformations (SS), reduced fetal weight (SS), increased incidence of some skeletal variations (mainly reduced ossification) (SS). 4,000 ppm: Reduced fetal weight (SS), increased incidence of some skeletal variations (mainly reduced ossification) (SS).	Bevan et al. 1997b, Tyl and Neepers-Bradley 1989

Study design⁽¹⁾	Reported effects⁽²⁾	Reference
Rabbit (New Zealand White) inhalation gd 6-18, 6 hours/day Target concentrations: 0, 1,000, 4,000, 8,000 ppm Analytical concentrations: 0, 1,021, 4,058, 8,021 ppm	No maternal death, reduced body weight, clinical signs of toxicity before or after daily exposure periods. 8,000 ppm: Reduced maternal weight gain (gd 6-18) (SS), reduced food consumption (gd 6-18) (SS), increased relative liver weight (SS). Clinical signs (group observations during daily exposure periods): hypoactivity, ataxia. 4,000 ppm: Reduced maternal weight gain (gd 6-9) (SS), reduced food consumption (gd 6-10) (SS). No increased pre- or post-implant loss, reduced litter size, altered sex ratio, reduced fetal weight, increased malformations, or increased skeletal variations.	Bevan et al. 1997b, Tyl 1989

Study design ⁽¹⁾	Reported effects ⁽²⁾	Reference
Rat (Sprague-Dawley) inhalation 2 generation reproductive Target concentrations: 0, 400, 3,000, 8,000 ppm Analytical concentrations: 0, 402, 3,019, 8,007 ppm Male: 6 hours/day, 10 weeks (5 days/week) + mating + gestation Female: 6 hours/day, 10 weeks (5 days/week) + mating + gestation (gd 1-19) + lactation (pnd 5-28) Exposures for F ₀ starting at pnd 42, and F ₁ starting on pnd 29-31. Pups not placed in inhalation chambers during lactation.	No adult male or female deaths (F ₀ or F ₁), reduced adult female body weight (F ₀), reduced adult female body weight gain (F ₁), reduced adult female food consumption (F ₀). 8,000 ppm: Reduced adult male body weight (F ₀ , F ₁) (SS), reduced adult male body weight gain (F ₀ : weeks 0-3, 5-7; F ₁ : weeks 0-2, 5-6), reduced adult female body weight (F ₁ : weeks 0-8, not gestation or lactation) (SS), reduced adult female body weight gain (F ₀ : weeks 0-1, 5-6, not gestation or lactation) (SS), increased female body weight gain during lactation (F ₀ , F ₁) (SS), increased adult male and female absolute and relative liver weights (F ₁) (SS), reduced adult female food consumption (F ₁ : lactation days 7-14, not pre-breed or gestation) (SS). Clinical signs (individual observations): adult male, perioral wetness (F ₀ , F ₁), perioral encrustation and salivation (F ₁); adult female, perioral wetness (F ₀ , F ₁), perioral encrustation, salivation and urine stains (F ₁). Clinical signs (group observations during daily exposure periods): adult male and female, ataxia (F ₀ , F ₁), hypoactivity (F ₀ , F ₁), blepharospasm (F ₀ , F ₁), lack of startle reflex (F ₀ , F ₁). 3,000 ppm: Increased adult male relative liver weights (F ₁) (SS), increased adult female body weight gain (F ₁ : lactation) (SS). Clinical signs (group observations during daily exposure periods): adult male and female, hypoactivity (F ₀ , F ₁), blepharospasm (F ₀ , F ₁), lack of startle reflex (F ₀ , F ₁). No reduced mating (F ₀ , F ₁), reduced fertility (F ₀ , F ₁), reduced live litter size (F ₁ , F ₂), reduced postnatal survival after pnd 4 (F ₁ , F ₂), reduced litter size at end of lactation (F ₁ , F ₂), reduced birth (lactation day one) weight (F ₁ , F ₂). 8,000 ppm: Increased dead pups pnd 4 (F ₁ , F ₂) (SS), reduced postnatal weight (F ₁ : pnd 14-28, F ₂ : pnd 7-28) (SS), reduced postnatal weight gain (F ₁ : pnd 7-21, F ₂ : pnd 1-21) (SS). 3,000 ppm: Reduced postnatal weight (F ₁ : pnd 14, F ₂ : pnd 14-28) (SS), reduced postnatal weight gain (F ₁ : pnd 7-14, F ₂ : pnd 7-21) (SS).	Bevan et al. 1997a, Neeper-Bradley 1991

Study design ⁽¹⁾	Reported effects ⁽²⁾	Reference
Rat (Sprague-Dawley) inhalation Reproductive: 1 generation, 2 litter Male: 6 hours/day, 12 weeks (5 days/week), + mating Female: 6 hours/day, 3 weeks (5 days/week), + mating + gestation (gd 0-20) + lactation (pnd 5-21) Male and female exposures repeated for second litter. Target concentrations in text: 0, 250, 1,000, 2,500 ppm. Target concentrations in abstract: 0, 300, 1,300, 3,400 ppm. Analytical exposures, Male/Female: 0/0, 290/300, 1,180/1,240, 2,860/2,980.	No male or female death, reduced male or female body weight, altered testes or ovary weight, adverse histopathological effects on ovaries or testes (F ₀). 2,500, 250 ppm: Increased incidence dilated renal pelvis in females (NOT 1,000 ppm). No reduced mating, reduced fertility (pregnancy rate), reduced litter size (live/total) (F _{1a} , F _{1b}), altered sex ratio (F _{1a} , F _{1b}), reduced pup viability at birth (live/total) (F _{1a}), reduced birth weight (F _{1a} , F _{1b}). reduced pup survival on pnd 4 (F _{1b}), reduced pup survival on pnd 21 (F _{1a} , F _{1b}). 2,500 ppm: Reduced pup viability at birth (live/total) (F _{1b}) (SS) (Note high in controls: control 99%, 1,000 and 2,500 ppm 95.5%. Authors discount biological significance), reduced postnatal weight on pnd 14, 21 (F _{1a} , F _{1b}) (NOT SS). 1,000 ppm: Reduced pup viability at birth (live/total) (F _{1b}) (SS) (Note high in controls: control 99%, 1,000 and 2,500 ppm 95.5%. Authors discount biological significance), reduced pup survival on pnd 4 (F _{1a}) (NOT 2,500 ppm), reduced postnatal weight on pnd 14, 21 (F _{1a} , F _{1b}) (NOT SS). 250 ppm: Reduced pup survival on pnd 4 (F _{1a}) (NOT 2,500 ppm) (SS).	Biles et al. 1987

Study design⁽¹⁾	Reported effects⁽²⁾	Reference
Mouse (CD-1) inhalation Male and female 6 hours/day, 5 days/week, 18 months 0, 400, 3,000, 8,000 ppm	Male: No alteration in testes weight, testicular (or other reproductive organ) histopathological effects. 8,000 ppm: Increased male death (SS), reduced weight (SS), increased liver weight (SS), blepharospasm, hypoactivity, ataxia, lack of startle reflex, prostration. 3,000 ppm: Increased liver weight (SS), blepharospasm, hypoactivity, ataxia, lack of startle reflex, stereotypy. 400 ppm: Increased liver weight (SS). Female: No ovarian (or other reproductive organ) histopathological effects. 8,000 ppm: Reduced weight (SS), increased liver weight (SS), blepharospasm, hypoactivity, ataxia, lack of startle reflex, prostration. 3,000 ppm: Increased liver weight (SS), blepharospasm, hypoactivity, ataxia, lack of startle reflex, stereotypy.	Burleigh- Flayer et al. 1992

Study design ⁽¹⁾	Reported effects ⁽²⁾	Reference
Rat (Fischer 344) inhalation Male and female 6 hours/day, 5 days/week Male: 0, 400 ppm; 104 weeks Male: 3,000 ppm; 97 weeks Male: 8,000 ppm; 82 weeks Female: 0, 400, 3,000, 8,000 ppm; 104 weeks	Male: No altered liver or testes weight to 400 ppm (see note). 8,000 ppm: Increased male death (SS), reduced weight (SS), (increased) nephropathy, ataxia, hypoactivity, blepharospasm, lack of startle reflex, testicular mineralization. 3,000 ppm: Increased male death (SS), nephropathy, ataxia, hypoactivity, blepharospasm, lack of startle reflex, testicular mineralization. 400 ppm: Increased male death (SS), nephropathy, testicular mineralization. Note: Remaining males in 8,000 and 3,000 ppm groups were sacrificed early due to high group mortality. Authors attribute mortality and mineralization of "numerous tissues" to nephropathy. No statistical evaluation of testes or other organ weight, or, apparently, histopathological changes, was performed by the authors for the 8,000 or 3,000 ppm groups. Female: No increased female death or ovarian (or other reproductive organ) histopathological effects. 8,000 ppm: Reduced female weight (SS), increased liver weight (SS), ataxia, hypoactivity, blepharospasm, lack of startle reflex, nephropathy. 3,000 ppm: Increased liver weight (SS), ataxia, hypoactivity, blepharospasm, lack of startle reflex, nephropathy.	Chun et al. 1992

Study design⁽¹⁾	Reported effects⁽²⁾	Reference
Rat (Sprague-Dawley) inhalation gd 6-15, 6 hours/day Target concentrations: 0, 250, 1,000, 2,500 ppm Analytical concentrations: 0, 250, 1,000, 2,430 ppm "Nominal" concentrations: 0, 260, 1,100, 3,300 ppm	No maternal death, reduced maternal weight, altered water consumption, or altered liver weight. 2,500, 1,000, 250 ppm: Reduced maternal food consumption on gd 9-12 (SS). No increased pre- or post-implant loss, reduced live litter size, reduced fetal weight, reduced crown-rump distance, altered sex ratio, increased malformations, increased ossification variations.	Conaway et al. 1985, Bio/dynamics, Inc. 1984a
Mouse (CD-1) inhalation gd 6-15, 6 hours/day Target concentrations: 0, 250, 1,000, 2,500 ppm Analytical concentrations: 0, 280, 1,110, 2,710 ppm "Nominal" concentrations: 0, 280, 1,200, 3,500 ppm	No maternal death, reduced maternal weight, altered food or water consumption, altered liver weight. No increased pre- or post-implant losses, reduced live litter size, reduced fetal weight, reduced crown-rump distance, altered sex ratio, increased malformations. [Fetuses with skeletal malformations: control, 1.6%; 250 ppm, 1.7%; 1,000 ppm, 2.4%; 2,500 ppm, 3.1% NOT SS]	Conaway et al. 1985, Bio/dynamics, Inc. 1984b

Study design ⁽¹⁾	Reported effects ⁽²⁾	Reference
Rat (Sprague-Dawley) oral (gavage) Male and female 14 days 0, 357, 714, 1,071, 1,428 mg/kg/day	Male: No increased male death, altered testes weight, or testicular histopathological effects. 1,428 mg/kg/day: anesthesia, reduced male weight gain (SS), diarrhea 1,071, 714 mg/kg/day: reduced male weight gain (SS), diarrhea 357 mg/kg/day: diarrhea Female: No increased female death, altered ovary weight, or ovarian histopathological effects. 1,428 mg/kg/day: anesthesia, reduced female weight gain (SS), diarrhea 1,071 mg/kg/day: reduced female weight gain (SS), diarrhea 714, 357 mg/kg/day: diarrhea	Robinson et al. 1990
Rat (Sprague-Dawley) oral (gavage) Male and female 90 days 0, 100, 300, 900, 1,200 mg/kg/day	Male: No increased male death, altered testes weight, or testicular histopathological effects. 1,200 mg/kg/day: anesthesia, reduced male weight (NOT SS), diarrhea 900, 300, 100 mg/kg/day: diarrhea Female: No increased female death, altered ovary weight, or ovarian histopathological effects. 1,200 mg/kg/day: anesthesia, reduced female weight (SS), diarrhea 900, 300 mg/kg/day: reduced female weight (NOT SS), diarrhea 100 mg/kg/day: diarrhea	Robinson et al. 1990

(1) Abbreviations: gd = gestation day, pnd = postnatal day.

(2) Effects reported by authors to be statistically significant (SS) or biologically noteworthy.

Immunotoxicity

Oral administration of 1,428 mg/kg/day MTBE for 14 days reduced absolute spleen weights and absolute and relative thymus weights in female rats but not in males and did not produce histopathological lesions in the spleen or thymus. Similar results were observed following 90 days treatment with an oral dose of 100 to 1,200 mg/kg/day MTBE (Robinson et al. 1990). An increased incidence of dysplastic proliferation of lymphoreticular tissues was observed in female rats gavaged with 250 or 1,000 mg/kg/day MTBE, four days per week for 104 weeks (Belpoggi et al. 1995). The authors discussed the possibility that these lesions had the potential to develop into the lymphomas and leukemias also observed in this study. Gavaged Sprague-Dawley male rats daily for 28 days with 40, 400, or 800 mg/kg/day of MTBE produced an overall increased percentage of apoptotic-type comets in peripheral blood lymphocytes but no dose

produced a statistical increase over vehicle controls. DNA strand breakage was significantly increased in the 800 mg/kg/day group and depressed body weight gain and high corticosterone levels were observed at 28 days (Lee et al. 1998).

Neurotoxicity

Acute oral exposure in rats caused marked CNS depression at doses greater than 1,900 mg/kg, ataxia at doses greater than 2,450 mg/kg, loss of righting reflex at doses greater than 3,160 mg/kg, and tremors and labored breathing at doses greater than 4,080 mg/kg. A no observed effect level (NOEL) of 40 mg/kg for adverse but reversible neurological effects for acute oral exposure was identified (Bioresearch Laboratories 1990b) and an acute oral MRL of 0.4 mg/kg/day was calculated by ATSDR (1996).

Scholl et al. (1996) measured the duration of ataxia and hypnosis in male Fischer 344 rats pretreated with P₄₅₀ inducers following a single sub-hypnotic (0.5 mg/kg) and hypnotic (1.2 mg/kg) i.p. dose of MTBE. Pretreatment with phenobarbital, and to a lesser extent clofibrate but not beta-naphthoflavone, prolonged the duration of ataxia or narcosis from MTBE compared with the vehicle control. The data suggested that the biotransformation status is a major potential determinant of sensitivity to the CNS depression effects of MTBE.

Two inhalation studies indicated that MTBE might be a weak neurotoxicant in adult rats with primary effects of acute impairment. A six-hour inhalation study and a 13-week repeated vapor inhalation study produced signs of reversible CNS depression following exposure to 8,000 ppm and, to a lesser extent, to 4,000 ppm vapor with a NOAEL of 800 ppm (Dodd and Kintigh 1989, Daughtrey et al. 1997). MTBE induced some mild and reversible CNS toxicity but did not appear to be a neurotoxicant under the conditions of these studies (Fueta et al. 1994).

Chronic Toxicity

Sprague-Dawley rats (60 animals per sex, per dose group) were given 0, 250 or 1,000 mg/kg/day MTBE in olive oil via gavage, four days per week, for 104 weeks. This dosing regimen gives a seven-day time-weighted average daily dose of 0, 143, and 571 mg/kg/day. Survival appeared to be decreased in female rats after 16 weeks, but no statistical treatments on data were reported. There was no reporting of hematological, clinical chemistry or urinalysis parameters, or any indication as to whether or not these endpoints were evaluated. The authors did not observe any differences in food consumption or final body weights in the various groups. In addition, they did not report any noncancer histopathological changes (Belpoggi et al. 1995, 1997). Due to the

limited scope, intermittent treatment schedule and scant data reporting on noncancer endpoints in this study, it is not possible to identify an adequate NOAEL or LOAEL.

Kidney toxicity was also observed in both males and females in the two-year inhalation study in Fischer 344 rats by Chun et al. (1992) discussed in the next section on carcinogenicity. U.S. EPA derived a RfC of three mg/m³ based on the kidney and liver effects of MTBE (U.S. EPA 1993, 1997c). These data support the conclusion that, after MTBE exposure, kidney toxicity is of toxicological concern. However, the use of the Robinson et al. (1990) study for evaluation of kidney effects, as detailed in the previous section on subchronic toxicity, has two significant uncertainties. One is that the study was for 90 days and not for a lifetime, and the second is the extrapolation of dose from a single daily bolus dose in corn oil to the continuous small doses from drinking water exposure. In general, it would be anticipated that a 90-day exposure period would tend to underestimate the toxicity, while the bolus dose (a NOAEL of 100 mg/kg/day) would be more likely to overestimate the toxic response. However, the relative effects of these two factors are uncertain.

Animal studies conducted at very high levels of exposure to MTBE, i.e., at greater than 1,000 ppm, through inhalation caused increased liver, kidney, spleen, and adrenal weights; decreased brain weight, body weight, and body weight gain; swollen periocular tissue; and ataxia in rodents. Increased prostration (lying flat) or exhaustion was reported in female rodents only.

Carcinogenicity

No data on long-term effects of human exposure to MTBE relevant to cancer risk were found in recent literature searches performed by OEHHA.

The carcinogenic activity of MTBE has been investigated in male and female Sprague-Dawley rats administered MTBE by gavage (Belpoggi et al. 1995, 1997) and in male and female Fischer 344 rats (Chun et al. 1992, Bird et al. 1997) and CD-1 mice (Burleigh-Flayer et al. 1992, Bird et al. 1997) exposed to MTBE by inhalation. In rats receiving MTBE by gavage for 24 months, statistically significant increases in Leydig interstitial cell tumors of the testes were observed in males, and statistically significant increases in lymphomas and leukemias (combined) were observed in females. An increase in the incidence of uterine sarcomas was also observed in MTBE-exposed female rats, but was not statistically significant at the $p < 0.05$ level. In rats exposed to MTBE by inhalation for up to 24 months, statistically significant increases in the incidences of renal tubular tumors and Leydig interstitial cell tumors of the testes were observed in males. In mice exposed to MTBE by inhalation for up to 18 months, statistically significant increases in the incidences of liver tumors were observed in females (hepatocellular adenomas; hepatocellular adenomas and carcinomas combined) and males (hepatocellular carcinomas; hepatocellular adenomas and carcinomas combined). These studies are described in more detail below.

Oral Exposure Studies

Rat gavage exposure studies: Belpoggi et al. (1995, 1997)

Groups of 60 male and 60 female eight-week old Sprague-Dawley rats were administered MTBE in olive oil by gavage at doses of 0 (oil only), 250 or 1,000 mg/kg body weight/day, four days per week for 104 weeks. Animals were maintained until natural death; the last animal died at

DRAFT

174 weeks of age. No difference in water or food consumption, or in mean body weights was observed between treated and control animals of either sex. A dose-related decrease in survival was observed in females. At 56 weeks of age, survival was approximately 98%, 85%, and 78% in controls, low- and high-dose females, respectively; at 88 weeks of age, survival in those same groups was approximately 76%, 60%, and 43%. In males, there was no difference in survival between the controls and the low-dose animals. However, after 88 weeks, survival in high-dose males exceeded that of low-dose and control males. At 104 weeks of age, survival was approximately 30% in low-dose and control males and 43% in high-dose males; at 120 weeks of age, survival in those same groups was approximately 11% and 32%. A dose-related increase in the combined incidence of lymphomas and leukemia was observed in female rats (Table 7). The authors reported that the increase was highly significant ($p < 0.01$) in the high-dose group and marginally significant in the low-dose group, when analyzed using a log-ranked test as described by Mantel (1966) and Cox (1972). When analyzed using the Fisher Exact test, the combined incidence of lymphomas and leukemia in high-dose females was significantly different from controls at the $p = 0.001$ level. Historical control incidence rates in this laboratory for lymphomas and leukemias (combined) was $< 10\%$ in female Sprague-Dawley rats (Belpoggi et al. 1995). The authors also noted an increase in uterine sarcomas in the low-dose females (5/60 versus 1/60 in controls), however, this increase did not reach statistical significance ($p = 0.1$ by Fisher's Exact test). In males, a statistically significant increased incidence of Leydig cell tumors of the testes was observed in the high-dose group (Table 7). The authors reported that this increase was significant at the $p = 0.05$ level using a prevalence analysis for nonlethal tumors (Hoel and Walburg 1972).

Table 7. Tumors in Sprague-Dawley Rats Receiving MTBE by Gavage, 250 or 1,000 mg/kg/day, Four days/week for 104 Weeks (Belpoggi et al. 1995, 1997)

Tumor site and type		Dose ^a (mg/kg/day)		
		0	250	1,000
Females				
Hemolymphoreticular tissues (including mesenteric lymph nodes)	Lymphomas and leukemias	2/58 ^b (3.4%)	6/51 ^b (11.8%)	12/47 ^{b,c,d} (25.5%)
Males				
Testes	Leydig interstitial cell tumors	2/26 ^e (7.7%)	2/25 ^e (8.0%)	11/32 ^{e,f} (34.4%)

^a Administered in olive oil, four days/week, for 104 weeks.

^b Number of lesion-bearing animals/total alive at 56 weeks of age, when the first leukemia was observed.

^c Incidence relative to control group was significant ($p < 0.01$) using a long-ranked test (Mantel 1966, Cox 1972), as reported by Belpoggi et al. (1995). Incidence relative to control group was significant by the Fisher Exact test ($p = 0.001$).

^d Dose-related trend was significant by the Cochran-Armitage trend test ($p < 0.01$).

^e Number of lesion-bearing animals/total alive at 96 weeks of age, when the first Leydig cell tumor was observed.

^f Incidence relative to control group was significant at the $p = 0.05$ level using prevalence analysis for non-lethal tumors (Hoel and Walburg 1972), as reported by Belpoggi et al. (1995). Incidence relative to control group was significant by the Fisher Exact test ($p = 0.015$).

Inhalation Exposure Studies

Rat inhalation exposure: Chun et al. (1992), Bird et al. (1997)

Groups of 50 male and 50 female eight-week old Fischer 344 rats were exposed to 0, 400, 3,000, or 8,000 ppm MTBE vapor by inhalation (corresponding to analytical mean concentrations of 403, 3,023, or 7,977 ppm, or 1,453, 10,899, 28,760 mg/m³ MTBE). The animals were exposed for six hours/day, five days/week for 24 months, except for the mid- and high-dose males, which were terminated at 97 and 82 weeks, respectively, due to a dose-dependent increased mortality rate from chronic progressive nephropathy. Low-dose males also experienced an increase in nephropathy that was associated with a slight increase in mortality and a decrease in survival. Survival times for females were not significantly different between exposed and control rats.

However, there were slightly more deaths due to chronic progressive nephropathy in the mid- and high-dose females than in the low-dose and control females. Body weight gain and absolute body weight were decreased in both sexes of the high-dose group. Exposure-related increases in kidney and liver weights were reported in mid- and high-dose females, but not in males. Chun et al. (1992) concluded that the maximum tolerated dose (MTD) was exceeded in both sexes at high- and mid-dose levels, based on increased mortality. Other observed effects of MTBE exposure included anesthetic effects in rats of both sexes in the mid- and high-dose groups.

A detailed histopathology examination was performed on all animals in the control and high-dose groups, and on all animals that died or were sacrificed moribund. Only a limited histopathology examination was performed on non-moribund animals from the low- and mid-dose groups that survived to terminal sacrifice; for males, only the liver, kidneys, testes and gross lesions were evaluated, while for females, only the liver and gross lesions were examined microscopically (Bird et al. 1997).

In males, a statistically significant increase in renal tubular adenoma and carcinoma (combined) was observed in the mid-dose group (Table 8). In high-dose males renal tubular adenomas were increased, however, this increase did not reach statistical significance (Table 8). The sensitivity of the bioassay to detect a dose-related increase in renal tumors in the high-dose group is likely to have been reduced by the high rate of early mortality, and the early termination of this treatment group at week 82. Despite the reduced sensitivity of the bioassay, a statistically significant increase in Leydig interstitial cell testicular tumors was observed in mid- and high-dose males, with a clear dose-response evident (Table 8).

Historical laboratory control values for Leydig testicular tumors in Fischer rats ranged from 64 to 98% (Bird et al. 1997).

In female Fischer 344 rats exposed to MTBE vapor, a single rare renal tubular cell adenoma was observed in one mid-dose animal; no treatment-related increases in tumor incidence were observed (Chun et al. 1992, Bird et al. 1997). MTBE treatment of females was associated with several nonneoplastic kidney lesions, however. Both female and male rats exposed to MTBE experienced a dose-related increase in mortality from chronic progressive nephropathy. Increases in microscopic kidney changes indicative of chronic nephropathy were seen in all treated males and in mid- and high-dose females. All treated males had increases in the severity of mineralization and interstitial fibrosis of the kidney, while increases in mild to moderate glomerulosclerosis, interstitial fibrosis, and tubular proteinosis were observed in females.

Table 8. Tumors in Male Fischer 344 Rats Receiving MTBE by Inhalation, 0, 400, 3,000, or 8,000 ppm, for up to 24 Months^a (Chun et al. 1995, Bird et al. 1997)

Tumor site and type		Concentration ^b (ppm)			
		0	400	3,000	8,000
Kidney	renal tubular adenoma	1/35 ^c	0/32 ^c	5/31 ^c	3/20 ^c
	renal tubular carcinoma	0/35 ^c	0/32 ^c	3/31 ^c	0/20 ^c
	renal tubular adenoma and carcinoma (combined)	1/35 ^c (3%)	0/32 ^c (0%)	8/31 ^{c,d} (26%)	3/20 ^c (15%)
Testes	Leydig interstitial cell tumors	32/50 (64%)	35/50 (70%)	41/50 ^e (82%)	47/50 ^f (94%)

^a Mid- and high-dose animals were terminated at 97 and 82 weeks, respectively, due to a dose-dependent increased mortality rate from chronic progressive nephropathy.

^b Administered as MTBE vapor six hours/day, five days/week.

^c Survival-adjusted tumor incidence rates were used to attempt to control for excess early mortality in the mid- and high-dose groups (U.S. EPA, 1995c).

^{d, e, f} Incidence relative to control group was significant by the Fisher Exact test (^dp < 0.01, ^ep < 0.05, ^fp < 0.001).

Mouse inhalation exposure: Burleigh-Flayer et al. (1992), Bird et al. (1997)

Groups of 50 male and 50 female eight-week old CD-1 mice were exposed to 0, 400, 3,000, or 8,000 ppm MTBE vapor by inhalation (corresponding to analytical mean concentrations of 402, 3,014, or 7,973 ppm or 1,442, 10,816, or 28,843 mg/m³ MTBE). The animals were exposed for six hours per day, five days per week, for 18 months. Increased mortality and decreased mean survival time were observed only for male mice in the high-dose group. A slightly increased frequency of obstructive uropathy, a condition which occurs spontaneously in this mouse strain, was observed in high-dose males, however, deaths due to the condition were within the range noted for historical controls. Body weight gain and absolute body weights were decreased in high-dose males and females. Dose-dependent increases in liver weights were observed in both sexes. Kidney weights were increased in high-dose females and in low- and mid-dose males. Burleigh-Flayer et al. (1992) concluded that the MTD was exceeded in both sexes at the high-dose level. Other observed effects of MTBE exposure included anesthetic effects in mice of both sexes in the mid- and high-dose groups.

DRAFT

A detailed histopathology examination was performed on all animals in the control and high-dose groups, and on all animals that died or were sacrificed moribund. Only a limited histopathology examination was performed on non-moribund animals from the low- and mid-dose groups that survived to terminal sacrifice; for males, only the liver, spleen and submandibular lymph nodes were evaluated, while for females, only the liver, uterus and stomach were examined microscopically (Bird et al. 1997).

In females, a statistically significant increased incidence of hepatocellular adenomas was observed in the high-dose group (Table 9). The incidence of hepatocellular adenomas and carcinomas (combined) was also increased in high-dose females, however, only two hepatocellular carcinomas were reported, one each in the low- and high-dose groups. In males, a statistically significant increase in hepatocellular carcinomas and hepatocellular adenomas and carcinomas (combined) was observed in the high-dose group (Table 9). Bird et al. (1997) noted that the combined incidence of adenomas and carcinomas in high-dose males was similar to the historical incidence for male CD-1 mice of 33%. However, after correcting for the number of animals alive at 49 weeks, when the first hepatocellular adenoma was observed in males, the incidence in the high-dose group was 43% (16/37, see Table 9), representing a clear increase above the cited historical incidence in male CD-1 mice. Burleigh-Flayer et al. (1992) concluded that the increased incidence of liver tumors in the high-dose groups (adenomas in females and carcinomas in males) could be attributed to MTBE exposure. The ability of this study to detect increases in tumor incidence was likely decreased by the shortened study length (18 versus 24 months).

Table 9. Tumors in CD-1 Mice Receiving MTBE by Inhalation, 0, 400, 3,000 or 8,000 ppm, for up to 18 Months^a (Burleigh-Flayer et al. 1992, Bird et al. 1997)

Tumor site and type		Dose ^b (ppm)			
		0	400	3,000	8,000
Females					
Liver	hepatocellular adenoma	2/50	1/50	2/50	10/50 ^c
	hepatocellular carcinoma	0/50	1/50	0/50	1/50
	hepatocellular adenoma and carcinoma (combined)	2/50	2/50	2/50	11/50 ^d
Males					
Liver	hepatocellular adenoma	11/47 ^e	11/47 ^e	9/46 ^e	12/37 ^e
	hepatocellular carcinoma	2/42 ^f	4/45 ^f	3/41 ^f	8/34 ^{c,f}
	hepatocellular adenoma and carcinoma (combined)	12/47 ^e	12/47 ^e	12/46 ^e	16/37 ^{c,e}

^a Male mice in the high-dose group experienced early mortality.

^b Administered as MTBE vapor six hours/day, five days/week.

^{c,d} Incidence relative to control group was significant by the Fisher Exact test (^cp < 0.05, ^dp < 0.01).

^e Number of lesion-bearing animals/total alive at 49 weeks, when the first hepatocellular adenoma was observed.

^f Number of lesion-bearing animals/total alive at 63 weeks, when the first hepatocellular carcinoma was observed.

Other Relevant Data

Structure-Activity Comparisons

MTBE and similar ethers generally undergo metabolism at the ethereal bond to form the corresponding alcohol and an aldehyde (Savolainen et al. 1985). Other structurally similar ethers include ETBE and tertiary-amyl methyl ether (TAME). No studies have been reported to date on the carcinogenicity of ETBE or TAME. Published data on the genotoxic potential of ETBE and TAME are few in number; ETBE and TAME tested negative in the Salmonella reverse mutation assay, and TAME did not induce micronuclei in mouse bone marrow cells following exposure in vivo (NSTC 1997). MTBE is made by isobutene and methanol, or TBA and methanol. NTP has documented some evidence of carcinogenic activity for isobutene in male rats (NTP 1997), and for TBA in male rats and female mice (NTP 1995).

Pathology

The tumors observed by Belpoggi et al. (1995, 1997) in hemolymphoreticular tissues in the female Sprague-Dawley rat were diagnosed as lymphomas and leukemias. Lymphomas and leukemias of the Sprague-Dawley rat commonly arise from a similar cellular origin, in which case the aggregation of these tumors for carcinogen identification and risk assessment purposes is appropriate.

The testicular tumors observed in both the Sprague-Dawley (Belpoggi et al. 1995, 1997) and Fischer 344 (Chun et al. 1992, Bird et al. 1997) rat strains were diagnosed as Leydig interstitial cell tumors. The spontaneous incidence of these tumors is typically much lower in the Sprague-Dawley rat, as compared to the Fischer 344 rat (approximately 5% and 88%, respectively at 24-months) (Clegg et al. 1997). The control incidence of these tumors reported by Belpoggi et al. (1995) is consistent with levels typically observed in the Sprague-Dawley strain. The control incidence observed by Chun et al. (1992) was lower than that typically observed in the Fischer 344 strain, but within the range reported for aged male rats of this strain (Bird et al. 1997, Haseman and Arnold 1990). The lower spontaneous Leydig cell tumor incidence observed in the Chun et al. (1992) study is likely to have facilitated the detection of the dose-dependent increase in Leydig cell tumors in MTBE-treated males, despite the early termination of the mid- and high-dose groups.

The tumors observed in male Fischer 344 rat kidney tissues (Chun et al. 1992, Bird et al. 1997) were diagnosed as renal tubular adenomas and carcinomas. These two tumor phenotypes are generally considered to be related in origin, with the possibility that adenomas may progress to carcinomas (Borghoff et al. 1996b). Therefore, they are normally aggregated for carcinogen identification and risk assessment purposes (U.S. EPA 1991). The possibility that the male rat-specific α_{2u} -globulin nephropathy plays a significant role in the pathogenesis of MTBE rat kidney tumors has been investigated, and reported to be unlikely (NSTC 1997, U.S. EPA 1997a). The data indicate that MTBE induces only mild accumulation of α_{2u} -globulin and mild or partial expression of α_{2u} -globulin associated nephropathy in male rats, while clearly exacerbating the expression of non- α_{2u} -globulin rat nephropathy in both males and females (NSTC 1997). Support for this conclusion includes the observation that a dose-dependent increase in mortality from chronic progressive nephropathy was observed in male rats at all dose levels, and in females at the mid- and high-dose levels in the rat inhalation bioassay (Bird et al. 1997). Observed microscopic kidney changes included increases in the severity of mineralization and interstitial fibrosis in all treated males, and increases in mild to moderate glomerulosclerosis, interstitial fibrosis, and tubular proteinosis in mid- and high-dose females (Chun et al. 1992). In addition, a rare renal tubular tumor was observed in one MTBE-treated female rat (Chun et al. 1992). In a separate analysis of a 13-week inhalation exposure study of male rats conducted at the Bushy Run Research Center laboratory, Swenberg and Dietrich (1991) measured the levels of α_{2u} -globulin associated with hyaline droplets in MTBE-treated and control kidneys. Although a slight increase in renal cortex staining for α_{2u} -globulin was observed in MTBE-treated animals, as compared with controls, there was no relationship between the level of α_{2u} -globulin staining and the dose of MTBE received (U.S. EPA 1997c, Swenberg and Dietrich 1991). In a study by Lington et al. (1997), inhalation of 4,000 and 8,000 ppm MTBE for 13 weeks resulted in a moderate increase in the size of hyaline droplets in male rat kidney, but no MTBE-associated increase in the area or intensity of α_{2u} -globulin immunostaining was observed, as reported by Bird et al. (1997). In a four-week inhalation study, exposure to 3,000 and 8,000 ppm MTBE increased the levels of protein accumulated in male rat kidney tubule epithelial cells, but not the levels of α_{2u} -globulin, as compared with controls (Bird et al. 1997).

The tumors observed by Burleigh-Flayer et al. (1992) and Bird et al. (1997) in mouse liver were diagnosed as hepatocellular adenomas and carcinomas. These two tumor phenotypes are generally considered to be related in origin, with the possibility that adenomas may progress to carcinomas. They are normally therefore aggregated for carcinogen identification and risk assessment purposes. The sensitivity of the study to detect treatment-related tumors, especially in the low- and mid-dose groups, may have been compromised by the less-than-lifetime length of the study (18 months).

Mechanism

The mechanism(s) by which MTBE induces tumors at multiple sites in rats and mice is unknown at this time. It is unclear whether MTBE itself plays a direct role in the observed tumorigenesis, or whether metabolism to one or more active metabolites is required. The two major metabolites of MTBE, HCHO (Kerns et al. 1983, Sellakumar et al. 1985, Til et al. 1989, Woutersen et al. 1989) and TBA (NTP 1995), have both been shown to possess tumorigenic activity in animal studies. Interestingly, there is a commonality of tumor sites observed for MTBE, HCHO, and TBA. Leukemias were observed in male and female Sprague-Dawley rats administered HCHO in drinking water (Soffritti et al. 1989), and renal tubular cell adenomas and carcinomas were observed in male Fischer 344 rats administered TBA in drinking water (NTP 1995, Cirvello et al. 1995). IARC (1995) concluded that the evidence on the carcinogenicity of HCHO was sufficient in animals and limited in humans, and classified the agent in Group 2A probably carcinogenic to humans. NTP (1995) in reviewing the results of two-year drinking water studies with TBA concluded that "there was 'some' evidence of carcinogenic activity of TBA in male Fischer 344/N rats based on increased incidences of renal tubule adenoma or carcinoma (combined)". It is presently unknown whether the nature or degree of MTBE metabolism is tissue- or sex-specific, or whether there is any relationship between the site of metabolism and target tumor sites. Comparison of the target tumor sites in rats administered MTBE by two different routes of administration is inherently limited by the use of different rat strains in these studies; however, these findings suggest that route-specific distribution and metabolism of MTBE may be of importance in the development of some (e.g., leukemias and lymphomas, renal tumors), but not all treatment-associated tumors (e.g., testicular tumors). It has also been suggested that sex-specific differences in metabolism may underlie the development of leukemias and lymphomas in female, but not male rats (Belpoggi et al. 1995, 1997). This hypothesis remains untested, however.

MTBE was negative in a number of genotoxicity assays as noted in the section on genetic toxicity in this document and by ATSDR (1996), testing positive only in the activated mouse lymphoma forward mutation assay (ARCO 1980, Mackerer et al. 1996). The MTBE metabolite TBA was not mutagenic in either the Salmonella assay (Zeiger et al. 1987) or the mouse lymphoma assay (McGregor et al. 1988). HCHO is genotoxic, testing positive in numerous assay systems (IARC 1995). Data on HCHO-related genotoxicity in MTBE tumorigenesis are too limited to draw any conclusions at this time. Studies conducted in freshly isolated mouse hepatocytes from female CD-1 mice (Casanova and Heck 1997) did not find any dose-related increase in HCHO-associated DNA-protein cross-links or RNA-HCHO adducts following MTBE-treatment. Similar results were obtained with freshly isolated hepatocytes from male B6C3F1 mice and male Fischer 344 rats (Casanova and Heck 1997). These data suggest that HCHO is not the active species responsible for MTBE liver tumorigenesis in the mouse. In studies using the mouse lymphoma assay, however, HCHO has been implicated as the active

species responsible for MTBE's mutagenic activity (Mackerer et al. 1996). DNA-protein cross-link data and RNA-HCHO adduct data are not available for the other tumor sites noted after MTBE exposure in laboratory animals.

Several hypotheses have been put forward suggesting that MTBE may act via a variety of nongenotoxic mechanisms, such as the involvement of endocrine modulation in mouse liver and rat testicular tumorigenesis (Bird et al. 1997, Moser et al. 1996b) and α_{2u} -globulin nephropathy in male rat kidney tumorigenesis (Poet and Borghoff 1997a, 1997b, Prescott-Mathews et al. 1997a). While MTBE exposure of the mouse is associated with various endocrine-related tissue and cellular responses (see the section on developmental and reproductive toxicity in this document), the available data are insufficient to support an endocrine-mediated mode of action for MTBE-associated liver (Moser et al. 1996a, 1996b, Moser et al. 1998, Okahara et al. 1998) or testicular tumors (Day et al. 1998) at this time.

Data which suggest that α_{2u} -globulin nephropathy may be involved in MTBE kidney tumorigenesis include the following:

- A mild to moderate increase in the number and size of hyaline droplets in the renal proximal tubule cells of MTBE-treated male rats has been observed.
 - ◊ In a 10-day inhalation study, MTBE increased the number of protein droplets within the renal proximal tubules of male rats with a statistically significant concentration-related positive trend (Prescott-Mathews et al. 1997a).
 - ◊ In a 14-day gavage study, MTBE increased the formation of hyaline droplets in male rat kidney proximal tubular epithelial cells at the highest dose tested (Robinson et al. 1990).
 - ◊ In a 13-week inhalation study, MTBE slightly increased hyaline droplet formation in male rat kidney (Swenberg and Dietrich 1991).
 - ◊ In another 13-week inhalation study, MTBE slightly increased the size of hyaline droplets in male rat kidney (Bird et al. 1997 reporting on findings of Lington et al. 1997).
 - ◊ In a 90-day gavage study, MTBE slightly increased the number of hyaline droplets in male rat kidney proximal tubular epithelial cells (Robinson et al. 1990).
- Protein in the renal proximal tubule cells of MTBE-treated male rats stains weakly for α_{2u} -globulin.
 - ◊ In a 13-week inhalation study, MTBE slightly increased hyaline droplet formation and staining for α_{2u} -globulin in male rat kidney but these increases were not dose-dependent (Swenberg and Dietrich 1991).
 - ◊ In a 10-day inhalation study, no increase in α_{2u} -globulin staining could be detected in MTBE-treated male rat kidney by immunohistochemical staining (Prescott-Mathews et al. 1997a).
- Using an ELISA-based method, a mild dose-dependent increase in α_{2u} -globulin-immunoreactivity (approximately 150 μg α_{2u} -globulin/mg total protein in controls versus 200 μg α_{2u} -globulin/mg total protein in the high-dose animals) has been observed in rat kidney cytosol of male rats exposed to MTBE by inhalation for 10 days (Prescott-Mathews et al. 1997a).
- MTBE binds weakly to α_{2u} -globulin in vitro. Using a kidney homogenate system, only a very weak interaction between MTBE and male rat renal proteins was detected (Prescott-

Mathews et al. 1997b). This interaction did not survive dialysis or anion exchange chromatography (Prescott-Mathews et al. 1997b).

- A dose-dependent increase in cell proliferation has been observed in the renal cortex of male rats exposed to MTBE by inhalation for 10 days (Prescott-Mathews et al. 1997a).
- Agents which are thought to induce renal tubular tumors via an α_{2u} -globulin-mediated mechanism are nongenotoxic. MTBE has demonstrated little or no genotoxicity in vitro or in vivo.

Data which argue against a significant role for α_{2u} -globulin nephropathy in MTBE kidney tumorigenesis include the following:

- Male rat specificity for nephropathy and renal tumorigenicity has not been observed.
 - ◊ In a two-year inhalation study, MTBE exacerbated chronic progressive nephropathy and increased mortality associated with chronic progressive nephropathy in a dose-dependent manner in both in female and male rats (Chun et al. 1992, Bird et al. 1997).
 - ◊ A rare kidney tumor was observed in one MTBE-treated female rat in the two-year inhalation study (Chun et al. 1992, Bird et al. 1997).
- A clear exposure-related increase in staining for α_{2u} -globulin, an effect typical of classical α_{2u} -globulin nephropathy-inducing agents, has not been observed in male rats treated with MTBE.
 - ◊ In a 13-week inhalation study, MTBE slightly increased hyaline droplet formation and staining for α_{2u} -globulin in male rat kidney but these increases were not dose-dependent (Swenberg and Dietrich 1991).
 - ◊ In another 13-week inhalation study, MTBE slightly increased the size of hyaline droplets in male rat kidney, but no increase in the area or intensity of α_{2u} -globulin staining was observed (Bird et al. 1997 reporting on findings of Lington et al. 1997).
 - ◊ In a four-week inhalation study, MTBE slightly increased protein accumulation in male rat kidney, but did not increase α_{2u} -globulin levels (Bird et al. 1997).
 - ◊ In a 10-day inhalation study, no increase in α_{2u} -globulin staining could be detected in MTBE-treated male rat kidney by immunohistochemical staining, but using a more sensitive ELISA-based assay a mild increase in the concentration of α_{2u} -globulin (approximately 150 μg α_{2u} -globulin/mg total protein in controls versus 200 μg α_{2u} -globulin/mg total protein in the high-dose animals) was observed (Prescott-Mathews et al. 1997a). This small increase is in contrast to the marked increase seen with classical α_{2u} -globulin nephropathy-inducing agents, such as 2,2,4-trimethylpentane (approximately 200 μg α_{2u} -globulin/mg total protein in controls versus 550 μg α_{2u} -globulin/mg total protein in treated animals) (Prescott-Mathews et al. 1997a).
- α_{2u} -globulin-positive proteinaceous casts, another effect typical of classical α_{2u} -globulin nephropathy-inducing agents, were not seen at the junction of the proximal tubules and the thin loop of Henle in a 13-week inhalation study (Swenberg and Dietrich 1991, U.S. EPA 1997c).
- Investigators have been unable to detect the binding of MTBE to α_{2u} -globulin or male rat renal proteins in vivo (Prescott-Mathews et al. 1997b), and only a very weak interaction between MTBE and male rat renal proteins has been detected in vitro, using a kidney homogenate system (Prescott-Mathews et al. 1997b). This interaction did not survive dialysis or anion exchange chromatography (Prescott-Mathews et al. 1997b), in contrast to

observations with classical α_{2u} -globulin nephropathy-inducing agents, where typically 20-40% of bound ligand is retained after dialysis (NSTC 1997).

The available data on renal tumorigenesis indicate that MTBE induces only mild accumulation of α_{2u} -globulin and mild or partial expression of α_{2u} -globulin associated nephropathy in male rats, while clearly exacerbating the expression of non- α_{2u} -globulin rat nephropathy in both males and females (NSTC 1997). The U.S. EPA (1991) established three criteria for causation of an α_{2u} -globulin effect:

- (1) increased number and size of hyaline droplets in renal proximal tubule cells of treated male rats;
- (2) accumulating protein in the hyaline droplets is α_{2u} -globulin; and
- (3) additional aspects of the pathological sequence of lesions associated with α_{2u} -globulin nephropathy are present.

If the response is mild all of the typical lesions may not be observed, however, some elements consistent with the pathological sequence must be demonstrated to be present. Evaluation of the available data indicates that the first criterion has been satisfied, but not the second or third (NSTC 1997, U.S. EPA 1997a). Thus, α_{2u} -globulin nephropathy does not appear to play a significant role in MTBE kidney tumorigenesis.

Summary of the Evidence

Carcinogenicity of MTBE has been observed in both sexes of the rat in a lifetime gavage study (Belpoggi et al. 1995, 1997), in male rats of a different strain in a 24-month inhalation study (Chun et al. 1992, Bird et al. 1997), and in male and female mice in an 18-month inhalation study (Burleigh-Flayer et al. 1992, Bird et al. 1997). Statistically significant increases in Leydig interstitial cell tumors of the testes were observed in two different strains of rats by two separate routes of administration. Other statistically significant increases in the rat were leukemias and lymphomas (combined) in females and renal tubular tumors in males. Statistically significant increases in hepatocellular adenomas and carcinomas (combined) were observed in both sexes of the mouse. MTBE has demonstrated little or no genotoxicity in vitro or in vivo. The mechanism by which MTBE induces tumors at multiple sites in animals remains unknown (Menear 1995, 1997a, 1997b). Additional supporting evidence is provided by the carcinogenic activity of HCHO and TBA, two primary metabolites of MTBE, which share target tumor sites in common with MTBE.

Conclusion

Based on the information reviewed in the preparation of this document, there is evidence for the carcinogenicity of MTBE at multiple sites in both sexes of the rat and the mouse; MTBE is a multi-species, multi-strain, multi-sex, and multi-route carcinogen. Positive animal carcinogenicity data for HCHO and TBA, metabolites of MTBE, provide support for this conclusion.

Ecotoxicity

Concern has been raised about the effects of MTBE in water on plants, animals and ecosystems. Rowe et al. (1997) summarized aquatic toxicity information and water quality criteria for VOCs

including MTBE being monitored in the NAWQA Program by the USGS. The species tested so far for toxic effects of MTBE have high thresholds in the parts per million (ppm or mg/L) range indicating that MTBE has limited acute and chronic toxicity for aquatic species (Mancini 1997, Stubblefield et al. 1997). Acute studies generated MTBE LC₅₀ values with the freshwater green algae of 184 ppm, the freshwater Ceriodaphnia fleas of 348 ppm, the freshwater Daphnia water fleas of 542 and 681 ppm, the freshwater fathead minnows of 672, 706, 929 and 979 ppm, the freshwater rainbow trouts of 887 and 1237 ppm, the freshwater tadpoles of 2,500 ppm, the marine mysid shrimps of 44 and 136 ppm, the marine inland silverside of 574 ppm, the marine bleak of > 1,000 ppm, the marine copepod of > 1,000 ppm, and the marine sheepshead minnows of > 2,500 ppm.

Toxicity of MTBE to *Daphnia magna* and *Photobacterium phosphoreum* was reported (Gupta and Lin 1995). A recent laboratory toxicity study with three unicellular algae suggests that the dissolved MTBE may alter algal community composition in the natural environment (Rousch and Sommerfeld 1998). Research by the API and others on ecological hazards of MTBE exposure is continuing. Because of the large amount of MTBE usage in California, high water and lipid solubility of MTBE, and lack of information on toxic effects of long-term exposure to low doses of MTBE (e.g., reproductive impairment in plants or animals), Cal/EPA (1998) has a continuing interest in reviewing current and proposed research to fill in these data gaps.

TOXICOLOGICAL EFFECTS IN HUMANS

No studies were located regarding toxic effects of MTBE in humans following ingestion or skin contact. No studies were located regarding toxic effects of ingested or inhaled or skin-contacted MTBE in drinking water in humans. No studies were located regarding acute effects, subchronic effects, chronic effects, death, systemic effects including respiratory, gastrointestinal, cardiovascular, hematological, musculoskeletal, hepatic, renal, endocrine, dermal, ocular, or body weight effects, immunological or lymphoreticular effects, neurological effects, developmental or reproductive effects, genotoxic effects, or cancer in humans after oral exposure to MTBE alone (ATSDR 1996).

No epidemiological study data on long-term effects and the carcinogenic effects of human exposure to MTBE were found in an earlier search by ATSDR (1996) or more recently by OEHHA. Neither U.S. EPA nor IARC has classified MTBE with respect to potential human carcinogenicity based on animal studies. The NSTC (1997) report concluded that "there is sufficient evidence to indicate that MTBE is an animal carcinogen and to regard MTBE as having a human hazard potential." Nevertheless, health complaints from the public have raised the concern of federal and state health agencies (Begley 1994, Begley and Rotman 1993, CDC 1993a, 1993b, 1993c, Drew 1995, Joseph 1995, Mehlman 1995, 1996).

No studies were located regarding death, cardiovascular effects, hematological effects, musculoskeletal effects, hepatic effects, renal effects, endocrine effects, body weight effects, developmental and reproductive effects, genotoxic effects, or cancer in humans after inhalation exposure to MTBE. No studies were located regarding death, respiratory effects, gastrointestinal effects, cardiovascular effects, hematological effects, musculoskeletal effects, hepatic effects, renal effects, endocrine effects, body weight effects, immunological or lymphoreticular effects, neurological effects, developmental and reproductive effects, genotoxic effects, or cancer in humans after dermal exposure to MTBE (ATSDR 1996).

Acute Toxicity

A recent literature review (Borak et al. 1998) summarizes the exposure to MTBE and acute human health effects including nine epidemiological studies, ten industrial hygiene studies, and 12 clinical studies. No studies were located regarding acute toxic effects of ingested or skin-contacted MTBE in humans. There are very limited data on the acute toxicity of MTBE in humans through inhalation exposure. Several studies undertaken over the past four to five years were unable to find any correlation between reported acute health effects and MTBE exposures experienced by the general public, mainly through inhalation, from the use of MTBE in gasoline (ATSDR 1996, Balter 1997, McCoy et al. 1995, NSTC 1996, 1997, U.S. EPA 1997a).

Ingestion of gasoline-MTBE mixtures may result in aspiration and pneumonitis. Complaints of acute effects from exposure to oxygenates such as MTBE in gasoline, mainly via inhalation, have been received by health authorities (Fiedler et al. 1994, McCoy et al. 1995, Raabe 1993). However, the limited epidemiological studies that have been conducted to date have not demonstrated a causal association between acute effects and inhalation exposure in a relatively small population (ATSDR 1996). Three human volunteer inhalation studies did not show increased symptoms among healthy adults (Cain et al. 1996, Johanson et al. 1995, Prah et al. 1994).

In 1993, the J. B. Pierce Laboratory of Yale University (Cain et al. 1996) and U.S. EPA (Prah et al. 1994), in two separate studies, exposed individuals to clean air and air mixed with MTBE. In cases where 37 or 43 human volunteers were exposed to low levels of MTBE in air (1.39 or 1.7 ppm) for one hour, there was no significant increase in symptoms of eye, nasal, or pulmonary irritation when the results for periods of exposure to MTBE were compared to results from exposure to ambient air. There were also no significant effects on mood or in the results from several performance-based neurobehavioral tests. In both studies, the females ranked the general quality of the air containing MTBE lower than the control atmosphere. However, in the study by Cain et al. (1996), where the subjects were also exposed to an atmosphere containing a total of 7.1 ppm mixture of 17 VOCs that are frequent air contaminants in areas around gasoline stations, the air quality of the MTBE-containing atmosphere ranked higher than that with the VOC mixture. No increase in acute symptoms was observed in individuals exposed to MTBE at concentrations that would be encountered while refueling a car.

The studies by Hakkola (1994), Hakkola et al. (1996, 1997) and White et al. (1995) compared the effects in two groups exposed to different concentrations of MTBE from treated gasoline because of their lifestyles. The moderately-exposed individuals either drove a gasoline delivery truck, worked in a gasoline station, or worked on car repairs. The minimally-exposed individuals merely used a gasoline-powered vehicle to go to and from work or as part of their job. In the study by White et al. (1995), the odds ratio was 8.9 (95% confidence interval = 1.2 to 75.6) for the reporting of one or more symptoms when 11 individuals with blood MTBE levels of $> 2.4 \mu\text{g/L}$ were compared with 33 individuals with lower levels. The odds ratio increased to 21 (95% confidence interval = 1.8 to 539) when commuters were excluded from the population studied and eight workers with blood levels of $> 3.8 \mu\text{g/L}$ were compared to 22 individuals with lower blood MTBE levels. All individuals lived and worked in the area around Stamford, Connecticut.

In a series of studies conducted in Finland where the gasoline contains 10% MTBE, Hakkola (1994) first evaluated neuropsychological symptoms among 61 male tanker drivers with exposure to organic solvents at work, and found that the differences between the exposed group and the two control groups (56 males with occasional exposure at work and 31 male with no exposure) were

not statistically significant. Hakkola et al. (1996) again found that there were no statistically significant differences between the signs and symptoms reported by 101 drivers of tanker trucks and 100 milk truck drivers. Blood concentrations of MTBE or its metabolites were not monitored. However, the latest Hakkola et al. (1997) study compared symptoms and moods among 101 road tanker drivers with 100 milk delivery drivers. The tanker drivers with long exposure to gasoline during the work week reported significantly higher changes in fatigue scores than drivers with short exposure, and 20% of tanker drivers reported acute symptoms connected to MTBE exposure.

In the winter of 1992, the state of Alaska began using 15% MTBE in wintertime oxygenated gasoline as part of the federal requirements to reduce emissions of CO in Fairbanks and Anchorage. There were reports of headaches, dizziness, nausea, and spaciness after refueling and/or working around oxygenated gasoline (Smith and Duffy 1995). The Centers for Disease Control (CDC), U.S. EPA, and the state of Alaska investigated these complaints but were unable to associate them with MTBE exposure. Instead, it was suggested that the increase in price of the new federal RFG, the odor of MTBE, and the harsh climate of Alaska resulted in some of the public associating changes in fuel with the reported symptoms. The state is now using ethanol in its gasoline during the winter (Beller et al. 1992, Chandler and Middaugh 1992, CDC 1993a). Gordian et al. (1995) reported no increase in claims for respiratory illness in Anchorage or Fairbanks after introduction of MTBE in Alaska.

A study in Alaska (Moolenaar et al. 1994) compared effects and blood levels of MTBE from a time period when oxygenated fuels were in use (Phase I) to those after the oxygenated fuels use had stopped (Phase II). The subjects were volunteers who were occupationally exposed to motor vehicle exhaust or gasoline fumes. Eighteen workers participated in Phase I and 22 in Phase II. Twelve of those that participated in Phase I of the study also participated in Phase II. A questionnaire was used to gather information on signs and symptoms and blood samples were collected for measurement of MTBE at the beginning and end of a typical work day. In Phase I, the median post-shift MTBE level was higher than the pre-shift value (1.80 versus 1.15 ppb). During Phase II, the values were more comparable (0.25 versus 0.21 ppb). Median post-shift blood measurements of TBA were higher during Phase I than in Phase II (5.6 versus 3.9 ppb). Signs and symptoms that could be associated with MTBE exposure were reported more frequently during Phase I than Phase II (Moolenaar et al. 1994). During Phase I, 50% or more of the participants reported headaches, eye irritations, and nose and throat irritations. Reporting of these symptoms occurred in less than 10% of the participants during Phase II. However, it is difficult to evaluate if psychosomatic factors and individual sensitivity had influenced these results. The volunteers may have chosen to participate because of their sensitivity to contaminants in the atmosphere. A follow-up survey of workers exposed to oxygenated fuel in Fairbanks, Alaska (Moolenaar et al. 1997) detected higher blood benzene concentrations in mechanics than drivers and other garage workers.

Milwaukee, Wisconsin began to use MTBE in its gasoline as part of the federal RFG program in November 1994. Similar health complaints, as voiced in Alaska (Beller et al. 1992), were registered in Wisconsin. U.S. EPA, the Wisconsin Department of Health, CDC, and the University of Wisconsin investigated complaints from approximately 1,500 people. They wrote two reports (May and September 1995) and concluded that they could find no relationship between reported health effects and MTBE exposure. It was suggested that the odor of MTBE, increase in price of wintertime gasoline, and negative media coverage were responsible for the reports of health problems associated with exposure to gasoline (Anderson et al. 1995).

National Institute for Working Life in Sweden (Nihlen et al. 1998a, 1998b) assessed acute effects up to the Swedish occupational exposure limit value both with objective measurements and questionnaires. The healthy male volunteers were exposed to MTBE vapor for two hours at five, 25, and 50 ppm during light physical work. In the questionnaire, only the ratings of solvent smell increased up to 50% of the scale as the volunteers entered the chamber and declined slowly with time. No ocular effects were observed.

Nasal airway resistance blockage index increased but was not related to exposure levels. Decreased nasal volume was seen but with no dose-effect relationship. The authors concluded no or minimal acute effects of MTBE vapor upon short-term exposure at these relatively high levels.

An interview questionnaire study (Fiedler et al. 1994) was conducted to assess exposure and the symptomatic responses of individuals with multiple chemical sensitivities (MCS) while using gasoline products with MTBE, and to compare their responses to individuals with chronic fatigue syndrome (CFS) which can not contribute to exposure to chemicals, and to compare with normal controls. Fourteen MCS, five CFS, and six normal control subjects of comparable age, education, gender, and ethnicity completed several structured interview and assessment sessions. It was concluded that while the sample was limited, MTBE symptoms were not uniquely associated with chemical sensitivity or with situations where MTBE was more prevalent.

Several additional major literature reviews on the acute health effects of MTBE have been conducted. Reviews from studies in Connecticut (CDC 1993b, White et al. 1995), Montana (MCCHD 1993), New Jersey (Mohr et al. 1994), New York (CDC 1993c), Illinois and Wisconsin (Anderson et al. 1995) and the HEI (1996) could find no evidence linking acute health effects with exposure to MTBE from gasoline use. In 1993, the Environmental and Occupational Health Sciences Institute (EOHSI) surveyed New Jersey garage workers and service station attendants, some of whom were exposed to MTBE, and some of whom were not. No significant differences in the frequency of reported symptoms were observed between the two groups (Hartle 1993, Mohr et al. 1994). EOHSI is conducting a study on individuals who have reported sensitivity to MTBE and were recruited from the "Oxybuster" group in New Jersey. The Oxybuster group is a citizens' group which claims their members experience acute health effects from breathing MTBE (Joseph 1995). Those individuals will be exposed to gasoline with and without MTBE. Results are expected later in 1998.

In response to the negative publicity associated with the use of federal wintertime oxygenated fuel, the White House OSTP through the NSTC in September 1995 directed federal agencies to review fuel economy and engine performance issues, water quality, air quality benefits, and health effects of oxygenates in fuel with a final report issued in June 1997. NSTC (1997) concluded that with the information collected to date there was no evidence that MTBE is causing increases in acute symptoms or illnesses at concentrations experienced by the general population, but anecdotal reports of acute health symptoms among some individuals cannot yet be explained or dismissed. NSTC also recommended that greater attention should be given to the potential for increased symptom reporting among workers exposed to high concentrations of oxygenated gasoline containing MTBE. Regarding the issue of acute sensitivity to MTBE, NRC which peer-reviewed an earlier draft of the NSTC report, concluded that there was no reason to believe that some people have extreme sensitivity to MTBE. The final NSTC report concluded that "an examination of possible predisposing factors might be useful to better understand the occurrence of various symptoms in the general public following exposure to MTBE-containing gasoline."

MTBE has had a limited use as a therapeutic drug for dissolving cholesterol gallbladder stones (ATSDR 1996, HSDB 1997). Perfusion of MTBE through the bile duct and gallbladder by a

percutaneous transhepatic catheter under local anesthesia was once used as a medical treatment to dissolve gallstones as an alternative to surgery (Diaz et al. 1992, Edison et al. 1993, Lin et al. 1994). Another solvent, ethyl propionate, has been suggested to be preferable to MTBE in this investigational procedure (Hofmann et al. 1997). Acute exposure of humans to MTBE has occurred via injection through the catheter into the gallbladder. During this procedure, some of the MTBE enters the blood stream and is distributed systemically. Side effects reported in patients treated by this procedure included nausea, vomiting, coughing, bronchitis, sleepiness, sedation, perspiration, bradycardia (slow heart beat), elevation of liver enzymes, apnea, CNS depression, and respiratory depression (Allen et al. 1985, Juliani et al. 1985, Wyngaarden 1986). A case of acute renal failure was also reported (Ponchon et al. 1988). These signs cannot be attributed totally to MTBE because of the confounding effects of anesthesia and the infusion process itself. Borak et al. (1998) reviewed 12 dissolution studies and reported that the peak MTBE blood levels averaged 40,000 µg/L in one study and ranged up to 10,000 µg/L in another study.

Immunotoxicity

There are very limited human studies available on the immunotoxicity of MTBE-added fuels through inhalation or MTBE-contaminated water. Duffy (1994) concluded that single day exposures to oxyfuel and its combustion products did not show an immediate effect on the immune system as measured by serum plasma interleukin 6 (IL-6) levels. In this study, blood samples from 22 individuals exposed to auto emissions derived from oxyfuel were analyzed for effects on the immune system by monitoring IL-6 levels at the beginning and at the end of the eight-hour workday during a four-week period in late November and early December 1992 (Duffy 1994).

Vojdani et al. (1997b) reported the detection of MTBE antibodies in seven out of 24 gasoline station attendants (six females and 18 males ranging in age from 21 to 58 years) who were employed for more than two years in service stations, and none out of the 12 healthy control subjects (four females and eight males 24 to 60 years of age). The results indicated that these IgG and IgM antibodies were produced against the methyl or tert-butyl group of MTBE. They also indicated that the immune reactions to MTBE occurred through hapten carrier reactions which could be related to airborne exposures to TBF. However, the antibody response did not correlate with claimed symptoms.

The same group (Mordechai et al. 1997, Vojdani et al. 1997a) also reported reversible but statistically significant increased rates of abnormal apoptosis (programmed cell death) and cell cycle progression in peripheral blood lymphocytes in 20 Southern California residents exposed to MTBE and benzene contaminated water as compared to ten healthy human controls. Apoptosis is an organism's way of maintaining healthy cell populations, the process can lead to the development of disease if it is unduly suppressed or stimulated. For example, cancer may be the result of a failure in the apoptotic process, in which mutant cells are allowed to proliferate freely rather than being recognized as damaged and destroyed.

Neurotoxicity

Burbacher (1993) reviewed gasoline and its constituents as neuroactive substances and recommended future studies to focus on examining the dose-response relationship between chronic low-level exposure and subtle toxic effects in CNS functions. The results from human

studies of neurological effects, e.g. headache, dizziness, disorientation, fatigue, emotional distress, gastrointestinal problems, e.g. nausea or diarrhea, and symptoms of respiratory irritation in individuals exposed to MTBE vapors through MTBE-containing fuels are inconclusive (Hakkola et al. 1996, Hakkola and Saarinen 1996, Moolenaar et al. 1994, White et al. 1995). The three studies cited were different in their design and utilized slightly different parameters for monitoring effects. All studies evaluated exposure to an MTBE-gasoline mixture and not MTBE alone.

However, in the most recent study by Hakkola et al. (1997) comparing neuropsychological symptoms and moods among 101 road tanker drivers from three Finnish oil companies with 100 milk delivery drivers from two milk companies, the tanker drivers with long exposure to gasoline during the work week reported significantly higher changes in fatigue scores than drivers with short exposure, and 20% of tanker drivers reported acute symptoms of headache, dizziness, nausea, dyspnoea, and irritation of saliva excretion. These symptoms have been connected to MTBE exposure. The authors suggested that exposure to MTBE during the work week could be reason for acute symptoms among the tanker drivers in this study.

dose-response assessment

INTERNAL DOSE ESTIMATION

Due to the lack of a clear mode of action of TBA or other MTBE metabolites in MTBE-induced carcinogenesis in experimental animals, OEHHA has necessarily had to treat the parent compound MTBE as the cause of the observed effects in animal studies for the purpose of determining dose metrics. In order to estimate internal doses of MTBE, in addition to simple continuous applied doses, a simplified PBPK model was employed. This model is based on both the Borghoff et al. (1996a) model, in that it has five compartments for MTBE and five compartments for TBA, and the Rao and Ginsberg (1997) model with its MTBE metabolic parameters and slowly perfused compartment/blood partition coefficient for TBA. The PBPK model employs compartments loosely representing "Fat, Liver, Kidneys, Muscle, and rapidly perfused tissues termed as Vessel Rich Group (VRG)". The model's fundamental structure is based on that developed by Hattis et al. (1986) for perchloroethylene and was formulated in Stella® software (ithink® v. 3.0.6b for the Power Macintosh, High Performance Systems Inc., Hanover, New Hampshire 03755). The model units for the whole animal are moles, L, moles/L, hour, moles/hour, L/hour, and ppm in alveolar air. Simulations of up to 32 hours were run at approximately 1,000 steps/simulated hour, using the Runge-Kutta 4 computation method on a Power Macintosh 7100/80. The model parameters were obtained from Borghoff et al. (1996a) or Rao and Ginsberg (1997) and are listed in Table 10. In addition to simulations of the pharmacokinetic data of Miller et al. (1997) with a model 0.22 kg rat, simulations of cancer bioassay doses were conducted assuming 0.35 kg for female and 0.5 kg for male lifetime average body weights. Physiological and metabolic parameters were scaled to these body weights as described in Borghoff et al. (1996a).

Table 10. Parameters Used in the PBPK Model Simulations for MTBE and TBA

Parameter	Female rat	Male rat	Source
Body weight (kg)	0.35	0.5	Estimated from Belpoggi et al. 1995
Compartment volumes (L)			
Liver	0.014	0.020	Borghoff et al. 1996a
Kidney	0.00245	0.0035	Borghoff et al. 1996a
Muscle	0.2625	0.375	Borghoff et al. 1996a
Fat	0.0245	0.035	Borghoff et al. 1996a
Vessel Rich Group (VRG)	0.01505	0.0215	Borghoff et al. 1996a
Flows (L/hour)			
Alveolar			
ventilation	6.4	8.32	Borghoff et al. 1996a
Cardiac output	6.4	8.32	Borghoff et al. 1996a
Liver	1.6	2.88	Borghoff et al. 1996a
Kidney	1.6	2.88	Borghoff et al. 1996a
Muscle	0.96	1.248	Borghoff et al. 1996a
Fat	0.576	0.7488	Borghoff et al. 1996a
VRG	1.664	2.1632	Borghoff et al. 1996a
Partition coefficients (MTBE)			
Blood/Air	11.5	11.5	Borghoff et al. 1996a
Liver/Blood	1.1826	1.1826	Borghoff et al. 1996a
Kidney/Blood	3.113	3.113	Borghoff et al. 1996a
Muscle/Blood	0.565	0.565	Borghoff et al. 1996a
Fat/Blood	10.05	10.05	Borghoff et al. 1996a
VRG/Blood	3.113	3.113	Borghoff et al. 1996a
Partition coefficients (TBA)			
Blood/Air	481-75	481-75	Borghoff et al. 1996a*
Liver/Blood	0.8316	0.8316	Borghoff et al. 1996a
Kidney/Blood	1.1289	1.1289	Borghoff et al. 1996a
Muscle/Blood	0.4	0.4	Rao & Ginsberg 1997
Fat/Blood	0.3971	0.3971	Borghoff et al. 1996a
VRG/Blood	1.1289	1.1289	Borghoff et al. 1996a
Metabolism (MTBE)			
Vmax ₁ (mole/hour)	2.05 × 10 ⁻⁶	2.66 × 10 ⁻⁶	Rao & Ginsberg 1997
Vmax ₂ (mole/hour)	2.27 × 10 ⁻⁴	2.94 × 10 ⁻⁴	Rao & Ginsberg 1997
Km ₁ (M)	2.27 × 10 ⁻⁶	2.27 × 10 ⁻⁶	Rao & Ginsberg 1997
Km ₂ (M)	1.25 × 10 ⁻³	1.25 × 10 ⁻³	Rao & Ginsberg 1997
Metabolism (TBA)			
Vmax (mole/hour)	2.46 × 10 ⁻⁵	3.21 × 10 ⁻⁵	Rao & Ginsberg 1997
Km (M)	3.79 × 10 ⁻⁴	3.79 × 10 ⁻⁴	Rao & Ginsberg 1997
GI absorption (hour ⁻¹)	0.8	0.8	Model assumption

* Note: see text

DRAFT

The PBPK model simulation results for oral exposures to MTBE are summarized in Table 11. The italic boldface values are observed experimental data from Miller et al. (1997). The simulated or predicted values for 0.215 kg, 0.35 kg female, and 0.5 kg male rats are shown in normal type. In general, better predictions were obtained for MTBE than for TBA both for maximum blood concentration and the area under the blood concentration x time curve, or AUC.

Adequate simulation of TBA blood kinetics became increasingly difficult with increased body size and lower TBA blood-air partition coefficients of 150 and 75 had to be employed to achieve stable simulations. In all cases MTBE doses were cleared within 24 hours and there was no need for multi-day simulations to estimate an average daily MTBE AUC for the bioassays. In all cases MTBE AUC was linear with applied dose for a particular body size.

Table 11. Comparison of PBPK Predictions with Experimental Data from Oral MTBE Administrations*

Oral dose/ Body weight	MTBE mM Cmax	TBA mM Cmax	MTBE AUC mM hour	TBA AUC mM hour	Blood:Air MTBE/TBA
40 mg/kg					
0.215 kg rat	0.068	0.176	0.150	0.863	11.5/481
<i>Observed</i>					
<i>Frat</i>	<i>0.127</i>	<i>0.12</i>	<i>0.142</i>	<i>0.495</i>	
<i>Mrat</i>	<i>0.195</i>	<i>0.135</i>	<i>0.193</i>	<i>0.526</i>	
250 mg/kg					
0.35 kg Frat	0.527	0.974	1.03	6.3	11.5/75
0.5 kg Mrat	0.813	1.42	2.32	10.7	
400 mg/kg					
0.215 kg rat	0.801	2.26	1.88	30.7	11.5/150
<i>Observed</i>					
<i>Frat</i>	<i>1.30</i>	<i>0.66</i>	<i>2.19</i>	<i>3.90</i>	
<i>Mrat</i>	<i>1.41</i>	<i>0.68</i>	<i>2.61</i>	<i>4.10</i>	
1,000 mg/kg					
0.35 kg Frat	2.36	3.03	6.08	30.9	11.5/75
0.5 kg Mrat	3.81	3.26	11.9	30.6	

*Note: Mrat = male rat; Frat = female rat, in both cases values are for assumed lifetime average body weights. Simulation values are single day results and not averaged over a week.

Table 12 gives the average daily doses based on the blood MTBE AUC values for male and female rat simulations and the linear relations for each with applied oral dose.

Table 12. MTBE AUC Based PBPK Doses

Nominal dose mg/kg/day	Average applied dose mg/kg/day	MTBE AUC females mg/kg/day	MTBE AUC males mg/kg/day
0	0	0	0
250	143	116.1	124.2
1,000	571	576.0	575.1

Males: $\text{mg/kg/day} = 26.28 + 82.36(\text{mM hour})$, $r = 0.998$;

females: $\text{mg/kg/day} = 38.95 + 159.37(\text{mM hour})$, $r = 0.996$.

Table 13 presents similar simulation results for inhalation exposures with the observed experimental values in *italic boldface*. The results are similar to the oral exposures with predictions of MTBE blood concentrations and AUCs being closer to observed values than TBA predictions. On the basis of comparison of MTBE AUC values, a 3,000 ppm \times six-hour exposure appeared to be equivalent to a 1,000 mg/kg oral gavage dose to a 0.5 kg rat. As seen in the oral exposures, the MTBE AUC in mM hour varied linearly with applied dose [$\text{ppm} \times \text{six-hour/day} = 145.84 + 255.17 (\text{mM hour})$, $r = 0.999$]. Also given in the lower part of Table 13 are dose conversions from MTBE AUC to oral mg/kg/day averaged for lifetime daily intake. This conversion assumes that the same relation exists between AUC and mg/kg/day as seen above in the oral simulations. If this assumption holds, the oral equivalent male doses from the inhalation bioassay would be 0, 82.9, 618.8, and 1,848.3 mg/kg/day. The male oral doses from the gavage bioassay study would be 0, 124.2, and 575.1 mg/kg/day.

**Table 13. Comparison of MTBE PBPK Predictions with Experimental Data:
Rat Inhalation**

Inhalation dose/ Body weight	MTBE mM Cmax	TBA mM Cmax	MTBE AUC mM hour	TBA AUC mM hour	Blood:Air MTBE/TBA
400 ppm × 6 hours 0.215 kg rat	0.219	1.34	1.31	15.8	11.5/350
Observed 400 ppm					
Mrat	0.169	0.535	0.956	5.45	
Frat	0.171	0.531	0.884	5.05	
400 ppm × 6 hours 0.5 kg Mrat	0.182	0.914	1.09	12.2	11.5/350
3,000 ppm × 6 hours 0.5 kg Mrat	1.7	5.4	10.2	125est	11.5/150
8,000 ppm × 6 hours 0.215 kg rat	5.65	9.83	33.9	22.6	11.5/150
Observed 8,000 ppm					
Mrat	6.3	7.2	33.6	81.0	
Frat	6.4	3.3	32.6	34.4	
8,000 ppm × 6 hours 0.5 kg Mrat	5.2	9.6	31.1	487est	11.5/150
Male rats	Nominal dose ppm × 6 hours	MTBE AUC mM hour	Dose from MTBE AUC ppm	Dose from MTBE AUC* mg/kg/day	
	400	1.09	424	82.9	
	3,000	10.2	2,749	618.8	
	8,000	31.1	8,082	1,848.3	

*Note: This conversion assumes the same relation between AUC and mg/kg/day as seen in oral studies or what single oral dose would give the predicted MTBE AUC seen during the 6-hour inhalation exposures. See also Dourson and Felter (1997) for alternative route-to-route extrapolation.

Overall, the PBPK pharmacokinetic correction for delivered dose when based on MTBE blood AUC is relatively modest compared to the simple applied dose. It is presently uncertain whether other dose metrics would be superior to MTBE AUC and will probably remain so until a more definitive mode(s) of action of MTBE carcinogenesis develops.

NONCARCINOGENIC EFFECTS

The most sensitive noncarcinogenic effect by oral route is in the kidney based on the Robinson et al. (1990) 90-day gavage study with a NOAEL of 100 mg/kg/day. As noted above this value was used by U.S. EPA (1996a) to derive a proposed lifetime HA of 70 ppb (or 0.07 mg/L) in drinking water for MTBE. In its more recent document (U.S. EPA 1997a), U.S. EPA employed this toxicity endpoint along with two other noncancer endpoints, neurological and reproductive/developmental, as well as three cancer endpoints in a margin of exposure (MOE) analysis to develop longer-term HAs. Other states also used this toxicity endpoint to develop regulatory guidelines for MTBE as described later in this document.

CARCINOGENIC EFFECTS

Possible Modes of Action

There are limited data available on the mechanism of action of MTBE. It remains unknown whether biotransformation is required for expression of MTBE's carcinogenic activity. The data from several in vitro and in vivo tests indicate that MTBE lacks significant genotoxic activity and suggest that a genotoxic mode of action is unlikely. It has been proposed that MTBE's induction of renal tubular cell tumors in the male rat is the result of α_{2u} -globulin nephropathy. Although some characteristic features of α_{2u} -globulin nephropathy have been associated with MTBE, the absence of others leads to the overall conclusion that α_{2u} -globulin nephropathy is not likely to account for the induction of kidney tumors by MTBE. Although endocrine-mediated modes of action have been suggested for MTBE's induction of testicular tumors in rats and liver tumors in mice, there are insufficient data to support these hypotheses. In summary, the data available at this time do not provide sufficient evidence in support of a specific mode of action of MTBE carcinogenicity.

Estimation of Carcinogenic Potency

According to the proposed guidelines for carcinogen risk assessment (U.S.EPA 1996f) the type of extrapolation employed for a given chemical depends on the existence of data supporting linearity or non-linearity or a biologically-based or case-specific model. When insufficient data are available supporting either approach the default is to use a linear extrapolation. MTBE seems to fit this category, since no mode of action is known (U.S. EPA 1994a, 1994c). Although the lack of genotoxicity and the nonlinearity of the carcinogenic response in some studies might be argued as supportive of a mechanism other than direct genotoxicity via covalent modification of DNA, attempts to identify positively an alternative mechanism have not so far succeeded. Dourson and Felton (1997) attempted to perform an extrapolation of the cancer potency of MTBE from inhalation route (Chun et al. 1992) to oral route. Cancer potency or cancer potency factor (CPF) is a slope derived from a mathematical function used to extrapolate the probability of incidence of cancer from a bioassay in animals using high doses to that expected to be observed at the low doses which are likely to be found in chronic human exposure. The mathematical model, such as the LMS model, is commonly used in quantitative carcinogenic risk assessments in which the chemical agent is assumed to be a

complete carcinogen and the risk is assumed to be proportional to the dose at very low doses. q_1^* is the upper 95% confidence limit on the cancer potency slope calculated by the LMS model. Or another cancer slope factor (CSF) is a potency derived from the lower 95% confidence limit on the 10% tumor dose (LED_{10}). LED_{10} is the 95% lower bound on the dose that is predicted to give a 10% tumor incidence. The CSF equals to 10% dividing by LED_{10} .

Earlier guidelines for cancer risk assessment, including those formerly used by OEHHA (DHS 1985) have required the use of the LMS model to estimate an upper bound on the low-dose potency (q_1^*). However, more recent OEHHA methodologies, and the draft proposed U.S. EPA (1996f) guidelines for carcinogen risk assessment, recommend a linear extrapolation approach based on the LED_{10} . A multistage polynomial is used to fit data in the observable range, unless some other dose-response curve is specifically indicated by the available data. Because adequate data do not exist for MTBE, the default curve-fitting approach is appropriate. Interspecies scaling for oral doses (and internal doses calculated from a single-species pharmacokinetic model) is based on (body weight)^{3/4} as proposed by U.S. EPA (1996f, 1992b) instead of the (body weight)^{1/3} used previously. For inhalation exposures U.S. EPA has in the past used an assumption of equivalence between different species of exposures to a given atmospheric concentration. This provides roughly similar scaling in effect, due to the way that breathing rate and related parameters affecting uptake scale with body weight. More recently PBPK modeling has been seen as a preferable approach to both dose estimation and interspecies scaling of inhalation exposures, where data are available to support this. Since pharmacokinetic data are available for MTBE in the rat, the modeling approach was feasible in this case for that species only.

Table 14 summarizes the cancer potency values derived by both the LED_{10} method and the LMS model (for comparison with earlier results) from the available statistically significant rodent cancer bioassay data sets for MTBE described earlier in the section on carcinogenicity. In all cases the Tox_Risk v.3.5 (Crump et al. 1993) program was used to fit the multistage model to the quantal data sets. The q_1^* cancer potencies or the 95% upper bound on the LMS linear slope at low dose were calculated directly by the program. CSF's are based on the LED_{10} . The CSF is $0.1/LED_{10}$, in units of $(\text{mg/kg/day})^{-1}$. For the curve fitting to estimate the LED_{10} , we have employed a $p \geq 0.05$ criterion for the Chi-squared goodness of fit statistic of the optimized polynomial. In order to obtain an adequate fit it was necessary to exclude the data for kidney tumors in the high dose (8,000 ppm) males rats in the study by Chun et al. (1992). As can be seen from Table 14, the potency estimates for all tumors are similar whether based on the q_1^* or the CSF. Results in the inhalation studies (Chun et al. 1992, Burleigh-Flayer et al. 1992) are effectively the same (within a factor of two) for the different sites in rats and mice, except that the potency for testicular interstitial cell tumors in male rats is about five times higher. Comparison between different routes and experiments for the rat is easiest by examining the data calculated using the pharmacokinetic model to convert the inhalation exposures to equivalent oral doses. In this case it is apparent that all the results are comparable, with the testicular interstitial cell tumors in the Chun et al. (1992) males again showing a slightly higher value than those found at other sites or in the testis in the Belpoggi et al. (1995, 1997) oral study.

Table 14. Dose Response Parameters for MTBE Carcinogenicity Studies

a) Inhalation studies - ppm in air as dose metric

Species	Sex	Tumor site and type	q_1^* (ppm ⁻¹)	LED ₁₀ (ppm)	CSF (ppm ⁻¹)
Mouse	Female	hepatocellular adenoma + carcinoma	3.2×10^{-4}	320	3.2×10^{-4}
	Male	hepatocellular adenoma + carcinoma	7.3×10^{-4}	140	7.0×10^{-4}
Rat	Male	renal tubular cell adenoma + carcinoma	4.4×10^{-4}	240	4.2×10^{-4}
		testicular interstitial cell tumors	2.3×10^{-3}	46	2.2×10^{-3}

Assumed:

Data reassessment by U.S. EPA (1994c, 1995c).

Duration correction based on $(t_e/t_1)^3$: $t_1 = 104$ weeks for both rats and mice.

Interspecies correction: ppm equivalency.

b) Rat oral - Administered dose as dose metric

Study	Sex	Tumor site and type	q_1^* (mg/kg/day) ⁻¹	LED ₁₀ mg/kg/day	CSF (mg/kg/day) ⁻¹
Belpoggi et al. 1995	Male	Leydig cell tumors	1.38×10^{-3}	76	1.38×10^{-3}
	Female	Leukemia/lymphoma	2.13×10^{-3}	49	2×10^{-3}

Assumed:

No duration correction: $t_e = t_1$.

Interspecies correction: $BW^{3/4}$.

c) Rat oral and inhalation studies - AUC as dose metric

Route	Sex	Tumor site and type	q_1^* (mM.hour/day) ⁻¹	LED ₁₀ mM.hour/day	CSF (mM.hour/day) ⁻¹
Inhalation (Chun et al. 1992)	Male	renal tubular cell adenoma + carcinoma	0.037	2.9	0.035
	Male	testicular interstitial cell tumors	0.16	0.66	0.15
Gavage (Belpoggi et al. 1995)	Male	Leydig cell tumors	0.044	2.4	0.041
	Female	Leukemia/lymphoma	0.051	2.1	0.048

Assumed:

Data reassessment by U.S. EPA (1994c, 1995c) for Chun et al. (1992) study.

Duration correction based on $(t_e/t_i)^3$: $t_i = 104$ weeks for rats.

Interspecies correction: AUC equivalency.

d) Rat oral and inhalation studies - Equivalent oral dose as dose metric

Route	Sex	Tumor site and type	q_1^* (mg/kg/day) ⁻¹	LED ₁₀ mg/kg/day	CSF (mg/kg/day) ⁻¹
Inhalation (Chun et al. 1992)	Male	renal tubular cell adenoma + carcinoma	1.9×10^{-3}	55	1.8×10^{-3}
	Male	testicular interstitial cell tumors	9.2×10^{-3}	11	8.7×10^{-3}
Gavage (Belpoggi et al. 1995)	Male	Leydig cell tumors	1.38×10^{-3}	76	1.38×10^{-3}
	Female	Leukemia/lymphoma	2.13×10^{-3}	49	2×10^{-3}

Assumed:

Data reassessment by U.S. EPA (1994c, 1995c) for Chun et al. (1992) study.

Duration correction based on $(t_e/t_i)^3$: $t_i = 104$ weeks for rats.

Interspecies correction: $BW^{3/4}$.

e) Oral and inhalation studies -Study design

Species	Route	Sex	Body weight	Study duration	Lifetime assumed	Dosing schedule	Concentrations	Study
Rat	Inhalation	Male	500 g	97 weeks	104 weeks	6 hour/day, 5 day/week	0, 400, 3,000, 8,000* ppm	Chun et al. 1992
Mouse	Inhalation	Male	35 g	68 weeks	104 weeks	6 hour/day, 5 day/week	0, 400, 3,000, 8,000* ppm	Chun et al. 1992
		Female	30 g	68 weeks	104 weeks	6 hour/day, 5 day/week	0, 400, 3,000, 8,000* ppm	
Rat	Gavage	Male	500 g	lifetime	104 weeks	4 day/week	0, 250, 1,000 mg/kg/day	Belpoggi et al. 1995
		Female	350 g	lifetime	104 weeks	4 day/week	0, 250, 1,000 mg/kg/day	

*8,000 ppm dose group not used in analysis of male rat renal tubule tumors due to inability of multistage polynomial to achieve adequate fit.

Carcinogen risk assessment guidelines used by OEHHA normally recommend selection of human cancer potency estimates based on the most sensitive site and species, unless there is evidence to indicate that the most sensitive site(s) are not relevant to human cancer induction, or represent data sets with unusually wide error bounds. As an alternative, where several equally plausible results are available and are sufficiently close to be regarded as concordant, the geometric mean of all such estimates may be used. The pharmacokinetic model which allows comparison of different routes and corrects for nonlinearities in the relationship between applied and internal dose, is not available for the mouse. Therefore, the potency estimates obtained in the rat are preferred for risk assessment purposes. Because the results in rats and mice are comparable, the use of the rat data is consistent with the policy of selecting appropriately sensitive species as the basis for the estimate of potency in humans.

In terms of the relevance to human cancer and the mechanism of the observed effects, the results of the studies by Chun et al. (1992) and Burleigh-Flayer et al. (1992) are limited by the relatively severe mortality seen in the highest dose groups, and the less-than lifetime exposure given the mice and the male rats. These experimental flaws are not so severe as to exclude the use of the data in risk assessment, nor more prohibitive than the experimental flaws associated with many studies on other compounds which have been successfully used for this purpose. There are, however, additional problems in the case of the testicular interstitial cell tumors observed in male rats by Chun et al. (1992). The study authors stated that the control incidence of these

tumors was lower than the historical incidence observed in animals from the colony from which these experimental animals were obtained. In view of this, the slightly divergent value for the potency estimate obtained with this data set is regarded with lower confidence than the other values obtained in this analysis.

An attempt was made to allow for the severe impact of mortality on the male rat kidney adenoma and carcinoma incidence in the study by Chun et al. (1992) by applying the time-dependent version of the LMS model to the individual time-to-tumor incidence data in this study. A suitable model available in the Tox_Risk program (multistage in dose, Weibull in time) was used, and an adequate fit was obtained. The program provided an estimate of $q_1^* = 7.6 \times 10^{-2} \text{ (mg/kg/day)}^{-1}$, which is substantially higher than the value estimated from the quantal data. The calculated end-of-life LED_{10} indicated a CSF of $7.2 \times 10^{-2} \text{ (mg/kg/day)}^{-1}$. However, the fit obtained involved a large Weibull exponent ($z = 8.7$, whereas more usual values are in the range of three to six), implying a very late appearance of this tumor. This observation may be of interest in addressing the unsolved question of the mechanism of induction of this tumor by MTBE. However it implies a marked reduction in the confidence which can be placed in the potency estimate using this model. Few tumor data were obtained during the final third of the expected lifetime of the exposed rats (due to the early death of all the rats dosed with 8,000 ppm, and most of the rats dosed with 3,000 ppm by this time). The potency estimate therefore involves a substantial extrapolation outside the range of the observed data, even using the LED_{10} /CSF methodology which is designed to avoid such problems. The extreme time dependency, deficiency in genotoxicity data, and other uncertainties described previously also raise the question of how appropriate it is to use this particular model to fit these data. Its use for extrapolation outside the range of observed data (as opposed to merely as a curve-fitting device within the range of observed data) imply an acceptance of the classic Armitage-Doll theory of action for genotoxic carcinogens, which may not be warranted in the case of MTBE. Because the mechanistic information and the technical resources which would be required to undertake a more appropriate analysis of these time-to-tumor data are lacking, it was decided not to include the results of the time-dependent analysis in the final risk estimate.

In view of the closeness of the other values obtained in the rat, and their similar confidence levels, the preferred value for the cancer potency is therefore the geometric mean of the potency estimates obtained for the male rat kidney adenomas and carcinomas combined (Chun et al. 1992), and the male rat Leydig interstitial cell tumors and the leukemia and lymphomas in female rats (Belpoggi et al. 1995). The combined use of these data yields an estimated CSF of $1.7 \times 10^{-3} \text{ (mg/kg/day)}^{-1}$. While it is theoretically possible that the true human CSF could exceed this value, that is considered unlikely. On the other hand it is plausible that the lower bound on the human CSF includes zero. This is a result of statistical uncertainty with a zero lower bound estimate on q_1 by the LMS method with some MTBE data sets and biological uncertainties due to interspecies extrapolation and mode of action.

A unit risk value is similarly derived from the geometric mean of the respective LED_{10} values for the blood MTBE AUC (Table 14c) as follows:

- a) the geometric mean of $2.1 \text{ mM} \times \text{hour}$ is converted to external concentration (in ppm) using the regression expression derived above i.e., $145.84 + 225.17(2.1) = 618.7 = 619 \text{ ppm}$;
- b) this value is converted to mg/m^3 using the $3.6 \text{ mg/m}^3/\text{ppm}$ conversion factor, or $619 \text{ ppm} \times 3.6 \text{ mg/m}^3/\text{ppm} = 2,230 \text{ mg/m}^3$,
- c) the unit risk is calculated as $0.1/2230 \text{ mg/m}^3$ or $4.5 \times 10^{-5} \text{ (mg/m}^3)^{-1}$ or $4.5 \times 10^{-8} \text{ (}\mu\text{g/m}^3)^{-1}$.

Since the LED values were in human equivalent doses no additional interspecies scaling is required. This unit risk would indicate negligible theoretical lifetime cancer risk at ambient MTBE air concentrations below about 6.2 ppbv (ppb by volume).

Calculation of phg

Calculations of public health-protective concentrations of chemical contaminants in drinking water associated with negligible risks for carcinogens or noncarcinogens must take into account the toxicity of the chemical itself, as well as the potential exposure of individuals using the water. Tap water is used directly as drinking water, for preparing foods and beverages. It is also used for bathing or showering, and in washing, flushing toilets, and other household uses resulting in potential dermal and inhalation exposures.

NONCARCINOGENIC EFFECTS

Calculation of a public health-protective concentration (C, in mg/L) for MTBE in drinking water for noncarcinogenic endpoints uses the following general equation adopted by U.S. EPA (1990, 1992a, 1996c):

$$C = \frac{\text{NOAEL/LOAEL} \times \text{BW} \times \text{RSC}}{\text{UF} \times \text{DWC}}$$

where,

- NOAEL/LOAEL = no observed adverse effect level or lowest observed adverse effect level.
- BW = body weight (a default of 70 kg for a male or 60 kg for a female adult).
- RSC = relative source contribution (a default of 20% to 80% as explained below).
- UF = Uncertainty factors (UFs) are included to account for gaps in our knowledge (uncertainty) about the toxicity of chemicals and for recognized variability in human responses to toxic chemicals.

In determining uncertainty factors for chronic effects it is conventional to apply an uncertainty factor where data are only available from short- or medium-term exposures of animals, rather than full lifetime exposures. In the case of MTBE noncarcinogenic effects, there is no adequate chronic study in experimental animals of the critical effect (increase in kidney weight in rats): the key study is of 90 days duration or about 10% the life span of a rat. Because of this, we consider that a 10-fold UF is justified.

For interspecies extrapolation of toxic effects seen in experimental animals to what might occur in exposed humans an UF of up to 10-fold is generally recommended. This is usually considered as consisting of two 3.1-fold parts: one that accounts for metabolic or pharmacokinetic differences between the species; and another that addresses pharmacodynamic differences, i.e. differences between the response of human and animal tissues to the chemical exposure. Based on the limited metabolic studies of MTBE in humans which indicate possible differences from rodents, and unresolved questions of its toxic

potential for neurological, immunological and endocrine effects we believe a 10-fold UF for interspecies differences is appropriate.

Exposed humans are known to vary considerably in their response to toxic chemical and drug exposures due to age, disease states, and genetic makeup, particularly in genetic polymorphisms for enzymes (isozymes) for detoxifying chemicals. While little is known about individual variation of MTBE metabolism and toxicity the use of a 10-fold UF seems prudent considering the widespread use of tap water in the population.

Finally an additional 10-fold UF is used to account for possible carcinogenicity. This follows a U.S.EPA policy applied to their Group C contaminants. OEHHA has previously employed this additional UF for other PHGs in situations where either a nonlinear dose response was applied to a carcinogen or where both linear and nonlinear approaches were used.

DWC = daily water consumption rate (a default of two L/day for an adult has been used by the U.S. EPA (1996b), or L equivalent/day (Leq/day) to account for additional inhalation and dermal exposures from household use of drinking water as explained below).

Based on the NOAEL of 100 mg/kg/day of the most sensitive noncarcinogenic effect in the kidney from the 90-day gavage (Robinson et al. 1990) study, the following calculation can be made:

$$C = \frac{100 \text{ mg/kg/day} \times 70 \text{ kg} \times 0.2}{10 \times 1,000 \times 3 \text{ Leq/day}} = 0.0467 \text{ mg/L} = 47 \text{ ppb (rounded)}$$

In this calculation an additional uncertainty factor of ten is employed to account for potential carcinogenicity and a DWC value of three Leq/day is used to account for inhalation exposures via typical household use as well as ingestion of tap water. The RSC addresses other non-drinking-water sources, principally airborne MTBE from vehicular exhaust. Support for these values is presented below in a discussion of exposure factors.

EXPOSURE FACTORS

The U.S. EPA (1994b) estimated scenarios of potential human exposure to MTBE related to RFG. In terms of the equation for calculating the public health-protective concentrations of chemical contaminants in drinking water as shown above, the first exposure factor to be considered is the RSC. The RSC is a factor which is based on an estimate of the contribution of drinking water exposure relative to other sources such as food, air, etc. While food is often a significant source of chronic chemical exposure, in the case of MTBE, airborne exposures are likely to be most significant, if highly variable. U.S. EPA typically uses 20% as the default RSC. Maine Department of Human Services used 10% RSC for their proposed MCL for MTBE of 35 ppb (Smith and Kemp 1998) based on the same renal toxicity (Robinson et al. 1990) NOAEL in the 90-day oral study.

Estimates for combined population's airborne exposures and occupational subpopulations' exposures vary by three orders of magnitude or more and include few California data sets. Some

of these estimates are collected in Table 15 where RSC values are calculated for a range of drinking water concentrations. The analyses of Brown (1997) include a combined population grand average of 0.00185 mg/kg/day for various activity associated airborne exposures and an average ambient water concentration of 0.36 ppb. The NSTC (1997) report gives MTBE concentrations in groundwater and surface water ranging from 0.2 to 8.7 ppb with a median value of 1.5 ppb, presumably resulting from nonpoint sources. Although the air exposure analysis of Brown (1997) is the most comprehensive it may underestimate MTBE exposures to the general public in local areas in California (e.g., the Los Angeles basin), possibly by a factor of two. Also due to the year-round and universal use of MTBE in California gasoline, commuters, other drivers, gasoline station customers and neighbors, and the general public are likely to receive greater exposures than elsewhere in the U.S. For this reason a health-protective value of 0.2 (or 20%), equal to the default value used by U.S. EPA (1994a, 1994b, 1996a), is used here for the RSC. The other exposure factor in the equation to calculate the public health-protective concentrations of chemical contaminants in drinking water as shown above is DWC, the daily water intake in Leq/day. This represents the amount of tap water consumed as drinking water as well as that mixed with beverages and used in cooking. The default for an adult is two L/day. For children a default of one Leq/day is used. For VOCs, additional exposures occur via the inhalation and dermal routes (i.e., multi-route) during and after showering, bathing, flushing of toilets, washing clothes and dishes, and other domestic uses. Estimates of inhalation and dermal exposure of MTBE relative to ingestion exposure vary from 15% at 0.36 ppb in water (Brown 1997) to 45 to 110% at 70 ppb in water based on predictions of the CalToxTM Model (DTSC 1994) assuming only 50% of inhaled MTBE is absorbed (Nihlen et al. 1998a). A value of 50% or three Leq/day for total exposure would appear to be a reasonable estimate for the purpose of calculating the PHG as shown in Table 16. The Henry's Law constant for MTBE is about 6×10^{-4} atm³/mole at 25°C which is approximately 1/4 that of benzene and 1/14 that of perchloroethylene, the two common VOCs that have been studied previously (Robbins et al. 1993). MTBE is less volatile and its solubility in water is significantly higher than these VOCs. Accordingly, the correction for showering and other activities for assumed daily water consumption for MTBE is smaller than these other common VOCs.

Table 15. Relative Source Contribution (RSC) Estimates (%) for Different Combinations of Air and Drinking Water Exposures to MTBE*

Air exposure estimate (mg/kg/day)	Air exposure scenario	RSC (%)				Reference
		0.36 ppb*	2 ppb*	12 ppb*	70 ppb*	
0.00185	Combined U. S. population grand average	0.6	3	16	52	Brown 1997
0.01	One million exposed U. S. nationwide	0.1	0.6	3.3	17	Brown 1997
0.002	Los Angeles basin at 4 ppbv ambient	0.5	2.8	15	50	ARB 1996
0.0093	Scenario I annual	0.1	0.6	3.6	18	NSTC 1996
0.0182	Scenario II annual	0.06	0.3	1.8	10	NSTC 1996
6.7×10^{-5}	Milwaukee, Wisconsin Air	13	46	84	97	HEI 1996
0.37	MTBE distribution of fuel mixture Time-Weighted-Average (TWA) for workers	0.003	0.02	0.09	27	HEI 1996
1.3×10^{-4}	Albany, New York air	7	30	72	94	NSTC 1997
Geometric mean		0.28	1.5	6.4	34	
Arithmetic mean		2.6	10.4	24.5	45.6	

Note:

$RSC = (I_{water} \times 100) / (I_{water} + I_{air})$. Food and soil sources are considered negligible for MTBE.

I_{water} = uptake by ingestion of tap water containing MTBE at the concentrations noted assuming two L/day and 100% intestinal absorption.

I_{air} = uptake by inhalation of airborne MTBE assuming 20 m³ air inhaled/day and 50% absorption.

Both I_{water} and I_{air} are assumed for a 70 kg human.

*The concentrations of MTBE in drinking water were taken from the reports noted rather than using arbitrary values: 0.36 ppb (Brown 1997); two ppb (NSTC 1997 rounded); 12 ppb (rounded 10⁻⁶ risk estimate, U.S. EPA 1996a); and 70 ppb (proposed Longer-Term and Lifetime HA, U.S. EPA 1996a). However, any plausible range could have been used, e.g., five, 10, 20, 40, etc.

DRAFT

Table 16. CalTox™ Predictions of Inhalation (I), Oral (O) and Dermal (D) Exposures (mg/kg/day) from 70 ppb MTBE Contaminated Tap Water: Effects of Varying Henry's Law Constant and Drinking Water Intake Level

Henry's Law constant (Pa m ³ /mole)	Water intake (mL/kg/day)			
	19.4	33.3	43.9	
66.5	I=	1.165×10^{-3}	1.165×10^{-3}	1.165×10^{-3}
	O=	1.11×10^{-3}	1.91×10^{-3}	2.52×10^{-3}
	D=	4.41×10^{-6}	4.41×10^{-6}	4.41×10^{-6}
		2.275×10^{-3}	3.08×10^{-3}	3.69×10^{-3}
	All	2.46 Leq/day	3.30 Leq/day	3.97 Leq/day
142	I=	1.17×10^{-3}	ND	ND
	O=	1.09×10^{-3}		
	D=	4.43×10^{-6}		
		2.26×10^{-3}		
	All	2.48 Leq/day		
228	I=	1.18×10^{-3}	1.18×10^{-3}	1.18×10^{-3}
	O=	1.09×10^{-3}	1.88×10^{-3}	2.47×10^{-3}
	D=	4.34×10^{-6}	4.3×10^{-6}	4.34×10^{-6}
		2.27×10^{-3}	3.06×10^{-3}	3.65×10^{-3}
	All	2.51 Leq/day	3.33 Leq/day	4.03 Leq/day

Note:

The CalTox™ model vadose and root zone compartments were loaded to predict 70 ppb MTBE in the groundwater used for residential drinking water. Various values for Henry's Law constant and water intake in mL/kg/day for a 62 kg female were used. MTBE parameters for molecular weight, octanol-water partition coefficient, melting point, vapor pressure, and water solubility were entered. Water intake values (mL/kg/day) correspond to median tap water for 20 to 64 year old females (19.4), median total water intake for 20 to 64 year old females (33.3), and average total water intake for all females (43.9) based on the Western Regional data (Ershow and Cantor 1989). Inhalation (I) value assumes 50% of inhaled MTBE is absorbed. Oral (O) and dermal (D) values assume 100% absorption. Total intakes by all routes are also expressed as L equivalents (Leq)/day.

CARCINOGENIC EFFECTS

For carcinogens, the following general equation can be used to calculate the public health-protective concentration (C) for a chemical in drinking water (in mg/L):

$$C = \frac{BW \times R}{q_1^* \text{ or CSF} \times DWC} = \text{mg/L}$$

where,

- BW = adult body weight (a default of 70 kg).
- R = de minimis level for lifetime excess individual cancer risk (a default of 10^{-6}).
- q_1^* or CSF = cancer slope factor. The q_1^* is the upper 95% confidence limit on the cancer potency slope calculated by the LMS model, and CSF is a potency derived from the lower 95% confidence limit on the 10% (0.1) tumor dose (LED₁₀).
CSF = $0.1 / \text{LED}_{10}$. Both potency estimates are converted to human equivalent [in (mg/kg/day)⁻¹] using BW^{3/4} scaling.
- DWC = daily volume of water consumed by an adult (a default of two L/day or other volume in Leq/day to account for additional inhalation and dermal exposures from household use of drinking water as explained above).

Two cancer potency estimates, q_1^* or CSF, were calculated because our current experience with the LMS model is extensive whereas the new methodology proposed by U.S. EPA (1996f) in its draft guidelines for carcinogen risk assessment is based on the LED₁₀ for which little is known about the problems and outcome of using this procedure. The LMS model focuses on the linear low dose extrapolation and analysts (e.g., U.S. EPA) have often accepted relatively poor fits to the observed tumor incidence data. The new method places a higher premium on fitting the observed data to estimate the ED₁₀ and the 95% lower bound LED₁₀, the point from which the low dose extrapolation is made (U.S. EPA 1996a). In the case of the estimates obtained for carcinogenic potency of MTBE, the values calculated using the LMS model are not significantly different from that obtained using the preferred LED₁₀ approach. The calculated public health-protective concentration accounting for carcinogenic effects of MTBE is based on a carcinogenic potency of 1.7×10^{-3} (mg/kg-day)⁻¹. This estimate is the geometric mean of the potency estimates (CSFs) obtained for the combined male rat kidney adenomas and carcinomas in the inhalation study by Chun et al. (1992), the male rat Leydig cell tumors in the oral study by Belpoggi et al. (1995), and the leukemia and lymphomas in female rats, also in the study by Belpoggi et al. (1995). It is consistent with potencies obtained at other sites in another species (mice). The estimate for the inhalation route was converted to an oral intake using the pharmacokinetic model described earlier. The public health-protective concentration was therefore calculated using the following values:

- BW = 70 kg (the default male adult human body weight).
- R = 10^{-6} (default de minimis lifetime excess individual cancer risk).
- q_1^* or CSF = 1.7×10^{-3} (mg/kg-day)⁻¹ (CSF estimated as above).
- DWC = 3 Leq/day (daily water consumption. As described previously in the section on RSCs, there are various probable routes of exposure in addition to ingestion which would result from contamination of water supplies. To

allow for these additional exposures the assumed daily volume of water consumed by an adult is increased from the default of two L/day to three Leq/day).

Thus,

$$C = \frac{70 \times 10^{-6}}{1.7 \times 10^{-3} \times 3} = 14 \times 10^{-3} \text{ mg/L} = 14 \text{ } \mu\text{g/L} = 14 \text{ ppb}$$

Since the calculated public health-protective concentration based on noncancer toxicity of 47 ppb is less protective of public health than the above cancer based value of 14 ppb, the recommended PHG level for MTBE is therefore 14 ppb (0.014 mg/L or 14 μ g/L). The proposed PHG is considered to contain an adequate margin of safety for the potential noncarcinogenic adverse effects including adverse effects on the renal and neurological systems.

RISK CHARACTERIZATION

MTBE is used as an additive in cleaner burning automotive fuel in California. This results in opportunities for airborne exposures as well as drinking water exposures through leaking USTs and to a lesser extent from certain powered water craft and air deposition. The public health risks of exposure to MTBE can be characterized as follows:

ACUTE HEALTH EFFECTS

Acute health effects are not expected to result from typical exposure to MTBE in drinking water. This includes household airborne exposures from showering, flushing toilets, etc. Reports of health complaints of various nonspecific symptoms (e.g., headache, nausea, cough) associated with exposure to gasoline containing MTBE have not been confirmed in controlled studies and remain to be fully evaluated.

CARCINOGENIC EFFECTS

Inhalation exposure to MTBE produced increased incidences of kidney and testicular tumors in male rats and liver tumors in mice. Oral administration of MTBE produced leukemia and lymphoma in female rats and testicular tumors in male rats. A summary of our evaluation is listed below.

- As a result of this assessment OEHHA considers MTBE to be an animal carcinogen and a possible human carcinogen.
- Three cancer bioassays have shown MTBE induced tumors at several sites, in two species, in both sexes, by oral and inhalation routes of exposure.
- Cancer study results exhibit consistency. For example, testicular tumors were induced in rats by both routes of MTBE administration.
- The oral rat study by Belpoggi et al. (1995, 1997) was found to be adequate for risk assessment purposes despite early mortality in the females.
- The inhalation studies in rats and mice were also considered adequate for risk assessment despite early mortality in both studies.

DRAFT

- In general the quality of the three studies was as good or better than those typically available for chemical risk assessment.
- While there are varying degrees of uncertainty as to the relevance to human cancer causation for each of the tumor types induced by MTBE in rodents (i.e., hepatocellular adenoma and carcinoma, renal tubular adenoma and carcinoma, Leydig interstitial cell tumors of the testes, leukemias and lymphomas), the occurrence of tumors at all of these sites adds considerably to the weight of evidence supporting the conclusion that MTBE should be considered a possible human carcinogen.
- MTBE genotoxicity data is mixed, and there is no clear evidence that genotoxicity of its metabolites is involved in the carcinogenicity observed.
- There is no evidence to support a specific nongenotoxic mode of action (e.g., hormone receptor binding) and no evidence that metabolism of MTBE is required for carcinogenicity. In the absence of sufficient evidence, dose metrics based on the parent compound, MTBE, were necessarily chosen for the dose-response assessment.
- Cancer potency estimates derived from different studies, sites, and routes of administration are similar.
- Cancer potency estimates are low compared to other known carcinogens despite the health conservative default assumptions employed.
- The proposed PHG of 14 ppb is based on an average of three quantitatively similar carcinogen slope factors (CSFs) for three sites (kidney tumors, testicular tumors, leukemia/lymphoma). If the PHG value was based on individual tumor sites instead of an average, the values would range from 2.5 to 17 ppb.
- The CSFs are upper-bound estimates defined by the 95% confidence limit on the ED₁₀. It is theoretically possible that the true value of the cancer potency of MTBE in humans could exceed these values, but that is considered unlikely. It is plausible that the true value of the human cancer potency for MTBE has a lower bound of zero based on statistical and biological uncertainties including interspecies extrapolation and mode of action.
- The estimate of multi-route exposure employed in the PHG calculation was three Leq/day. The range of exposure estimates based on different Henry's Law constants and water ingestion rates was 2.3 to four Leq/day. The range of possible PHGs based on this range and the average CSF of 0.0017 (mg/kg-d)⁻¹ is 10 to 18 ppb.
- Additional peer review of all the cancer bioassays would be useful as would be a separate bioassay of MTBE in drinking water. However, these desiderata should be seen in the context of the data already available, which are substantial and of better quality than is available for some other compounds for which risk assessments have been undertaken.
- Lack of knowledge of the mode(s) of action of MTBE or its metabolites is a major limitation of this risk assessment.
- Lack of evidence of cancer causation in humans is also a significant limitation, although widespread use and potential exposure is relatively recent in California and the rest of the U.S.
- Additional pharmacokinetic data in humans and improved PBPK models in animals and humans are desirable.
- Lack of information on the role that interindividual variability (i.e., stemming from metabolic polymorphisms, age-related differences, and concurrent disease conditions) may

play in determining susceptibility to the carcinogenicity of MTBE severely hinders identification of sensitive subgroups in the California population.

The cancer potency estimate derived from the geometric mean of the CSFs of the combined male rat kidney adenomas and carcinomas, the male rat Leydig cell tumors, and the leukemia and lymphomas in female rats was $1.7 \times 10^{-3} \text{ (mg/kg/day)}^{-1}$. Individual tumor endpoint CSFs ranged from $1.4 \times 10^{-3} \text{ (mg/kg/day)}^{-1}$ to $8.7 \times 10^{-3} \text{ (mg/kg/day)}^{-1}$, or a range of about six-fold. Potencies based on the LMS were similar ranging from $1.4 \times 10^{-3} \text{ (mg/kg/day)}^{-1}$ to $9.2 \times 10^{-3} \text{ (mg/kg/day)}^{-1}$, or a range of seven-fold. A time-to-tumor analysis gave much higher values of $0.076 \text{ (mg/kg/day)}^{-1}$ and $0.072 \text{ (mg/kg/day)}^{-1}$ for the LMS and LED_{10} approaches, respectively. However this latter estimate has a low degree of confidence. The water concentration associated with a 10^{-6} negligible theoretical extra lifetime cancer risk calculated from this analysis is 14 ppb. This includes an estimate of inhalation exposure from showering in MTBE contaminated water, flushing toilets, and other household activities involving tap water. The estimate of one Leq/day of additional exposure via the inhalation route is lower than the default value of two Leq/day suggested by U.S. EPA (1996b) based on average estimated showering exposures of a number of typical VOCs. This reflects the fact that MTBE is less volatile and more water soluble than other VOCs commonly found in drinking water. The proposed PHG value of 14 ppb also compares favorably with the Provisional Health and Consumer Acceptability Advisory range of 20 to 40 ppb established by U.S. EPA (1997a) using a MOE approach. Since the proposed value of 14 ppb was calculated for a 1×10^{-6} theoretical lifetime extra risk from a linear extrapolation, the values of 140 ppb and 1,400 ppb (1.4 ppm or 1.4 mg/L) would be associated with the higher risk estimates of 1×10^{-5} and 1×10^{-4} , respectively. For PHGs, our use of the RSC has, with a few exceptions, followed U.S. EPA drinking water risk assessment methodology. U.S. EPA has treated carcinogens differently from noncarcinogens with respect to the use of RSCs. For noncarcinogens, RfDs (in mg/kg/day), DWELs (in mg/L) and MCLGs (in mg/L) are calculated using uncertainty factors (UFs), body weights and DWC (in Leq/day) and RSC, respectively. The typical RSC range is 20% to 80% (0.2 to 0.8), depending on the scientific evidence.

U.S. EPA follows a general procedure in promulgating MCLGs:

- if Group A and B carcinogens (i.e., strong evidence of carcinogenicity) MCLGs are set to zero;
 - if Group C (i.e., limited evidence of carcinogenicity), either an RfD approach is used (as with a noncarcinogen) but an additional UF of 1 to 10 (usually 10) is applied to account for the limited evidence of carcinogenicity, or a quantitative method (potency and low-dose extrapolation) is used and the MCLG is set in the 10^{-5} to 10^{-6} cancer risk range;
 - if Group D (i.e., inadequate or no animal evidence) an RfD approach is used to promulgate the MCLG.
- For approaches that use low-dose extrapolation based on quantitative risk assessment, U.S. EPA does not factor in a RSC. The use of low-dose extrapolation is considered by U.S. EPA to be adequately health-protective without the additional source contributions. In developing PHGs, we have used the assumption that RSCs should not be factored in for carcinogens grouped in U.S. EPA categories A and B, and for C carcinogens for which we have calculated a cancer

potency based on low-dose extrapolation. This is an area of uncertainty and scientific debate and it is not clear how this assumption impacts the overall health risk assessment.

other regulatory standards

The World Health Organization (WHO) has recently issued the second draft of an environmental health criteria document on MTBE. In it they state that carcinogenic findings in animal bioassays seem to warrant some concern of potential carcinogenic risk to humans, but the document does not contain a risk characterization. The Dutch Expert Committee on Occupational Standards (Wibowo 1994) recommended a health-based eight hour-Time-Weighted Average (TWA) exposure limit for MTBE of 180 mg/m³ or 50 ppm to be averaged over an eight-hour working day, and a short-term 15-minute-TWA limit of 360 mg/m³ or 100 ppm in the Netherlands. Czechoslovakia has an Occupational Exposure Limit (OEL) TWA of 100 mg/m³ and a Short-Term OEL (STEL) of 200 mg/m³ since January 1993. Russia has a STEL of 100 mg/m³ since January 1993 (RTECS 1997). Sweden established a TWA of 50 ppm and a 15-minute STEL of 75 ppm in 1988 (ACGIH 1996). The British Industrial Biological Research Association (BIBRA) compiled a toxicological profile on MTBE in 1990. The Danish EPA is considering to set a 30 ppb limit of MTBE in groundwater.

In the U.S., the OSHA and NIOSH established the TLV-TWA as 40 ppm in air (144 mg/m³) in 1994 as proposed by ACGIH in 1993. ACGIH (1996) also lists MTBE as an A3 animal carcinogen in 1995 as proposed in 1994. MTBE is on the Emergency Preparedness and Community Right-to-Know Section of the Superfund Amendments and Reauthorization Act of 1986 (SARA Title III) Extremely Hazardous Substances (EHS) list and in the TSCA Test Submission (TSCATS) Database. MTBE is on the Hazardous Air Pollutant (HAP) list with 189 other chemicals to be regulated under the Air Toxics Program of the 1990 CAAA and therefore a California TAC. Article 211(b) of Title III of the CAAA requires that oil companies conduct gasoline inhalation studies and U.S. EPA sent the testing requirement notification on August 20, 1997. Negotiations with industry on the extent of these studies are ongoing. Animal research will focus on short and long-term inhalation effects of conventional gasoline and gasoline with MTBE. The Article 211 studies will also include human exposure research. The research will be completed at varying intervals over the next five years. HEI is funding three new studies designed to answer key questions on the metabolism of MTBE and other ethers in animals and humans. Texas established a half-hour limit in ambient air of 0.6 mg/m³ and an annual limit of 0.288 mg/m³ in 1992 (Sittig 1994).

MTBE is not a priority pollutant under the Clean Water Act and is not a target analyte in routine water quality monitoring and assessment programs. MTBE is included in the draft and final Drinking Water Contaminant Candidate List required by the Safe Drinking Water Act (U.S. EPA 1997b, 1997d, 1998). The final list is published on March 2, 1998, with decisions on whether to establish a standard on at least five contaminants by August 2003. MTBE is proposed for inclusion on the federal "National Drinking Water Contaminant Occurrence Data Base". In the interim, the Office of Water has initiated a database based on voluntary reporting from some states, USGS data, and other available sources. MTBE is on the U.S. EPA Drinking Water Priority List for future regulation. The U.S. EPA's Office of Research and Development is working to identify MTBE research needs, including monitoring, exposure, health effects, and remediation. A workshop is planned for early September to present an initial assessment of research needs to industry and academic groups. A draft report will be issued for public comment in early 1998. Other U.S. EPA activities include development of a protocol to collect

data on potential CO reductions using federal oxygenated gasoline. USGS is conducting urban land use studies this year to characterize VOCs, including MTBE contamination as a part of the larger national NAWQA program.

Since the early 1990s, U.S. EPA has evaluated MTBE to quantify its toxic effects (Farland 1990, Hiremath and Parker 1994, Klan and Carpenter 1994, Gomez-Taylor et al. 1997). U.S. EPA (1996a) proposed a 70 ppb HA for MTBE in its December 1996 draft report based on noncarcinogenic kidney and liver effects in laboratory animals with large uncertainty factors (U.S. EPA 1996f). U.S. EPA also included an extra uncertainty factor in its draft report to account for the possible carcinogenicity of the substance. The laboratory animal cancer bioassays of MTBE by the oral route were performed in Italy (Belpoggi et al. 1995, 1997), and U.S. EPA has not had an opportunity to audit the studies. Nevertheless, in the 1996 draft, U.S. EPA indicated that the animal studies would suggest that 12.5 ppb would equate to a theoretical risk level of one excess fatal case of cancer per million people per 70-year lifetime (a 10^{-6} risk), a level usually viewed as *de minimis*, for MTBE as a probable human carcinogen.

The U.S. EPA (1997c) IRIS database lists the RfC for inhalation of MTBE in air as three mg/m³ as last revised on September 1, 1993. The RfC is based on increased liver and kidney weights, increased prostration in females, and swollen periocular tissues in male and female rats. The RfD for oral exposure to MTBE is under review by U.S. EPA (1997c). In 1992, U.S. EPA derived a draft long-term HA range for MTBE in drinking water of 20 to 200 ppb (or 0.02 to 0.2 mg/L) based on a RfD of 0.1 mg/kg/day from a 90-day rat drinking water study with dose-related increases in relative kidney weights in both sexes (Robinson et al. 1990). The range is due to the uncertainty for the carcinogen classification. The guideline would be either 20 ppb if MTBE were classified as a Group B2 or C carcinogen, or 200 ppb if MTBE is a Group D carcinogen. In 1994, U.S. EPA drafted a proposal in reviewing data from animal studies for the possibility of listing MTBE as a Group B2 carcinogen, and derived an oral cancer potency estimate of

8.6×10^{-3} (mg/kg/day)⁻¹ and a HA of 4 ppb for a 10^{-6} risk.

The States of Vermont and Florida established drinking water standards for MTBE of 40 ppb and 50 ppb, respectively. The New York Department of Public Water promulgated a MCL of 50 ppb in 1988. The New Jersey Department of Environmental Protection proposed in 1994 and established in 1996 a health-based MCL for MTBE in drinking water of 70 ppb, reducing from 700 ppb, based on the 1993 evaluation of the U.S. EPA but using an uncertainty factor of 10,000 instead of the 30,000 applied by the U.S. EPA (NJDWQI 1994, Post 1994). The Illinois Environmental Protection Agency listed a human threshold toxicant advisory concentration of 230 ppb in 1994 and has proposed a health-based MCL for MTBE in drinking water ranging from 70 to 2,000 ppb. The Massachusetts Department of Environmental Protection in 1995 proposed to decrease the guideline for MTBE in drinking water from 700 ppb to 70 ppb (MORS 1995). The Maine Department of Human Services listed a drinking water threshold of 50 ppb in 1995 and is considering to adopt 35 ppb based on noncancer health effects with a RSC of 10% (Smith and Kemp 1998). NCDEHNR has proposed a primary MCL of 70 ppb. The Wisconsin Department of Natural Resources in 1995 established a groundwater enforcement standard for MTBE of 60 ppb (WDOH 1995). The guideline for MTBE in drinking water is 35 ppb in Arizona, 40 ppb in Michigan, 50 ppb in Rhode Island, and 100 ppb in Connecticut and New Hampshire (ATSDR 1996, HSDB 1997, Sittig 1994).

DHS has added MTBE to a list of non-regulated chemicals which require monitoring by drinking water suppliers in California in compliance with the California Safe Drinking Water Act, Sections 116300 to 116750. An interim Action Level of 35 ppb or 0.035 mg/L for drinking water was proposed by the Cal/EPA in 1991 which was based on the oral RfD of

DRAFT

0.005 mg/kg/day then reported on the U.S. EPA IRIS database for an anesthetic effect in rats in a 13-week inhalation study performed in Europe (Greenough et al. 1980). DHS is proceeding with establishing drinking water standards for MTBE in California.

The initial standard to be developed for MTBE is a secondary MCL, which is to be adopted by DHS as a regulation by July 1998. Secondary MCLs address aesthetic qualities of drinking water supplies. In the case of MTBE, the focus is on its organoleptic qualities, that is, its odor and taste. The purpose of the secondary MCL is to protect the public from exposure to MTBE in drinking water at levels that can be smelled or tasted. Secondary MCLs in California are enforceable standards, which means that drinking water should not be served by public water systems if it contains MTBE higher than the secondary standard. Enforceable secondary standards are unique to California. The proposed secondary MCL for MTBE will be proposed based on data from experiments that have been performed by researchers, using panels of subjects who were exposed to varying concentrations of MTBE in water to determine levels at which it could be smelled or tasted. As part of the process by which regulations are adopted under California's Administrative Procedures Act, the proposed regulation will be available for public comment in early 1998.

The next standard to be developed is a primary MCL that protects the public from MTBE at levels that can affect public health. A primary MCL for MTBE will include consideration of the health risk assessment, the technical feasibility of meeting the MCL (in terms of monitoring and water treatment requirements for MTBE) and costs associated with compliance. DHS has requested the OEHHA to provide a risk assessment for MTBE that is required for the development of the primary standard. DHS requested that the risk assessment be completed by July 1998, in order to meet the scheduled adoption of this regulation by July 1999. The proposed primary MCL is anticipated to be available for public comment in early 1999.

REFERENCES

- Aarstad K, Zahlsten K, Nilsen OG (1985). Inhalation of butanols: Changes in the cytochrome P₄₅₀ enzyme system. *Arch. Toxicol. Suppl* 8: 418-421
- Abel EL, Bilitzke PJ (1992). Effects of prenatal exposure to methanol and t-butanol in Long Evans rats. *Am. J. Obstet. Gynecol.* 166(1, pt 2): 433.
- ACGIH (1996). Methyl Tert-Butyl Ether. In: Documentation of the Threshold Limit Values and Biological Exposure Indices. Supplements to the Sixth Edition: Methyl Tert-Butyl Ether. pp. 1 - 12. Cincinnati, Ohio: American Conference of Governmental Industrial Hygienists, Inc. (ACGIH).
- Allen MJ, Borody TJ, Bugliosi TF, May GR, Thistle JL, Etzel AL (1985). Rapid dissolution of gallstones by methyl tertiary-butyl ether: Preliminary observations. *New Engl. J. Med.* 312(4): 217-220. January 24.
- Anderson HA, Hanrahan L, Goldring J, Delaney B (1995). An Investigation of Health Concerns Attributed to Reformulated Gasoline (RFG) Use in Southeastern Wisconsin. Final Report. May 30. Addendum. September 18. 99 pp. Milwaukee, Wisconsin: Section of Environmental Epidemiology and Prevention, Bureau of Public Health, Division of Health, Wisconsin Department of Health and Social Services.
- Anderson RAJ, Reddy JM, Joyce C, Willis BR, Van der Ven H, Zaneveld LJ (1982). Inhibition of mouse sperm capacitation by ethanol. *Biol. Reprod.* 27(4): 833-40.
- Angle CR (1991). If the tap water smells foul, think MTBE. Letter to the Editor. *JAMA* 266(21): 2985-2986. December 4.
- Anonymous (1995). USGS reports MTBE in groundwater. *Oil & Gas J.* 93(16): 21-22. April 17. PennWell Publishing Company.
- API (1994). Odor threshold studies performed with gasoline and gasoline combined with MTBE, ETBE and TAME. Prepared by TRC Environmental Corporation, Windsor, Connecticut, for American Petroleum Institute (API). January. API # 4592. Submitted to the U.S. Environmental Protection Agency with cover letter dated February 22, 1995. EPA/OTS DOC#86950000131. Washington, D.C.: API.
- Arashidani K, Katoh T, Yoshikawa M, Kikuchi M, Kawamoto T, Kodama Y (1993). LD₅₀ and weight change in organs of mice following intraperitoneal administration of methyl tertiary-butyl ether. *Sangyo Igaku (Japan)* 35(5): 404-405. September.
- ARB (1996). Preliminary Ambient Oxygenates Data of the California ARB Laboratory Information System, and San Francisco Bay Area Air Quality Management District (BAAQMD) Ambient Air Toxics Data on MTBE. Laboratory and Monitoring Division, California Air Resources Board (ARB), California Environmental Protection Agency. Sacramento, California: ARB, Cal/EPA.
- ARCO (1995a). MTBE Use and Possible Occurrences in Water Supplies. Prepared by William J. Piel. May 10. 4 pp. Newton Square, Pennsylvania: ARCO Chemical Company.

- ARCO (1995b). Methyl t-Butyl Ether (MtBE): A status report of its presence and significance in U.S. drinking water. Presented by ARCO Chemical Company to the Office of Water, U.S. Environmental Protection Agency on June 8. Glenolden, Pennsylvania: ARCO Chemical Company.
- ARCO (1980). Methyl tertiary-butyl ether: acute toxicological studies. Unpublished study for ARCO Research and Development for ARCO Chemical Company. Glenolden, Pennsylvania: ARCO Chemical Company.
- ATSDR (1996). Toxicological profile for methyl tert-butyl ether. August. Prepared by Research Triangle Institute for the Agency for Toxic Substances and Disease Registry (ATSDR), Centers for Disease Control (CDC), Public Health Service (PHS), United States Department of Health and Human Services (USDHHS). Atlanta, Georgia: ATSDR, CDC.
- Baehr AL, Baker RJ, Lahvis MA (1997). Transport of methyl tert-butyl ether across the water table to the unsaturated zone at a gasoline-spill site in Beaufort, South Carolina. Paper ENVR 230. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 417-418.
- Balter NJ (1997). Casualty assessment of the acute health complaints reported in association with oxygenated fuels. *Risk Anal.* 17(6): 705-715. December.
- Barker JF, Hubbard CE, Lemon LA (1990). The influence of methanol and MTBE on the fate and persistence of monoaromatic hydrocarbons in ground water. *Ground Water Manag.* 113-117.
- Begley R (1994). MTBE high demand time looms as health questions linger. *Chemical Week* 155: 13. August 31 - September 7.
- Begley R, Rotman D (1993). Health complaints fuel federal concern over MTBE. *Chemical Week* 152(10): 7. March 17.
- Beller M, Schloss M, Middaugh J (1992). Potential Illness Due to Exposure to Oxyfuels. Evaluation of Health Effects from Exposure to Oxygenated Fuel in Fairbanks, Alaska. December 11. Bulletin Number 26. Anchorage, Alaska: Alaska Department of Health and Social Services.
- Belpoggi F, Soffritti M, Maltoni C (1995). Methyl tertiary-butyl ether (MtBE) - a gasoline additive - causes testicular and lymphohaematopoietic cancers in rats. *Toxicol. Ind. Hlth.* 11(2): 119-149. March.
- Belpoggi F, Soffritti M, Filippini F, Maltoni C (1997). Results of long-term experimental studies on the carcinogenicity of methyl tert-butyl ether. *Annals N. Y. Acad. Sci.* 837: 77-95. December 26.
- Bevan C, Neepier-Bradley TL, Tyl RW, Fischer LC, Panson RD, Kneiss JJ, Andrews LS (1997a). Two-generation reproductive study of methyl tertiary-butyl ether (MTBE) in rats. *J. Appl. Toxicol.* 17(S1): S13-S19. May.
- Bevan C, Tyl RW, Neepier-Bradley TL, Fischer LC, Panson RD, Douglas JF, Andrews LS (1997b). Developmental toxicity evaluation of methyl tertiary-butyl ether (MTBE) by inhalation in mice and rabbits. *J. Appl. Toxicol.* 17(S1): S21-S29. May.
- BIBRA (1990). Methyl t-butyl ether (MTBE) toxicology profile. The British Industrial Biological Research Association (BIBRA) Working Group. 6 pp. London, Great Britain: BIBRA Toxicology International.
- Biles RW, Schroeder RE, Holdsworth CE (1987). Methyl tert-butyl ether inhalation in rats: a single generation reproduction study. *Toxicol. Ind. Hlth.* 3(4): 519-534. December.
- Bio/dynamics, Inc. (1984a). An inhalation teratology study in rats with methyl tertiary-butyl ether (MtBE). Project No. 80-2515. Unpublished report submitted to API, Washington, D.C. East Millstone, New Jersey: Bio/dynamics, Inc.
- Bio/dynamics, Inc. (1984b). An inhalation teratology study in mice with methyl tertiary-butyl ether (MtBE). Project No. 80-2516. Unpublished report submitted to API, Washington, D.C. East Millstone, New Jersey: Bio/dynamics, Inc.
- Bio/dynamics, Inc. (1984c). The metabolic fate of methyl tertiary-butyl ether (MtBE) following acute intraperitoneal injection. Project No. 80089. Unpublished report submitted to API, Washington, D.C. 150 pp. East Millstone, New Jersey: Bio/dynamics, Inc.
- Bio/dynamics, Inc. (1981). A nine day inhalation toxicity study of MTBE in the rat. Project No. 40-1452. Unpublished report submitted to API, Washington, D.C. 352 pp. East Millstone, New Jersey: Bio/dynamics, Inc.

DRAFT

- Bioresearch Laboratories (1990a). Pharmacokinetics of methyl tertiary-butyl ether (MtBE) and tert-butyl alcohol (TBA) in male and female Fischer-344 rats after administration of ^{14}C -MtBE by intravenous, oral, and dermal routes. Report #38842. Senneville, Quebec, Canada: Bioresearch Laboratories.
- Bioresearch Laboratories (1990b). Mass balance of radioactivity and metabolism of methyl tert-butyl ether (MtBE) in male and female Fischer-344 rats after administration of ^{14}C MtBE by intravenous, oral, and dermal routes. Report #38843. Senneville, Quebec, Canada: Bioresearch Laboratories.
- Bioresearch Laboratories (1990c). Pharmacokinetics of methyl tert-butyl ether (MtBE) and tert-butyl alcohol (TBA) in male and female Fischer-344 rats after single and repeat inhalation nose-only exposure to ^{14}C -MtBE. Report #38844. Senneville, Quebec, Canada: Bioresearch Laboratories.
- Bioresearch Laboratories (1990d). Disposition of radioactivity of methyl tertiary- butyl ether (MtBE) in male and female Fischer-344 rats after nose-only inhalation exposure to ^{14}C -MtBE. Report #38845. Senneville, Quebec, Canada: Bioresearch Laboratories.
- Bird MG, Burleigh-Flayer HD, Chun JS, Douglas JF, Kneiss JJ, Andrews LS (1997). Oncogenicity studies of inhaled methyl tertiary-butyl ether (MTBE) in CD-1 mice and F-344 rats. *J. Appl. Toxicol.* 17 (S1): S45-S55. May.
- Bonin MA, Ashly DL, Cardinali FL, Mccraw JM, Wooten JV (1995). Measurement of methyl tert-butyl ether and tert-butyl alcohol in human blood by purge-and-trap gas chromatography-mass spectrometry using an isotope-dilution method. *J. Analyt. Toxicol.* 19(3): 187-191. May - June.
- Borak J, Pastides H, Van Ert M, Russi M, Herastein J (1998). Exposure to MTBE and acute human health effects: A critical literature review. *Human and Ecological Risk Assess.* 4(1): 177-200. February.
- Borden RC, Daniel RA, LeBrun LE IV, Davis CW (1997). Intrinsic biodegradation of MTBE and BTEX in a gasoline-contaminated aquifer. *Water Resources Res.* 33(5): 1105-1115. May.

- Borghoff SJ, Murphy JE, Medinsky MA (1996a). Development of a physiologically based pharmacokinetic model for methyl tertiary-butyl ether and tertiary-butanol in male Fischer-344 rats. *Fundam. Appl. Toxicol.* 30(2): 264-275. April.
- Borghoff SJ, Prescott-Mathews JS, Poet TS (1996b). The mechanism of male rat kidney tumors induced by methyl tert-butyl ether (MTBE) and its relevance in assessing human risk. *CIIT Activities.* 16(10): 1-10. October. Research Triangle Park, North Carolina: Chemical Industry Institute of Toxicology (CIIT).
- Brady JF, Xiao F, Ning WJ, Yang CS (1990). Metabolism of methyl tertiary-butyl ether by rat hepatic microsomes. *Arch. Toxicol.* 64(2): 157-160.
- Brown A, Devinsky JS, Davis MK, Browne TE, Rodriguez RA (1997). A review of potential technologies for the treatment of methyl tertiary-butyl ether (MTBE) in drinking water. In: *Proceedings of the NGWA-API Conference on Petroleum Hydrocarbons and Organic Chemicals in Ground Water: Prevention, Detection, and Remediation.* 16 pp. November 12 - 14. Houston, Texas. Dublin, Ohio: National Ground Water Association (NGWA).
- Brown SL (1997). Atmospheric and potable water exposures to methyl tertiary-butyl ether (MTBE). *Reg. Toxicol. Pharmacol.* 25(3): 256-276. June.
- Bruce BW, McMahon PB (1996). Shallow groundwater quality beneath a major urban center: Denver, Colorado, USA. *J. Hydrology (Amsterdam, Netherlands)* 186(1-4): 129-151.
- Buckley TJ, Prah JD, Ashley D, Zweidinger RA, Wallace LA (1997). Body burden measurements and models to assess inhalation exposure to methyl tertiary-butyl ether (MTBE). *J. Air Waste Manag. Assoc.* 47(7): 739-752. July.
- Burbacher TM (1993). Neurotoxic effects of gasoline and gasoline constituents. *Environ. Hlth. Perspect.* 101(S6): 133-141. December.
- Burleigh-Flayer HD, Chun JS, Kintigh WJ (1992). Methyl tertiary butyl ether: vapor inhalation oncogenicity study in CD-1 mice. Bushy Run Research Center Report No. 91N0013A. October 15. Submitted to the U.S. EPA under TSCA Section 4 Testing Consent Order 40 CFR 799.5000 with cover letter dated October 29, 1992. EPA/OPTS#42098. Export, Pennsylvania: Bushy Run Research Center.
- Burleigh-Flayer HD, Doss DE, Bird MG, Ridlon SA (1991). Oncogenicity study of inhaled methyl tertiary butyl ether (MTBE) in CD-1 mouse. MTBE Task Force. Washington, D.C.: Oxygenated Fuels Association.
- Buxton HT, Landmeyer JE, Baehr AL, Church CD, Tratnyek PG (1997). Interdisciplinary investigation of subsurface contaminant transport and fate at point-source releases of gasoline containing methyl tert-butyl ether (MTBE). In: *Proceedings of the Petroleum Hydrocarbons and Organic Chemicals in Ground Water: Prevention, Detection, and Remediation Conference.* pp. 15. November 11 - 14. Houston, Texas. Dublin, Ohio: NGWA.
- Cain WS, Leaderer BP, Ginsberg GL, Andrews LS, Cometto-Muniz JE, Gent JF, Buck M, Berglund LG, Mohsenin V, Monahan E, Kjaergaard S (1996). Acute exposure to low-level methyl tertiary-butyl ether (MTBE): Human reactions and pharmacokinetic responses. *Inhalation Toxicol.* 8(1): 21-49. January.
- Cal/EPA (1998). MTBE (methyl tertiary butyl ether) briefing paper by the California Environmental Protection Agency (Cal/EPA). April 24, 1997, updated March 30, 1998. 32 pp. Sacramento, California: Cal/EPA.

- Casanova M, Heck HA (1997). Lack of evidence for the involvement of formaldehyde in the hepatocarcinogenicity of methyl tertiary-butyl ether in CD-1 mice. *Chem. Biol. Interact.* 105(2): 131-143. July 11.
- CDC (1993a). An investigation of exposure to methyl tertiary-butyl ether (MTBE) in oxygenated fuel in Fairbanks, Alaska. October 22. National Center for Environmental Health (NCEH), Centers for Disease Control and Prevention (CDC), PHS, USDHHS. Atlanta, Georgia: CDC.
- CDC (1993b). An investigation of exposure to MTBE among motorists and exposed workers in Stamford, Connecticut. September 14. NCEH, CDC, PHS, USDHHS. Atlanta, Georgia: CDC.
- CDC (1993c). An investigation of exposure to MTBE and gasoline among motorists and exposed workers in Albany, New York. August 4. NCEH, CDC, PHS, USDHHS. Atlanta, Georgia: CDC.
- Cederbaum AI, Cohen G (1980). Oxidative demethylation of t-butyl alcohol by rat liver microsomes. *Biochem. Biophys. Res. Comm.* 97: 730-736.
- Chandler B, Middaugh J (1992). Potential illness due to exposure to oxyfuels: Anchorage, Alaska. Bulletin Number 1. Middaugh J ed. Anchorage, Alaska: Alaska Department of Health and Social Services.
- Chun JS, Burleigh-Flayer HD, Kintigh WJ (1992). Methyl tertiary ether: vapor inhalation oncogenicity study in Fisher 344 rats. Bushy Run Research Center Report No. 91N0013B. November 13. Submitted to the U.S. EPA under TSCA Section 4 Testing Consent Order 40 CFR 799.5000 with cover letter dated November 19, 1992. EPA/OPTS#42098. Export, Pennsylvania: Bushy Run Research Center.
- Church CD, Isabelle LM, Pankow JF, Rose DL, Tratnyek PG (1997). Method for determination of methyl tert-butyl ether and its degradation products in water. *Environ. Sci. Technol.* 31(12): 3723-3726.
- Cinelli S, Ciliutti P, Falezza A, Meli C, Caserta L, Marchetti S, Seeberg AH, Vericat JA (1992). Absence of mutagenicity of methyl-tertiary-butyl ether [abstract No. P36/P10]. *Toxicol. Lett. Suppl.* 1-356: 300.
- Cirvello JD, Radovsky JE, Heath DR, Farnell DR, Lindamood, III C (1995). Toxicity and carcinogenicity of t-butyl alcohol in rats and mice following chronic exposure in drinking water. *Toxicol. Ind. Hlth.* 11(2): 151-165.
- Clary JJ (1997). Methyl tertiary butyl ether systemic toxicity. *Risk Analy.* 17(6): 661-672. December.
- Clegg ED, Cook JC, Chapin RE, Foster PMD, Daston GP (1997). Leydig cell hyperplasia and adenoma formation: Mechanisms and relevance to humans. *Reproductive Toxicol.* 11(1): 107-121.
- Cohen Y (1998). Multimedia distribution of oxygenated fuel additives. Presented at the MTBE Workshop, co-sponsored by the UCLA Center for Environmental Risk Reduction, UC Toxic Substances Research and Teaching Program, and Southern California Society for Risk Analysis. February 28. UCLA, Los Angeles, California. 29 pp.

DRAFT

- Cohen Y, Allen DT, Clay R, Tsai WT, Rosselot K, Klee H, Blewitt D (1991). Multimedia assessment of refinery emissions. Paper No. 91-84.6 presented at the 84th Annual Meeting of the Air and Waste Management Association. June 16 - 21. Vancouver, British Columbia, Canada.
- Conaway CC, Schroeder RE, Snyder NK (1985). Teratology evaluation of methyl tertiary-butyl ether in rats and mice. *J. Toxicol. Environ. Hlth.* 16(6): 797-809.
- Cooney CM (1997). California struggles with presence of MTBE in public drinking water wells. *Environ. Sci. Technol.* 31(6): A269. June.
- Cornitius T (1996). California air rules foster MTBE demand. *Chemical Week* 158(27): 33. July 17.
- Cox DR (1972). Regression models and life-tables (with discussion). *J. R. Stat. Soc. B.* 24: 187-220.
- Crump KS, Howe RB, van Landingham C, Fuller WG (1993). TOX_RISK, a Toxicology Risk Assessment Program. Version 3.5. October. Ruston, Louisiana: K.S. Crump Division, Clement International Corp.
- Dale MS, Losee RF, Crofts EW, Davis MK (1997a). MTBE: Occurrence and fate in source-water supplies. Paper ENVR 101. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 376-377.
- Dale MS, Moylan MS, Koch B, Davis MK (1997b). MTBE: Taste and odor threshold determinations using the flavor profile method. Presented at the Water Quality Technology Conference. November 9 - 13. Denver, Colorado. American Water Works Association (AWWA).
- Daly MH, Lindsey BD (1996). Occurrence and concentrations of volatile organic compounds in shallow groundwater in the Lower Susquehanna River basin, Pennsylvania and Maryland. USGS Water Resources Investigation Report. WRIR 96-4141. 8 pp. USGS.
- Daniel MA, Evans MA (1982). Quantitative comparison of maternal ethanol and maternal tertiary butanol diet on postnatal development. *J. Pharmacol. Exp. Ther.* 222(2): 294-300.
- Daniel RA (1995). Intrinsic Bioremediation of BTEX and MTBE: Field, Laboratory and Computer Modeling Studies. Graduate Thesis, Department of Civil Engineering, NCU. Raleigh, North Carolina: North Carolina State University.
- Daughtrey WC, Gill MW, Pritts IM, Douglas JF, Kneiss JJ, Andrews LS (1997). Neurotoxicological evaluation of methyl tertiary-butyl ether in rats. *J. Appl. Toxicol.* 17(S1): S57-S64. May.
- Davidson JM (1995). Fate and transport of MTBE - The latest data. In: Proceedings of the Petroleum Hydrocarbons and Organic Chemicals in Ground Water: Prevention, Detection, and Remediation Conference. pp. 285-301. November 29 - December 1. Houston, Texas. Dublin, Ohio: NGWA.
- Davidson JM, Parsons R (1996). Remediating MTBE with current and emerging technologies. In: Proceedings of the Petroleum Hydrocarbons and Organic Chemicals in Ground Water: Prevention, Detection, and Remediation Conference. pp. 15-29. November 13 - 25. Houston, Texas. Dublin, Ohio: NGWA.

DRAFT

- Day KJ, de Peyster A, Allgaier BS, Luong A, MacGregor JA (1998). Methyl t-butyl ether (MTBE) effects on the male rat reproductive endocrine axis [Abstract No. 861 presented at the 1998 Society of Toxicology Annual Meeting]. *The Toxicologist* 42(1-S): 174. March.
- Delzer GC, Zogorski JS, Lopes TJ (1997). Occurrence of the gasoline oxygenate MTBE and BTEX compounds in municipal stormwater in the United States, 1991-1995. Paper ENVR 100. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 374-376.
- Denton J, Masur L (1996). California's cleaner burning gasoline and methyl tertiary-butyl ether [Abstract]. In: the Proceedings of the Society of Environmental Toxicology and Chemistry (SETAC) 17th Annual Meeting. p. 115. November 17 - 21. Washington D.C.
- DHS (1997). Proposed Regulation for a Revision to the Secondary MCL List and Revisions to Unregulated Chemical Monitoring List and Associated Requirements for Drinking Water. Division of Drinking Water Environmental Management. October and December. Sacramento, California: California Department of Health Services (DHS), Health and Welfare Agency, State of California.
- DHS (1985). Guidelines for Chemical Carcinogen Risk Assessments and Their Scientific Rationale. Sacramento, California: DHS, Health and Welfare Agency, State of California.
- Diaz, D, Bories P, Ampelas M, Larrey D, Michel H (1992). Methyl tert-butyl ether in the endoscopic treatment of common bile duct radiolucent stones in elderly patients with nasobiliary tube. *Digest. Dis. Sci.* 37: 97-100.
- Dodd DE, Kintigh WJ (1989). Methyl tertiary butyl ether (MTBE) repeated 13-week vapor inhalation study in rats with neurotoxicity evaluation. Bushy Run Research Center Project Report 52-507. Final Report submitted to the U.S. EPA with cover letter dated September 27, 1989 with draft submitted dated August 11, 1989. TSCATS/403186. EPA/OTS #40-8913440. No. FYI-OTS-0889-0689. WPA/PTS # FYI-OTS-0889-0689. Export, Pennsylvania: Bushy Run Research Center.
- Dourson ML, Felter SP (1997). Route-to-route extrapolation of the toxic potency of MTBE. *Risk Analy.* 17(6): 717-725. December.
- Drew RT (1995). Misunderstand MTBE. Letter to the Editor. *Environ. Hlth. Perspect.* 103(5): 420. May.
- DTSC (1994). CalTox™, A Multimedia Total Exposure Model for Hazardous Waste Sites. Spreadsheet User's Guide. Version 1.5. Prepared by the University of California, Davis, in corporation with Lawrence Livermore National Laboratory for the Department of Toxic Substances Control., California Environmental Protection Agency. Sacramento, California: DTSC, Cal/EPA.
- Du JT, Abernathy CO, Donohue J, Mahfouz A, Khanna K (1998). Provisional health and consumer acceptability advisory for methyl t-butyl ether (MTBE) [Abstract no. 1123 presented at the 1998 Society of Toxicology Annual Meeting]. *The Toxicologist* 42(1-S): 228. March.
- Duffy JS, Del Pup JA, Kneiss JJ (1992). Toxicological evaluation of methyl tertiary-butyl ether (MTBE): Testing performed under TSCA consent agreement. *J. Soil Contam.* 1(1): 29-37. and *Hydrocarbon Contam. Soils* 2: 757-765. Washington, D.C.: American Chemical Society.

- Duffy LK (1994). Oxyfuel in Alaska: Use of interleukins to monitor effects on the immune system. *Sci. Total Environ.* 151(3): 253-256. July 18.
- Edison SA, Maier M, Kohler B, Schlauch D, Buttmann A, Gauer E, Riemann JF (1993). Direct dissolution of gallstones with MTBE by endoscopic cannulation of the gallbladder. *Am. J. Gastroenterol.* 88(8): 1242-1248.
- Environmental Canada (1993). Canadian Environmental Protection Act Priority Substances List Supporting Document, Methyl Tertiary Butyl Ether (MTBE). January. 28 pp. Ottawa, Canada: Canada Communication Group Publisher, Government of Canada.
- Environmental Canada (1992). Canadian Environmental Protection Act Priority Substances List Assessment Report No. 5, Methyl Tertiary Butyl Ether (MTBE). 19 pp. Ottawa, Canada: Canada Communication Group Publisher, Government of Canada.
- Erdal S, Gong H Jr., Linn WS, Rykowski R (1997). Projection of health benefits from ambient ozone reduction related to the use of methyl tertiary butyl ether (MTBE) in the reformulated gasoline program. *Risk Anal.* 17(6): 693-704. December.
- Ershow AG, Cantor KP (1989). Total Water and Tap Water Intake in the United States: Population Based Estimates of Quantities and Sources. Life Sciences Research Office, Federation of American Societies for Experimental Biology (FASEB). Baltimore, Maryland: FASEB.
- Farland WH (1990). Memorandum on Drinking Water Quantification of Toxicologic Effects (QTE) for MTBE. Final Draft. ECAO-CIN-DO23. Office of Health and Environmental Assessment. Washington, D.C.: U.S. EPA.
- Faulkner TP, Wiechart JD, Hartman DM, Hussain AS (1989). The effects of prenatal tertiary butanol administration in CBA/J and C57BL/6J mice. *Life Sci.* 45(21): 1989-95.
- Fenelon JM, Moore RC (1996). Occurrence of volatile organic compounds in groundwater in the White River basin, Indiana, 1994 - 1995. USGS Fact Sheet FS 138-96. 4 pp. USGS.
- Fiedler N, Mohr SN, Kelly-McNeil K, Kipen HM (1994). Response of sensitive groups to methyl tertiary-butyl ether (MTBE). In: Proceedings of the 1993 Conference on MTBE and Other Oxygenates, pp. D65-D84. EPA/600-R95-134. *Inhalation Toxicol.* 6 (6): 539-552. November - December.
- Freed CN (1997). EPA fuel programs. Paper ENVR 95. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 366-368.
- Fueta Y, Arashidani K, Katoh T, Fukata K, Kodama Y (1994). Effects of methyl tertiary-butyl ether on the central nervous system. *Sangyo Igaku* 36(1): 26-27.
- Fujita EM, Bowen JL, Green MC, Moosmuller H (1997). Southern California Ozone Study (SCOS97) Quality Assurance Plan. Report prepared for California Air Resources Board, Desert Research Institute, Reno, Nevada. April 4, 1997. Sacramento, California: ARB, Cal/EPA.
- Fujiwara U, Kinoshita T, Sato H, Kojima I (1984). Biodegradation and bioconcentration of alkyl ethers. *Yukagaken* 33: 111-114.

- Garrett P, Moreau M, Lowry JD (1986). MTBE as a groundwater contaminant. In: Proceedings of the NWWA-API Conference on Petroleum Hydrocarbons and Organic Chemicals in Ground Water - Prevention, Detection and Restoration. pp. 227-238. November 12 - 14. Houston, Texas. Dublin, Ohio: National Water Well Association.
- Gilbert CE, Calabrese EJ (1992). Developing a standard for methyl tertiary-butyl ether in drinking water. In: Regulating Drinking Water Quality. pp. 231-252. Gilbert CE, Calabrese EJ eds. Ann Arbor, Michigan: Lewis Publishers, Inc.
- Gomez-Taylor MM, Abernathy CO, Du JT (1997). Drinking water health advisory for methyl tertiary-butyl ether. Paper ENVR 98. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 370-372.
- Gordian ME, Huelsman MD, Brecht ML, Fisher DG (1995). Health effects of MTBE in gasoline in Alaska. *Alaska Med.* 37(3): 101-103, 119. July - September.
- Grady SJ (1997). Distribution of MTBE in groundwater in New England by aquifer and land use. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 392-394.
- Grant KA, Samson HH (1982). Ethanol and tertiary butanol induced microcephaly in the neonatal rat: Comparison of brain growth parameters. *Neurobehav. Toxicol. Teratol.* 4(3): 315-321.
- Greenough RJ, McDonald P, Robinson P, Cowie J, Maule W, Macnaughtan F, Rushton A (1980). MTBE (Driveron) three month inhalation toxicity in rats. Project No. 413038. Inveresk Research International. Unpublished report to Chemise Werke Hols AG, Marl, West Germany. Report No. 1596. 230 pp. Edinburgh. As cited in the U.S. EPA's IRIS, MTBE, 1991. NTIS Publication No. NTIS/OTS 0513483.
- Grosjean E, Grosjean D, Gunawardena R, Rasmussen RA (1998). Ambient concentrations of ethanol and methyl tert-butyl ether (MTBE) in Porto Alegre, Brazil, March 1996 - April 1997. *Environ. Sci. Technol.* 32(6): 736-742. March 15.
- Gupta G, Lin YJ (1995). Toxicity of methyl tertiary butyl ether to *Daphnia magna* and *Photobacterium phosphoreum*. *Bull. Environ. Contam. Toxicol.* 55(4): 618-620. October.
- Hakkola M, Honkasalo ML, Pulkkinen P (1997). Changes in neuropsychological symptoms and moods among tanker drivers exposed to gasoline during a work week. *Occup. Med. (Oxford, London, England)* 47(6): 344-348. August.
- Hakkola M, Honkasalo ML, Pulkkinen P (1996). Neuropsychological symptoms among tanker drivers exposed to gasoline. *Occup. Med. (Oxford)* 46(2): 125-130. April.
- Hakkola M, Saarinen L (1996). Exposure of tanker drivers to gasoline and some of its components. *Annals Occup. Hyg.* 40(1): 1-10. February.
- Hakkola M (1994). Neuropsychological symptoms among tanker drivers with exposure to solvents. *Occup. Med. (Oxford)* 44(5): 243-246. December.
- Hall AH, Rumack BH, eds. (1998). TOMES PLUS System. Micromedex, Inc., Englewood, Colorado.

- Hartle R (1993). Exposure to methyl tertiary-butyl ether and benzene among service station attendants and operators. *Environ. Hlth. Perspect.* 101(S6): 23-26. December.
- Hartley WR, Englande SJ Jr. (1992). Health risk assessment of the migration of unleaded gasoline - a model for petroleum products. *Water Sci. Technol.* 25(3): 65-72.
- Haseman JK, Arnold J. (1990). Tumor incidence in Fischer 344 rats: NTP historical data. In: *Pathology of the Fischer Rat*. Boorman GA, Eustis SL, Elwell MR, Montgomery CA Jr, MacKenzie WF eds. pp. 555-564. San Diego, California: Academic Press.
- Hattis D, Tuler S, Finkelstein L, Luo Z-Q (1986). *A Pharmacokinetic/Mechanism-based Analysis of the Carcinogenic Risk of Perchloroethylene*. Center for Technology, Policy and Industrial Development. Cambridge, Massachusetts: Massachusetts Institute of Technology.
- HEI (1996). *The Potential Health Effects of Oxygenates Added to Gasoline: A Review of the Current Literature*. April. A special report of the Health Effect Institute's Oxygenates Evaluation Committee. Cambridge, Massachusetts: Health Effects Institute (HEI).
- Hiremath CB, Parker JC (1994). Methyl tertiary butyl ether: cancer risk assessment issues (U.S. EPA, Office of Research and Development, Office of Health and Environmental Assessment) [Abstract No. 540 presented at the 1994 Society of Toxicology Annual Meeting]. *The Toxicologist* 14(1): 152. March. 25 pp.
- Hoel DG, Walburg HE (1972). Statistical analysis of survival experiments. *J. Natl. Cancer Inst.* 49: 361-372.
- Hofmann AF, Amelsberg A, Esch O, Schteingart CD, Lyche K, Jinich H, Vanssonenberg E, D'Agostino HB (1997). Successful topical dissolution of cholesterol gallbladder stones using ethyl propionate. *Digest. Dis. Sci.* 42(6): 1274-1282. June.
- Hong JY, Wang YY, Bondoc FY, Yang CS, Lee M, Huang WQ (1997a). Rat olfactory mucosa displays a high activity in metabolizing methyl tert-butyl ether and other gasoline ethers. *Fund. Appl. Toxicol.* 40(2): 205-210. December.
- Hong JY, Yang CS, Lee M, Wang YY, Huang WQ, Tan Y, Patten CJ, Bondoc FY (1997b). Role of cytochrome P₄₅₀ in the metabolism of methyl tert-butyl ether in human livers. *Arch. Toxicol.* 71(4): 266-269.
- Howard PH ed. (1993). *Handbook of Environmental Fate and Exposure Data for Organic Chemicals*. Volume IV. pp. 71-75. Ann Arbor, Michigan: Lewis Publishers, Inc.
- Howard PH, Boethling RS, Jarvis WF, Meylan WM, Michalenko EM (1991). *Handbook of Environmental Degradation Rates*. pp. 653-654. Chelsea, Michigan: Lewis Publishers, Inc.
- HSDB (1997). *Hazardous Substances Data Bank (HSDB): t-Butyl Methyl Ether*. Last revised on April 23, 1997. Bethesda, Maryland: National Library of Medicine (NLM), National Toxicology Program (NTP).
- Hubbard, CE, Barker JF, O'Hannesin SF, Vandegriendt M, Gillham RW (1994). Transport and fate of dissolved methanol, MTBE, and monoaromatic hydrocarbons in a shallow sand aquifer. API Publication 4601, Appendix H. Ontario, Canada: Institute for Groundwater Research, University of Waterloo.
- Hutcheon DE, Arnold JD, ten Hove W, Boyle J III (1996). Disposition, metabolism and toxicity of methyl tertiary-butyl ether, an oxygenate for reformulated gasoline. *J. Toxicol. Environ. Hlth.* 47(5): 453-464. April 5.

- IARC (1995). Formaldehyde. In: International Agency for Research on Cancer (IARC) monographs on the evaluation of carcinogenic risks to humans: wood dust and formaldehyde. 62: 217-362. Lyon, France: IARC, World Health Organization (WHO).
- IARC (1987). Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans: An Updating of IARC Monographs Volumes 1 to 42, Supplement 7. Lyons, France: IARC, WHO.
- Imbriani M, Ghittori S, Pezzagno G (1997). Partition coefficients of methyl tert-butyl ether (MTBE). *G. Ital. Med. Lav. Ergon. (Occupational Medicine, Pisa, Italy)* 19(3): 63-65. July - September.
- Jeffrey D (1997). Physical-chemical properties of MTBE and preferred environmental fate and compartmentalization. Paper ENVR 209. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 397-399.
- Jensen HM, Arvin E (1990). Solubility and degradability of the gasoline additive MTBE and gasoline compounds in water. In: *Contaminated Soils*, 90: 445-448. Netherlands: Kluwer Academic Publishers.
- Johanson G, Nihlen A, Lof A (1995). Toxicokinetics and acute effects of MTBE and ETBE in male volunteers. *Toxicol. Lett. (Shannon, Ireland)* 82/83: 713-718. December.
- Johnson W, Boyne RA (1992). 28-day oral (gavage) toxicity study of methyl tert-butyl ether in rats. ITT Research Institute Project No. L08100. Chicago, Illinois: ITT Research Institute.
- Joseph P (1995). Illness due to methyl tertiary butyl ether. Letter to the Editor. *Arch. Environ. Hlth.* 50(5): 395-396. September - October.
- Juliani G, Gandini G, Gabasio S, Bonardi L, Fascetti E, Gremo L (1985). Colelitolisi chimica transcutanea con metil-ter-butyl etere (MtBE). *La Radiol. Med.* 71: 569-574.
- Kado NY, Kuzmicky PA, Loarca-Pina G, Mumtaz MM (1998). Genotoxicity testing of methyl tertiary-butyl ether (MTBE) in the Salmonella microsuspension assay and mouse bone marrow micronucleus test. *Mutation Res.* In press.
- Katoh T, Arashidani K, Kikuchi M, Yoshikawa M, Kodama Y (1993). Effects of methyl tertiary-butyl ether on hepatic lipid peroxidation in mice. *Nippon Eiseigaku Zasshi (Japan)* 48(4): 873-878. October.
- Kerns KD, Pavkov KL, Donofrio DJ, Gralla EJ, Swenberg JA (1983). Carcinogenicity of formaldehyde in rats and mice after long-term inhalation exposure. *Cancer Res.* 43: 4382-4392.
- Killian W (1998). Oxyfuel advertisement offends readers. Letter to the Editor. *J. Am. Water Works Assoc.* 90(2): 4-4. February.
- Kirschner EM (1996). Growth of top 50 chemicals slowed in 1995 from very high 1994 rate. *C&EN (Chemical and Engineering News)* 74(15): 16-22. April 8.
- Klan MJ, Carpenter MJ (1994). A risk-based drinking water concentration for methyl tertiary-butyl ether (MTBE). U.S. EPA. In: Proceedings of the Petroleum Hydrocarbons and Organic Chemicals in Ground Water: Prevention, Detection and Remediation Conference. pp. 107-115. November 2 - 4. Houston, Texas. Dublin, Ohio: National Water Well Association.

- Klan MJ, Johnson W, Hatoum NS, Yermakoff JK (1992). 28-day oral (gavage) toxicity study of methyl tert-butyl ether (MTBE) in rats [Abstract No. 382 presented at the 1992 Society of Toxicology Annual Meeting]. *The Toxicologist* 12(1): 117. February.
- Kneiss J (1995). MTBE - the headache of clean air. *Environ. Hlth. Perspect.* 103(7-8): 666-670. July - August.
- Koenigsberg S (1997). MTBE: Wild card in groundwater cleanup. *Environ. Protection* 8(11): 26-28. November.
- Landmeyer JE, Pankow JF, Church CD (1997). Occurrence of MTBE and tert-butyl alcohol in a gasoline-contaminated aquifer. Paper ENVR 228. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 413-415.
- Lapham WW, Neitzert KM, Moran MM, Zogorski JS (1997). USGS compiles data set for national assessment of VOCs in groundwater. *Ground Water Monitoring & Remediation*. XVII(4): 147-157. Fall.
- Lee LC, Quintana PJE, de Peyster A (1998). Comet assay evaluation of the effect of methyl t-butyl ether (MTBE) on rat lymphocytes [Abstract No. 923 presented at the 1998 Society of Toxicology Annual Meeting]. *The Toxicologist* 42(1-S): 187. March.
- Lee CW, Weisel CP (1998). Determination of methyl tert-butyl ether and tert-butyl alcohol in human urine by high-temperature purge-and-trap gas chromatography-mass spectrometry. *J. Analyt. Toxicol.* 22(1): 1-5. January/February.
- Leuschner U, Hellstern A, Ansell A, Gatzen M, Guldutuna S, Leuschner M (1994). Manual and automatic gallstone dissolution with methyl tert-butyl ether. *Digest. Dis. Sci.* 39(6): 1302-1308. June.
- Li D, Zhu M, Yang XY, Cheng WB, Mo GQ, Wang XF (1991). Pharmacokinetics of methyl tert-butyl ether (MTBE) in rats. *Zhong Guo Yao Li Xue Yu Du Li Xue Za Zhi (Chinese J. Pharmacol. Toxicol.)* 5(4): 287-290. November.
- Life Science Research Roma Toxicology Centre S.P.A. (1989a). Reverse mutation in *Salmonella typhimurium*, test substance: MtBE. Report No. 216001-M-03489. Rome, Italy: Roma Toxicology Centre S.P.A.
- Life Science Research Roma Toxicology Centre S.P.A. (1989b). Gene mutation in Chinese hamster V79 cells, test substance: MtBE. Report No. 216002-M-03589. Rome, Italy: Roma Toxicology Centre S.P.A.
- Life Science Research Roma Toxicology Centre S.P.A. (1989c). Unscheduled DNA synthesis (UDS) in primary rat hepatocytes (autoradiographic method), test substance: MtBE. Report No. 216003-M-03689. Rome, Italy: Roma Toxicology Centre S.P.A.
- Lin XZ, Chou TC, Lin PW, Chow YL, Li CC, Chen SK (1994). Chemical dissolution of gallstones in Taiwan: An in vitro study. *J. Gastroenterol. Hepatol.* 9(2): 143-147.
- Lindsey BD, Breen KJ, Daly MH (1997). MTBE in water from fractured-bedrock aquifers, southcentral Pennsylvania. Paper ENVR 210. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 399-400.

- Lindstrom SB Pleil JD (1996). Alveolar breath sampling and analysis to exposures to methyl tertiary-butyl ether (MTBE) during motor vehicle refueling. *J. Air Waste Manag. Assoc.* 46: 676-682. July.
- Lington AW, Dodd DE, Ridlon SA, Douglas JF, Kneiss JJ, Andrew LS (1997). Evaluation of 13-week inhalation toxicity study on methyl t-butyl ether (MTBE) in Fischer 344 rats. *J. Appl. Toxicol.* 17(S1): S37-S44. May.
- Little CJ, Dale AD, Whatley JA (1979). Methyl tert-butyl ether: A new chromatographic effluent. *J. Chromatogr.* 169: 381-385.
- Livo KB (1995). Overview of public's perspective of health effects from oxyfuels in Colorado. Presented at the HEI Workshop on Acute Health Effects on Oxygenates and Oxygenated Gasolines. July 27. Chicago, Illinois.
- Long G, Meek ME, Savard S (1994). Methyl tertiary-butyl ether (MTBE): Evaluation of risks to health from environmental exposure in Canada. *J. Environ. Sci. Hlth., Part C, Environ. Carcinog. Ecotoxicol. Rev.* C12(2): 389-395.
- Lucier G, Genter MB, Lao YJ, Stopford W, Starr T (1995). Summary of the carcinogenicity assessment of MTBE conducted by the Secretary's Scientific Advisory Board on Toxic Air Pollutants. Report to the North Carolina Department of Environment, Health, and Natural Resources. *Environ. Hlth. Perspect.* 103(5): 420-422. May.
- MacGregor JA, Richter WR, Magaw RI (1993). Uterine changes in female mice following lifetime inhalation of wholly vaporized unleaded gasoline: A possible relationship to liver tumors? *Am. Coll. Toxicol.* 12: 119-126.
- Mackay D, Shiu WY, Ma KC (1993). *Illustrated Handbook of Physical-Chemical Properties and Environmental Fate for Organic Chemicals.* V. III. Volatile Organic Chemicals. p. 756. Chelsea, Michigan: Lewis Publishers.
- Mackerer CR, Angelosanto FA, Blackburn GR, Schreiner CA (1996). Identification of formaldehyde as the metabolite responsible for the mutagenicity of methyl tertiary-butyl ether in the activated mouse lymphoma assay. *Proc. Soc. Exp. Biol. Med.* 212(4): 338-341. September.
- Mancini ER (1997). Aquatic toxicity data for methyl tertiary-butyl ether (MTBE): Current status, future research. Paper ENVR 245. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 427-429.
- Mantel N (1966). Evaluation of survival data on two new rank order statistics arising in its consideration. *Cancer Chemother. Rep.* 50: 163-170.
- MCCHD (1993). Oxygenated Fuel Data Collection: Missoula Physician Screening. Missoula, Montana: Missoula City-County Health Department (MCCHD).
- McCoy M Jr, Abernethy J, Johnson T (1995). Anecdotal Health-Related Complaint Data Pertaining to Possible Exposures to MTBE: 1993 and 1994 Follow-up Survey (1984 - 1994). API Publication 4623. Washington D.C.: API.
- McGregor DB, Brown A, Cattnach P, Edwards I, McBride D, Caspony WJ (1988). Responses of the LS178y tk⁺/tk⁻ mouse lymphoma cell forward mutation assay. II: 18 coded chemicals. *Environ. Mol. Mutagen.* 11: 91-118.

- McKee RH, Vergnes JS, Galvin JB, Douglas JF, Kneiss JJ, Andrews LS (1997). Assessment of the in vivo mutagenic potential of methyl tertiary-butyl ether. *J. Appl. Toxicol.* 17(S1): S31-S36. May.
- McKinnon RJ, Dyksen JE (1984). Removing organics from groundwater through aeration plus GAC. *J. Am. Water Works Assoc.* 76: 42-47. May.
- Medlin J (1995). MTBE: The headache of clean air. *Environ. Hlth. Perspect.* 103: 666-670. July - August.
- Mehlman MA (1996). Dangerous and cancer-causing properties of products and chemicals in the oil refining and petrochemical industry: Part XXII, health hazards from exposure to gasoline containing methyl tertiary butyl ether, study of New Jersey residents. *Toxicol. Ind. Hlth.* 12(5): 613-627. September - October.
- Mehlman MA (1995). Dangerous and cancer-causing properties of products and chemicals in the oil refining and petrochemical industry: Part XV, health hazards and health risks from oxygenated automobile fuels (MTBE), lessons not heeded. *Int. J. Occup. Med. Toxicol.* 4(2): 219-236.
- Menear JH (1997a). Carcinogenicity studies on Methyl tertiary butyl ether (MTBE): Critical review and interpretation. *Risk Anal.* 17(6): 673-681. December.
- Menear JH (1997b). Methyl tertiary butyl ether (MTBE) carcinogenesis: Hazard identification and safety assessment. *Toxicol. Ecotoxicol. News (TEN)* 4(5): 139-147.
- Menear JH (1995). MTBE: Not carcinogenic. Letter to the Editor. *Environ. Hlth. Perspect.* 103(11): 985-986. November.
- Merck (1989). *The Merck Index: An encyclopedia of Chemicals, Drugs, and Biologicals.* 11th ed. Budavari S, O'Neil MJ, Smith A, Heckelman PE eds. Rahway, New Jersey: Merck & Co., Inc.
- Miller MJ, Ferdinandi ES, Klan M, Andrews LS, Douglas JF, Kneiss JJ (1997). Pharmacokinetics and disposition of methyl t-butyl ether in Fischer-344 rats. *J. Appl. Toxicol.* 17(S1): S3-S12. May.
- Mohr S, Fiedler N, Weisel C, Kelly-McNeil K (1994). Health effects of MTBE among New Jersey garage workers. *Inhalation Toxicol.* 6(6): 553-562. November - December.
- Moolenaar RL, Hefflin BJ, Ashley DL, Middaugh JP, Etzel RA (1997). Blood benzene concentrations in workers exposed to oxygenated fuel in Fairbanks, Alaska. *Int. Arch. Occup. Environ. Hlth.* 69(2): 139-143.
- Moolenaar RL, Hefflin BJ, Ashley DL, Middaugh JP, Etzel RA (1994). MTBE in human blood after exposure to oxygenated fuel in Fairbanks, Alaska. *Arch. Environ. Hlth.* 49(5): 402-409. September - October.
- Mordechai E, Vojdani A, Magtoto L, Choppa PC, Brautbar N (1997). Induction of apoptosis in individuals exposed to methyl tertiary butyl ether (MTBE) [Abstract No. 1727 presented at the 1997 Society of Toxicology Annual Meeting]. *The Toxicologist* 36(1, pt. 2): 340. March.
- Mormile MR, Liu S, Suflita JM (1994). Anaerobic biodegradation of gasoline oxygenates: Extrapolation of information to multiple sites and redox conditions. *Environ. Sci. Technol.* 28(9): 1727-1732. September.

- MORS (1995). Justification for Decreasing the ORS Guideline for MTBE in Drinking Water from 0.7 mg/L to 0.07 mg/L. Office of Research and Standards (ORS), Department of Environmental Protection, the Commonwealth of Massachusetts. Boston, Massachusetts: MORS.
- Moser GJ, Wolf DC, Sar M, Gaido KW, Janszen D, Goldsworthy TL (1998). Methyl tertiary butyl ether-induced endocrine alterations in mice are not mediated through the estrogen receptor. *Toxicol. Sci.* (formerly *Fund. Appl. Toxicol.*) 41(1): 77-87. January.
- Moser GJ, Wong BA, Wolf DC, Fransson-Steen RL, Goldsworthy TL (1996a). Methyl tertiary butyl ether lacks tumor-promoting activity in N-nitrosodiethylamine-initiated B6C3F1 female mouse liver. *Carcinogenesis* 17(12): 2753-2761. December.
- Moser GJ, Wong BA, Wolf DC, Moss OR, Goldsworthy TL (1996b). Comparative short-term effects of methyl tertiary-butyl ether and unleaded gasoline vapor in female B6C3F1 mice. *Fundam. Appl. Toxicol.* 31(2): 173-183. June.
- Neeper-Bradley TL (1991). Two-generation reproduction study of inhaled methyl tert-butyl ether (MTBE) in CD Sprague-Dawley rats. Final Report. August 13. Bushy Run Research Center Project ID 53-594. Submitted to the U.S. EPA with cover letter dated August 16, 1991. EPA/OTS #40-9113465. Export, Pennsylvania: Bushy Run Research Center.
- Nelson BK, Brightwell WS, Khan A, Burg JR, Goa PT (1989a). Lack of selective developmental toxicity of three Butanol isomers administered by inhalation to rats. *Fundam. Appl. Toxicol.* 12(3): 469-79.
- Nelson BK, Brightwell WS, Khan A, Kreig EFJ, Massari VJ (1989b). Behavioral teratology investigation of tertiary-butanol administered by inhalation to rats. *Teratology* 39(5): 504.
- Nihlen A, Lof A, Johanson G (1998a). Experimental exposure to methyl tertiary-butyl ether I. Toxicokinetics in humans. *Toxicol. Appl. Pharmacol.* 148(2): 274-280. February.
- Nihlen A, Walinder R, Lof A., Johanson G (1998b). Experimental exposure to methyl tertiary-butyl ether II. Acute effects in humans. *Toxicol. Appl. Pharmacol.* 148(2): 281-287. February.
- Nihlen A, Lof A, Johanson G (1997). Liquid/air partition coefficients of methyl and ethyl t-butyl ethers, t-amyl methyl ether, and t-butyl alcohol. *J. Clean Technol., Environ. Toxicol., Occup. Med.* 6(2): 205-213.
- Nihlen A, Lof A, Johanson G (1995). Liquid/air partition coefficients of methyl and ethyl t-butyl ethers, t-amyl methyl ether, and t-butyl alcohol. *J. Exposure Analyt. Environ. Epidemiol.* 5(4): 573-592.
- Nihlen A, Walinder R, Lof A, Johanson G (1994). Toxicokinetics and irritative effects of MTBE in man. Presented at the International Society for Environmental Epidemiology meeting. Durham, North Carolina.
- NJDWQI (1994). Maximum Contaminant Level Recommendations for Hazardous Contaminants in Drinking Water, Appendix A: Health-Based Maximum Contaminant Level Support Documents and Addenda. September 26. Newark, New Jersey: New Jersey Drinking Water Quality Institute (NJDWQI).

- NRC (1996). Toxicological and Performance Aspects of Oxygenated Motor Vehicle Fuels. 160 pp. Committee on Toxicological and Performance Aspects of Oxygenated Motor Vehicle Fuels, Board on Environmental Studies and Toxicology, Commission on Life Sciences, National Research Council (NRC), National Academy of Sciences (NAS). Washington, D.C.: National Academy Press.
- NSTC (1997). Interagency Assessment of Oxygenated Fuels. June. National Science and Technology Council (NSTC), Committee on Environment and Natural Resources (CENR) and Interagency Oxygenated Fuels Assessment Steering Committee. White House Office of Science and Technology Policy (OSTP) through the CENR of the Executive Office of the President. Washington, D.C.: NSTC.
- NSTC (1996). Interagency Assessment of Potential Health Risks Associated with Oxygenated Gasoline. February. OSTP through the President's NSTC, CENR and Interagency Oxygenated Fuels Assessment Steering Committee. Washington, D.C.: NSTC.
- NTP (1998). National Toxicology Program January 1998 Update. National Toxicology Program (NTP), National Institute of Environmental Health Sciences (NIEHS), National Institutes of Health (NIH). Research Triangle Park, North Carolina: NTP. January.
- NTP (1997). Toxicology and carcinogenesis studies of isobutene (CAS No. 115-11-7) in Fischer 344/N rats and B6C3F1 mice (inhalation studies). Board Draft. National Toxicology Program Technical Report Series No. 487, NIH Publication No. 97-3977. NTP, NIEHS, NIH. Research Triangle Park, North Carolina: NTP.
- NTP (1995). Toxicology and carcinogenesis studies of t-butyl alcohol (CAS No. 76-65-0) in Fischer 344/N rats and B6C3F1 mice (drinking water studies). National Toxicology Program Technical Report Series No. 436, NIH Publication No. 94-3167. NTP, NIEHS, NIH. Research Triangle Park, North Carolina: NTP.
- OEHHA (1991). MTBE Interim Action Level. February 19. Memorandum from the Pesticide and Environmental Toxicology Section (PETS), Office of Environmental Health Hazard Assessment (OEHHA) to the Office of Drinking Water, DHS. Berkeley, California: PETS, OEHHA, DHS.
- Okahara N, de Peyster A, McPherson SE, MacGregor JA (1998). Effect of MTBE on estrogen-sensitive tissues of immature female CD-1 mice [Abstract No. 862 presented at the 1998 Society of Toxicology Annual Meeting]. *The Toxicologist* 42(1-S): 174-175. March.
- Pankow JF, Thomson NR, Johnson RL, Baehr AL, Zogorski JS (1997). The urban atmosphere as a non-point source for the transport of MTBE and other VOCs to shallow groundwater. *Environ. Sci. Technol.* 31(10): 2821-2828.
- Poet TS, Borghoff SJ (1998). Metabolism of methyl t-butyl ether in human liver microsomes [Abstract No. 464 presented at the 1998 Society of Toxicology Annual Meeting]. *The Toxicologist* 42(1-S): 94. March.
- Poet TS, Borghoff SJ (1997a). In vitro uptake of methyl tert-butyl ether (MTBE) in male rat kidney: Use of a two-compartment model to describe protein interactions. *Toxicol. Appl. Pharmacol.* 145(2): 340-348. August.
- Poet TS, Borghoff SJ (1997b). Metabolism of methyl t-butyl ether in methyl tert-butyl ether (MTBE) in F-344 rats liver microsomes [Abstract No. 1717 presented at the 1997 Society of Toxicology Annual Meeting]. *The Toxicologist* 36(1, pt. 2): 338. March.

- Poet TS, Lalentine JL, Borghoff SJ (1997c). Pharmacokinetics of tertiary butyl alcohol in male and female Fischer 344 rats. *Toxicol. Lett.* 92: 179-186.
- Poet TS, Murphy JE, Borghoff SJ (1996). In vitro uptake of MTBE in male and female rat kidney homogenate: Solubility and protein interactions [Abstract No. 1563 presented at the 1996 Society of Toxicology Annual Meeting]. *The Toxicologist* 30(1, pt. 2): 305. March.
- Ponchon T, Baroud J, Pujol B, Valette PJ, Perrot D (1988). Renal failure during dissolution of gallstone by methyl tert-butyl ether. Letter to the Editor. *Lancet* 2: 276-277. July 30.
- Poore M, Chang B, Niyati F, Madden S (1997). Sampling and analysis of methyl t-butyl ether (MTBE) in ambient air at selected locations in California. Paper ENVR 215. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 407.
- Post G (1994). Methyl Tertiary Butyl Ether, Health-Based Maximum Contaminant Level Support Document. July. Division of Science and Research, New Jersey Department of Environmental Protection (NJDEP). Trenton, New Jersey: NJDEP.
- Prah JD, Goldstein GM, Devlin R, Otto D, Ashley D, House D, Cohen KL, Gerrity T (1994). Sensory, symptomatic, inflammatory, and ocular responses to and the metabolism of MTBE in a controlled human exposure experiment. *Inhalation Toxicol.* 6(6): 521-538. Erratum. (1995) *Inhalation Toxicol.* 7(4): 575.
- Prescott-Mathews JS, Wolf DC, Wong BA, Borghoff SJ (1997a). Methyl tert-butyl ether causes α_{2u} -globulin nephropathy and enhanced renal cell proliferation in male F-344 rats. *Toxicol. Appl. Pharmacol.* 143(2): 301-314. April.
- Prescott-Mathews JS, Garrett MJ, Borghoff SJ (1997b). Chemical binding to α_{2u} -globulin following gavage dosing of ^{14}C - methyl tert-Butyl Ether (MTBE) in F-344 rats [Abstract No. 1726 presented at the 1997 Society of Toxicology Annual Meeting]. *The Toxicologist* 36(1, pt. 2): 340. March.
- Prescott-Mathews JS, Wolf DC, Wong BA, Borghoff SJ (1996). MTBE-induced protein droplet nephropathy and cell proliferation in male F-344 rats [Abstract No. 1559 presented at the 1996 Society of Toxicology Annual Meeting]. *The Toxicologist* 30(1, pt. 2): 304. March.
- Raabe GK (1993). API Health Complaint Survey. EPA/600/R-95/134. In: Proceedings of the Conference on MTBE and Other Oxygenates: A Research Update. July 26-28. Falls Church, Virginia.
- Rao HV, Ginsberg GL (1997). A physiologically-based pharmacokinetic model assessment of methyl t-butyl ether in groundwater for a bathing and showering determination. *Risk Anal.* 17(5): 583-598. October.
- Reese E, Kimbrough RD (1993). Acute toxicity of gasoline and some additives. *Environ. Hlth. Perspect.* 101(S6): 115-131.
- Reisch MS (1994). Top 50 chemicals production rose modestly in the U.S. *C&EN* 72: 12-15. April 8.
- Reynolds G (1998). Oxyfuel advertisement offends readers. Letter to the Editor. *J. Am. Water Works Assoc.* 90(2): 4-4. February.

DRAFT

- Robbins GA, Wang S, Stuart JD (1993). Using the static headspace method to determine Henry's Law Constants. *Analyt. Chemistry* 65(21): 3113-3118.
- Robinson M, Bruner RH, Olson GR (1990). Fourteen- and ninety-day oral toxicity studies of methyl tertiary-butyl ether (MTBE) in Sprague-Dawley rats. *J. Am. Coll. Toxicol.* 9(5): 525-540.
- Rousch JM, Sommerfeld MR (1998). Liquid-gas partitioning of the gasoline oxygenate methyl tert-butyl ether (MTBE) under laboratory conditions and its effect on growth of selected algae. *Arch. Environ. Contam. Toxicol.* 34(1): 6-11. January.
- Rowe BL, Landrigan SJ, Lopes TJ (1997). Summary of published aquatic toxicity information and water-quality criteria for selected volatile organic compounds. USGS Open-File Report. OFR 97-563. 60 pp. USGS.
- RTECS (1997). Registry of Toxic Effects of Chemical Substances (RTECS): Ether, Tert-Butyl Methyl. Last revision date: 1997. National Institute for Occupational Safety and Health (NIOSH). Cincinnati, Ohio: NIOSH.
- Rudo KM (1995). Methyl tertiary butyl ether (MTBE) - evaluation of MTBE carcinogenicity studies. *Toxicol. Ind. Hlth.* 11(2): 167-173. March - April.
- Saarinen L, Hakkola M, Pekari K, Lappalainen K, Aitio A (1998). Exposure of gasoline road-tanker drivers to methyl tert-butyl ether and methyl tert-amyl ether. *Int. Arch. Occup. Environ. Hlth.* 71(2): 143-147. March.
- Savolainen H, Pfaffli P, Elovaara E (1985). Biochemical effects of methyl tertiary-butyl ether (MTBE) in extended vapor exposure in rats. *Arch. Toxicol.* 57(4): 285-288.
- SCDEM (1997). Groundwater Monitoring Report: Project Status, Quarterly Monitoring, and NPDES Report for July to September 1997. October. Fairfield, California: Solano County Department of Environmental Management (SCDEM).
- Scheible MH (1997). California's cleaner-burning gasoline regulations. Paper ENVR 96. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 368-369.
- Scholl HR, Seelig B, Thomas PE, Iba MM (1996). Methyl t-butyl ether (MTBE) induced central nervous system (CNS) depression: effects of metabolism of the compound [Abstract No. 122 presented at the 1996 Society of Toxicology Annual Meeting]. *The Toxicologist* 30(1, Part 2): 24. March.
- Sellakumar AR., Snyder CA, Solomon JJ, Albert RE (1985). Carcinogenicity of formaldehyde and hydrogen chloride in rats. *Toxicol. Appl. Pharmacol.* 81: 401-406.
- Sernau RC (1989). Mutagenicity test on methyl tertiary butyl ether in *Drosophila melanogaster* sex-linked recessive lethal test. Final Report. HLA study no. 10484-0-461. Fiche # OYSO528039. Submitted to the U.S. Environmental Protection Agency with cover letter dated April 14, 1989. EPA/OTS DOC#40-8913430. Kensington, Maryland: Hazelton Laboratories America, Inc.
- Shaffer KL, Uchrin CG (1997). Uptake of methyl tertiary butyl ether (MTBE) by groundwater solids. *Bull. Environ. Contam. Toxicol.* 59(5): 744-749. November.

- Shen YF, Yoo LY, Fitzsimmons SR, Yamamoto MK (1997). Threshold odor concentrations of MTBE and other fuel oxygenates. Paper ENVR 216. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 407-409.
- Siljeholm J (1997). A hazard ranking of organic contaminants in refinery effluents. *Toxicol. Ind. Hlth.* 13(4): 527-551. July - August.
- Sittig (1994). T-Butyl Methyl Ether. In: *World-wide Limits for Toxic and Hazardous Chemicals in Air, Water and Soil.* p. 135. Park Ridge, New Jersey: Noyes Publications.
- Smith SL, Duffy LK (1995). Odor and health complaints with Alaska gasolines. *Chemical Hlth. Safety* 32(3): 2-38.
- Smith AE, Kemp PA (1998). Maximum Contaminant Level for Methyl Tertiary Butyl Ether (MTBE). Technical Support Document. February 1. Environmental Toxicology Program, Bureau of Health. Augusta, Maine: Maine Department of Human Services.
- Soffritti M, Maltoni C, Maffei F, Biagi R (1989). Formaldehyde: an experimental multipotential carcinogen. *Toxicol. Ind. Hlth.* 5: 699-730.
- Sonawane B (1994). Assessment of potential developmental toxicity risk of methyl tertiary butyl ether (MTBE). *Teratology* 49: 388A. May.
- Spitzer HL (1997). An analysis of the health benefits associated with the use of MTBE reformulated gasoline and oxygenated fuels in reducing atmosphere concentrations of selected volatile organic compounds. *Risk Analy.* 17(6): 683-691. December.
- Squillace PJ, Pope DA, Price CV (1995). Occurrence of the Gasoline Additive MTBE in Shallow Groundwater in Urban and Agricultural Areas. USGS Fact Sheet FS-114-95. 4 pp. Washington D.C.: Department of Interior, USGS.
- Squillace PJ, Zogorski JS, Wilber WG, Price CV (1996). Preliminary assessment of the occurrence and possible sources of MTBE in groundwater in the United States, 1993 - 1994. *Environ. Sci. Technol.* 30(5): 1721-1730. May.
- Squillace PJ, Pankow JF, Korte NE, Zogorski JS (1997a). Review of the environmental behavior and fate of methyl tert-butyl ether (MTBE). *J. Environ. Toxicol. Chemistry* 16(9): 1836-1844. September.
- Squillace PJ, Zogorski JS, Wilber WG, Price CV (1997b). Preliminary assessment of the occurrence and possible sources of MTBE in groundwater in the United States, 1993 - 1994. Paper ENVR 99. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 372-374.
- Stackelberg PE, O'Brien AK, Terracciano SA (1997). Occurrence of MTBE in surface and groundwater, Long Island, New York, and New Jersey. Paper ENVR 208. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 394-397.
- Standeven AM, Blazer T, Goldsworthy TL (1994). Investigation of antiestrogenic properties of unleaded gasoline in female mice. *Toxicol. Appl. Pharmacol.* 127: 233-240.

- Steffan RJ, McClay K, Vainberg S, Condee CW, Zhang D (1997). Biodegradation of the gasoline oxygenates methyl tert-butyl ether (MTBE), ethyl tert butyl ether (ETBE), and tert-amyl methyl ether (TAME) by propane-oxidizing bacteria. *Appl. Environ. Microbiol.* 63(11): 4216-4222. November.
- Stelljes ME (1997). Issues associated with the toxicological data on MTBE Paper ENVR 212. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 401-403.
- Stern BR, Kneiss JJ (1997). Methyl tertiary-butyl ether (MTBE): Use as an oxygenate in fuels. *J. Appl. Toxicol.* 17(S1): S1-S2. May.
- Stern BR, Tardiff RG (1997). Risk characterization of methyl tertiary butyl ether (MTBE) in tap water. *Risk Analy.* 17(6): 727-743. December.
- Stoneybrook Laboratories Inc. (1993). Activated mouse lymphoma (L5178Y/tk⁺/tk⁻) mutagenicity assay supplemented with formaldehyde dehydrogenase for methyl tertiary butyl ether. Status Report 65579. Princeton, New Jersey: Stoneybrook Laboratories Inc.
- Stubblefield WA, Burnett SL, Hockett JR, Naddy R, Mancini ER (1997). Evaluation of the acute and chronic aquatic toxicity of methyl tertiary-butyl ether (MTBE). Paper ENVR 246. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 429-430.
- Swenberg JA, Dietrich DR (1991). Immunohistochemical localization of α_{2u} -globulin in kidneys of treated and control rats of a 13-week vapor inhalation study undertaken with methyl tertiary butyl ether (MTBE). Report to John Kneiss, Manager, MTBE Task Force. July 26, 1991. Chapel Hill, North Carolina: University of North Carolina.
- Tardiff RG, Stern BR (1997). Estimating the risks and safety of methyl tertiary butyl ether (MTBE) and tertiary butyl alcohol (TBA) in tap water for exposures of varying duration. Paper ENVR 247. In: Proceedings of the 213th American Chemical Society National Meeting, Division of Environmental Chemistry, Environmental Fate and Effects of Gasoline Oxygenates. April 13 - 17. San Francisco, California. 37(1): 430-432.
- Terracciano SA, O'Brien AK (1997). Occurrence of volatile organic compounds in streams on Long Island, New York, and New Jersey. USGS Fact Sheet. FS 063-97. 4 pp. USGS.
- Til HP, Woutersen RA., Feron VJ, Hollanders VMH, Falke HE (1989). Two-year drinking-water study for formaldehyde in rats. *Food Chem. Toxicol.* 27: 77-87.
- Tepper JS, Jackson MC, McGee JK, Costa DL, Graham JA (1994). Estimation of respiratory irritancy from inhaled MTBE in mice. *Inhalation Toxicol.* 6(6): 563-569. November - December.
- Tyl RW, Neeper-Bradley TL (1989). Developmental toxicity study of inhaled methyl tertiary butyl ether (MTBE) in CD-1 mice. Final Report. July 20. Bushy Run Research Center Report No. 51-266. Submitted to the U.S. EPA with cover letter dated July 26, 1989. TSCATS/403186. EPA/OTS #40-8913432. No. FYI-OTS-0889-0689. Export, Pennsylvania: Bushy Run Research Center.

DRAFT

- Tyl RW (1989). Developmental toxicity study of inhaled methyl tertiary butyl ether (MTBE) in New Zealand white rabbits. Final Report. May 12. Bushy Run Research Center Report No. 51-268. Submitted to the U.S. EPA with cover letter dated July 12, 1989. EPA/OTS #40-8913426. No. FYI-OTS-0889-0689. Export, Pennsylvania: Bushy Run Research Center.
- U.S. EPA (1998). 63 FR 10274. 40 CFR Parts 141 and 142, Announcement of the Drinking Water Contaminant Candidate List; Notice. Federal Register 63(40): 10274-10287. Monday, March 2. Washington, D.C.: U.S. EPA.
- U.S. EPA (1997a). Drinking Water Advisory: Consumer Acceptability Advice and Health Effects Analysis on Methyl Tertiary-Butyl Ether (MtBE). December. Fact Sheet 4 pp. and Advisory 42 pp. EPA-822-F-97-009. ODW 4304. Health and Ecological Criteria Division, Office of Science and Technology, Office of Water. Washington, D.C.: U.S. Environmental Protection Agency (U.S. EPA).
- U.S. EPA (1997b). 62 FR 52193 40 CFR Parts 141 and 142, Announcement of the Draft Drinking Water Contaminant Candidate List; Notice. Federal Register 62(128): 52193-52219. Monday, October 6. Washington, D.C.: U.S. EPA.
- U.S. EPA (1997c). Integrated Risk Information System (IRIS): Methyl Tertiary-Butyl Ether (MTBE). Last revised on September 1, 1993 for Reference Concentration (RfC) assessment. Washington, D.C.: U.S. EPA.
- U.S. EPA (1997d). 40 CFR Parts 141 and 142, Drinking Water Monitoring Requirements for Certain Chemical Contaminants - Chemical Monitoring Reform (CMR) and Permanent Monitoring Relief (PMR); Proposed Rule. Federal Register 62(128): 36100-36136. Thursday, July 3. Washington, D.C.: U.S. EPA.
- U.S. EPA (1996a). Methyl-t-Butyl Ether (MTBE) Drinking Water Health Advisory (draft). December. Health and Ecological Criteria Division, Office of Science and Technology, Office of Water. Washington, D.C.: U.S. EPA.
- U.S. EPA (1996b). 40 CFR Part 131, National Toxics Rule: Remand of Water Quality Criteria for Dioxin and Pentachlorophenol to EPA for Response to Comments -- Response to Comments from American Forest and Paper Association on Two of the Exposure Assumptions Used by EPA In Developing the Human Health Water Quality Criteria for Dioxin and Pentachlorophenol. Federal Register 61(239): 65183-65185. Wednesday, December 11. Washington, D.C.: U.S. EPA.
- U.S. EPA (1996c). Benchmark Dose Technical Guidance Document. August 6. External Review Draft. EPA/600/P-96/002A. Risk Assessment Forum. Office of Health and Environmental Assessment. Washington, D.C.: U.S. EPA.
- U.S. EPA (1996d). Fact Sheet on MTBE in Water. June. Office of Water. Washington, D.C.: U.S. EPA.
- U.S. EPA (1996e). Questions and Answers Fact Sheet on MTBE in Water. June. Office of Water. Washington, D.C.: U.S. EPA.
- U.S. EPA (1996f). 40 CFR Part 131, Proposed Guidelines for Carcinogen Risk Assessment. Federal Register 61(79): 17959-18011. Wednesday, April 23. Washington, D.C.: U.S. EPA.
- U.S. EPA (1995a). Toxics Release Inventory. An online database maintained by the U.S. EPA and accessible via TOXLINE. 1993 data accessed June 1995. Office of Toxic Substances Control. Washington, D.C.: U.S. EPA.

DRAFT

- U.S. EPA (1995b). MTBE in Water. Draft by Evelyn Washington. August 1. Office of Ground Water and Drinking Water. Washington, D.C.: U.S. EPA.
- U.S. EPA (1995c). Summary of cancer risk derivations for MTBE, a screening evaluation. Appendix A of the NSTC 1996 report. November 29. By J. Jinot. Quantitative Risk Methods Group, National Center for Environmental Assessment. Washington, D.C.: U.S. EPA.
- U.S. EPA (1994a). Health Risk Perspectives on Fuel Oxygenates. December. Office of Research and Development (ORD). EPA 600/R-94/217. Washington, D.C.: U.S. EPA.
- U.S. EPA (1994b). Estimates of MTBE Exposures Related to Reformulated Gasoline. December 8. Environmental Criteria and Assessment Office, Office of Health and Environmental Assessment. Research Triangle Park, North Carolina: U.S. EPA.
- U.S. EPA (1994c). Summary of Cancer Risk Derivatives for MTBE, A Screening Evaluation for Internal Consideration Related to Gasoline Oxygenates. December 2. Cancer Assessment Statistics and Epidemiology Branch, Office of Health and Environmental Assessment. Washington, D.C.: U.S. EPA.
- U.S. EPA (1994d). Reproductive and Developmental Effects from Gasoline Vapors, A Screening Evaluation for Internal Consideration Related to Gasoline Oxygenates. November 7. Reproductive and Developmental Toxicology Branch, Office of Health and Environmental Assessment. Washington, D.C.: U.S. EPA.
- U.S. EPA (1993). Assessment of Potential Health Risks of Gasoline Oxygenated with Methyl Tertiary-Butyl Ether (MtBE). November. Office of Research and Development, United States Environmental Protection Agency. EPA/600/R-93/206. Washington, D.C.: U.S. EPA.
- U.S. EPA (1992a). 40 CFR Parts 141 and 142, National Primary Drinking Water Regulations (NPDWR); Synthetic Organic Chemicals and Inorganic Chemicals, Final Rule. Federal Register 57(138): 31776-31849. Friday, July 17. Washington, D.C.: U.S. EPA.
- U.S. EPA (1992b). Draft Report: A Cross-Species Scaling Factor for Carcinogen Risk Assessment Based on Equivalence of mg/kg 3/4/Day. Federal Register 57(109): 24152-24173. June 5. Washington, D.C.: U.S. EPA.
- U.S. EPA (1992c). Methyl t-Butyl Ether (MtBE) Drinking Water Health Advisory. January. Health and Ecological Criteria Division, Office of Science and Technology, Office of Water. Washington, D.C.: U.S. EPA.
- U.S. EPA (1991). α_{2u} -globulin association with chemically induced renal toxicity and neoplasia in the male rat. Risk Assessment Forum. United States Environmental Protection Agency. EPA/625/3-91/019F. Washington, D.C.: U.S. EPA.
- U.S. EPA (1990). 40 CFR Parts 141, 142 and 143, National Primary and Secondary Drinking Water Regulations; Synthetic Organic Chemicals and Inorganic Chemicals, Proposed Rule. Federal Register 55(143): 30370-30448. Wednesday, July 25. Washington, D.C.: U.S. EPA.
- U.S. EPA (1987a). Technical Support Document - MtBE. Prepared for Test Rules Development Branch, Existing Chemicals Assessment Division. OTS Publication SRC RT-86-394. Washington, D.C.: U.S. EPA.
- U.S. EPA (1987b). Reference Dose (RfD): Description and use in health risk assessments. IRIS, Appendix A. United States Environmental Protection Agency. Integrated risk information system documentation, vol. 1. EPA/600/8-66/032a. Washington, D.C.: U.S. EPA.

DRAFT

- USGS (1996). Occurrence of the gasoline additive MTBE in shallow ground water in urban and agricultural areas. October. United States Geological Survey Fact Sheet 114.95. Washington, D.C.: USGS.
- Vergnes JS, Chun JS (1994). Methyl tertiary butyl ether: In vivo in vitro hepatocyte unscheduled DNA synthesis assay in mice. Bushy Run Research Center Laboratory project ID 93N1316. Submitted to the U.S. EPA with cover letter dated June 13, 1994 with draft submitted dated October 15, 1993. EPA/OTS DOC#86940000975 and #86940000009. Export, Pennsylvania: Bushy Run Research Center.
- Vergnes JS, Kintigh WJ (1993). Methyl tertiary butyl ether: Bone marrow micronucleus test in mice. Bushy Run Research Center Laboratory project ID 93N1244. Submitted to the U.S. EPA with cover letter dated November 15, 1993. EPA/OTS #86940000031. Export, Pennsylvania: Bushy Run Research Center.
- Vergnes JS, Morabit ER (1989). Methyl tertiary butyl ether repeated exposure vapor inhalation study in rats: In vivo cytogenetic evaluation. Bushy Run Research Center Project report 51-635. Fiche # OTSO528040 Doc # 40-8913431. Export, Pennsylvania: Bushy Run Research Center.
- Vojdani A, Mordechai E, Brautbar N (1997a). Abnormal apoptosis and cell cycle progression in humans exposed to methyl tertiary butyl ether and benzene contaminating water. *Human Exp. Toxicol.* 16(9): 485-494. September.
- Vojdani A, Namatalla G, Brautbar N (1997b). Methyl tertiary butyl ether antibodies among gasoline service station attendants. *Annals N. Y. Acad. Sci.* 837: 96-104. December 26.
- Von Burg R (1992). Toxicological update: Methyl tertiary butyl ether (MTBE). *J. Appl. Toxicol.* 12(1): 73-74.
- Ward Jr. JB, Au WW, Whorton EB, et al. (1994). Genetic toxicity of methyl-tertiary butyl ether. Division of Environmental Toxicology, Department of Preventive Medicine and Community Health. Galveston, Texas: University of Texas Medical Branch.
- Ward Jr. JB, Daiker DH, Hastings DA, Ammenheuser MM, Legator MS (1995). Assessment of the mutagenicity of methyl-tertiary butyl ether at the HPRT gene in CD-1 mice [Abstract No. 417 presented at the 1995 Society of Toxicology Annual Meeting]. *The Toxicologist* 15: 79.
- WDOH (1995). Methyl Tert-Butyl Ether (MTBE). February. Wisconsin Division of Health (WDOH), Department of Natural Resources. Madison, Wisconsin: WDOH.
- Weil CS (1970). Significance of organ-weight changes in food safety evaluation. In: *Metabolic Aspects of Food Safety*. Roe FJ eds. pp. 419-454. New York, New York: Academic Press.
- White MC, Johnson CA, Ashley DL, Buchta TM, Pelletier DJ (1995). Exposure to methyl tertiary-butyl ether from oxygenated gasoline in Stamford, Connecticut. *Arch. Environ. Hlth.* 50(3): 183-189. May - June.
- Wibowo AAE (1994). Methyl t-Butyl Ether: Health Based Recommended Occupational Exposure Limit. Dutch Expert Committee on Occupational Standards: Werkgroep van Deskundigen ter Vaststelling van MAL-waarden. 75 pp. *Arbete. Och. Hals.* No. 22. 24 pp. Gezondheidsrad, Postbus 90517, 2509 Lm Den Haag, Netherlands.
- Wiley K (1998). Clean air versus clean water: Does California need MTBE? California Senate position paper on MTBE. February. 15 pp. Sacramento, California: Senate Office of Research.

DRAFT

- Woutersen RA, van Garderen-Hoetmer A, Bruijntjes JP, Swart A, Feron VJ (1989). Nasal tumors in rats after severe injury to the nasal mucosa and prolonged exposure to 10 ppm formaldehyde. *J. Appl. Toxicol.* 9: 39-46.
- Wyngaarden JB (1986). New nonsurgical treatment removes gallstones. *JAMA* 256(13): 1692. October 3.
- Yoshikawa M, Arashidani K, Katoh T, Kawamoto T, Kodama Y (1994). Pulmonary elimination of methyl tertiary-butyl ether after intraperitoneal administration in mice. *Arch. Toxicol.* 68(8): 517-519.
- Young WF, Horth H, Crane R, Ogden T, Arnott M (1996). Taste and odour threshold concentrations of potential potable water contaminants. *Water Res.* 30(2): 331-340.
- Zakko SF, Scirica JC, Guttermuth MC, Dodge J, Hajjar JJ (1997). Ethyl propionate is more effective and less cytotoxic than methyl tert-butyl ether for topical gallstone dissolution. *Gastroenterology* 113(1): 232-237.
- Zeiger E, Anderson B, Haworth S, Lawlor T, Mortelmans K, Speck W (1987). Salmonella mutagenicity tests: III. Results from the testing of 255 chemicals. *Environ. Mutagen.* 9(S9): 1-110.
- Zhang YP, Macina OT, Rosenkranz HS, Karol MH, Mattison DR, Klopman G (1997). Prediction of the metabolism and toxicological profiles of gasoline oxygenates. *Inhalation Toxicol.* 9(3): 237-254. April.
- Zielinska B, Shire J, Harshfield G, Pasek R (1997). Paper 97-RP139.06. In: Proceedings of the 90th Annual Meeting of the Air and Waste Management Association (AWMA). June 8 - 13. Toronto, Canada.
- Zogorski JS, Baehr AL, Bauman BJ, Conrad DL, Drew RT, Korte NE, Lapham WW, Morduchowitz A, Pankow JF, Washington ER (1997). Significant findings and water-quality recommendations of the Interagency Oxygenated Fuel Assessment. In: Proceedings of the 11th Conference on Contaminated Soils. In press.
- Zogorski JS, Morduchowitz A, Baehr AL, Bauman BJ, Conrad DL, Drew RT, Korte NE, Lapham WW, Pankow JF, Washington ER (1996). Fuel Oxygenates and Water Quality: Current Understanding of Sources, Occurrence in Natural Waters, Environmental Behavior, Fate and Significance. Final Report. September. Prepared for Interagency Oxygenated Fuel Assessment as Chapter 2 of the NSTC 1997 final report, pp. 2-1 - 2-80. Coordinated by OSTP. Washington, D.C.: NSTC.