

**Malignant Lymphomas found in female rats administered glyphosate in the diet. Data extracted from Monsanto 2-year study
Lankas & Hogan (1981)**

Glyphosate Dose mg/kg/day number affected / number of animals	0 control	5 treated---->	10	30
BRAIN	0/50	0/50	0/50	1/50 (2%)
SPINE	0/50	0/50	0/50	1/50 (2%)
HEART	0/49	1/50 (2%)	0/50	1/50 (2%)
LUNG	0/49	1/50 (2%)	0/49	1/50 (2%)
LIVER	0/50	0/50	1/50 (2%)	2/50 (4%)
THYMUS	0/25	0/32	1/37 (3%)	1/34 (3%)
SPLEEN	0/50	0/50	1/50 (2%)	2/50 (4%)
STOMACH	0/50	0/50	0/50	1/50 (2%)
BONE MARROW	0/46	0/44	1/46 (2%)	1/45 (2%)
ADRENAL GLAND	0/50	0/50	0/50	1/49 (2%)
HARDERIAN GLAND	0/47	0/45	0/47	1/44 (2%)
LYMPH NODE MESENTERIC	0/42	0/39	0/48	1/47 (2%)
LYMPH NODE MANDIBULAR	0/2	0/3	0/6	1/6 (17%)
LYMPH NODE MEDIASTINAL	0/33	0/29	1/37 (3%)	2/30 (7%)

Anthony Samsel, Research Scientist
SEAPHS
Samsel Environmental & Public Health Services
Deerfield, NH 03037