



# Health Advisory and Guidelines for Eating Fish from Taylor Lake (Imperial County)

November 2019



Fish, Ecotoxicology, and Water Section  
Pesticide and Environmental Toxicology Branch  
Office of Environmental Health Hazard Assessment  
California Environmental Protection Agency

# LIST OF CONTRIBUTORS

## ***Office of Environmental Health Hazard Assessment***

### ***Authors***

Lori A. Chumney, M.S.  
Huyen Tran Pham, M.P.H.

### ***Primary Reviewers***

Susan A. Klasing, Ph.D., Section Chief  
Shannon R. Murphy, Ph.D.  
Wesley Smith, Ph.D.

### ***Final Reviewers***

David Ting, Ph.D., Branch Chief  
David Siegel, Ph.D., Assistant to the Deputy Director  
Allan Hirsch, Chief Deputy Director

### ***Director***

Lauren Zeise, Ph.D.

# ACKNOWLEDGMENTS

Developing fish consumption advisories depends on sampling and analysis of fish. The Office of Environmental Health Hazard Assessment acknowledges the contribution of information from the following entities: the State Water Resources Control Board, the California Department of Fish and Wildlife and its analytical resources, the Moss Landing Marine Laboratories and the Water Pollution Control Laboratory. Data were obtained from the California Environmental Data Exchange Network (<http://ceden.waterboards.ca.gov/AdvancedQueryTool>). The map was created using ArcMap (10.5) from Environmental Systems Resource Institute (ESRI, Redlands, California).

### ***For further information, contact:***

Pesticide and Environmental Toxicology Branch  
Office of Environmental Health Hazard Assessment  
California Environmental Protection Agency

1515 Clay Street, 16<sup>th</sup> Floor  
Oakland, California 94612  
Telephone: (510) 622-3170  
Email address: fish@oehha.ca.gov

1001 I Street, P.O. Box 4010  
Sacramento, CA 95812-4010  
Telephone: (916) 324-7572

## LIST OF ACRONYMS AND ABBREVIATIONS

ATL	Advisory Tissue Level
CDFW	California Department of Fish and Wildlife
DDT(s)	dichlorodiphenyltrichloroethane (DDT) and its metabolites dichlorodiphenyldichloroethane (DDD) and dichlorodiphenyldichloroethylene (DDE)
DHA	docosahexaenoic acid
EPA	eicosapentaenoic acid
FDA	Food and Drug Administration
Hg	mercury
MDL	method detection limit
MLML	Moss Landing Marine Laboratories
mm	millimeters
OEHHA	Office of Environmental Health Hazard Assessment
PCBs	polychlorinated biphenyls
ppb	parts per billion
RL	reporting limit
RWB7	Regional Water Board 7 (Colorado River)
Se	selenium
SWAMP	Surface Water Ambient Monitoring Program
SWRCB	State Water Resources Control Board
USDA	United States Department of Agriculture
USDHHS	United States Department of Health and Human Services
US EPA	United States Environmental Protection Agency

## PREFACE

The Office of Environmental Health Hazard Assessment (OEHHA), a department in the California Environmental Protection Agency, is responsible for evaluating potential public health risks from chemical contamination of sport fish. This includes issuing fish consumption advisories, when appropriate, for the State of California. OEHHA's authorities to conduct these activities are based on mandates in the:

- California Health and Safety Code
  - Section 59009, to protect public health
  - Section 59011, to advise local health authorities
- California Water Code
  - Section 13177.5, to issue health advisories

The health advisories are published in the California Department of Fish and Wildlife Sport Fishing Regulations in the section on public health advisories.


This report presents guidelines for eating fish from Taylor Lake in Imperial County. The report provides background information and a technical description of how the guidelines were developed. The resulting advice is summarized in the illustrations after the Table of Contents and List of Figures and Tables.

## TABLE OF CONTENTS

A GUIDE TO EATING FISH FROM TAYLOR LAKE.....	5
INTRODUCTION .....	6
<i>Location</i> .....	6
<i>Approach Used</i> .....	7
CHEMICALS OF POTENTIAL CONCERN.....	7
DATA SOURCES.....	8
<i>Colorado River Basin Fish Study, 2014 (RWB7)</i> .....	9
<i>Monitoring of Contaminants in Fish from California Lakes and Reservoirs, 2016 (SWAMP)</i> .....	9
FISH SAMPLED FROM TAYLOR LAKE .....	9
CHEMICAL CONCENTRATIONS .....	10
<i>Mercury</i> .....	10
<i>PCBs and Pesticides</i> .....	11
<i>Selenium</i> .....	11
DEVELOPMENT OF GUIDELINES FOR EATING FISH FROM TAYLOR LAKE.....	12
CONSUMPTION ADVICE FOR FISH FROM TAYLOR LAKE .....	14
<i>Black Bass Species (Largemouth Bass)</i> .....	15
<i>Common Carp</i> .....	15
<i>Sunfish Species (Bluegill, Redear Sunfish)</i> .....	15
RECOMMENDED MAXIMUM NUMBER OF SERVINGS.....	16
REFERENCES .....	17
APPENDIX I. Advisory Tissue Levels.....	19


## LIST OF FIGURES AND TABLES

Figure 1. Location of Taylor Lake.....	6
Table 1. Fish Samples Evaluated for the Taylor Lake Advisory .....	10
Table 2. Mercury and Selenium Concentrations in Fish from Taylor Lake .....	12
Table 3. Recommended Maximum Number of Servings per Week for Fish from Taylor Lake .....	16
Advisory Tissue Levels for Selected Analytes.....	19



**Women**  
(18-49 Years)

**Children**  
(1-17 Years)



**Women**  
(50+ Years)

**Men**  
(18+ Years)

## A GUIDE TO EATING FISH


*from*

# TAYLOR LAKE

(IMPERIAL COUNTY)


**Eat the Good Fish**

Eating fish that are low in chemicals may provide health benefits to children and adults.




**Avoid the Bad Fish**

Eating fish with higher levels of chemicals like mercury or PCBs may cause health problems in children and adults.




**Choose the Right Fish**


Chemicals may be more harmful to unborn babies and children.



**Common Carp**



**Black Bass Species**  
♥ *high in omega-3s*




**Sunfish Species**


**Serving Size**

A serving of fish is about the size and thickness of your hand. Give children smaller servings.

**For Adults**




**For Children**




**California Office of Environmental Health Hazard Assessment**

web [www.oehha.ca.gov/fish](http://www.oehha.ca.gov/fish)  
 email [fish@oehha.ca.gov](mailto:fish@oehha.ca.gov)  
 phone (916) 324-7572

**Eat only the skinless fillet**



**Eat only the meat**



Some chemicals are higher in the skin, fat, and guts.

## INTRODUCTION

This report presents guidelines for eating fish from Taylor Lake (Figure 1) in Imperial County, near the California-Arizona border. Taylor Lake is adjacent to the Colorado River in the Picacho State Recreation Area, and is under the jurisdiction of the California Department of Parks and Recreation.<sup>1</sup>

### *LOCATION*

Taylor Lake is located about 40 miles south of Blythe, and 20 miles north of Winterhaven, in southeastern California.

FIGURE 1. LOCATION OF TAYLOR LAKE



<sup>1</sup> Information regarding Taylor Lake was obtained from the California State Parks Division of Boating and Waterways. Online at: <http://www.dbw.ca.gov/BoatingFacilities/Details/1259>



### *APPROACH USED*

The Office of Environmental Health Hazard Assessment (OEHHA) used the results from two monitoring studies described in this report to develop the Taylor Lake Advisory. OEHHA uses the following general process in developing consumption advice for sport fish:

- 1) Evaluation of all fish contaminant data available from a water body and selection of appropriate data that meet data quality criteria and sampling plan guidelines.
- 2) Determination of fish species for which adequate data are available to issue fish consumption advice.
- 3) Calculation of an appropriate measure of central tendency (often a weighted arithmetic mean<sup>2</sup>) and other descriptive statistics of the contaminant data, as appropriate, for a chemical of potential concern for the selected fish species.
- 4) Comparison of the chemical concentrations with the OEHHA Advisory Tissue Levels (ATLs) for each chemical of potential concern.
- 5) Development of final advice based on a thorough review of the data and best professional judgment relating to the benefits and risks of consuming a particular fish species.

The ATLs (discussed further in a subsequent section of this report) are chemical levels in fish tissue that are considered acceptable, based on chemical toxicity, for a range of consumption rates. Development of the ATLs also included consideration of health benefits associated with including fish in the diet (OEHHA, 2008). The ATLs should not be interpreted as static “bright lines,” but one component of a complex process of data evaluation and interpretation used by OEHHA in the assessment and communication of the benefits and risks of consuming sport fish.

## CHEMICALS OF POTENTIAL CONCERN

Certain chemicals are considered to be of potential concern for people who eat fish because of their toxicity and their ability to accumulate in fish tissue. The majority of fish consumption advisories in California are issued because of mercury (Hg), followed by polychlorinated biphenyls (PCBs) and, in a few cases, selenium (Se) or some legacy pesticides (pesticides that are no longer used but remain in the environment).

Mercury is a natural element found in some rock and soil. Human activities, such as burning coal and the historic use of mercury to mine gold, also add mercury to the environment. If mercury enters waterways, it can be converted to a more toxic form

---

<sup>2</sup> Means are an arithmetic average of individual values and/or a weighted average of composites. A weighted average of composites is calculated by multiplying the chemical concentration in each composite by the number of fish in that composite for each species. Products are then summed and divided by the total number of fish in all composites for that species, combined.

known as methylmercury – which can pass into and build up in fish. High levels of methylmercury can harm the brain, especially in fetuses and children.

PCBs are industrial chemicals previously used in electrical transformers, plastics, and lubricating oils, often as flame retardants or electrical insulators. Their use was banned in the 1970s, but they persist in the environment because they do not break down easily and can accumulate in fish. Depending on the exposure level, PCBs may cause cancer or other health effects, including neurotoxicity, in humans.

Selenium is a naturally occurring metalloid and at low doses is an essential nutrient for many important human health processes, including thyroid regulation and vitamin C metabolism. Higher doses cause selenium toxicity, which can include symptoms ranging from hair loss and gastrointestinal distress to dizziness and tremors.

Chlordanes, dichlorodiphenyltrichloroethane (DDT), dieldrin, and toxaphene are pesticides that were banned from use in 1973 (DDT), the late 1980s (chlordanes and dieldrin) and 1990 (toxaphene), but are still found in some fish in certain California water bodies. Depending on the exposure level, these chemicals may cause cancer or adverse effects on the nervous system.

Detailed discussion of the toxicity of these chemicals and references are presented in “Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Chlordane, DDTs, dieldrin, methylmercury, PCBs, selenium, and toxaphene” (OEHHA, 2008).

All fish species collected from Taylor Lake and used in advisory development were analyzed for mercury (as a measure of methylmercury) and selenium. Common Carp were also analyzed for PCBs and the legacy pesticides chlordanes (cis-chlordane, trans-chlordane, cis-nonachlor, trans-nonachlor, and oxychlordane), dieldrin, and DDTs (DDT and its metabolites dichlorodiphenyldichloroethane [DDD] and dichlorodiphenyldichloroethylene [DDE]). Fish species that do not normally accumulate PCBs or other organic chemicals may not be analyzed for those contaminants in a particular monitoring study. Additionally, some studies do not analyze these chemicals and instead focus only on mercury.

## DATA SOURCES

The guidelines for eating fish from Taylor Lake are based on the chemicals detected in the fish collected for the two monitoring studies described below. These studies met OEHHA’s data quality criteria, including adequate documentation of sample collection, fish preparation methods (e.g., skinning or filleting), chemical analyses, quality assurance, and sufficiently low detection limits. “Sample,” as used in this report, refers to an individual fish or a composite of multiple fish for which contaminant data were reported. “Sampling” or “sampled” refers to the act of collecting fish for chemical analysis.

*COLORADO RIVER BASIN FISH STUDY, 2014 (RWB7)*

The Colorado River Regional Water Quality Control Board (RWB7)<sup>3</sup> staff, in cooperation with the State Water Resources Control Board (SWRCB), monitors water quality in California's surface waters. In 2014, the program performed regional monitoring surveys of Taylor Lake to evaluate contaminants in commonly consumed sport fish and to collect information about contamination in the greater aquatic food web. The surveys collected Bluegill, Common Carp, Largemouth Bass, and Redear Sunfish from Taylor Lake. All species were analyzed for selenium and mercury. Common Carp were additionally analyzed for chlordanes, dieldrin, DDTs, and PCBs.

*MONITORING OF CONTAMINANTS IN FISH FROM CALIFORNIA LAKES AND RESERVOIRS, 2016 (SWAMP)*

The Surface Water Ambient Monitoring Program (SWAMP), operated by the SWRCB, monitors water quality in California's surface waters. In 2016, SWAMP collected Common Carp, Largemouth Bass, and Redear Sunfish from Taylor Lake, which were analyzed for mercury and selenium. The purpose of the study was to supplement long-term monitoring data that document bioaccumulation impacts on the beneficial uses of California waters. The study focused on water bodies that provide beneficial uses through fishing and had either not been previously sampled or were previously sampled, but needed data gaps filled to determine impairment or develop consumption advisories (Davis et al., 2019).

**FISH SAMPLED FROM TAYLOR LAKE**

The fish sampling data used in this advisory were retrieved from the California Environmental Data Exchange Network (CEDEN), the state's repository for environmental data. Samples were excluded when the fish were not legal size to take or did not meet OEHHA's criteria for minimum "edible" size based on species size at maturity, and professional judgment (as described in OEHHA, 2005). A summary of all fish species evaluated for this advisory is shown in Table 1, including the name of the species, number of samples collected, total number of fish, project name, year sampled, and contaminants analyzed.

---

<sup>3</sup> Information on the Colorado River Regional Water Quality Control Board can be found online at: <https://www.waterboards.ca.gov/coloradoriver/>.

TABLE 1. FISH SAMPLES EVALUATED FOR THE TAYLOR LAKE ADVISORY

Common Name	Scientific Name	Number of Samples	Total Number of Fish	Project	Year Collected	Contaminants Analyzed
Bluegill	<i>Lepomis macrochirus</i>	1	6	RWB7	2014	Hg, Se
Common Carp	<i>Cyprinus carpio</i>	1	5	RWB7	2014	Chlordanes, DDTs, Dieldrin, Hg, PCBs, Se
		2	10	SWAMP	2016	Hg, Se
Largemouth Bass	<i>Micropterus salmoides</i>	8	8	RWB7	2014	Hg
		1	5	RWB7	2014	Se
		7	7	SWAMP	2016	Hg
		1	5	SWAMP	2016	Se
Redear Sunfish	<i>Lepomis microlophus</i>	2	10	RWB7	2014	Hg, Se
		2	10	SWAMP	2016	Hg, Se

## CHEMICAL CONCENTRATIONS

As shown in Table 1, samples were analyzed for one or more of the following: total mercury, selenium, chlordanes, DDTs, dieldrin, and PCBs (50 congeners<sup>4</sup>). Among the chemicals analyzed in fish tissue samples from Taylor Lake, only mercury and selenium levels were sufficiently high to impact consumption advice.

All fish samples were prepared as skinless fillets, and samples were analyzed as individual fish or composites. For this advisory, OEHHA used the weighted (by the number of individual fish) arithmetic mean (average) of the chemical concentrations (in wet weight) for each fish species to estimate average human exposure.

### MERCURY

Samples were analyzed for total mercury, as either individual fish or composite samples, using a direct mercury analyzer (DMA) at the California Department of Fish and Wildlife (CDFW) Moss Landing Marine Laboratories (MLML). The DMA method utilizes thermal decomposition and atomic absorption. OEHHA assumed all mercury

<sup>4</sup> Congeners are related compounds with similar chemical forms. Of the 209 possible PCB congeners, 54-55 are generally reported.

detected was methylmercury, which is the most common form found in fish and is also the more toxic form (Bloom, 1992). Table 2 shows the averages and ranges for total length<sup>5</sup>, as well as mercury concentrations in each fish species. The DMA method detection limit (MDL)<sup>6</sup> and the reporting limit (RL)<sup>7</sup> for total mercury were reported at 4 and 12 parts per billion (ppb), respectively.

#### *PCBS AND PESTICIDES*

Some composite samples were analyzed for PCBs and the legacy pesticides (chlordanes, DDTs, and dieldrin). PCBs and pesticides were analyzed by gas chromatography at the CDFW Water Pollution Control Laboratory. For chlordanes, DDTs, and PCBs, each of the concentrations presented was the sum of the detected parent compound, congeners, or metabolites, where applicable. Since the MDLs or RLs were relatively low ( $\leq 1$  ppb), individual congeners or metabolites with concentrations reported as non-detects were assumed to be zero. This is a standard method of handling non-detect values for PCBs and other chemicals with multiple congeners or metabolites in a given sample when detection levels are adequate (US EPA, 2000a).

Concentrations of chlordanes, dieldrin, DDTs, and PCBs were lower than the corresponding ATL threshold values for daily consumption (OEHHA, 2008 and 2011). These chemicals were therefore not considered further for developing consumption advice and are not shown in this report.

#### *SELENIUM*

The CDFW MLML analyzed species collected from Taylor Lake for selenium, as composite samples, using inductively coupled plasma-mass spectrometry (ICP-MS). The ICP-MS method utilizes desolvation, atomization and ionization with ion separation based on a mass-to-charge ratio to detect the total selenium concentration in a sample. The ICP-MS method detection limit (MDL) and the reporting limit (RL) for total selenium were reported at 150 and 400 ppb, respectively.

---

<sup>5</sup> Total length is the maximum length of the fish, measured from the tip of the closed mouth to the tip of the pinched tail fin.

<sup>6</sup> The MDL is the lowest quantity of a chemical that can be distinguished (as greater than zero) in a sample.

<sup>7</sup> The RL is the lowest quantity of a chemical that can be accurately quantified in a sample.

TABLE 2. MERCURY AND SELENIUM CONCENTRATIONS IN FISH FROM TAYLOR LAKE

Species from Taylor Lake	Number of Samples	Total Number of Fish	Mean* Total Length (mm)	Range of Total Lengths** (mm)	Mercury (ppb)	
					Mean*	Range**
Common Carp	3	15	526	465 - 588	9	4 - 12
Largemouth Bass	15	15	382	330 - 431	56	30 - 137
Sunfish Species	5	26	190	126 - 280	15	0 - 27
Bluegill	1	6	142	126 - 170	17	n/a
Redear Sunfish	4	20	204	150 - 280	14	0 - 27
					Selenium (ppb)	
Common Carp	3	15	526	465 - 588	1517	1440 - 1640
Largemouth Bass	2	10	368	330 - 400	1795	1580 - 2010
Sunfish Species	5	26	190	126 - 280	1882	1480 - 2170
Bluegill	1	6	142	126 - 170	1480	n/a
Redear Sunfish	4	20	204	150 - 280	2003	1770 - 2170

\*Means are an arithmetic average of individual values and/or a weighted average of composites.

\*\*Range of individuals and/or range of the composites.

n/a = not applicable due to a single sample

## DEVELOPMENT OF GUIDELINES FOR EATING FISH FROM TAYLOR LAKE

The OEHHA fish advisory process considers the health benefits of fish consumption as well as the risk from exposure to the chemical contaminants found in fish. Benefits are included in the advisory process because there is considerable evidence and scientific consensus that fish should be part of a healthy, well-balanced diet. Fish contain many nutrients that are important for general health and, in particular, help promote optimal growth and development of babies and young children, and may reduce the incidence of heart disease in adults (FDA/US EPA, 2017; American Heart Association, 2016; OEHHA, 2008; Institute of Medicine, 2007; Kris-Etherton et al., 2002). Fish are a significant source of the specific omega-3 fatty acids, docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA), thought to be associated with these beneficial health effects (USDA/USDHHS, 2015; Weaver et al., 2008).

The 2015-2020 US Dietary Guidelines recommend that 1) the general population “consume eight or more ounces per week (less for young children)” of a variety of

seafood<sup>8</sup> “for the total package of nutrients that seafood provides, including its EPA and DHA content” and 2) “women who are pregnant or breastfeeding should consume at least eight and up to twelve ounces of a variety of seafood per week from choices that are lower in methylmercury” (USDA/USDHHS, 2015). The particular fish that people eat is an important factor in determining the net beneficial effects of fish consumption. For example, studies have shown that children of mothers who ate low-mercury fish during pregnancy scored better on cognitive tests compared to children of mothers who did not eat fish or ate high-mercury fish (Oken et al., 2005 and 2008). Accordingly, because of the high mercury content of certain fish species, the US Food and Drug Administration (FDA) and the US Environmental Protection Agency (US EPA) recommend that women who are pregnant (or might become pregnant) or breastfeeding, and young children avoid consuming shark, swordfish, tilefish (Gulf of Mexico), bigeye tuna, marlin, orange roughy, and king mackerel (FDA/US EPA, 2017).

In order to address the potential health concerns associated with exposure to contaminants in sport fish, OEHHA has established ATLS for chemicals that are known to accumulate in the edible tissues of fish. ATLS consider both the toxicity of the chemical and potential benefits of eating fish. OEHHA uses the ATLS to determine the maximum number of servings per week that consumers can eat, for each species and at each location, to limit their exposure to these contaminants. Consumers can use OEHHA’s guidance when choosing which fish and how much to eat as part of an overall healthy diet.

There are two sets of ATLS for methylmercury in fish because of the age-related toxicity of this chemical (OEHHA, 2008). The fetus and children are more sensitive to the toxic effects of methylmercury. Thus, the ATLS for the sensitive population, including women who might become pregnant (typically 18 to 49 years of age) and children 1-17 years, are lower than those for women 50 years and older, and men 18 years and older. The lower ATL values for the sensitive population provide additional protection to allow for normal growth and development of the brain and nervous system of unborn babies and children. Detailed discussion about the toxicity of common fish contaminants and health benefits of fish consumption, as well as derivation of the ATLS, are provided in “Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Chlordane, DDTs, dieldrin, methylmercury, PCBs, selenium, and toxaphene” (OEHHA, 2008). A list of the ATLS used in this report is presented in Appendix I.

For each fish species in this advisory, OEHHA compared the mean mercury and selenium concentrations detected in the fillet to the corresponding ATLS to establish the maximum number of servings per week that could be consumed (see Appendix I). A serving size is considered to be 8 ounces, prior to cooking, or about the size and

---

<sup>8</sup> “Marine animals that live in the sea and in freshwater lakes and rivers. Seafood includes fish, such as salmon, tuna, trout, and tilapia, and shellfish, such as shrimp, crab, and oysters” (USDHHS/USDA, 2015).

thickness of a hand for fish fillets. Children should be given smaller servings. For smaller fish species, several individuals may be required to yield a serving.

The consumption advice for a fish species is initially based on the chemical with the lowest allowable number of servings per week. Because some chemicals, such as mercury and PCBs, are known to have similar adverse effects, additivity of toxicity is assumed in such cases and may be assessed using multiple chemical exposure methodology (US EPA, 1989 and 2000b). If two or more chemicals with similar adverse effects are present in fish tissue at levels above the corresponding ATL values for daily consumption, multiple chemical exposure methodology is employed. This may result in advising the sensitive population to consume fewer meals per week than would be the case for the presence of one chemical alone, in a similar concentration. The potential effect of multiple chemical exposures (mercury and PCBs) was not assessed in Common Carp, the only species for which both contaminants were analyzed, because PCB levels were non-detectable. Advice for other species in this advisory was based solely on mercury or selenium concentrations.

OEHHA recommends that individuals strive to meet the US Dietary Guidelines seafood consumption recommendations, while also adhering to federal and OEHHA recommendations to limit the consumption of fish with higher contaminant levels. The advice discussed in the following section represents the maximum recommended number of servings per week for different fish species. People should eat no more than the recommended number of servings for each fish species or species group. OEHHA's consumption advice for a particular fish species can be extended to other closely related fish species<sup>9</sup> known to accumulate similar levels of contaminants.

Consumption advice should not be combined. That is, if a person chooses to eat a fish from the "one-serving-a-week" category, then they should not eat any other fish from any source (including commercial) until the next week. If a person chooses to eat a fish from the "two-servings-per-week" category, they can combine fish species from that category, or eat one fish from that category and one from a category that recommends more than two-servings-per-week (if available), for a total of two servings in that week. Then they should not eat any other fish from any source (including commercial) until the following week.

## CONSUMPTION ADVICE FOR FISH FROM TAYLOR LAKE

OEHHA's advisory protocol requires at least nine fish of a species to be collected from a water body before an advisory can be developed for the primary contaminant of concern. This is to ensure the sample dataset is representative of the fish species population in the water body. In some cases, an exception is made for species that are commonly caught and consumed from a given water body but where available data may be limited. For Taylor Lake, the sample size criterion was met for the following species:

---

<sup>9</sup> Fish species within the same genus are most closely related, and family is the next level of relationship.



Black bass species (Largemouth Bass), Common Carp, and sunfish species (Bluegill, Redear). There were not sufficient data to evaluate other species that may be found in this water body. For fish species found in Taylor Lake that are not included in this advisory, OEHHA recommends following the [statewide advisory for lakes and reservoirs without site-specific advice](#).

#### *BLACK BASS SPECIES (LARGEMOUTH BASS)*

The mean mercury and selenium concentrations in Largemouth Bass from Taylor Lake were 56 and 1795 ppb, respectively. OEHHA recommends a maximum of three servings a week of black bass species for the sensitive population (women 18 to 49 years and children 1 to 17 years) based on mercury, and a maximum of four servings a week for the general population (women 50 years and older, and men 18 years and older), based on selenium.

OEHHA has evaluated mercury concentrations in black bass species in many water bodies in California and has found a similar range of mercury concentrations when two or more of these species were caught from the same water body. Selenium levels are presumed to show a similar pattern with comparable concentrations expected to be observed in individual black bass species within the same water body. Therefore, OEHHA extends the consumption advice for Largemouth Bass to other black bass species, including Redeye, Smallmouth, and Spotted Bass.

#### *COMMON CARP*

The mean mercury and selenium concentrations in Common Carp from Taylor Lake were 9 and 1517 ppb, respectively. OEHHA recommends a maximum of four servings a week of Common Carp for both the sensitive population (women 18 to 49 years and children 1 to 17 years) and the general population (women 50 years and older and men 18 years and older), based on selenium.

#### *SUNFISH SPECIES (BLUEGILL, REDEAR SUNFISH)*

The mean mercury and selenium concentrations in sunfish species from Taylor Lake were 15 and 1882 ppb, respectively. Mercury and selenium concentrations for individual sunfish species were as follows, Bluegill (Hg: 17 ppb, Se: 1480 ppb), and Redear Sunfish (Hg: 14, Se: 2003 ppb). Based on the concentration of selenium in these sunfish species, OEHHA recommends a maximum of three servings a week of sunfish species for both the sensitive population (women 18 to 49 years and children 1 to 17 years) and the general population (women 50 years and older and men 18 years and older).

OEHHA has evaluated mercury concentrations in sunfish species in many water bodies in California and has found a similar range of mercury concentrations when two or more of these species were caught from the same water body. Selenium concentrations in sunfish species in this region of California are generally similar within the same water

body. Therefore, OEHHA extends the consumption advice for sunfish species (Bluegill, Redear Sunfish) to other sunfish species, including Green Sunfish and Pumpkinseed.

## RECOMMENDED MAXIMUM NUMBER OF SERVINGS

The recommended maximum numbers of servings per week for fish from Taylor Lake are shown in Table 3.

TABLE 3. RECOMMENDED MAXIMUM NUMBER OF SERVINGS PER WEEK FOR FISH FROM TAYLOR LAKE

Fish Species from Taylor Lake	Women 18–49 years and Children 1-17 years	Women 50 years and older and Men 18 years and older
Black Bass Species	3	4
Common Carp	4	4
Sunfish Species	3	3

## REFERENCES

- American Heart Association. 2016. Fish and Omega-3 Fatty Acids. Online at: [http://www.heart.org/HEARTORG/HealthyLiving/HealthyEating/HealthyDietGoals/Fish-and-Omega-3-Fatty-Acids\\_UCM\\_303248\\_Article.jsp#.Wl57BnlG2Uk](http://www.heart.org/HEARTORG/HealthyLiving/HealthyEating/HealthyDietGoals/Fish-and-Omega-3-Fatty-Acids_UCM_303248_Article.jsp#.Wl57BnlG2Uk).
- Bloom, N.S. 1992. On the chemical form of mercury in edible fish and marine invertebrate tissue. *Can. J. Fish. Aquat. Sci.* 49(5):1010-1017.
- Davis, J.A., J.R.M. Ross, S.N. Bezalel, A. Bonnema, G. Ichikawa, B. Jakl, and W.A. Heim. 2019. Monitoring of Contaminants in Fish from California Lakes and Reservoirs: 2016 Data Report. A Report of the Surface Water Ambient Monitoring Program (SWAMP). California State Water Resources Control Board, Sacramento, CA.
- FDA/US EPA. 2017. Eating Fish: What pregnant women and parents should know. Advice by FDA and US EPA/January, 2017. Online at: <http://www.fda.gov/downloads/Food/FoodborneIllnessContaminants/Metals/UCM537120.pdf>.
- Institute of Medicine. 2007. Seafood choices, balancing benefits and risks. Committee on Nutrient Relationships in Seafood: Selections to Balance Benefits and Risks. Institute of Medicine, Food and Nutrition Board. The National Academies Press, Washington, D.C.
- Kris-Etherton, P.M., W.S. Harris, and L.J. Appel. 2002. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circ.* 106:2747-2757.
- OEHHA. 2005. General Protocol for Sport Fish Sampling and Analysis. Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Sacramento, California. Online at: <http://oehha.ca.gov/media/downloads/fish/document/fishsamplingprotocol2005.pdf>.
- OEHHA. 2008. Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Chlordane, DDTs, dieldrin, methylmercury, PCBs, selenium, and toxaphene. Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Sacramento, California. Online at: <http://oehha.ca.gov/media/downloads/fish/report/atlmhgandothers2008c.pdf>.
- Oken, E., R.O. Wright, K.P. Kleinman, D. Bellinger, C.J. Amarasiriwardena, H. Hu, J.W. Rich-Edwards, and M.W. Gillman. 2005. Maternal fish consumption, hair mercury, and infant cognition in a U.S. cohort. *Environ. Health Perspect.* 113(10):1376-1380.
- Oken, E., J.S. Radesky, R.O. Wright, D. Bellinger, C.J. Amarasiriwardena, K.P. Kleinman, H. Hu, J.W. Rich-Edwards, and M.W. Gillman. 2008. Maternal fish intake

during pregnancy, blood mercury levels, and infant cognition at age 3 years in a U.S. cohort. *Am. J. Epidemiol.* 167(10):1171-1181.

USDA/USDHHS. 2015. 2015-2020 Dietary Guidelines for Americans. 8<sup>th</sup> Edition. U.S. Government Printing Office, Washington, D.C. December. Online at: <http://health.gov/dietaryguidelines/2015/guidelines/>.

US EPA. 1989. Risk Assessment Guidance for Superfund Volume I: Human Health Evaluation Manual (Part A) Interim Final. EPA/5401-89/002, December 1989. Office of Emergency and Remedial Response, U.S. Environmental Protection Agency, Washington, D.C. Online at: <https://rais.ornl.gov/documents/HHEMA.pdf>.

US EPA. 2000a. Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories: Volume 1. Fish Sampling and Analysis. 3<sup>rd</sup> Ed. EPA 823-B00-007. Office of Water, U.S. Environmental Protection Agency, Washington, D.C.

US EPA. 2000b. Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories: Volume 2. Risk Assessment and Fish Consumption Limits, 3<sup>rd</sup> Edition. EPA 823-B-00-007. Office of Water, U.S. Environmental Protection Agency, Washington, D.C.

Weaver, K.L., P. Ivester, J.A. Chilton, M.D. Wilson, P. Pandey, and F.H. Chilton. 2008. The content of favorable and unfavorable polyunsaturated fatty acids found in commonly eaten fish. *J. American Dietetic Assoc.* 108:1178-1185.

## APPENDIX I. ADVISORY TISSUE LEVELS

Advisory Tissue Levels (ATLs) guide the development of advice for people eating sport fish. ATLs are levels of contaminants found in fish that correspond to the maximum numbers of recommended fish servings. OEHHA uses ATLs to provide advice to prevent consumers from being exposed to:

- More than the average daily reference dose<sup>10</sup> for chemicals not known to cause cancer, such as methylmercury, or
- For cancer-causing chemicals, a risk level greater than one additional cancer case in a population of 10,000 people consuming fish at the given consumption rate over a lifetime. This cancer endpoint is the maximum acceptable risk level recommended by the US EPA (2000b) for fish advisories.

For each chemical, ATLs were determined for both cancer and non-cancer risk, if appropriate, for one to seven eight-ounce servings per week. The most health-protective ATLs for each chemical, selected from either cancer or non-cancer based risk, are shown in the table below for zero to seven servings per week. When the guidelines for eating fish from Taylor Lake are followed, exposure to chemicals in fish from Taylor Lake would be at or below the average daily reference dose or the cancer risk probability of one in 10,000.

## ADVISORY TISSUE LEVELS FOR SELECTED ANALYTES

Contaminant	Consumption Frequency Categories (8-ounce servings/week) <sup>a</sup> and ATLs (in ppb)							
	7	6	5	4	3	2	1	0
Chlordanes	≤ 80	>80-90	>90-110	>110-140	>140-190	>190-280	>280-560	>560
DDTs	≤ 220	>220-260	>260-310	>310-390	>390-520	>520-1,000	>1,000-2,100	>2,100
Dieldrin	≤ 7	>7-8	>8-9	>9-11	>11-15	>15-23	>23-46	>46
MeHg (Women 18-49 and children 1-17)	≤ 31	>31-36	>36-44	>44-55	>55-70	>70-150	>150-440	>440
MeHg (Women > 49 and men)	≤ 94	>94-109	>109-130	>130-160	>160-220	>220-440	>440-1,310	>1,310
PBDEs	≤ 45	>45-52	>52-63	>63-78	>78-100	>100-210	>210-630	>630
PCBs	≤ 9	>9-10	>10-13	>13-16	>16-21	>21-42	>42-120	>120
Selenium	≤ 1000	>1,000-1200	>1,200-1,400	>1,400-1,800	>1,800-2,500	>2,500-4,900	>4,900-15,000	>15,000
Toxaphene	≤ 87	>87-100	>100-120	>120-150	>150-200	>200-300	>300-610	>610

<sup>a</sup> Serving sizes (prior to cooking, wet weight) are based on an average 160-pound person. Individuals weighing less than 160 pounds should eat proportionately smaller amounts.

<sup>10</sup> The reference dose is an estimate of the maximum daily exposure to a chemical likely to be without significant risk of harmful health effects over a lifetime.