

Cal/Ecotox Exposure Factors for Stephens' Kangaroo Rat (*Dipodomys stephensi*)*

Page 1

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Age at Fledging, Metamorphosis, Weaning			18 - 22	d	B	Juvenile	Lab	a	1
Age at Sexual Maturity	3			mo	F	Adult	Riverside; CA	b	2
Body Weight - Mean	67.26			g	B	Adult	San Diego; CA	c	1
Body Weight - Mean	62.30	2.40 SD	59.9 - 64.4	g	F	Adult	Lab	d	3
Body Weight - Mean	65.83	1.19 SD	64.5 - 66.8	g	M	Adult	Lab	e	3
Body Weight - Mean	40			g	B	Juvenile	Lab	f	1
Body Weight - Mean	30			g	B	Juvenile	Lab	g	1
Body Weight - Mean	46.1			g	B	Juvenile	Lab	h	1
Body Weight - Mean	20			g	B	Juvenile	Lab	i	1
Body Weight - Mean	10			g	B	Juvenile	Lab	j	1
Body Weight - Mean	4.4		3.8 - 4.7	g	B	Neonate	Lab	k	1
Body Weight - Mean	62.2		45 - 73	g	NR	NR	Riverside; CA	l	4
Clutch or Litter Size	2.67			young/litter	F	Adult	Riverside; CA	m	2
Clutches or Litters per year			1.50 - 1.93	litters/yr	F	Adult	Riverside; CA	n	2
Dietary Composition	Erodium cicutarium leaf (19.4%) and seed (18.8%); Schismus barbatus non-seed (15.3%) and seed (2.8%); unidentified grass (14.1%); Bromus madritensis (7.1%); insect (8.1%); Hirschfeldia incana seed (4.7%); Eriogonum fasciculatum seed (3.5%); Salsola tragus seed (3.4%); Amsinckia menziesii seed (2.4%); Artemisia californica seed (0.9%)			%	NR	NR	Riverside; CA	o	5
Growth Rate	see citation			g	B	Juvenile	Lab	p	1
Growth Rate	19.00 (20-29g); 8.82 (30-39g); 8.80 (40-49g); 0.94 (60-69g)			g/mo	B	NR	Riverside; CA	q	2
Home Range			30.5 - 45.0	m	F	Adult	Riverside; CA	r	6
Home Range			22.7 - 30.0	m	F	Adult	Riverside; CA	s	6
Home Range			78.8 - 86.9	m	M	Adult	Riverside; CA	t	6
Home Range			30.0 - 33.5	m	M	Adult	Riverside; CA	u	6
Home Range			15.0 - 16.5	m	F	Juvenile	Riverside; CA	v	6
Home Range			22.4 - 30.0	m	M	Juvenile	Riverside; CA	w	6
Home Range	review				NR	NR		x	7
Longevity	19			mo	B	Adult	Riverside; CA	y	2
Population Density	36.1			#/ha	B	NR	Riverside; CA	z	8
Population Density	26.2			#/ha	B	NR	Riverside; CA	aa	8
Population Density	19.7			#/ha	B	NR	Riverside; CA	ab	8
Population Density	see citation			#/ha	B	NR	Riverside; CA	ac	2
Population Density	review				NR	NR		ad	7
Population Density			4.9 - 5.2	#/ha	NR	NR	Riverside; CA	ae	9
Population Density			< 1 - >10	#/ha	NR	NR	CA	af	10
Survival/ Mortality			0.79 - 0.85		B	Adult	Riverside; CA	ag	8
Survival/ Mortality			0.52 - 0.81		B	Juvenile	Riverside; CA	ah	8

Notes

a N=NR

04/20/99



Exposure Factors for Stephens' Kangaroo Rat (*Dipodomys stephensi*)

Page 2

b	N=NR; Motte Rimrock Reserve and San Jacinto Wildlife Area
c	N=14; Bonsall and San Luis Rey
d	N=5; captured 3 days prior at Lake Mathews, Riverside County, CA
e	N=3; captured 3 days prior at Lake Mathews, Riverside County, CA
f	N=16; Age=31 d
g	N=16; Age=22 d
h	N=16; Age=63 d
i	N=16; Age=15 d
j	N=16; Age=5 d
k	N=16; Age=0 d
l	N=40
m	N=6; Motte Rimrock Reserve and San Jacinto Wildlife Area
n	range over 2 yrs; N=8-14/yr; Motte Rimrock Reserve and San Jacinto Wildlife Area
o	annual % frequency of occurrence in fecal pellets; N=5-10 pellets/mo; Aug. - Nov.; Motte Rimrock Reserve and San Jacinto Wildlife Area; see citation for monthly changes in diet
p	figure relating body weight or length to age; N=16; Age=0 - 63 d
q	mass specific growth rates pooled by sex and site expressed as change in body mass per mo for 4 weight classes (20-69g); N=1-125/site and sex; Motte Rimrock Reserve and San Jacinto Wildlife Area
r	range of median maximum distance moved based on telemetry at 2 sites; 21 - 91 d observation; N=9; Motte Rock Ecological Reserve [lat., 33°o53'N; long., 117°o6'W] and San Jacinto Wildlife Area [lat., 33°o51'N; long., 116°o54'W]
s	range of median maximum distance moved based on monthly live trapping capture data at 2 sites; N=94; Motte Rock Ecological Reserve [lat., 33°o53'N; long., 117°o6'W] and San Jacinto Wildlife Area [lat., 33°o51'N; long., 116°o54'W]
t	range of median maximum distance moved based on telemetry at 2 sites; 15 - 127 d of observation; N=4-9; Motte Rock Ecological Reserve [lat., 33°o53'N; long., 117°o6'W] and San Jacinto Wildlife Area [lat., 33°o51'N; long., 116°o54'W]
u	range of median maximum distance moved based on monthly live trapping capture data at 2 sites; N=97 - 121; Motte Rock Ecological Reserve [lat., 33°o53'N; long., 117°o6'W] and San Jacinto Wildlife Area [lat., 33°o51'N; long., 116°o54'W]
v	range of median maximum distance moved based on monthly live trapping capture data at 2 sites; N=21-54; Motte Rock Ecological Reserve [lat., 33°o53'N; long., 117°o6'W] and San Jacinto Wildlife Area [lat., 33°o51'N; long., 116°o54'W]
w	range of median maximum distance moved based on monthly live trapping capture data at 2 sites; N=17-59; Motte Rock Ecological Reserve [lat., 33°o53'N; long., 117°o6'W] and San Jacinto Wildlife Area [lat., 33°o51'N; long., 116°o54'W]
x	N=NR
y	maximum time of persistence at site; N=NR; Motte Rimrock Reserve and San Jacinto Wildlife Area
z	average density over 19 mo study; N=125; Lake Mathews [lat., 33°o50'N; long., 117°o27'30"W]; see citation for figure of seasonal changes in population size
aa	average density over 19 mo study; N=96; Motte Rimrock Reserve [lat., 33°o47'30"N; long., 117°o15'W]; see citation for figure of seasonal changes in population size
ab	average density over 19 mo study; N=87; San Jacinto Wildlife Area [lat., 33°o52'N; long., 117°o7'15"W]; see citation for figure of seasonal changes in population size
ac	figure of monthly population densities at two sites; N=NR; Motte Rimrock Reserve and San Jacinto Wildlife Area
ad	N=NR
ae	range of densities estimated over 2 yrs; N=320 - 640 trap nights; Age=Feb. - Mar.; San Jacinto Wildlife Refuge
af	N=NR; Aug. - Dec.
ag	range of minimum rate of survival of individuals in the trappable population per mo, measured over 19 mo at 3 sites; N=110-234; Lake Mathews [lat., 33°o50'N; long., 117°o27'30"W]; Motte Rimrock Reserve [lat., 33°o47'30"N; long., 117°o15'W] and San Jacinto Wildlife Area [lat., 33°o52'N; long., 117°o7'15"W]
ah	range of minimum rate of survival of individuals in the trappable population per mo, measured over 19 mo at 3 sites; N=23-49; Lake Mathews [lat., 33°o50'N; long., 117°o27'30"W]; Motte Rimrock Reserve [lat., 33°o47'30"N; long., 117°o15'W] and San Jacinto Wildlife Area [lat., 33°o52'N; long., 117°o7'15"W]

References

- 1 Lackey, James A. 1967. Growth and development of *Dipodomys stephensi*. *J. Mammal.* 48:624-632.
- 2 Price, Mary V. and Patrick A. Kelly. 1994. An age-structured demographic model for endangered Stephens' kangaroo rat. *Conserv. Biol.* 8:810-821.
- 3 Bleich, Vernon C. and Mary V. Price. 1995. Aggressive behavior of *Dipodomys stephensi*, an endangered species, and *Dipodomys agilis*, a sympatric congener. *J. Mammal.* 76:646-651.
- 4 Price, Mary V., William S. Longland and Ross L. Goldingay. 1991. Niche relationships of *Dipodomys agilis* and *D. stephensi*: two sympatric kangaroo rats of similar size. *Am. Midl. Nat.* 126:172-186.
- 5 Lowe, Margot. 1997. Diet of Stephens' Kangaroo rat, *Dipodomys stephensi*. *Southwest. Nat.* 42:358-361.
- 6 Price, Mary V., Patrick A. Kelly and Ross L. Goldingay. 1994. Distances moved by Stephens' kangaroo rat (*Dipodomys stephensi* Merriam) and implications for conservation. *J. Mammal.* 75:929-939.
- 7 Bleich, Vernon C. 1977. *Dipodomys stephensi*. *Mamm. Species.* 73:1-3.
- 8 McClenaghan, Leroy R. Jr. and Ernest Taylor. 1993. Temporal and spatial demographic patterns in *Dipodomys stephensi* from Riverside County, California. *J. Mammal.* 74:636-645.
- 9 Price, M.V. and P.R. Endo. 1989. Estimating the distribution and abundance of a cryptic species, *Dipodomys stephensi* (Rodentia: Heteromyidae), and implications for management. *Conserv. Biol.* 3:293-301.
- 10 O'Farrell, Michael J. and Curt Uptain. Jul. 1989. Assessment of population and habitat status of the Stephens' kangaroo rat (*Dipodomys stephensi*). Sacramento, CA: California Department of Fish and Game, Nongame Bird and Mammal Section. 198 p.

*Cal/EPA, OEHHA and the University of California Regents are not responsible for damages of any kind resulting from the use of or reliance on information in this report. Users are encouraged to consult the original data. Updated: February 1999.