



# Health Advisory and Guidelines for Eating Fish from Gene Wash Reservoir (San Bernardino County)

June 2024



Fish, Ecotoxicology, and Water Section  
Pesticide and Environmental Toxicology Branch  
Office of Environmental Health Hazard Assessment  
California Environmental Protection Agency

# LIST OF CONTRIBUTORS

## ***Office of Environmental Health Hazard Assessment***

### ***Project Leads***

Loren Chumney, M.S.  
Huyen Tran Pham, M.P.H.

### ***Primary Reviewers***

Wesley Smith, Ph.D., Section Chief  
Shannon R. Murphy, Ph.D.  
Susan A. Klasing, Ph.D. (Retired Annuitant)

### ***Final Reviewers***

Elaine Khan, Ph.D., Pesticide and Environmental Toxicology Branch Chief  
Martha Sandy, Ph.D., Acting Deputy Director for Scientific Programs  
David Edwards, Ph.D., Chief Deputy Director

### ***Director***

Lauren Zeise, Ph.D.

# ACKNOWLEDGMENTS

Developing fish consumption advisories depends on sampling and analysis of fish. The Office of Environmental Health Hazard Assessment acknowledges the contribution of information from the following entities: the State Water Resources Control Board, the California Department of Fish and Wildlife, and the Marine Pollution Studies Laboratory at Moss Landing Marine Laboratories. Data were obtained from the [California Environmental Data Exchange Network](#). The map was created using ArcMap (10.5) from Environmental Systems Resource Institute (ESRI, Redlands, California).

### ***For further information, contact:***

Pesticide and Environmental Toxicology Branch  
Office of Environmental Health Hazard Assessment  
California Environmental Protection Agency

1001 I Street, P.O. Box 4010  
Sacramento, CA 95812-4010  
Telephone: (916) 324-7572  
Email address: [fish@oehha.ca.gov](mailto:fish@oehha.ca.gov)

1515 Clay Street, 16<sup>th</sup> Floor  
Oakland, CA 94612  
Telephone: (510) 622-3170

## LIST OF ACRONYMS AND ABBREVIATIONS

ATL	Advisory Tissue Level
CDFW	California Department of Fish and Wildlife
CEDEN	California Environmental Data Exchange Network
CRRWQCB	Colorado River Regional Water Quality Control Board (Region 7)
DDT(s)	dichlorodiphenyltrichloroethane (DDT) and its metabolites dichlorodiphenyldichloroethane (DDD) and dichlorodiphenyldichloroethylene (DDE)
DHA	docosahexaenoic acid
DMA	direct mercury analyzer
EPA	eicosapentaenoic acid
FDA	United States Food and Drug Administration
Hg	mercury
ICP-MS	inductively coupled plasma-mass spectrometry
MDL	method detection limit
MeHg	methylmercury
mm	millimeters
MPSL	Marine Pollution Studies Laboratory at Moss Landing Marine Laboratories
OEHHA	Office of Environmental Health Hazard Assessment
PBDEs	polybrominated diphenyl ethers
PCBs	polychlorinated biphenyls
ppb	parts per billion
RL	reporting limit
Se	selenium
SWAMP	Surface Water Ambient Monitoring Program
SWRCB	State Water Resources Control Board
USDA	United States Department of Agriculture
USDHHS	United States Department of Health and Human Services
US EPA	United States Environmental Protection Agency

## PREFACE

The Office of Environmental Health Hazard Assessment (OEHHA), a department in the California Environmental Protection Agency, is responsible for evaluating potential public health risks from chemical contamination of sport fish.<sup>1</sup> This includes issuing fish consumption advisories, when appropriate, for the State of California. OEHHA's authorities to conduct these activities are based on mandates in the:

- California Health and Safety Code
  - Section 59009, to protect public health
  - Section 59011, to advise local health authorities
- California Water Code
  - Section 13177.5, to issue health advisories.

The health advisories are published in the California Department of Fish and Wildlife's (CDFW) Inland and Ocean Sport Fishing Regulations in their respective sections on public health advisories.<sup>2</sup>

This report presents guidelines for eating fish from Gene Wash Reservoir in San Bernardino County. The report provides background information and a technical description of how the guidelines were developed. The resulting advice is summarized in the illustrations after the Table of Contents and the List of Figures and Tables.

---

<sup>1</sup> Sport fish includes all fish and shellfish caught from California waters for non-commercial purposes (e.g., recreational, tribal/cultural, and subsistence practices).


<sup>2</sup> CDFW's Inland and Ocean Sport Fishing Regulations can be found online at: <https://wildlife.ca.gov/Fishing/Inland> and <https://wildlife.ca.gov/Fishing/Ocean>, respectively.

## TABLE OF CONTENTS

A GUIDE TO EATING FISH FROM GENE WASH RESERVOIR .....	5
INTRODUCTION .....	6
<i>Location</i> .....	6
<i>Approach Used</i> .....	7
CHEMICALS OF CONCERN.....	7
DATA SOURCES.....	8
<i>Contaminants in Fish From California Lakes and Reservoirs, 2007–2008 (SWAMP)</i> .....	9
<i>Long-Term Monitoring of Bass Lakes and Reservoirs in California, 2015–ongoing (SWAMP)</i> .....	9
FISH SAMPLED FROM GENE WASH RESERVOIR.....	9
CHEMICAL CONCENTRATIONS.....	10
<i>Mercury</i> .....	10
<i>PCBs, PBDEs, and Pesticides</i> .....	11
<i>Selenium</i> .....	11
DEVELOPMENT OF GUIDELINES FOR EATING FISH FROM GENE WASH RESERVOIR ...	12
CONSUMPTION ADVICE FOR FISH FROM GENE WASH RESERVOIR .....	15
<i>Black Bass Species (Based on Largemouth Bass)</i> .....	16
<i>Common Carp</i> .....	16
<i>Flathead Catfish</i> .....	16
<i>Sunfish Species (Based on Redear Sunfish)</i> .....	16
RECOMMENDED MAXIMUM NUMBER OF SERVINGS.....	17
REFERENCES .....	18
APPENDIX. Advisory Tissue Levels .....	21


## LIST OF FIGURES AND TABLES

Figure 1. Location of Gene Wash Reservoir .....	6
Table 1. Fish Samples Evaluated for the Gene Wash Reservoir Advisory.....	10
Table 2. Mercury Concentrations in Fish from Gene Wash Reservoir.....	12
Table 3. Selenium Concentrations in Fish from Gene Wash Reservoir.....	12
Table 4. Recommended Maximum Number of Servings per Week for Fish from Gene Wash Reservoir.....	17
Advisory Tissue Levels for Selected Analytes.....	21



**Women**  
(18 – 49 Years)

**Children**  
(1 – 17 Years)



**Women**  
(50+ Years)

**Men**  
(18+ Years)


## A GUIDE TO EATING FISH

*from*


# GENE WASH RESERVOIR

(SAN BERNARDINO COUNTY)


**Eat the Good Fish**  
Eating fish that are low in chemicals may provide health benefits to children and adults.




**Avoid the Bad Fish**  
Eating fish with higher levels of chemicals like mercury or PCBs may cause health problems in children and adults.



**Choose the Right Fish**  
Chemicals may be more harmful to unborn babies and children.




Common Carp




Sunfish Species

---



**Black Bass Species**  
*♥ high in omega-3s*


---




Flathead Catfish

**Serving Size**  
A serving of fish is about the size and thickness of your hand. Give children smaller servings.

**For Adults**




**For Children**




California Office of Environmental Health Hazard Assessment

web [www.oehha.ca.gov/fish](http://www.oehha.ca.gov/fish)  
email [fish@oehha.ca.gov](mailto:fish@oehha.ca.gov)  
phone (916) 324-7572

**Eat only the skinless fillet**



**Eat only the meat**



Some chemicals are higher in the skin, fat, and guts.

## INTRODUCTION

This report presents guidelines for eating black bass species, Common Carp, Flathead Catfish, and sunfish species from Gene Wash Reservoir (Figure 1). Consumption advice is based on levels of mercury and selenium found in these species.

### LOCATION

Gene Wash Reservoir is located near the California-Arizona border, by the Colorado River. The reservoir was formed by construction of Gene Wash Dam in 1937, which is owned and operated by the Metropolitan Water District of Southern California. As part of the Colorado River Aqueduct System, the reservoir stores water for use in the Los Angeles and San Diego areas.<sup>3</sup>

FIGURE 1. LOCATION OF GENE WASH RESERVOIR



<sup>3</sup> Information regarding Gene Wash Reservoir was obtained from the US Department of Energy, online at: <https://www.osti.gov/biblio/192168>.



### *APPROACH USED*

The Office of Environmental Health Hazard Assessment (OEHHA) used the results from two monitoring studies described in this report to develop the Gene Wash Reservoir Advisory. OEHHA uses the following general process in developing consumption advice for sport fish:

- 1) Evaluation of all fish contaminant data available from a water body and selection of appropriate data that meet data quality criteria and sampling plan guidelines.
- 2) Determination of fish species for which adequate data are available to issue fish consumption advice.
- 3) Calculation of an appropriate measure of central tendency (often a weighted arithmetic mean)<sup>4</sup> and other descriptive statistics of the contaminant data, as appropriate, for a chemical of concern for the selected fish species.
- 4) Comparison of the chemical concentrations with the OEHHA Advisory Tissue Levels (ATLs) for each chemical of concern.
- 5) Development of final advice based on a thorough review of the data and best professional judgment relating to the benefits and risks of consuming a particular fish species.

The ATLs (discussed further in a subsequent section of this report) are chemical levels in fish tissue that are considered acceptable, based on chemical toxicity, for a range of consumption rates. Development of the ATLs also includes consideration of health benefits associated with including fish in the diet (OEHHA, 2008). The ATLs should not be interpreted as static “bright lines,” but as one component of a complex process of data evaluation and interpretation used by OEHHA in the assessment and communication of the benefits and risks of consuming sport fish.

## CHEMICALS OF CONCERN

Certain chemicals, because of their toxicity and their ability to accumulate in fish tissue, are of concern for people who eat fish. The majority of fish consumption advisories in California are issued because of mercury (Hg), followed by polychlorinated biphenyls (PCBs) and, in a few cases, selenium (Se), polybrominated diphenyl ethers (PBDEs), or some legacy pesticides (pesticides that are no longer used but remain in the environment).

Mercury is an element found in some rocks and soil. Human activities, such as burning coal and the historical use of mercury to mine gold, also add mercury to the environment. If mercury enters waterways, it can be converted to a more toxic form known as methylmercury (MeHg) – which can pass into and build up in fish. High levels

---

<sup>4</sup> Means are an arithmetic average of individual values and/or composites weighted by number of fish. A weighted average of composites is calculated by multiplying the chemical concentration in each composite by the number of fish in that composite for each species. Products are then summed and divided by the total number of fish in all composites for that species.

of methylmercury can harm the brain, especially in fetuses and children, whose brains are still developing.

PCBs are industrial chemicals previously used in electrical transformers, plastics, and lubricating oils, and were often used as flame retardants or electrical insulators. Their use was banned in the 1970s, but they can accumulate in fish because they do not break down easily and they persist in the environment. Depending on the exposure level, PCBs may cause cancer or other health effects, including neurotoxicity, in humans.

Selenium is an element and at low doses is an essential nutrient for many important human health processes, including thyroid regulation and vitamin C metabolism. Higher doses cause selenium toxicity, which can include symptoms ranging from hair loss and gastrointestinal distress to dizziness and tremors.

PBDEs are a class of flame retardants historically used in a variety of consumer products, including furniture, textiles, automotive parts, and electronics. The use of PBDEs in new products was largely phased out by 2013 but, due to their wide usage and persistence in the environment, they are still being detected in fish tissues. PBDEs may affect hormone levels or learning and behavior in children.

Chlordanes, dichlorodiphenyltrichloroethane (DDT), dieldrin, and toxaphene are pesticides that were banned from use in 1973 (DDT), the late 1980s (chlordanes and dieldrin) and 1990 (toxaphene), but are still found in some fish in certain California water bodies. Depending on the exposure level, these chemicals may cause cancer or adverse effects on the nervous system.

A detailed discussion of the toxicity of these chemicals is presented in “Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Chlordane, DDTs, Dieldrin, Methylmercury, PCBs, Selenium, and Toxaphene” (OEHHA, 2008) and “Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Polybrominated Diphenyl Ethers (PBDEs)” (OEHHA, 2011).

All fish species collected from Gene Wash Reservoir and used in advisory development were analyzed for mercury and selenium. Common Carp were additionally analyzed for PCBs, PBDEs, and legacy pesticides as indicated in Table 1.

## DATA SOURCES

The guidelines for eating fish from Gene Wash Reservoir are based on the chemicals detected in the fish collected for the two monitoring studies described below. These studies met OEHHA’s data quality criteria, including adequate documentation of sample collection, fish preparation methods (e.g., skinning or filleting), chemical analyses, quality assurance, and sufficiently low detection limits. “Sample,” as used in this report, refers to an individual fish or a composite of multiple fish for which contaminant data

were reported. “Sampling” or “sampled” refers to the act of collecting fish for chemical analysis. The studies or entities contributing data to this advisory are described below.

*CONTAMINANTS IN FISH FROM CALIFORNIA LAKES AND RESERVOIRS, 2007–2008 (SWAMP)*

The Surface Water Ambient Monitoring Program (SWAMP), operated by the State Water Resources Control Board (SWRCB) in cooperation with the Colorado River Regional Water Quality Control Board (CRRWQCB), monitors water quality in California’s surface waters. This survey of inland water bodies, conducted by SWAMP, was the State’s largest survey of chemical contaminants in sport fish. The survey sampled popular fishing sites at 272 lakes and reservoirs from 2007 to 2008 (SWRCB, 2010). The SWRCB used the data from this survey to characterize statewide water quality conditions. The program collected Common Carp and Largemouth Bass from Gene Wash Reservoir in 2007, which were analyzed for mercury. Common Carp were also analyzed for chlordanes, DDTs, dieldrin, PBDEs, PCBs, and selenium.

*LONG-TERM MONITORING OF BASS LAKES AND RESERVOIRS IN CALIFORNIA, 2015–ONGOING (SWAMP)*

This monitoring study is a multi-year effort initiated in 2015 to document the status and trends related to contamination in sport fish from California lakes and reservoirs where bass species reside.<sup>5</sup> In 2021, the program collected Flathead Catfish, Largemouth Bass, and Redear Sunfish from Gene Wash Reservoir, which were analyzed for mercury and selenium.

## FISH SAMPLED FROM GENE WASH RESERVOIR

The fish sampling data used in this advisory were retrieved from the California Environmental Data Exchange Network (CEDEN),<sup>6</sup> the state’s repository for environmental data. Samples were excluded when the fish were not legal size to take or did not meet OEHHA’s criteria for minimum “edible” size, based on species size at maturity and professional judgment (as described in OEHHA, 2022). A summary of all fish species evaluated for this advisory is shown in Table 1, including the name of the species, number of samples collected, total number of fish, project name, year sampled, and contaminants analyzed.

---

<sup>5</sup> Information on SWAMP’s Lake and Reservoir Bioaccumulation Monitoring Surveys can be found online at: [https://www.waterboards.ca.gov/water\\_issues/programs/swamp/lakes\\_study.html](https://www.waterboards.ca.gov/water_issues/programs/swamp/lakes_study.html).

<sup>6</sup> Online at: <http://ceden.waterboards.ca.gov/AdvancedQueryTool>.

TABLE 1. FISH SAMPLES EVALUATED FOR THE GENE WASH RESERVOIR ADVISORY

Common Name	Scientific Name	Number of Samples	Total Number of Fish	Project	Year Collected	Contaminants Analyzed
Common Carp	<i>Cyprinus carpio</i>	1	5	SWAMP	2007	Chlordanes, DDTs, Dieldrin, PBDEs, PCBs
		2	10	SWAMP	2007	Hg, Se
Flathead Catfish	<i>Pylodictis olivaris</i>	2	9	SWAMP	2021	Hg, Se
Largemouth Bass	<i>Micropterus salmoides</i>	7	7	SWAMP	2007	Hg
		6	6	SWAMP	2021	Hg
		1	7	SWAMP	2021	Se
Redear Sunfish	<i>Lepomis microlophus</i>	2	10	SWAMP	2021	Hg, Se

All fish were analyzed as skinless fillets.

## CHEMICAL CONCENTRATIONS

As shown in Table 1, samples were analyzed for one or more of the following: total mercury, selenium, chlordanes (5 congeners), DDTs (6 congeners), dieldrin, PBDEs (7 congeners), and PCBs (54 congeners).<sup>7</sup> Among the chemicals analyzed in fish tissue samples from Gene Wash Reservoir, only mercury and selenium levels were sufficiently high to impact consumption advice.

All fish samples were prepared as skinless fillets and analyzed as individual fish or composites.

For this advisory, OEHHA used the weighted (by the number of individual fish) average (arithmetic mean) of the chemical concentrations (in wet weight) for each fish species to estimate average human exposure.

### MERCURY

Samples were analyzed for total mercury, as either individual fish or composite samples, using a direct mercury analyzer (DMA) at the Marine Pollution Studies

<sup>7</sup> Congeners are related compounds with similar chemical forms. Five and six congeners are typically analyzed for chlordanes and DDTs, respectively. Of the 209 possible PBDE and PCB congeners, 6–7 and 48–54 are generally analyzed, respectively. See the OEHHA (2022) Sampling Protocol available online at

<https://oehha.ca.gov/media/downloads/fish/report/fishadvisorysamplinganalysisprotocolreport2022.pdf>.

Laboratory (MPSL) at Moss Landing Marine Laboratories. The DMA method utilizes thermal decomposition and atomic absorption. OEHHA assumed all mercury detected was methylmercury, which is the most common form found in fish and is also the more toxic form (Bloom, 1992). Table 2 shows the averages and ranges for total length,<sup>8</sup> as well as mercury concentrations in each fish species. Depending on the study, the DMA method detection limits (MDLs)<sup>9</sup> for total mercury were reported at 3 or 12 parts per billion (ppb), and the reporting limits (RLs)<sup>10</sup> were 10 or 12 ppb.

### *PCBs, PBDEs, AND PESTICIDES*

Pesticides, PBDEs, and PCBs in either individual fish or composite samples were analyzed by gas chromatography at the CDFW Water Pollution Control Laboratory. Where applicable, the concentrations presented were the sum of the detected analytes (parent compound, congeners, or metabolites) for chlordanes, DDTs, PCBs, and PBDEs. Because the MDLs or RLs were relatively low ( $\leq 5$  ppb), individual congeners or metabolites with concentrations reported as non-detects were assumed to be zero. This is a standard method of handling non-detect values for PCBs and other chemicals with multiple congeners or metabolites in a given sample when detection levels are adequate (US EPA, 2000a).

### *SELENIUM*

The MPSL analyzed species collected from Gene Wash Reservoir for selenium as composite samples, using inductively coupled plasma-mass spectrometry (ICP-MS). The ICP-MS method uses desolvation, atomization, and ionization with ion separation based on a mass-to-charge ratio to detect the total selenium concentration in a sample. Depending on the study, the MDLs for selenium were reported at 100, 150, or 230 ppb, and the RLs were 300, 400, or 700 ppb. Table 3 shows the averages and ranges for total length, as well as selenium concentrations in each fish species.

Concentrations of chlordanes, dieldrin, DDTs, PBDEs, and PCBs were lower than the corresponding ATL threshold values for daily consumption (OEHHA, 2008 and 2011). With the exception of assessing for multiple chemical exposures, these chemicals were not considered further for developing consumption advice and are not shown in this report.

---

<sup>8</sup> Total length is the maximum length of the fish, measured from the tip of the closed mouth to the tip of the pinched tail fin.

<sup>9</sup> The MDL is the lowest quantity of a chemical that can be distinguished (as greater than zero) in a sample.

<sup>10</sup> The RL is the lowest quantity of a chemical that can be accurately quantified in a sample.

TABLE 2. MERCURY CONCENTRATIONS IN FISH FROM GENE WASH RESERVOIR

Species from Gene Wash Reservoir	Number of Samples <sup>a</sup>	Total Number of Fish	Mean <sup>b</sup> Total Length (mm)	Range of Total Lengths <sup>c</sup> (mm)	Mercury (ppb)	
					Mean <sup>b</sup>	Range <sup>c</sup>
Common Carp	2	10	625	540 – 682	14	13 – 15
Flathead Catfish	2	9	614	505 – 825	200	133 – 254
Largemouth Bass	13	13	449	346 – 521	119	32 – 248
Redear Sunfish	2	10	167	152 – 186	20	19 – 21

<sup>a</sup>Samples were prepared as skinless fillets.

<sup>b</sup>Means are an arithmetic average of individual values and/or a weighted average of composites.

<sup>c</sup>Range of individuals and/or range of the composites.

TABLE 3. SELENIUM CONCENTRATIONS IN FISH FROM GENE WASH RESERVOIR

Species from Gene Wash Reservoir	Number of Samples <sup>a</sup>	Total Number of Fish	Mean <sup>b</sup> Total Length (mm)	Range of Total Lengths <sup>c</sup> (mm)	Selenium (ppb)	
					Mean <sup>b</sup>	Range <sup>c</sup>
Common Carp	2	10	625	540 – 682	2,135	1,600 – 2,670
Flathead Catfish	2	9	614	505 – 825	892	870 – 910
Largemouth Bass	1	7	466	308 – 521 <sup>d</sup>	1,540	n/a
Redear Sunfish	2	10	167	152 – 186	2,165	2,100 – 2,230

<sup>a</sup>Samples were prepared as skinless fillets.

<sup>b</sup>Means are an arithmetic average of individual values and/or a weighted average of composites.

<sup>c</sup>Range of individuals and/or range of the composites.

<sup>d</sup>Composite includes fish below legal size for black bass in the Colorado River District (330 mm), but above legal size (305 mm) for most other California water bodies. These data were included in the evaluation because it was health-protective to consider the contribution of selenium to the toxicity of this species.

n/a = not applicable due to a single sample

## DEVELOPMENT OF GUIDELINES FOR EATING FISH FROM GENE WASH RESERVOIR

The OEHHA fish advisory process considers the health benefits of fish consumption as well as the risk from exposure to the chemical contaminants found in fish. Benefits are included in the advisory process because there is considerable evidence and scientific consensus that fish should be part of a healthy well-balanced diet. Fish contain many nutrients that are important for general health and, in particular, help promote optimal growth and development of babies and young children, and may reduce the incidence of heart disease in adults (FDA/US EPA, 2017; American Heart Association, 2016; OEHHA, 2008; Institute of Medicine, 2007; Kris-Etherton et al., 2002). Fish are a

significant source of the beneficial omega-3 fatty acids, docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA) (USDA/USDHHS, 2020; Weaver et al., 2008).

As part of a healthy US-style dietary pattern at the 2,000-calorie level, the “Dietary Guidelines for Americans, 2020 – 2025” (USDA/USDHHS, 2020) recommends consuming eight ounces of seafood<sup>11</sup> per week. Young children are advised to eat proportionately smaller amounts. “Women who are pregnant or lactating should consume at least 8 and up to 12 ounces of a variety of seafood per week from choices that are lower in methylmercury.” Additionally, “based on FDA [US Food and Drug Administration] and EPA’s [US Environmental Protection Agency] advice,<sup>12</sup> depending on body weight, some women should choose seafood lowest in methylmercury or eat less seafood than the amounts in the Healthy U.S.-Style Dietary Pattern” and avoid certain species (USDA/USDHHS, 2020). The species of fish that people eat is an important factor in determining the net beneficial effects of fish consumption. For example, studies have shown that children of mothers who ate low-mercury fish during pregnancy scored better on cognitive tests compared to children of mothers who did not eat fish or ate high-mercury fish (Oken et al., 2005 and 2008). Accordingly, because of the high mercury content of certain fish species, the FDA and the US EPA recommend that women who are pregnant (or might become pregnant) or breastfeeding, and young children avoid consuming shark, Swordfish, tilefish (Gulf of Mexico), Bigeye Tuna, marlin, Orange Roughy, and King Mackerel (FDA/US EPA, 2017).

To address the potential health concerns associated with exposure to contaminants in sport fish, OEHHA has established ATLS for chemicals that are known to accumulate in the edible tissues of fish. ATLS consider both the toxicity of the chemical and potential benefits of eating fish. OEHHA uses the ATLS to determine the maximum number of servings per week that consumers can eat safely, for each species and from each location, to limit their exposure to these contaminants. Consumers can use OEHHA’s guidance when choosing which fish and how much to eat as part of an overall healthy diet.

There are two sets of ATLS for methylmercury in fish because of the age-related toxicity of this chemical (OEHHA, 2008). The fetus and children are more sensitive to the toxic effects of methylmercury. Thus, the ATLS for the sensitive population, including women who might become pregnant (typically 18–49 years of age) and children 1–17 years of age, are lower than those for women 50 years and older and men 18 years and older. The lower ATL values for the sensitive population provide additional protection to allow for normal growth and development of the brain and nervous system of unborn babies and children. Detailed discussion about the toxicity of common fish contaminants and health benefits of fish consumption, as well as derivation of the ATLS, are provided in “Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Chlordane, DDTs, Dieldrin, Methylmercury, PCBs, Selenium, and Toxaphene” (OEHHA, 2008) and “Development of Fish

---

<sup>11</sup> Seafood as used here refers to fish and shellfish from freshwater and marine environments.

<sup>12</sup> Online at: <https://www.fda.gov/food/consumers/advice-about-eating-fish>.

Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Polybrominated Diphenyl Ethers (PBDEs)” (OEHHA, 2011). A list of the ATLs used in this report is presented in the Appendix.

For each fish species in this advisory, OEHHA compared the mean chemical concentrations detected in the fillet to the corresponding ATLs to establish the maximum number of servings per week that can be safely consumed (see Appendix). For fish fillets, a serving size is considered to be 8 ounces, prior to cooking, or about the size and thickness of a hand. Children should be given smaller servings. For smaller fish species, several individual fish may be required to yield a serving.

The consumption advice for a fish species is initially based on the chemical with the lowest allowable number of servings per week. Because some chemicals, such as mercury and PCBs, are known to have similar adverse effects, additivity of toxicity is assumed in such cases and may be assessed using multiple chemical exposure methodology (US EPA, 1989 and 2000b). If two or more chemicals with similar adverse effects are present in fish tissue, multiple chemical exposure methodology involving hazard index calculations is employed. This may result in advising fewer servings per week than would be the case for the presence of either chemical alone, in a similar concentration. The potential effect of multiple chemical exposures (mercury and PCBs) was assessed in Common Carp, the only species for which PCBs were analyzed, and did not affect advice. Advice for all species in this advisory was based solely on mercury or selenium concentrations.

OEHHA recommends that individuals strive to meet the US dietary guidelines’ seafood consumption recommendations,<sup>13</sup> while also adhering to federal and OEHHA recommendations to limit the consumption of fish with higher contaminant levels. The advice discussed in the following section represents the maximum recommended number of servings per week for listed fish species. People should eat no more than the recommended number of servings for each fish species or species group. When noted, OEHHA’s consumption advice for a particular fish species can be extended to other closely related fish species<sup>14</sup> known to accumulate similar levels of contaminants.

Consumption advice should not be combined. That is, if a person chooses to eat a serving of fish from the “one-serving-a-week” category, then they should not eat any other fish from any source (including commercial) until the next week. If a person chooses to eat a serving of fish from the “two-servings-per-week” category, they can combine fish species from that category, or eat one serving of fish from that category and one from a category that recommends more than two servings per week (if available), for a total of two servings in that week. Then they should not eat any other fish from any source (including commercial) until the following week.

---

<sup>13</sup> Online at: <https://www.dietaryguidelines.gov/>.

<sup>14</sup> Fish species within the same genus are most closely related, and family is the next level of relationship.



## CONSUMPTION ADVICE FOR FISH FROM GENE WASH RESERVOIR

OEHHA's sampling and analysis protocol (OEHHA, 2022) requires that a minimum of nine edible-size fish of a species that may be legally caught are collected and analyzed before an advisory can be developed for the primary contaminant of concern. Additional fish beyond this number will increase confidence that the sample dataset is representative of the fish species population in the water body (OEHHA, 2022). The majority of fish consumption advisories in California are based on mercury, which is typically analyzed in individual fish, rather than as composites. Mercury analysis is relatively inexpensive and mercury concentrations in fish are more likely to be positively correlated with fish size than other contaminants. Other contaminants, such as PCBs, pesticides, and selenium, may also impact advice. These contaminants are often analyzed as a composite of a smaller subset of fish (usually at least five individuals) as a cost-saving mechanism, a common practice that is considered acceptable. In some cases, an exception is made regarding the minimum sample size. This is particularly true if the advice is based on a chemical other than mercury where sample size is often limited, and/or if doing so leads to more health-protective advice than would otherwise be provided.

For Gene Wash Reservoir, the sample size criterion was met for mercury for all species. However, Largemouth Bass advice for the general population was based on a selenium analysis using data from only one composite of seven fish. The composite included fish below the CDFW minimum legal take length 13 inches (330 mm) within the Colorado River District, but above the typical legal size of 12 inches (305 mm) for black bass in most other water bodies in the state. Although the sample contained slightly fewer than the preferred number of fish, the data were included because they reduced the number of recommended servings of Largemouth Bass for the general population from five to four per week, compared to the advice based on mercury, and thus provided more health protective advice.

There were not sufficient data to evaluate other species that may be found in this water body. For fish species found in Gene Wash Reservoir that are not included in this advisory, OEHHA recommends following the statewide advisory for lakes and reservoirs without site-specific advice.<sup>15</sup>

The following advice is based solely on mercury or selenium concentrations. The sensitive population is defined as women ages 18 to 49 years and children ages 1 to 17 years, and the general population is defined as women 50 years and older and men 18 years and older.

---

<sup>15</sup> Online at: <https://oehha.ca.gov/advisories/statewide-advisory-eating-fish-californias-lakes-and-reservoirs-without-site-specific>.

*BLACK BASS SPECIES (BASED ON LARGEMOUTH BASS)*

Black bass species are one of the most targeted species of freshwater game fish in California. OEHHA groups black bass species because they have similar predatory diets which suggests a comparable chemical uptake (Long and Fisher, 2000). They are also known to hybridize (Pierce and Van Den Avyle, 1997), largely due to species introductions for angling purposes and weak genetic barriers between members of the genus (Thongda et al., 2020). OEHHA has also evaluated mercury concentrations in black bass species in many water bodies in California and has found a similar range of mercury concentrations when two or more of these species were caught from the same water body. Selenium levels are also presumed to show a similar pattern with comparable concentrations expected to be observed in individual black bass species within the same water body. Therefore, OEHHA extends the consumption advice for Largemouth Bass to other black bass species, including Redeye, Smallmouth, and Spotted Bass.

The mean mercury and selenium concentrations in Largemouth Bass from Gene Wash Reservoir were 119 and 1,540 parts per billion (ppb), respectively. OEHHA recommends a maximum of two servings per week of Largemouth Bass for the sensitive population based on mercury, and a maximum of four servings per week for the general population, based on selenium.

*COMMON CARP*

The mean mercury and selenium concentrations in Common Carp from Gene Wash Reservoir were 14 and 2,135 ppb, respectively. Based on selenium, OEHHA recommends a maximum of three servings per week of Common Carp for both the sensitive and general populations.

*FLATHEAD CATFISH*

The mean mercury and selenium concentrations in Flathead Catfish from Gene Wash Reservoir were 200 and 892 ppb, respectively. Based on mercury, OEHHA recommends a maximum of one serving per week of Flathead Catfish for the sensitive population, and a maximum of three servings per week for general population.

*SUNFISH SPECIES (BASED ON REDEAR SUNFISH)*

OEHHA groups sunfish species due to a known ability to hybridize (Avisé and Smith, 1974) and extensive dietary overlap (Kirby, 1982), which suggests a similar contaminant uptake. OEHHA has evaluated mercury concentrations in sunfish species from many water bodies in California and has found a similar range of mercury concentrations when two or more of these species were caught from the same water body. Selenium concentrations in sunfish species in this region of California are generally similar within the same water body. Therefore, OEHHA extends the consumption advice for Redear Sunfish to other sunfish species, including Bluegill, Green Sunfish, and Pumpkinseed.

The mean mercury and selenium concentrations in Redear Sunfish from Gene Wash Reservoir were 20 and 2,165 ppb, respectively. Based on selenium, OEHHA recommends a maximum of three servings per week of sunfish species for both the sensitive and general populations.

## RECOMMENDED MAXIMUM NUMBER OF SERVINGS

The recommended maximum numbers of servings per week for fish from Gene Wash Reservoir are shown in Table 4.

TABLE 4. RECOMMENDED MAXIMUM NUMBER OF SERVINGS PER WEEK FOR FISH FROM GENE WASH RESERVOIR

Species from Gene Wash Reservoir	Women 18–49 years and Children 1–17 years		Women 50 years and older and Men 18 years and older	
	Number of Servings	Risk Driver	Number of Servings	Risk Driver
Black Bass Species	2	Hg	4	Se
Common Carp	3	Se	3	Se
Flathead Catfish	1	Hg	3	Hg
Sunfish Species	3	Se	3	Se

## REFERENCES

- American Heart Association. 2016. Fish and Omega-3 Fatty Acids. Online at: <https://www.heart.org/en/healthy-living/healthy-eating/eat-smart/fats/fish-and-omega-3-fatty-acids>.
- Awise, J.C. and M.H. Smith. 1974. Biochemical Genetics of Sunfish II. Genic Similarity between Hybridizing Species. *The American Naturalist*. 108 (962). Online at: <https://www.journals.uchicago.edu/doi/pdf/10.1086/282926>.
- Bloom, N.S. 1992. On the chemical form of mercury in edible fish and marine invertebrate tissue. *Can. J. Fish. Aquat. Sci.* 49(5):1010–1017.
- FDA/US EPA. 2017. Eating Fish: What pregnant women and parents should know. Advice by FDA and US EPA/January, 2017. Online at: <https://www.fda.gov/food/consumers/advice-about-eating-fish>.
- Institute of Medicine. 2007. Seafood choices, balancing benefits and risks. Committee on Nutrient Relationships in Seafood: Selections to Balance Benefits and Risks. Institute of Medicine, Food and Nutrition Board. The National Academies Press, Washington, D.C.
- Kirby, J.M. 1982. Prey Utilization among Four Sympatric Species of Sunfish. *Proc. PA. Acad. of Sci.* 56(2):147–150. Online at: <https://www.jstor.org/stable/44111415>.
- Kris-Etherton, P.M., W.S. Harris, and L.J. Appel. 2002. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circ.* 106:2747–2757.
- Long, J.M. and W.L. Fisher. 2000. Inter-Annual and Size-Related Differences in the Diets of Three Sympatric Black Bass in an Oklahoma Reservoir. *J. Freshw. Ecol.* 15(4): 465–474. Online at: <https://www.tandfonline.com/doi/abs/10.1080/02705060.2000.9663768>.
- OEHHA. 2008. Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Chlordane, DDTs, Dieldrin, Methylmercury, PCBs, Selenium, and Toxaphene. Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Sacramento, California. Online at: <http://oehha.ca.gov/media/downloads/fish/report/atlmhgandothers2008c.pdf>.
- OEHHA. 2011. Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Polybrominated diphenyl ethers (PBDEs). Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Sacramento, California. Online at: <http://oehha.ca.gov/media/downloads/fish/report/pbdes052311.pdf>.

OEHHA. 2022. General Protocol for Sport Fish Sampling and Analysis. Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Sacramento, California. Online at: <https://oehha.ca.gov/media/downloads/fish/report/fishadvisorysamplinganalysisprotocolreport2022.pdf>.

Oken, E., R.O. Wright, K.P. Kleinman, D. Bellinger, C.J. Amarasiriwardena, H. Hu, J.W. Rich-Edwards, and M.W. Gillman. 2005. Maternal fish consumption, hair mercury, and infant cognition in a U.S. cohort. *Environ. Health Perspect.* 113(10):1376–1380.

Oken, E., J.S. Radesky, R.O. Wright, D. Bellinger, C.J. Amarasiriwardena, K.P. Kleinman, H. Hu, J.W. Rich-Edwards, and M.W. Gillman. 2008. Maternal fish intake during pregnancy, blood mercury levels, and infant cognition at age 3 years in a U.S. cohort. *Am. J. Epidemiol.* 167(10):1171–1181.

Pierce, P.C., and M.J. Van Den Avyle. 1997. Hybridization between Introduced Spotted Bass and Smallmouth Bass in Reservoirs. *Trans. Am. Fish. Soc.* 126(6):939–947. Available online at: <https://www.tandfonline.com/doi/abs/10.1577/1548-8659%281997%29126%3C0939%3AHBISBA%3E2.3.CO%3B2>.

SWRCB. 2010. Contaminants in Fish from California Lakes and Reservoirs, 2007–2008: Summary Report on a Two-Year Screening Survey. State Water Resources Control Board, California Environmental Protection Agency, Sacramento, California. Online at: [http://www.waterboards.ca.gov/water\\_issues/programs/swamp/docs/lakes\\_study/lake\\_survey\\_yr2\\_no\\_app.pdf](http://www.waterboards.ca.gov/water_issues/programs/swamp/docs/lakes_study/lake_survey_yr2_no_app.pdf).

Thongda, W., M. Lewis, H. Zhao, B. Bowen, D.J. Lutz-Carrillo, B.K. Peoples, and E. Peatman. 2020. Species-diagnostic SNP markers for the black basses (*Micropterus spp.*): a new tool for black bass conservation and management. *Cons. Genet. Resour.* 12:319–328. Available at: <https://link.springer.com/article/10.1007%2Fs12686-019-01109-8>.

USDA/USDHHS. 2020. Dietary Guidelines for Americans, 2020–2025. 9<sup>th</sup> Edition. U.S. Department of Health and Human Services and U.S. Department of Agriculture. Online at: <https://www.dietaryguidelines.gov/>.

US EPA. 1989. Risk Assessment Guidance for Superfund Volume I: Human Health Evaluation Manual (Part A) Interim Final. EPA/5401-89/002, December 1989. Office of Emergency and Remedial Response, U.S. Environmental Protection Agency, Washington, D.C. Online at: <https://rais.ornl.gov/documents/HHEMA.pdf>.

US EPA. 2000a. Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories: Volume 1. Fish Sampling and Analysis, 3<sup>rd</sup> Edition. EPA 823-B00-007. Office of Water, U.S. Environmental Protection Agency, Washington, D.C.

US EPA. 2000b. Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories: Volume 2. Risk Assessment and Fish Consumption Limits, 3<sup>rd</sup> Edition. EPA 823-B-00-007. Office of Water, U.S. Environmental Protection Agency, Washington, D.C.

Weaver, K.L., P. Ivester, J.A. Chilton, M.D. Wilson, P. Pandey, and F.H. Chilton. 2008. The content of favorable and unfavorable polyunsaturated fatty acids found in commonly eaten fish. J. American Dietetic Assoc. 108:1178–1185.

## APPENDIX. ADVISORY TISSUE LEVELS

Advisory Tissue Levels (ATLs; OEHHA, 2008 and 2011) guide the development of advice for people eating sport fish. ATLs are levels of contaminants found in fish that correspond to the maximum numbers of recommended fish servings. OEHHA uses ATLs to provide advice to prevent consumers from being exposed to:

- More than the reference dose<sup>16</sup> on an average daily basis for chemicals not known to cause cancer, such as methylmercury, or
- For cancer-causing chemicals, a risk level greater than one additional cancer case in a population of 10,000 people consuming fish at the given consumption rate over a lifetime. This cancer risk level is the maximum acceptable risk level recommended by the US EPA (2000b) for fish advisories.

For each chemical, ATLs were determined for both cancer and non-cancer risk, if appropriate, for one to seven eight-ounce servings per week. The most health-protective ATLs for each chemical, selected from either cancer or non-cancer based risk, are shown in the table below for zero to seven servings per week. When the guidelines for eating fish from a water body are followed, exposure to chemicals in fish from that water body would be at or below the average daily reference dose or the cancer risk probability of one in 10,000.

## ADVISORY TISSUE LEVELS FOR SELECTED ANALYTES

Contaminant	Consumption Frequency Categories (8-ounce servings/week) <sup>a</sup> and ATLs (in ppb)							
	7	6	5	4	3	2	1	0
Chlordanes	≤ 80	>80–90	>90–110	>110–140	>140–190	>190–280	>280–560	>560
DDTs	≤ 220	>220–260	>260–310	>310–390	>390–520	>520–1,000	>1,000–2,100	>2,100
Dieldrin	≤ 7	>7–8	>8–9	>9–11	>11–15	>15–23	>23–46	>46
MeHg <sup>b</sup> (Women 18–49 and children 1–17)	≤ 31	>31–36	>36–44	>44–55	>55–70	>70–150	>150–440	>440
MeHg (Women ≥ 50 and men ≥ 18)	≤ 94	>94–109	>109–130	>130–160	>160–220	>220–440	>440–1,310	>1,310
PBDEs	≤ 45	>45–52	>52–63	>63–78	>78–100	>100–210	>210–630	>630
PCBs	≤ 9	>9–10	>10–13	>13–16	>16–21	>21–42	>42–120	>120
Selenium	≤ 1,000	>1,000–1,200	>1,200–1,400	>1,400–1,800	>1,800–2,500	>2,500–4,900	>4,900–15,000	>15,000
Toxaphene	≤ 87	>87–100	>100–120	>120–150	>150–200	>200–300	>300–610	>610

<sup>a</sup>Serving sizes (prior to cooking, wet weight) are based on an average 160-pound person. Individuals weighing less than 160 pounds should eat proportionately smaller amounts.

<sup>b</sup>All mercury detected is assumed to be methylmercury, which is the most common form found in fish and is also the more toxic form (Bloom, 1992).

<sup>16</sup> The reference dose is an estimate of the maximum daily exposure to a chemical likely to be without significant risk of harmful health effects over a lifetime.