

Appendix F

OEHHA Synthetic Turf Study

March 2026

Prepared by

**Pesticide and Environmental Toxicology Branch
Office of Environmental Health Hazard Assessment
California Environmental Protection Agency**

Page intentionally left blank



Appendix F. Exposure Analyses, Studies, Concentrations, and Doses

F.1. Assessing Human Time-Activity Exposure Patterns Occurring on Synthetic Turf Fields – by The University of California, Berkeley



Draft Report: Assessing Human Time-Activity Exposure Patterns Occurring on Synthetic Turf Fields

Agreement #16-E0019

Principal Investigator: Asa Bradman, PhD, MS
Center for Environmental Research and Children's Health
University of California – Berkeley

Prepared for the California Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Report Authors

Asa Bradman, PhD

Carly Hyland, MS

Rosemary Castorina, PhD

Center for Environmental Research and Children's Health
University of California - Berkeley
1995 University Avenue, Suite 265
Berkeley, CA 94704



Disclaimer

The statements and conclusions in this Report are those of the contractor and not necessarily those of the California Office of Environmental Health Hazard Assessment.



Acknowledgements

The authors wish to thank the California Office of Environmental Health Hazard Assessment (OEHHA) for sponsoring this study and especially OEHHA staff members Dr. Patty Wong, Dr. Jocelyn Claude, Rebecca Beloso, and Ladan Khandel for their ongoing support, including review of the final report. We also thank our colleagues Dr. Paloma Beamer and Nicolas Lopez-Galvez at the University of Arizona, College of Public Health for their expertise in time activity exposure assessment and on-going support over the course of this project. We would also like to thank members of the Scientific Advisory Panel for their input at various stages throughout this project. We thank OEHHA and CalRecycle staff Bennett Lock, Dr. David Ting, Carolyn Flowers, Jill Hayashida, Emily Wang, Tamara Kadir, Huyen Tran Pham, Dr. Ida Flores-Avila, Karen Randles, Karla Mondo, and Dr. Keita Ebisu and UC Berkeley students Hannah Luong, Laura Fouquette, Tessa Oliaro, Samantha Schildroth, Anna Smith, and Kevin Soun who participated in the fieldwork for this project. We also thank Arti Kundu for help with literature reviews and formatting. Finally, we thank all the soccer players and their parents who participated in this project.

This Report was submitted in fulfillment of OEHHA Contract No. 16-E0019 “Assessing human time-activity exposure patterns occurring on synthetic turf fields and development of studies to assess synthetic turf related exposures”, by the Center for Environmental Research and Children’s Health, School of Public Health, UC Berkeley, in collaboration with OEHHA.



Table of Contents

F.1.2.3.1. Survey Development.....	F-13
F.1.2.3.2. Recruitment.....	F-13
F.1.2.3.3. Data Collection	F-14
F.1.2.3.4. Data Management and Analysis.....	F-14
F.1.2.4.1. Questionnaire Development.....	F-14
F.1.2.4.2. Recruitment.....	F-14
F.1.2.4.3. Data Collection	F-15
F.1.2.4.4. Data Management and Analysis.....	F-15
F.1.3.4.1. Demographic Information.....	F-19
F.1.3.4.2. Player History	F-22
F.1.3.4.3. Longest Amount of Time Spent on Synthetic Turf Field in Past Year ..F-25	
F.1.3.4.4. Reported Exertion During Practices and Games.....	F-26
F.1.3.4.5. Contact with Crumb Rubber During Practices and Games	F-27
F.1.3.4.6. Food and Water Consumption Before and During Practices and Games F-36	
F.1.3.4.7. Observation of Crumb Rubber in Home	F-37
F.2.2.2.1. Children Playing on Turf.....	F-49
F.2.2.2.2. Children Playing on Playgrounds	F-49
F.2.3.1.1. Amount of Footage with Turf and Demographics of Children Who Played on Turf.....	F-51
F.2.3.1.2. Hand Contact Activity While Playing on Turf.....	F-52
F.2.3.1.3. Mouthing Activity While Playing on Turf.....	F-64
F.2.3.2.1. Children’s Description and Footage Length	F-74
F.2.3.2.2. Hands Object Activity While Playing on Playground Structures	F-74
F.2.3.2.3. Mouthing Activity While Playing on Playground	F-86
F.3.2.2.1. Training	F-105
F.3.2.2.2. Quality Control and Quality Assurance.....	F-105
F.3.2.3.1. Analysis for Both Hands and Mouthing Events.....	F-106
F.3.2.3.2. Analysis for Macro-Activities, Actions and Intensity	F-107



F.3.2.3.3. Statistical Analysis	F-108
F.3.3.6.1. Event Schedule (Moment/Time)	F-119
F.3.3.6.2. Location (On or Off Field) (Supplemental Material Sections S3.2.2.1 and S3.2.2.2) F-120	
F.3.3.6.3. Use Of Gloves (On or Off) (Supplemental Material Sections S3.2.3.1 and S3.2.3.2) F-121	
F.3.3.6.4. Player Age (Supplemental Material Section S3.3)	F-121
F.3.3.6.5. Player Gender	F-126
F.3.3.6.6. Player Position (Goalie or Others)	F-131
F.3.3.6.7. Event Type (Game or Practice).....	F-136



Abbreviations

COPC	Contaminants of Potential Concern
CPHS	Committee for the Protection of Human Subjects
NBI	Norwegian Building Institute
OEHHA	Office of Environmental Health Hazard Assessment
PAH	Polycyclic Aromatic Hydrocarbon
STFCR	Synthetic Turf Field with Crumb Rubber
UA	University of Arizona
VOC	Volatile Organic Compound



Overview

Due to their low water and maintenance requirements, the use of synthetic turf fields has become popular in California in recent years; however, concerns have emerged that playing soccer and other sports on synthetic turf fields with crumb rubber (STFCR) may result in exposure to chemicals present in recycled tires that pose risks to human health.¹ For example, crumb rubber infill made from recycled tires contains various toxicants such as PAHs and other organic contaminants.² Athletes and bystanders that spend time on or near synthetic turf fields may be exposed to these chemicals through oral, dermal, and inhalation routes.²

Exposure to chemicals from synthetic turf fields are of particular concern among soccer players in California. In addition to being the most popular sport among high school students in California, there is an abundance of recreational and competitive soccer leagues throughout the state that attract diverse participants of varying ages.³

While some health risk assessments have not found significantly increased risk from activity on synthetic turf fields,^{1 2, 4} the majority of studies to date have focused on short-term exposures. This study aims to provide information necessary to assess potential risks resulting from lifetime exposures to synthetic turf fields among California soccer players.³

This project includes three main components: i) administration of an online survey to obtain statewide information regarding players' soccer history and exposure-related behaviors; ii) videotaping players at soccer practices and games on synthetic turf fields containing crumb rubber in the San Francisco Bay Area and Sacramento area; and iii) administration of in-person questionnaires to players or parents of players participating in the videotaping. For the online survey, we recruited male and female recreational and competitive soccer players over the age of 18 and invited parents/guardians to complete the survey for soccer players under 18. For the videotaping and in-person questionnaires, we recruited male and female soccer players aged 7 to 25 years old.

Using publicly available datasets, we acquired email addresses for coaches, managers, club presidents, and other affiliates of youth, high school, collegiate, and adult recreational and competitive soccer teams throughout California. Between December, 2017 and April, 2018 we sent emails to soccer coaches and managers i) throughout California to recruit participants for the online survey, and ii) in the San Francisco Bay Area and Sacramento area to recruit participants for the videotaping and in-person questionnaires.

A total of 1,069 participants completed the online or in-person surveys and 40 soccer players were videotaped during five practices and five games on STFCRs. Information from these studies will inform the California Environmental Protection Agency's Office of Environmental Health Hazard Assessment (OEHHA) efforts to assess potential chemical exposures health risks resulting from use of synthetic turf fields.



F.1.1. Introduction

Concerns have been raised about the safety of using recycled tire crumb rubber as infill in synthetic turf playing fields due to the potential for exposure of athletes and bystanders to harmful chemicals, including known and suspected carcinogens, neurotoxins, and potential endocrine disruptors.^{5, 6} Numerous laboratory and field studies have found that volatile organic compounds (VOCs), semi-volatile organic compounds (SVOCs), metals, particulates, and other substances may be released from recycled tires and tire crumb under various conditions.⁷

While some risk assessments have concluded that the use of tire crumb rubber in synthetic turf playing fields presents minimal risk to athletes and bystanders, there is limited information from quantitative exposure assessments characterizing exposure to compounds released from crumb rubber via oral, inhalation, and dermal routes of exposure. Current risk assessments have also largely focused on short-term exposures and there are data gaps regarding exposures and risks associated with synthetic turf field use over a lifetime.⁶

Although no epidemiological studies have been published that have shown associations between use of synthetic turf fields with crumb rubber (STFCRs) and adverse health outcomes, there have been case reports of cancer among soccer players, particularly for goalies, who may have greater contact with synthetic turf.⁸

In this report, we describe time-activity data collected from Californians that play soccer on STFCRs. This information about players' histories and exposure-related behaviors will inform the development of exposure scenarios and modeling by OEHHA that will be used to evaluate potential health risks associated resulting from use of synthetic turf fields with crumb rubber.

F.1.1.1. Routes of Exposure

Soccer players, coaches, referees, bystanders, and others spending time on or near STFCRs may be exposed to chemicals present in the crumb rubber via inhalation, ingestion, and dermal contact.

Inhalation

Inhalation exposures occur when soccer players or bystanders inhale gases emitted from crumb rubber granules or suspended particles generally ≤ 10 microns. The greatest exposures are expected among athletes, particularly while they are engaged in high intensity activities that increase breathing rates. The resuspension of small crumb rubber particles into air from activities such as running, falling, or sliding can also result in higher inhalation exposures among players, coaches, referees, and other personnel located on the field. Bystanders near the field are likely to be engaged in sedentary activities, such as sitting or standing, and may have lower exposures via inhalation.



Ingestion

Non-dietary Ingestion

- Can be intentional (i.e., infant/toddler mouthing, pica behavior) or inadvertent (non-dietary ingestion);
- May occur during activities where hands, fingers, or other objects come in contact with the turf surface and then touch the face or mouth of an individual (i.e., when eating, scratching, etc.);
- May also occur when large dust particles are trapped in the upper respiratory system before entering the lungs and are then transported to the throat and swallowed.

Ingestion of crumb rubber may be inadvertent or intentional. Inadvertent ingestion may occur when a player or bystander's hands contact the crumb rubber or field surface and then their mouth. For example, soccer players may fall or dive during games or practice, touch the field surface and then touch their mouth with their hands, transferring chemicals into the oral cavity. Soccer players and bystanders may also be exposed via indirect hand-to-mouth activity when their hands contact an object that has contacted the field or other contaminated surface, such as a water bottle or toy, which ultimately comes into contact with their mouth. Inadvertent ingestion can also occur if crumb rubber granules directly enter the mouth when diving, slipping, tackling, or other strenuous contact with the field surface.⁹ Among soccer players, goalkeepers are expected to have higher ingestion exposures than other positions because they repeatedly dive and make contact with the field in practices and games.

Young bystanders, including infants and children, may crawl on the sidelines and then put their fingers in their mouth, also resulting in inadvertent ingestion. Young children may also be more likely to have a snack or play with toys on the field, which could result in indirect ingestion of crumb rubber particles, particularly if they do not wash their hands.

Intentional ingestion, in which crumb rubber is directly consumed, may be an important exposure pathway for infants and young children who crawl on the field sidelines and put crumb rubber directly in their mouth and eat it.

Dermal Absorption

Dermal exposure occurs when chemicals on skin are absorbed into the body. Soccer players routinely come in contact with the field during games and practices when diving, sliding, or slipping. Players may also push off the field with their hands to maintain balance during warm-up activities or to get themselves up after falling. Kicking the turf, running, and diving may temporarily dislodge crumb rubber from the field surface and suspend it into the air where it may contact skin. Falling or making contact with the turf can also cause crumb rubber to adhere to clothing or exposed skin, potentially resulting in more prolonged exposure if players do not remove their clothes and take a shower immediately following a practice or game. Anecdotal reports from parents often describe



extensive crumb rubber dust on children's skin or granules and dust in shoes, clothing, or hair. Bystanders may also be exposed by dermal contact with field surfaces, as adults may sit on the turf to watch practices or games and young children may play on or crawl around the turf.

Dermal exposure may also occur when gases emitted from crumb rubber granules are directly absorbed through the skin.

F.1.1.2. Relevant Research

Several laboratory studies, exposure assessments, and risk assessments conducted in the United States and Europe have aimed to assess exposure to chemicals released from recycled tires and crumb rubber and their potential health risks among users of STFCRs. Key studies include:

- The State of Connecticut collected personal air samples from 3-4 volunteers playing soccer for a 2-hour sampling event at one indoor and four outdoor STFCRs.¹⁰⁻¹² Investigators identified 22 potentially field-related chemicals of potential concern (COPC) that were elevated at the synthetic turf fields compared to background samples, including 10 VOCs, 11 PAHs, and the SVOC benzothiazole.¹⁰⁻¹²
- In a 2009 study in New York, investigators compared contaminants in the air above synthetic turf fields in the breathing zone of children collected under simulated playing conditions with background samples collected from grass fields and upwind locations.¹³ The investigators did not identify an increased risk for human health effects resulting from ingestion, dermal, or inhalation exposure to crumb rubber COPCs.¹³
- The Norwegian Building Institute (NBI) conducted two studies in which they measured VOCs in air samples from indoor artificial turf fields¹⁴ and analyzed PAHs in rubber granule and turf fiber samples from different distributors of artificial turf systems.¹⁵
- A French study measured the concentrations of VOCs emitted from crumb rubber under laboratory conditions.¹⁶ The data were used by the French National Institute for Industrial Environment and Risks to evaluate possible health risks from inhaling VOCs released from synthetic turf.¹⁶ The study authors concluded that the concentrations of organic compounds emitted did not pose a health concern for athletes, officials or spectators.¹⁶

Previous exposure assessment studies of synthetic turf with crumb rubber have not adequately addressed multiple scenarios of exposure frequency/duration and routes of exposure. Despite significant research on the topic, questions still remain about the safety of long-term exposure to crumb rubber in synthetic turf fields.

F.1.1.3. Research Objectives

Objective 1: Exposure Scenario Development - Estimate Nature, Duration, and Frequency of Exposures



- a. Use available databases to characterize the number and age of soccer players in California.
- b. Develop and administer online and in-person questionnaires to soccer players (and their parents, for players under 18) throughout California to characterize time-activity patterns of soccer participants.
- c. Videotape soccer players at practices and games taking place on synthetic turf fields containing crumb rubber in the San Francisco Bay Area and Sacramento area to capture information needed to identify exposure-related sport and non-sport activities that occur on synthetic turf fields.

F.1.2. Materials and Methods

F.1.2.1. Approach

This project focused on youth and adult competitive and recreational soccer players using STFCRs throughout California. The project included three primary components: i) administration of an online survey to collect statewide information about players' soccer history and behaviors while playing soccer; ii) videotaping soccer players during scheduled practices and games in the San Francisco Bay Area and Sacramento area to characterize activity patterns while playing soccer and characterize exposure-related behaviors among soccer players using STFCRs; and, iii) administration of an in-person questionnaire to players participating in the videotaping to collect information on their soccer history and behaviors while using STFCRs.

Our goal was to recruit 1,000 participants of various ages throughout California to participate in the online survey. For the in-person videotaping and questionnaire, our goal was to videotape four soccer players at up to 10 practices or games taking place on a STFCR, for a maximum of 40 participants. At each event, we attempted to videotape one soccer player for each player position on the team (i.e., forward, midfielder, defender, and goalkeeper). In order to capture exposure-related behaviors among a range of soccer players, we aimed to enroll one male and one female soccer team in each of the following age categories: under 9 years, 9-12 years, 12-15 years, 15-18 years, and 18-25 years.

Video data were sent to the University of Arizona (UA) for analysis. Investigators at UA transcribed the video data and coded the exposure-related time-activity behaviors using VirtualTimingDevice™. VirtualTimingDevice™ is a software program developed by UA to track the duration of contact of a body part (e.g., hands, legs) with surfaces (e.g., artificial turf) and other objects, such as a water bottle, food, or a mouth guard.

Data from the online surveys, videotaping and in-person questionnaires will be used by OEHHA to develop exposure scenarios for soccer players using STFCRs.

F.1.2.2. Human Subjects Review and Approval

Human subjects approval for this project was received from the UC Berkeley Committee for the Protection of Human Subjects (CPHS) on November 13, 2017 and from the



State of California CPHS on December 4, 2017. The UA CPHS relied on the UC Berkeley CPHS for this project.

F.1.2.3. Online Survey

F.1.2.3.1. Survey Development

The online survey was developed by study staff and OEHHA after review of instruments developed by U.S. EPA¹⁷ and consultation with soccer players and their parents. Additional input was provided by OEHHA management. An independent soccer coach also reviewed the questionnaire. The survey focused solely on use of STFCRs and included questions about general demographic information, frequency of use, frequency of contact with the field during games and practices, usual activity levels during games and practices, potential dermal and non-dietary ingestion exposures during games and practices, hygiene practices, and take-home of crumb rubber particles and dust (i.e., on shoes or clothing) after playing soccer on STFCRs. The online survey also included a detailed player history section. For the parents/guardians completing the survey for their child under the age of 18, the player history section addressed the use of STFCRs for soccer when their child was between the ages of 4-8, 9-12, and 13-17 years old. For soccer players over the age of 18 completing the survey themselves, the player history section addressed the use of STFCRs during college, high school, and youth soccer. The online survey is provided in Supplemental Material Section S1.1.

F.1.2.3.2. Recruitment

We primarily used email to recruit participants for the online survey. We obtained publicly available email addresses for coaches, managers, and other soccer club affiliates in California from websites for youth/club recreational and competitive soccer, high school soccer, adult recreational soccer, and collegiate competitive and intramural soccer. We obtained email addresses for collegiate competitive and recreational soccer teams by manually searching for coaches and managers associated with teams from all colleges and universities in California.

NorCal Premier, Cal North Soccer, and Cal South Soccer were the primary organizations that were used to obtain email addresses for youth and adult recreational and competitive soccer teams. Together, these organizations have over 600 member clubs and their websites were used to obtain the email addresses for coaches, managers, and other contacts associated with youth and adult soccer throughout California.

In December 2017 and January 2018, we sent a recruitment email for the online survey to over 10,000 email addresses for coaches and managers associated with youth, high school, collegiate, and adult recreational and competitive soccer teams throughout California. The email briefly outlined the study and asked coaches/managers to forward the link to the online survey to parents and players on the team. Links to the email in English and in Spanish were included in the email. The recruitment email also asked the coach/manager to complete the survey if they currently play soccer or had played



soccer in the past. Parents of soccer players under the age of 18 were invited to complete the survey for their child.

Between February and April 2018, we sent follow-up emails asking soccer coaches and managers to forward the link to the survey to the parents and players on their team if they had not done so already. Staff also conducted follow up calls to key clubs and organizations in different regions of the state to encourage participation.

In addition to sending emails to soccer coaches and clubs in California, study staff distributed recruitment flyers with a link to the online survey to bystanders and parents at the in-person videotaping fieldwork events. The flyer also included a link to the Facebook page that we maintained that allowed potential participants to obtain more information about the study and contact study investigators.

F.1.2.3.3. Data Collection

The UC Berkeley version of Qualtrics was used to administer the online surveys.

F.1.2.3.4. Data Management and Analysis

Data from the English and Spanish surveys were downloaded from Qualtrics in an Excel file and transferred to STATA statistical software Version 14, which was used for all data analysis.

F.1.2.4. In-Person Survey and Videotaping Time-Activity Assessment

F.1.2.4.1. Questionnaire Development

A questionnaire nearly identical to the online survey addressing exposure-related behaviors was administered to each of the players 14 years or older being videotaped. A parent/guardian was asked to complete the questionnaire for participants younger than 14 years. The questionnaire focused on use of STFCRS and included questions about demographics, use frequency of STFCRs, frequency of contact with the fields during games and practices, activity levels during games and practices, potential dermal and non-dietary ingestion exposures during games and practices, hygiene practices, and take-home of crumb rubber particles and dust (i.e., on shoes or clothing) after playing soccer on STFCRs. Information was also collected about the soccer history and the use of STFCRs at younger ages (4-8, 9-12, 13-17, and 18-25 years old, depending on the age of the participant). The in-person questionnaire is provided in Supplemental Material Section S1.1.

F.1.2.4.2. Recruitment

Beginning in December 2017, we sent a recruitment email to managers and coaches of recreational and competitive soccer teams in the San Francisco Bay Area and Sacramento area with players ages 7-25 years. The email detailed the goals of the study, what potential participation would involve for the team, and asked the coach and/or manager to respond to study staff via phone or email if they were potentially interested in participating. Study staff communicated with coaches and managers that



demonstrated interest in the study and provided more details about the study. Most coaches were able and willing to reach out to players and parents and help with forwarding study recruitment and consent information. Study staff then interacted with parents and players to obtain consent and assent (from players <18 years).

F.1.2.4.3. Data Collection

Videotape: Teams of two study staff were assigned a player to videotape from the time the participant stepped on the field until the time the participant left the field, including breaks. Study staff completed an “Object Palette Log Sheet” and “Taper’s Log Time Sheet” to record additional behaviors of the player they were videotaping. The purpose of the Object Palette Log Sheet was to record objects the participant touched to provide more detail for the staff transcribing the video footage. Examples of details study staff recorded on the Object Palette Log Sheet include “clear plastic water bottle” or “yellow polyester jersey”. Field staff also recorded other events that caused them to turn off or move the camera, such as during half-time or a camera malfunction. The videotape observation forms (Object Palette Log Sheet and Taper’s Log Time Sheet) are provided in Supplemental Material Section S1.1.

If enough staff were available, 1-2 high definition wide-angle cameras were set up on the edge of the field in order to capture the entire field to provide back-up footage in case participants were not in view for a segment of the primary video recording.

Questionnaire: Study staff administered the questionnaire to the soccer player, or a parent/guardian of the participant, as described above. A parent/guardian was asked to complete the questionnaire if the soccer player was younger than 14 years and the soccer player was asked to complete the questionnaire if they were 14 years or older. The UC Berkeley version of Qualtrics was used to administer the in-person questionnaires on electronic tablets.

F.1.2.4.4. Data Management and Analysis

Questionnaire Data

Data from the questionnaires collected on the Qualtrics offline application were uploaded to the Qualtrics server in the field or upon returning to the office.

Video Data

Video footage recorded on SD cards were transferred to i) an external hard drive device, and ii) Box file sharing service. A link to the password-protected and encrypted video files on Box was shared with collaborators at UA so they could download and analyze the data.

Video translation was completed using VirtualTimingDevice™ software, as described in previous studies.¹⁸⁻²⁰ Briefly, study staff from CERCH, OEHHA, and UA collaborated to tailor a video palette for this project containing potential locations of soccer players (e.g., field, sideline), time (e.g., before practice/game, during practice/game), activities



(e.g., diving), and object surface categories (e.g., artificial turf, water bottle). Translators at UA would transcribe the videos in 30-minute segments and activate cells in each of the location, time, activity, and object categories that correlated with the activity and contact occurring in the video. Once a cell was activated, a timer started that recorded the length of each activity and contact. If a new cell was activated, a new timer began. Through this process, data is collected that translates to the types of contacts being made and the contact frequencies and durations. This process was repeated for the left hand, right hand, left foot, and right foot to gather data for various body parts of interest.

Additionally, existing video footage from 56 children collected in 1998-2000 by Stanford's Exposure Research Group was reviewed and translated to gather micro-level data regarding the frequency of hand-to-mouth activity of young children playing outdoors.²¹ These data were used to develop exposure parameters for bystanders who may play on the sidelines of synthetic turf fields during practices and games.

F.1.3. Results and Discussion

F.1.3.1. Number and Age of Soccer Players in California (*Objective 1a*)

Soccer is often a lifelong sport and has been increasing in popularity in California in recent years. The abundance of recreational and competitive soccer leagues in California attracts diverse participants of varying ages. Soccer is also popular among both boys and girls in elementary, middle, and high school. While data are not readily available regarding how often soccer practices and games take place on STFCRs, factors such as minimal water requirements have resulted in increased installation of STFCRs across the United States and California.¹⁷ In 2016, California had over 900 STFCRs that were located primarily in the San Francisco Bay Area and Los Angeles County.²²

According to the US Youth Soccer Organization, over 320,000 boys and girls ages 18 and under were involved in recreational or competitive soccer in California between 2013 and 2014.²³ Among the sports most often played on STFCRs, soccer has one of the highest number of high school participants in California. The California Interscholastic Federation reported that nearly 100,000 males and females participated on high school soccer teams in 2016, representing nearly 13% of California's population of high school student athletes and over 5% of the total high school population in the state.^{3, 24} While participation in competitive soccer teams decreases at the collegiate, semi-professional, and professional levels, college intramural and adult recreational soccer leagues are also common in California.

In addition to being a favorite sport for recreational and competitive players of all ages, the Latino population has grown significantly in recent years and soccer has long been an important part of Latino culture in California. Data from the March 2016 Census Bureau Current Population Survey indicate that Hispanic/Latinos made up 38% of California's population in 2015.^{25, 26} In 2014, the estimated Latino population in



California was nearly 15 million, compared to 7.7 million in 1990 and 2.4 million in 1970.²⁷ An online blog titled “Hispanic Media” described soccer as an “outlet of cultural pride for Hispanics”²⁸ and a study on the impact of soccer in the Latin American community in Richmond, CA indicated that the sport plays a “central role” within the Latino community and creates “social networks through the community’s relationship with teams and clubs centered in Richmond”.²⁹ While data are not currently available regarding soccer enrollment among different ethnic groups, there are indications that soccer continues to grow in popularity in California due various factors, including the increasing Latino population in the state.

Factors such as increasing participation at the high school level, the widespread availability of recreational and competitive leagues for adults, and popularity among groups of diverse demographic backgrounds make soccer one of the most popular sports in California. We estimate that there are approximately 440,000 individuals involved with some form of recreational or competitive soccer in California that may have contact with synthetic turf fields containing crumb rubber (Table F-1).

Table F-1. Overview of Soccer Participation in California (Total=440,721)

Competitive Level	Age Range (Years)	Gender	Estimated Players (n)	Year
Youth Recreational and Competitive	4-18	Both	162,297 ^{a,b}	2013-2014 seasonal year (Northern CA)
Youth Recreational and Competitive	4-18	Both	159,278 ^{a,b}	2013-2014 seasonal year (Southern CA)
High School Competitive	14-18	Boys	52,266 ^{b,c}	2016
High School Competitive	14-18	Girls	46,778 ^{b,c}	2016
College Competitive (Divisions I-III)	18-22	Men	1,614 ^d	2016-2017
College Competitive (Divisions I-III)	18-22	Women	1,681 ^d	2016-2017
College Recreational (Intramural)	18-22	Both	5,000 ^e	2017
Adult Recreational	18+	Both	11,000 ^f	2017
Professional and Semi-Professional Competitive	18+	Men	566 ^g	2016-2017
Professional and Semi-Professional Competitive	18+	Women	241 ^g	2016-2017

^aRetrieved from US Youth Soccer Association at <http://www.usyouthsoccer.org/media kit/keystatistics/>.

^bHigh school players may participate in US Youth Soccer club teams and high school teams



^cRetrieved from California Interscholastic Federation at http://www.cifstate.org/coaches-admin/census/2016_CIF_Participation_Census.pdf.

^dCalculated by tallying rosters of all NCAA and NAIA collegiate teams in CA for 2016-2017 season

^eEstimate based on number of universities and colleges in CA, assuming two intramural teams per school and 15 players per team

^fEstimate based on number of teams affiliated with California Soccer Association-North (Cal North), California Soccer Association-South (Cal South), and other adult soccer leagues in California. Estimated 15 members per team for teams that did not provide number of participants.

^gCalculated by tallying rosters of all semi-professional and professional teams in Men's Major League Soccer, United Women's Soccer, and Premier Leagues in CA

F.1.3.2. Online Survey Overview (*Objective 1b*)

A total of 1,028 individuals completed the online survey (1,019 in English and nine in Spanish). Responses were collected from 763 parents/guardians who completed the questionnaire for their child and 265 soccer players 18 years or older completing the questionnaire for themselves. One individual completed an in-person questionnaire (virtually identical to the online survey) but was not videotaped, for a total of 1,029 questionnaire-only respondents.




Figure F-1. Heat Map of Zip Code Reported by Online Survey Participants

Figure F-1 is a heat map of the residential zip code reported in the online survey. While the majority of the survey respondents were located in the San Francisco, Sacramento, Los Angeles, and San Diego metropolitan areas, it is important to note that these are the California regions with the greatest population. Furthermore, California's STFCRs are located primarily in the San Francisco Bay Area and the Greater Los Angeles Area.²²



F.1.3.3. In-Person Questionnaire and Videotaping Time-Activity Assessment Overview (*Objective 1c*)

Forty individuals participated in the videotaping and completed the in-person questionnaire. Events were evenly split by gender (i.e., five events with males and five events with females) and event type (i.e., five games and five practices) (Table F-2). We also enrolled players with a range of ages and positions (Table F-2).

Table F-2. Videotaping Player and Event Summary (Total Players=40)

Event Type	Gender	Age (Years)	Positions Videotaped	Number Videotaped
Game	Male	9	Defender, Goalie, Variable (2)	4
Game	Female	9	Defender/Goalie, Midfielder (2)	3
Practice	Female	11-12	Defender, Defender/Goalie, Forward, Midfielder	4
Practice	Male	11-12	Forward, Goalie, Midfielder	4
Practice	Female	14	Defender/Midfielder, Forward, Forward/Goalie, Forward/Midfielder	4
Game	Male	14-15	Defender, Forward, Goalie (2), Midfielder	5
Game	Female	16-17	Defender (2), Defender/Forward/Goalie, Forward	4
Game	Male	16-18	Defender (2), Goalie, Midfielder	4
Practice	Male	19-22	Defender, Defender/Midfielder, Goalie, Midfielder	4
Practice	Female	19-21	Defender, Goalie, Midfielder (2)	4

F.1.3.4. Questionnaire Analysis

In the sections below, we provide overall population summaries of key variables collected using the online survey and in-person questionnaires. To support population-specific exposure modeling (for example, girls playing competitive soccer through age 25), we have prepared extensive tables stratifying player information by gender, participation in recreational or competitive soccer, player position, and age for all variables (data available upon request). This stratification results in hundreds of tables. For exposure scenario building, key parameters can be extracted to identify specific parameters for model building.

F.1.3.4.1. Demographic Information

Table F-3 presents demographic information of the online and in-person survey respondents combined (n=1,069). The median age of soccer players whose parent/guardian completed the survey was 14 years (n=763) and the median age of soccer players completing the questionnaire themselves was 38 years (n=264).



Approximately 60% of online survey respondents were Caucasian and 15% of respondents were Hispanic/Latino. Seventy-eight of the participants were between 9 and 25 years old and the respondents were almost evenly balanced on gender (50% male versus 49% female).

Table F-4 shows soccer characteristics reported by the survey respondents, including soccer position and percent of soccer practices and games that take place on a synthetic turf field with crumb rubber. Approximately 11% of the survey population mainly played goalie, 11% mainly played forward, 24% mainly played midfielder, and 25% mainly played defender. The majority (>75%) of the survey population only played competitive soccer, with approximately 11% playing recreational soccer and approximately 13% playing both recreational and competitive soccer.

Table F-3. Demographic Characteristics of Online and In-Person Survey Respondents (n=1,069)^a

Characteristic	Group	No. of Respondent, n (%)
Age of player	0 - <2	1 (0.1)
Age of player	2 - <9	11 (1.0)
Age of player	9 - <16	546 (51.1)
Age of player	16 - <25	296 (27.7)
Age of player	25 - <30	28 (2.6)
Age of player	30 - <40	64 (6.0)
Age of player	40 - <50	82 (7.7)
Age of player	≥ 50	30 (2.8)
Age of player	Prefer not to answer	11 (1.0)
Gender	Male	539 (50.4)
Gender	Female	522 (48.8)
Gender	Prefer not to answer	8 (0.8)
Ethnicity	Asian/Pacific Islander	55 (5.1)
Ethnicity	Black/African American	19 (1.8)
Ethnicity	Caucasian	640 (59.9)
Ethnicity	Hispanic/Latino	158 (14.8)
Ethnicity	Native American	5 (0.5)
Ethnicity	Mixed	139 (13.0)



Characteristic	Group	No. of Respondent, n (%)
Ethnicity	Other	17 (1.6)
Ethnicity	Prefer not to identify	36 (3.4)
Survey Language	English	1,060 (99.2)
Survey Language	Spanish	9 (0.8)

^aOnline survey completed by parent/guardian of soccer player (n=763) and players 18 years or older (n=265)

Table F-4. Soccer Player Characteristics (n=1,069)

Characteristic	Group	No. of Respondent, n (%)
Soccer Position	Goalie	120 (11.2)
Soccer Position	Forward	117 (10.9)
Soccer Position	Midfielder	258 (24.1)
Soccer Position	Defender	263 (24.6)
Soccer Position	Multiple Positions	300 (28.1)
Soccer Position	DK	11 (1.0)
Recreational/Competitive Soccer Player	Recreational	115 (10.8)
Recreational/Competitive Soccer Player	Competitive	815 (76.2)
Recreational/Competitive Soccer Player	Both	134 (12.5)
Recreational/Competitive Soccer Player	DK	5 (0.5)
Plays Soccer Year-Round	No	118 (11.0)
Plays Soccer Year-Round	Yes	946 (88.5)
Plays Soccer Year-Round	DK	5 (0.5)
Percent practices on synthetic turf with crumb rubber	0%	132 (12.4)
Percent practices on synthetic turf with crumb rubber	> 0 – 25%	175 (16.4)
Percent practices on synthetic turf with crumb rubber	> 25 – 50%	155 (14.5)
Percent practices on synthetic turf with crumb rubber	> 50 – 75%	157 (14.7)



Characteristic	Group	No. of Respondent, n (%)
Percent practices on synthetic turf with crumb rubber	>75%	443 (41.4)
Percent practices on synthetic turf with crumb rubber	DK	7 (0.7)
Percent games on synthetic turf with crumb rubber	0%	19 (1.8)
Percent practices on synthetic turf with crumb rubber	> 0 – 25%	168 (15.7)
Percent practices on synthetic turf with crumb rubber	> 25 – 50%	216 (20.2)
Percent practices on synthetic turf with crumb rubber	> 50 – 75%	243 (22.7)
Percent practices on synthetic turf with crumb rubber	>75%	418 (39.1)
Percent practices on synthetic turf with crumb rubber	DK	5 (0.5)

DK: Don't Know/Prefer not to answer

F.1.3.4.2. Player History

Information on player history collected in the questionnaires was categorized by pre-determined age groups (4-8; 9-12; 13-17; and 18-25 years, respectively). All other stratified information in this report is reported according to OEHHHA risk assessment age categories (0-<2; 2-<9; 9-<16; 16+ years, respectively).

Table F-5 to Table F-8 present overall player history results for study participants. Table F-5 and Table F-6 show the number of weeks per year and hours per week for players with more detailed age-specific history (i.e., for online subjects under age 18 and players who participated in the in-person questionnaire and videotaping). Note: because adults, many in middle age, completing the online survey may not have been able to recall detailed player history at younger ages, we asked participants 18 years and older detailed information about youth, high school, and college soccer (Table F-7 and Table F-8).

Table F-5 to Table F-8 show that with increasing age, soccer players play more weeks and more hours on STFCRs. Within each of the age groups in Table F-5, we did not find significant differences in the number of weeks participants reported playing soccer on STFCRs by gender (Kruskal-Wallis: $p=0.14$ for 4-8 age group; $p=0.36$ for 9-12 age group; $p=0.65$ for 13-17 age group; $p=0.64$ for 18-25 age group). Among adults who completed the online questionnaire, represented in Table F-7, females tended to report playing soccer on STFCRs more weeks per year than males ($p=0.10$ for youth soccer; $p=0.05$ for high school soccer; $p=0.06$ for college soccer).



Note: The number of players in each group in Table F-5 to Table F-10 do not equal the total number of surveys completed because many participants reported information for multiple age groups over their lifetime.

Table F-5. Player History – Weeks Per Year Child Played on Synthetic Turf Field with Crumb Rubber between the Ages of 4-8, 9-12, 13-17 and 18-25

Age Range (years)	n ^a	p25 ^b	p50	p75	p95	Mean	SD	Range ^b
4-8	705 ^c	0 ^b	2	12	40	9.3	13.4	0-52
9-12	692 ^d	4	18	36	48	20.5	16.7	0-52
13-17	402 ^e	15	30	45	52	29.5	16.4	0-52
18-25	7 ^f	20	36	52	52	34.3	17.8	6-52

^aTotal responses by age range > 785 because many children played in multiple age groups

^b"0" represent participants who played soccer but not on synthetic turf

^cNumber of responses from parent/guardian whose child played soccer between ages of 4 and 8 years

^dNumber of responses from parent/guardian whose child played soccer between ages of 9 and 12 years

^eNumber of responses from parent/guardian whose child played soccer between ages of 13 and 17 years

^fNumber of responses from in-person questionnaire and videotaping who played soccer between the ages of 18 and 25 years

SD: standard deviation

Table F-6. Player History – Hours Per Week Child Played on Synthetic Turf Field with Crumb Rubber between the Ages of 4-8, 9-12, 13-17 and 18-25

Age Range (years)	n ^a	p25 ^b	p50	p75	p95	Mean	SD	Range ^b
4-8	705 ^c	0	1	2	6	1.8	4.6	0-104
9-12	692 ^d	1.5	3	4.5	10	3.6	4.1	0-52
13-17	402 ^e	3	5	7	14	6.1	6.9	0-100
18-25	7 ^f	7	9	10	10	8.6	1.6	0-10

^aTotal responses by age range > 785 because many children played in multiple age groups

^b"0" represent participants who played soccer but not on synthetic turf

^cNumber of responses from parent/guardian whose child played soccer between ages of 4 and 8 years

^dNumber of responses from parent/guardian whose child played soccer between ages of 9 and 12 years

^eNumber of responses from parent/guardian whose child played soccer between ages of 13 and 17 years

^fNumber of responses from in-person questionnaire and videotaping who played soccer between the ages of 18 and 25 years

SD: standard deviation

Table F-7. Player History Reported by Older Adults – Weeks Per Year Played on Synthetic Turf Field with Crumb Rubber During Youth, High School, and College Soccer

Period	n ^a	p25 ^b	p50 ^b	p75	p95	Mean	SD	Range ^b
Youth	223 ^c	0	0	8	40	6.9	13.0	0-52



Period	n ^a	p25 ^b	p50 ^b	p75	p95	Mean	SD	Range ^b
High School	230 ^d	0	0	12	40	7.5	12.5	0-52
College	189 ^e	0	2	20	42	10.0	13.8	0-52

^aTotal responses by age range > 284 because many children played in multiple age groups

^b"0" represent participants who played soccer but not on synthetic turf

^cNumber of responses from participants who played youth soccer

^dNumber of responses from participants who played high school soccer

^eNumber of responses from participants who played college soccer

SD: standard deviation

Table F-8. Player History Reported by Older Adults – Hours Per Week Played on Synthetic Turf Field with Crumb Rubber During Youth, High School, and College Soccer

Period	n ^a	p25 ^b	p50 ^b	p75	p95	Mean	SD	Range ^b
Youth	223 ^c	0	0	3	9	2.0	4.1	0-25
High School	230 ^d	0	0	6	15	3.3	5.6	0-35
College	189 ^e	0	2	6	18	4.3	6.4	0-42

^aTotal responses by age range > 284 because many children played in multiple age groups

^b"0" represent participants who played soccer but not on synthetic turf

^cNumber of responses from participants who played youth soccer

^dNumber of responses from participants who played high school soccer

^eNumber of responses from participants who played college soccer

SD: standard deviation

Table F-9. Days Per Year Practiced on Synthetic Turf Field

Age Range (Years)	n	Mean	SD	p25	p50	p75	p95	p99	Max
0 - <2	1	32.5	-	32.5	32.5	32.5	32.5	32.5	32.5
2 - <9	8	71.5	54.7	26.0	45.5	123.5	156.0	156.0	156.0
9 - <16	470	91.6	51.1	52.0	91.0	117.0	182.0	221.0	273.0
16+	404	128.1	69.7	78.0	117.0	169.0	247.0	325.0	351.0

Table F-10. Days Per Year Played Games on Synthetic Turf Field

Age Range (Years)	n	Mean	SD	p25	p50	p75	p95	p99	Max
0 - <2	1	39.0	-	39.0	39.0	39.0	39.0	39.0	39.0
2 - <9	8	65.0	10.8	52.0	71.5	78.0	91.0	91.0	91.0
9 - <16	490	57.1	38.8	26.0	52.0	78.0	130.0	195.0	273.0



Age Range (Years)	n	Mean	SD	p25	p50	p75	p95	p99	Max
16+	460	80.3	44.0	52.0	65.0	104.0	156.0	221.0	286.0

F.1.3.4.3. Longest Amount of Time Spent on Synthetic Turf Field in Past Year

Table F-11 and Table F-12 summarize overall population information on the longest amount of time spent practicing and playing games on STFCRs in the past year.

There was a slight increase in the longest amount of time players reported practicing or playing soccer games on STFCRs as the age of the player increased, however, across all age groups, the most common responses were 1-2 or 2-4 hours. Among all ages combined, we did not observe differences in the longest amount of time reported spent practicing or playing games on STFCRs by gender (Chi-squared test: $p=0.87$ for practices; $p=0.46$ for games). We also did not find gender differences after stratifying by age for practices (Chi-squared test: $p=0.23$ for participants <9 years; $p=0.81$ for 9-<16 age group; $p=0.91$ for ≥ 16 age group; $p=0.44$ for those who prefer not to answer age) or games (Chi-squared test: $p=0.39$ for participants <9; $p=0.09$ for 9-<16 age group; $p=0.80$ for ≥ 16 age group; $p=0.44$ for those who prefer not to answer age).

Table F-11. Number of Players, n (percent), with the Corresponding Longest Amount of Time (Hours) Practiced on a Single Day On Synthetic Turf Field with Crumb Rubber in Past Year

Age Range (Years) ^a	<1	>1-2	>2-4	>4-5	>5	DK
0 - <2	0 (0)	0 (0)	1 (100.0)	0 (0)	0 (0)	0 (0)
2 - <9	0 (0)	4 (50.0)	3 (37.5)	0 (0)	1 (12.5)	0 (0)
9 - <16	11 (2.2)	215 (43.7)	204 (41.5)	35 (7.1)	27 (5.5)	0 (0)
≥ 16	6 (1.4)	137 (32.2)	194 (45.7)	43 (10.1)	39 (9.2)	6 (1.4)
Prefer not to answer	0 (0)	5 (45.5)	4 (36.4)	2 (18.2)	0 (0)	0 (0)

^aResponses collected from 1 player 0-<2 years, 8 players 2-<9 years, 492 players 9-<16 years, 425 players ≥ 16 years, and 11 players who preferred not to answer age or entered birth year incorrectly. Values in table are player in an age group with the corresponding longest amount of time (hours) practiced on a single day on synthetic turf field, n. Values in parathenesis are percent of players in an age group, percent.

DK: don't know/prefer not to answer

Table F-12. Number of Players, n (percent), with the Corresponding Longest Amount of Time (Hours) on a Single Day Players Played Games on Synthetic Turf Field with Crumb Rubber in Past Year

Age Range (Years) ^a	<1	>1-2	>2-4	>4-5	>5	DK



0 - <2	0 (0)	1 (100.0)	0 (0)	0 (0)	0 (0)	0 (0)
2 - <9	0 (0)	8 (88.9)	0 (0)	1 (11.1)	0 (0)	0 (0)
9 - <16	21 (3.9)	176 (32.8)	231 (43.1)	75 (14.0)	31 (5.8)	2 (0.4)
≥ 16	10 (2.0)	138 (28.0)	209 (42.4)	80 (16.2)	51 (10.3)	5 (1.0)
Prefer not to answer	0 (0)	4 (36.4)	5 (45.6)	2 (18.2)	0 (0)	0 (0)

^aResponses collected from 1 player 0-<2 years, 9 players 2-<9 years, 536 players 9-<16 years, 493 players > 16 years, and 11 players who preferred not to answer age or entered birth year incorrectly. Values in table are player in an age group with the corresponding longest amount of time (hours) of games on a single day on synthetic turf field, n. Values in parathensis are percent of players in an age group, percent.

DK: don't know/prefer not to answer

F.1.3.4.4. Reported Exertion During Practices and Games

Table F-13 and Table F-14 illustrate the overall population activity levels players reported during practices and games. Participants reported the proportion of the time they typically are: a) resting; b) lightly active; c) moderately active; and d) highly active during practices and games. Table F-13 and Table F-14 only include the results from participants whose responses totaled 100% across the four activity levels (n=774 for practices and 886 for games).

Overall, survey respondents reported a high level of activity. The median percent time participants reported being moderately or highly active during practices and games was 65% and 70%, respectively.

Males reported being lightly active slightly more percent of the time during practices than females (mean=17.6 vs. 16.2; Kruskal-Wallis test: $p=0.04$). We did not observe any differences in reported exertion levels during games by gender.

Table F-13. Reported Percent of Time at Varying Activity Levels During Practice (n=774)

Activity Level	Median (IQR)	Max
Highly Active	35 (25, 50)	100
Moderately Active	30 (25, 45)	80
Lightly Active	15 (10, 20)	80
Resting	10 (5, 15)	55

IQR: interquartile range (25th, 75th percentiles)



Table F-14. Reported Percent of Time at Varying Activity Levels During Games (n=886)

Activity Level	Median (IQR)	Max
Highly Active	40 (25, 55)	100
Moderately Active	30 (20, 40)	80
Lightly Active	10 (10, 20)	60
Resting	10 (5, 20)	90

IQR: interquartile range (25th, 75th percentiles)

F.1.3.4.5. Contact with Crumb Rubber During Practices and Games

Figure F-2 to Figure F-7 illustrate the frequency survey respondents reported diving, sliding/tackling, and slipping/falling at practices and games taking place on STFCRs, stratified by the percent of respondents per soccer position.

Goalies reported diving, sliding/tackling, and slipping/sliding on field during practices and games significantly more frequently than forwards, midfielders, and defenders combined (Chi-squared test: $p < 0.01$). For example, 50.8% and 58.7% of goalies reported diving more than 10 times at practices and games, respectively, whereas less than 5% of forwards, midfielders, or defenders reported diving more than 10 times at practices or games. Additionally, males reported diving and sliding/tackling more frequently than females during practices and games (Chi-squared test: $p < 0.01$).

Figure F-8 and Figure F-9 illustrate the frequency that players reported crumb rubber in their mouth and eyes and played with crumb rubber during practices and games. The majority of soccer players reported “Rarely” getting crumb rubber in their mouths or eyes. There were no large differences in the frequency of these contacts with crumb rubber during practices and games (Chi-squared test: $p < 0.01$).

Figure F-10 to Figure F-13 illustrate the frequency players reported crumb rubber in their mouths and eyes during practices and games, stratified by player position (goalie vs. midfielder, forwards, and defenders combined). Goalies reported getting crumb rubber in their mouths and eyes during practices and games more frequently than players of other positions (Chi-squared test: $p < 0.01$). For example, almost 43% of goalies reported getting crumb rubber in their mouth “sometimes”, “often”, or “always”, compared with 16% of midfielders, forwards, and defenders.

Figure F-14 illustrates the frequency players and parents/guardians of players reported observing crumb rubber on their body or personal objects after playing soccer on a synthetic turf field with crumb rubber. Only 8% of participants reported observing crumb rubber on their water bottles >25% of the time after playing soccer on a synthetic turf field. However, 51% and 36% of participants reported observing crumb rubber on their clothes and body >25% of the time after playing soccer on a synthetic turf field, respectively.




Figure F-2. Dive Frequency at Practices on Synthetic Turf Fields




Figure F-3. Dive Frequency at Games on Synthetic Turf Fields




Figure F-4. Slide/Tackle Frequency at Practices on Synthetic Turf Fields




Figure F-5. Slide/Tackle Frequency at Games on Synthetic Turf Fields




Figure F-6. Slip/Fall Frequency at Practices on Synthetic Turf Fields




Figure F-7. Slip/Slide Frequency at Games on Synthetic Turf Fields




Figure F-8. Frequency of Contact with Crumb Rubber During Practice




Figure F-9 Frequency of Contact with Crumb Rubber During Game




Figure F-10. Frequency of Crumb Rubber in Mouth During Practice - Goalies vs Other Positions




Figure F-11. Frequency of Crumb Rubber in Eyes During Practice - Goalies vs Other Positions






Figure F-12. Frequency of Crumb Rubber in Mouth During Games - Goalies vs Other Positions




Figure F-13. Frequency of Crumb Rubber in Eyes During Games - Goalies vs Other Positions




Figure F-14. Frequency Crumb Rubber Observed on Player or Personal Objects After Game or Practice



F.1.3.4.6. Food and Water Consumption Before and During Practices and Games

Table F-15 summarizes the amount of time before practices and games participants reported eating or having a snack/meal. While the most common response was 1-3 hours, more participants reported having eaten <1 hour before practices than before games (31.3% vs. 19.0%, respectively).

Table F-16 summarizes how often participants reported having a snack during practices and games. The majority of participants indicated they never had a snack during practices or games. Less than 25% of participants reported consuming one or more snacks during practices or games (Table F-16).

The majority of participants reported consuming between 16 and 32 ounces of water during practices and games (Table F-17). Only 4.6% and 2.3% of players reported drinking more than 64 ounces of water during practices and games, respectively.

Table F-15. Number of Participants, n (Percent), Reported Eating or Having a Snack Before Practices and Games

Time Eating or Having a Snack Before an Event (Hour)	Practice	Game
<1	333 (35.5)	204 (19.4)
> 1 -3	537 (57.3)	777 (74.0)
> 3	41 (4.4)	50 (4.8)
DK	26 (2.8)	19 (1.8)

DK: don't know/prefer not to answer

Table F-16. Number of Participants, n (Percent), Reported Consumption of Snack During Practices and Games

Frequency of Snack Consumption (unitless)	Practice	Game
Never	789 (84.2)	735 (70.0)
1	90 (9.6)	260 (24.8)
2	14 (1.5)	31 (3.0)
>2	6 (0.6)	6 (0.6)
DK	38 (4.0)	18 (1.7)

DK: don't know/prefer not to answer



Table F-17. Number of Participants, n (Percent), Reported Consumption of Water During Practices and Games

Amount of Water Consumed (Ounce)	Practice	Game
8	43 (4.6)	24 (2.3)
16	251 (26.8)	218 (20.8)
24	352 (37.8)	370 (25.2)
32	230 (24.6)	210 (29.5)
48	25 (2.7)	60 (5.7)
64	20 (2.1)	44 (4.2)
> 64	7 (0.8)	13 (1.3)
DK	9 (1.0)	10 (1.0)

DK: Don't Know/Prefer not to answer

F.1.3.4.7. Observation of Crumb Rubber in Home

Figure F-15 illustrates the frequency players and parents/guardians reported observing crumb rubber in various locations around the home after playing soccer on a synthetic turf field with crumb rubber. Nearly 46% of participants reported “always” observing crumb rubber in the garage/mudroom/entrance to their home after playing soccer, and 73% reported observing crumb rubber at least “sometimes” (i.e., “sometimes”, “often”, or “always”). While participants reported observing crumb rubber in other locations of the home less frequently, 59% reported finding crumb rubber in the laundry room at least “sometimes” and 43% reported finding crumb rubber in their bedrooms and bathrooms at least “sometimes”. Only 21% and 32% of participants reported wiping, cleaning, or removing socks, shoes, shin guards, or other equipment more than 75% of the time before entering their car and their house, respectively (data not shown).

When asked about the quantity of crumb rubber participants observed in their home each time after playing soccer on a synthetic turf field with crumb rubber, the majority of participants said they find some crumb rubber, but less than one tablespoon (Figure F-16).




Figure F-15. Frequency of Crumb Rubber Observed in Home After Playing Soccer




Figure F-16. Quantity of Crumb Rubber Observed in Home After Playing Soccer



F.1.3.5. Comparisons Between Survey Responses and Videotaping Data

For the 40 individuals that participated in the in-person questionnaire and videotaping, we compared the frequency of diving, sliding/tackling, and slipping/falling during the event that was videotaped with the average number of times they reported doing these activities during a typical practice or game in the questionnaire. We also compared the percent of time participants were resting, lightly active, moderately active, and highly active in the event videotaped with the percent of time they reported these exertion levels at a typical practice or game. We only compared the responses from the questionnaire with the type of event videotaped (i.e., if a participant was videotaped at a practice, we used responses to questions regarding practices).

To evaluate the frequency of diving, sliding/tackling and slipping/falling during a typical practice or game, we observed these activities in the event videotaped for that participant. We first categorized the frequency of these events into four categories corresponding to the options provided to participants in the questionnaire, where 0 refers to never, 1 refers to 1-3 times per event, 2 refers to 4-10 times per event, and 3 refers to >10 times per event. For 19 of the 40 (48%) participants, there was no difference between the frequency of diving reported in the questionnaires versus observed in videotaping, however participants tended to report more frequent diving in the questionnaire versus during a typical practice or game (Chi-squared test: $p=0.01$). There was concurrence between the questionnaire responses and videotape information for only 25% and 30% of the participants for slipping/falling and sliding/tackling, respectively. These differences were not statistically significant, although given relatively low p-values (0.10 and 0.12, respectively), the overall findings suggest that the self-reported information is directly aligned with the videotaped observations.

Figure F-17 presents the mean percent of time all 40 participants reported being highly active, moderately active, lightly active, and resting during a typical practice or game in the questionnaire compared with the mean percent of time we observed at these activity levels in the videotapes. We found that there were significant differences between the questionnaire responses and videotape (p -value <0.01 for each activity level). For example, compared to the event we videotaped, we found that participants reported being more highly and moderately active in the questionnaires, and conversely, we observed more moderate and resting activity they reported for a typical practice or game.

It is important to note that we only videotaped one practice or one game for each participant, and in the questionnaire we asked participants to report the average frequency of various activities and exertion levels in a typical practice or game. It is likely that their behaviors and exertion levels during the event videotaped are not representative of all practices or games. Our findings suggest that players may overestimate exertion levels, but further research is needed to appropriately compare self-report exertion against videotaping during the same time period.




Figure F-17. Mean Percent of Time Activity Levels Reported in Questionnaire and Observed in Videotaping

F.1.3.6. Population-Specific Exposure Parameter Estimates

This report presents an overview of population characteristics and key variables that determine short and long-term exposures resulting from use of STFCRs (e.g., player history, practice and game frequency and duration, exertion, contact with fields, etc). To estimate exposures to specific sub-groups, it is necessary to stratify the information and identify parameters unique to each population. For example, young adult women playing competitive soccer through age 25. In this case, information on player history specific to girls and young women combined with practice and game frequency, exertion, and contact with crumb rubber can be used to develop a lifetime exposure scenario and model exposures. Given the number of variables collected by the questionnaire (90+), and the key factors requiring stratification (gender, participation in recreational or competitive soccer, player position, and age), hundreds of tables are required to identify specific parameters for developing scenarios and building exposure models, resulting in several thousand pages of output.

To create a manageable process to identify key parameters for exposure modeling, we stratified responses from the online and in-person surveys by age (i.e., 0-<2 years, 2-<9 years, 9-<16 years, 16 years and over), gender (i.e., female, male, prefer not to identify), player level (i.e., recreational, competitive, both), and player position (i.e., goalie, forward, midfielder, defender, multiple) and produced output in a text format. Using Python, we replaced code with a descriptor for each stratification, and then



produced HTML format files with a linked table of contents for each question (for example, “*Approximately what percent of [your/your child’s] soccer PRACTICES take place on synthetic turf fields? (Questions 12 and 99)*”). We also produced an overall table of contents (index file) that automatically links to the appropriate HTML files with question-specific information. These are available upon request. This procedure simplifies access to the stratified information, eliminates the potential for data entry errors, and can be used for each question from the questionnaire, thereby providing a mechanism to extract key parameters from the output to inform exposure modeling when specific exposure scenarios are developed.

F.1.4. Strengths and Limitations

This study combines the results of a large online survey with player interviews and video data obtained through direct observation. One of the primary strengths of this study is that we were able to recruit diverse soccer players of both genders and of various ages and competitive levels living throughout the state of California for the online survey. With over 1000 survey participants and videotaping of 40 soccer players in the San Francisco Bay Area and greater Sacramento, this study is the most extensive to date examining exposure-related activity of individuals using STFCRs and it is unique to California. Combined with extensive environmental sampling testing by the Lawrence Berkeley National Laboratory (LBNL), these data will support OEHHHA’s efforts to assess chemical exposures and potential health risks resulting from use of STFCRs.

As with all surveys, it is possible that recall bias may have affected the accuracy of participants’ responses to the in-person and online questionnaires. To comply with the UC Berkeley CPHS and State of California CPHS human subjects guidelines, parents or guardians rather than the soccer players themselves were required to complete the online survey for all players under the age of 18. Because some parents or guardians are not familiar with their child’s activities, some responses may not represent actual practices. To the extent possible for this report, we have presented comparisons of the time activity video data from the 40 participating soccer players and their corresponding questionnaire data. It is important to note that the timeframe of the in-person questionnaires covered a longer timeframe (e.g., weeks, months) and did not match the timeframe of the particular game or practice that was videotaped for this study. In addition, we have presented information on key variables stratified by player age, gender, field position and level of play (i.e., competitive vs. recreational). For some of these stratifications, there are few or no observations in many cells, making it impossible in these cases to interpret the distributions or make statistical comparisons. For example, there was only one player 0-<2 years of age.

F.1.5. References

1. Schiliro, T., et al., *Artificial turf football fields: environmental and mutagenicity assessment*. Arch Environ Contam Toxicol, 2013. **64**(1): p. 1-11.



2. Cheng, H., Y. Hu, and M. Reinhard, *Environmental and Health Impacts of Artificial Turf: A Review*. Environmental Science & Technology, 2014. **48**(4): p. 2114-2129.
3. California Interscholastic Federation. *California High School Sports Participation at All-Time High For Fourth Consecutive Year*. 2016; Available from: http://www.cifstate.org/coaches-admin/census/2016_CIF_Participation_Census.pdf.
4. Pavilonis, B.T., et al., *Bioaccessibility and Risk of Exposure to Metals and SVOCs in Artificial Turf Field Fill Materials and Fibers*. Risk Anal, 2014. **34**(1): p. 44-55.
5. Dorsey, M.J.A., A; Ardo, O; Chou, M; Farrow, E; Glassman, EL; Manley, M; Meisner, H; Meyers, C; Morley, N; Rominger, K; Sena, M; Stiefbold, M; Stites, B; Tash, M; Weber, E; Counts, P., *Mutagenic Potential of Artificial Athletic Field Crumb Rubber at Increased Temperatures*. The Ohio Journal of Science, 2015. **115**(2).
6. Watterson, A., *Artificial Turf: Contested Terrains for Precautionary Public Health with Particular Reference to Europe?* International Journal of Environmental Research and Public Health, 2017. **14**(9): p. 1050.
7. California Office of Environmental Health Hazard Assessment (OEHHA), *Evaluation of Health Effects of Recycled Waste Tires in Playground and Track Products*. 2007.
8. Washington State Department of Health, *Investigation of Reported Cancer among Soccer Players in Washington State*. 2017.
9. Office of Environmental Health Hazard Assessment, *Air Toxics Hot Spots Program Risk Assessment Guidelines. Technical Support Document for Exposure Assessment and Stochastic Analysis*. 2012, California Environmental Protection Agency Sacramento, CA.
10. Simcox, N.J., A. Bracker, and J. Meyer, *Artificial Turf Field Investigation in Connecticut Final Report* 2010, University of Connecticut Health Center, Section of Occupational and Environmental Medicine.
11. Ginsberg, G., et al., *Human health risk assessment of synthetic turf fields based upon investigation of five fields in Connecticut*. J Toxicol Environ Health A, 2011. **74**(17): p. 1150-74.
12. Simcox, N.J., et al., *Synthetic turf field investigation in Connecticut*. J Toxicol Environ Health A, 2011. **74**(17): p. 1133-49.
13. Vetrano, K. and G. Ritter, *Air quality survey of synthetic turf fields containing crumb rubber infill*. 2009, New York City Department of Health and Mental Hygiene: New York, NY.



14. Dye, C., A. Bjerke, and S. Schmidbauer, *Measurement of air pollution in indoor artificial turf halls*. 2006, Norwegian Pollution Control Authority/Norwegian Institute for Air Research.
15. Plesser, T.S. and O.J. Lund, *Potential health and environmental effects linked to artificial turf systems-final report*. 2004, Norwegian Building Institute: Trondheim, Norway.
16. Moretto, R. *Environmental and Health Assessment of the Use of Elastomer Granulates (Virgin and from Used Tyres) as Filling in Third-Generation Artificial Pitches*. 2007; Available from: https://www.aliapur.fr/pdf/1_environmental_evaluation_0.pdf
17. USEPA, CDC, and ATSDR, *Research Protocol: Collections Related to Synthetic Turf Fields with Crumb Rubber Infill*. 2016, U.S. Environmental Protection Agency, Centers for Disease Control and Prevention, Agency for Toxic Substances and Disease Registry.
18. Ferguson, A., et al., *Video methods in the quantification of children's exposures*. J Expos Sci Environ Epidemiol, 2005. **16**(3): p. 287-298.
19. Beamer, P., et al., *Quantified activity pattern data from 6 to 27-month-old farmworker children for use in exposure assessment*. Environ Res, 2008. **108**(2): p. 239-46.
20. Beamer, P.I., et al., *Quantified outdoor micro-activity data for children aged 7-12-years old*. J Expos Sci Environ Epidemiol, 2012. **22**(1): p. 82-92.
21. Beamer, P.I., et al., *Quantified outdoor micro-activity data for children aged 7-12-years old*. Journal Of Exposure Science And Environmental Epidemiology, 2011. **22**: p. 82.
22. Gutierrez, M., *Federal study to examine safety of synthetic turf fields*. 2016: The Orange County Register.
23. US Youth Soccer. *Key Statistics Membership Statistics* May 22, 2017]; Available from: http://www.usyouthsoccer.org/media_kit/keystatistics/.
24. California Department of Education. *Fingertip Facts on Education in California - CalEdFacts*. 2015-2016; Available from: <http://www.cde.ca.gov/ds/sd/cb/ceffingertipfacts.asp>.
25. The Henry J. Kaiser Family Foundation. *Population Distribution by Race/Ethnicity*. State Health Facts [cited 2017 May 24]; Available from: <http://kff.org/other/state-indicator/distribution-by-raceethnicity/?currentTimeframe=0&selectedRows=%7B%22nested%22:%7B%22california%22:%7B%7D%7D%7D&sortModel=%7B%22colld%22:%22Location%22,%22sort%22:%22asc%22%7D>.
26. United States Census Bureau. *QuickFacts: California*. [cited 2017.
27. Panzar, J., *It's official: Latinos now outnumber whites in California*. 2015: LA Times.



28. de la Torre, E. *Copa America 2016: Importance of Soccer to US Hispanics* Learning the Hispanic Market 2016; Available from:
<http://www.echispanicmedia.com/learning-the-hispanic-market/copa-america-2016-importance-of-soccer-to-us-hispanics/>.
29. Messeri, I., *Vamos, Vamos Aceirteros: soccer and the Latino community in Richmond, California*. Soccer & Society, 2008. **9**(3): p. 416-427.



F.2. Quantification of Micro-level Activities from a Pre-existing Dataset of Children Playing on Turf/Playgrounds – by The University of Arizona



Part I: Final Report on the Quantification of Micro-level Activities from a Pre-existing Dataset of Children Playing on Turf/Playgrounds

Prepared for the California Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Principal Investigator: Paloma Beamer

Prepared by: Nicolas Lopez-Galvez

University of Arizona, College of Public Health

August 4, 2018



Table of Contents

F.2. Quantification of Micro-level Activities from a Pre-existing Dataset of Children Playing on Turf/Playgrounds – by The University of Arizona.....	Error!
Bookmark not defined.	
F.2.1 Introduction	Error! Bookmark not defined.
F.2.2 Methods	Error! Bookmark not defined.
F.2.3 Results	Error! Bookmark not defined.
F.2.4 Attachments	Error! Bookmark not defined.
F.2.5 References.....	Error! Bookmark not defined.



F.2.1. Introduction

Non-dietary ingestion and dermal routes of exposure are important to consider when evaluating children's aggregate exposure to artificial turf or other tire-derived surfaces like playground mats. The objective of this study was to quantify dermal and mouthing contact behaviors of children playing on turf and playground structures. While these children were playing on natural turf, this data could be used to represent potential exposures of children bystanders at sporting events on artificial turf. Similarly, while most of the children were playing on playgrounds that were not above playground mats made from recycled tires or from chopped up recycled tires, we quantified contacts with floor surfaces to provide data that could be used in estimating these potential exposures. This task consisted of analyzing a pre-existing micro-level activity time series (MLATS) dataset originally collected by Stanford's Exposure Research Group. The dataset was derived by transcribing video footage of children playing outdoors by using a video-translation software (Virtual Timing Device). The activity patterns were quantified and analyzed to understand differences in behavior by age and gender of the children playing outdoors on turf and playground structures.

F.2.2. Methods

F.2.2.1. Data Collection

Children MLATS data and videotapes were utilized from two previous studies, the Outdoor Residential Exposure Task Force (ORETF) project and the EPA study from the years 1998-2000. The videotaped activities have already been transcribed into computer text files using Virtual Timing Device by the Stanford's Exposure Research Group. For ORETF and EPA, the Virtual Timing Device palette presented in Figure F-18 was used for translation of microlevel activity data. This palette allows the translator to record the child's location, contact type and object/surface. Details on ORTEF and EPA methods have been previously described (AuYeung et al., 2004; AuYeung et al., 2006; Ferguson et al., 2006; Beamer et al., 2012).

For the ORTEF project, MLATS data was collected for 15 body parts including hands and mouth by videotaping 36 children, aged 1-12 years of age while they were playing outdoors. Meanwhile, for the EPA study, MLATS data was collected for mouth and hands by videotaping 20 children, aged 1-6 years of age while they were playing outdoors. For both studies, children were videotaped for 2 hours each. We combined these data to quantify the mouthing and dermal contact behavior of 56 children while playing on turf and playground.



Activity Monitor

Location	Yard	Indoor	Grass	Carpet/ Mat	Vegetation	Food_ Cont	Wood_ Tool/Appl	Paper/ Wrapper
	Patio	Street/ Sidewalk	Dirt	Wood_ Wall/Furn	Animal	Sticky_ Food	Plast_ Tool/Appl	Head
	Garden	Park	Asphalt/ Sidewalk	Rck/Brk_ Wall/Furn	Porous_ Plast_ Toy	Other_ Food	Metal_ Tool/Appl	Skin
	Garage	Other	Rck/Brk_ Floor	Fabric_ Wall/Furn	Fabric_ Toy	Water/ Beverage	Towel/ Washcloth	Other
Contact Type	Constant	Repetitive	Wood_Floor	Plastic_ Wall/Furn	Wood_Toy	Pool_Water	Clothes	Not_ In_View
			Tile/Linol_ Floor	Metal_ Wall/Furn	Hard_Toy	Puddle_ Water	Footwear	Nothing

Object/surface categories

PAUSE
OFF ON

Time: 10:00:00 AM
Counter:
Index: 0

Video ID: New Video
Subject ID: New Subject
Coder ID: New Coder
Boundary: Right_Hand

Figure F-18. Virtual Timing Device palette used by Stanford's Exposure Research Group for translation.

F.2.2.2. Data Processing

F.2.2.2.1. Children Playing on Turf

The translated plain text files, obtained from ORETF and EPA, for all 56 children for right hand, left hand, and mouthing activities were used for the analysis. The object/surfaces were selected as presented in Table F-18. Participants' contacts while playing in the location categories of yard, garden, and park were extracted and grouped as one turf-like location by using RStudio V1.1 (RStudio Team, 2016). Videotapes were reviewed to confirm turf like locations.

F.2.2.2.2. Children Playing on Playgrounds

Since playground was not categorized in the previous studies, all of the existing videotapes (n=56) were re-watched to only select footage where a playground structure was observed. The exact footage time and duration when children played on or near playground structures was recorded and then a specific "playground" location was added to each corresponding data file. The objects/surfaces selected for analysis are presented in Table F-19. All floor surfaces were grouped in one category for the playground locations (Table F-19) because we are assuming that playgrounds are on rubber mats made of tire or of chopped up tire.



Table F-18. Selected Categories for Object/Surfaces on Turf

Super Categories	Virtual Timing Device Palette Categories (Figure F-18)
<i>Location: Turf</i>	Yard, park, garden
<i>Object/Surface: Grass</i>	Grass
<i>Object/Surface: Dietary objects</i>	Water/beverage, sticky food, other food, food container
<i>Object/Surface: Non-dietary objects</i>	Everything, but dietary objects
<i>Object/Surface: Hands*</i>	Hands
<i>Object/Surface: All objects/surfaces</i>	Wood wall, wood tools, wood toy, vegetation, hard toys, porous plastic toys, fabric toys, plastic tool, plastic wall, paper, pool water, puddle water, metal wall, metal tool, footwear, carpet, wood floor, tile floor, rock floor, sidewalk, dirt

* Only analyzed for mouth contact files

Table F-19. Selected Categories for Object/Surfaces on Playgrounds

Super Categories	Virtual Timing Device Palette Categories (Figure F-18)
<i>Location: Playground</i>	Child near playground in the following locations: Yard, Park, Garden, Patio, Street, Garage
<i>Object/Surface: Floors</i>	Dirt, Asphalt, Rock floor, wood floor, tile, carpet/mat
<i>Object/Surface: Dietary objects</i>	Water/beverage, sticky food, other food, food container
<i>Object/Surface: Non-dietary objects</i>	Everything, but dietary objects
<i>Object/Surface: Hands*</i>	Hands
<i>Object/Surface: All objects/surfaces</i>	Wood wall, wood tools, wood toy, vegetation, hard toys, porous plastic toys, fabric toys, plastic tool, plastic wall, paper, pool water, puddle water, metal wall, metal tool, footwear, carpet, wood floor, tile floor, rock floor, sidewalk, dirt

* Only analyzed for mouth contact files

F.2.2.3. Data Analysis

Using RStudio V1.1, right hand, left hand, and mouth contact frequency, hourly contact duration and median contact duration with the selected object/surface categories (Table F-18 and Table F-19) were calculated for each child while playing on turf and playground structures, respectively.



Contact frequency (events/hour) was calculated by summing the total number of contact events by hands or mouth with a specific object category divided by the total time that the child was in view.

Hourly duration (minutes/hour) was calculated by summing the total time in minutes that hands or mouth were in contact with the specific object category divided by the total time in view (hours).

Median duration (seconds) was obtained from the contact duration for the hand or mouth contact events with a specific object/category during the time in view for each child. It is the median value of all contacts of the body part (mouth/hand) with the object. Only children who contacted the object were included in these calculations.

Since Wilcoxon signed rank is for paired data on the same person (dependent data), we used it to compare to assess differences in the activity variables (contact frequency, hourly duration and median duration) between right and left hand. If no differences were observed, then data for the hands were combined and reported as “both hands.”

The data was then summarized based on age and gender. The activity variables (contact frequency, hourly duration and median duration) were evaluated for significant differences between genders and by age groups using non-parametric tests. To determine differences between male and female participants, we used the two tailed Wilcoxon-rank sum test (non-parametric test, similar to a t-test, for independent data).

To determine if contact activities correlated with age, Spearman’s rank correlation coefficient was computed using STATA Version 12.0. (College Station, TX: StataCorp). The data for each contact activity was summarized into three age groups: 0 <2 years, 2 <9 years, and 9 <16 years. The Kruskal-Wallis test (non-parametric test similar to ANOVA) was used to assess if there are differences across the three age groups. Also, as presented in Supplemental Material Sections S2.1 and S2.2, the data was summarized into additional age groups by following the US EPA guidance (U.S. EPA, 2005). Differences between the activities of the 1-6 year old children and the 7-12 year old children was assessed using the two tailed Wilcoxon-rank sum test (see Supplemental Material Sections S2.1 and S2.2).

F.2.3. Results

A total of 43 hours (2548 minutes) of footage was collected from 56 children with a median footage time per child of 112 min (range: 60 – 133 min). This was the total amount of footage that was available for the retrospective analysis.

F.2.3.1. Children Playing on Turf:

F.2.3.1.1. Amount of Footage with Turf and Demographics of Children Who Played on Turf

All 56 children played on turf at some point during their footage, with a median time on



turf of 84 min (Table F-20). Of the footage of children playing on turf, 3.9% of the time hands and/or mouth was not in view and excluded from analysis.

There were comparable numbers of male (n=27) and female (n=29) participants (Table F-21). While among both male and females there were more children within the 2 <9 years of age than the other two age groups. As presented in Supplemental Material Section S2.1, for the EPA age groups, the male children had an even distribution among the age groups, while female children were disproportionally distributed between the ages of 3 and <11 years old. However, there are similar proportions of female children below and above the age of 6.

Table F-20. Time Spent on Turf (N=56)

Time on turf	Hand and/or mouth in view	Hand and/or mouth not in view
Total turf time (min)	1812.0	99.5
Median turf time (min per child)	84.0	2.5
Percentage of time spent on turf (%)	71.1*	3.9*

* Calculated from total footage time of 2,548 minutes

Table F-21. Number of Children Playing on Turf Grouped by Age Groups and Gender

Gender	0 <2 years	2 <9 years	9 <16 years	Total
Male	5	15	7	27
Female	3	21	5	29
Total	8	36	12	56

F.2.3.1.2. Hand Contact Activity While Playing on Turf

There were no significant differences in the contact frequency with object/surfaces between right hand and left hand (Table F-22 and Table F-23). Therefore, both hands were combined and summarized (Table F-24). This means that contact frequency for the left and right hands were summed, however it does not mean that both hands were in contact with the object at the same time. The median grass contact frequency for both hands combined was 4.14 events/hour. Although the median contact frequency with “grass” was relatively low (less than once every 10 minutes), the child with the greatest potential exposure was contacting the “grass” very frequently (over 4 times per minute).

Table F-22. Right Hand Contact Frequency (Event/Hour) While Playing on the Turf (n=56)

Distribution	Grass	Dietary	Non-Dietary	All objects
Mean	7.20	3.10	129.40	132.50



Distribution	Grass	Dietary	Non-Dietary	All objects
SD	17.50	5.40	54.60	54.30
Min	0.00	0.00	57.30	57.30
p25	1.51	0.00	93.92	96.00
Median	3.73	0.14	112.50	115.68
p75	10.00	3.28	151.30	155.68
p95	26.50	15.34	221.12	221.12
p99	89.65	20.68	297.78	298.40
Max	122.33	20.72	371.99	373.36

Table F-23. Left Hand Contact Frequency (Event/Hour) While Playing on the Turf (n=56)

Distribution	Grass	Dietary	Non-Dietary	All objects
Mean	7.00	3.10	134.20	137.30
SD	18.20	5.20	68.10	67.40
Min	0.00	0.00	9.60	9.60
p25	1.22	0.00	101.04	103.84
Median	3.55	0.15	118.70	127.17
p75	8.03	4.80	145.58	149.58
p95	30.91	14.92	242.44	242.72
p99	90.08	19.62	376.70	377.20
Max	129.54	19.64	393.85	394.96

Table F-24. Both Hands Object/Surface Contact Frequency (Event/Hour) (n=56)

Distribution	Grass	Dietary	Non-Dietary	All objects
Mean	14.50	6.10	276.80	282.90
SD	35.50	10.30	126.70	125.30
Min	0.00	0.00	107.70	109.70
p25	0.78	0.00	192.78	205.57
Median	4.14	0.71	236.31	246.32



Distribution	Grass	Dietary	Non-Dietary	All objects
p75	14.93	8.82	308.21	324.38
p95	48.57	27.03	503.25	504.34
p99	151.84	40.30	718.79	719.90
Max	251.87	40.33	765.84	768.32
p-value*	0.7249	0.935	0.535	0.546

*Wilcoxon Signed Rank test comparison of left- and right-hand contact frequency

There were no significant differences in hourly contact duration with object/surfaces between right hand and left hand (Table F-25 and Table F-26). Therefore, both hands were combined and summarized (Table F-27). This means that contact hourly duration for the left and right hands were summed, however it does not mean that both hands were in contact with the object at the same time. If both hands were in contact with object for the entire time, the maximum value would be 120 min/hour. The median hourly contact duration for both hands with grass was 0.24 min/hour. However, one child was in contact with grass 12 min/hour.

Table F-25. Right Hand Hourly Contact Duration (Min/Hour) While Playing on the Turf (n=56)

Distribution	Grass	Dietary	Non-Dietary	All Objects
Mean	0.44	1.75	33.19	34.94
SD	0.97	3.08	9.61	9.04
Min	0.00	0.00	11.88	11.88
p25	0.00	0.00	26.33	29.04
Median	0.09	0.01	32.21	34.22
p75	0.36	2.41	39.14	40.28
p95	2.40	8.62	48.93	50.22
p99	5.87	11.99	53.06	55.06
Max	5.87	11.99	55.06	55.06

Table F-26. Left Hand Hourly Contact Duration (Min/Hour) While Playing on the Turf (n=56)

Distribution	Grass	Dietary	Non-Dietary	All objects
Mean	0.45	1.61	31.94	33.55



Distribution	Grass	Dietary	Non-Dietary	All objects
SD	0.97	2.77	10.19	9.90
Min	0.00	0.00	5.68	5.68
p25	0.01	0.00	26.48	27.20
Median	0.11	0.01	31.23	32.94
p75	0.37	2.07	38.01	41.04
p95	2.20	7.46	48.49	48.49
p99	6.09	9.93	57.64	57.64
Max	6.09	9.93	57.64	57.64

Table F-27. Hourly Object/Surface Contact Duration for Both Hands While Playing on Turf (Min/Hour) (n=56)

Distribution	Grass	Dietary	Non-Dietary	All objects
Mean	0.93	1.81	36.57	38.38
SD	1.97	3.08	13.71	14.60
Min	0.00	0.00	20.65	22.36
p25	0.05	0.00	29.26	31.55
Median	0.24	0.12	33.33	34.83
p75	1.22	2.47	35.98	36.97
p95	4.63	8.10	64.64	67.61
p99	8.02	11.90	81.66	87.12
Max	11.96	12.14	92.96	105.10
p-value*	0.876	0.265	0.167	0.071

*Wilcoxon Signed Rank test comparison of left- and right-hand contact frequency

There were no significant differences in median contact duration with object/surfaces between right hand and left hand (Table F-28 and Table F-29). Therefore, both hands were combined, re-analyzed, and summarized (Table F-30). This means that for each object category the durations of all the discrete contacts with an object for both the right and left hands were grouped and reranked to determine the median value of contact duration across all the “hand” contacts with that object. Note that the median for aggregate categories like “All Objects” was calculated as the median duration of contacts with any object. The median contact duration of both hands combined and



grass was 2.00 seconds. However, one child had a median contact duration of both hands with grass of 13.00 seconds.

Table F-28. Right Hand Median Contact Duration While Playing on the Turf (Second)

Distribution	Grass	Dietary	Non-Dietary	All Objects
n*	38	28	56	56
Mean	2.54	8.07	3.37	3.44
SD	1.48	6.11	1.08	1.10
Min	0.50	1.00	1.00	1.00
p25	1.50	4.00	3.00	3.00
Median	2.00	6.50	3.00	3.00
p75	3.00	11.63	4.00	4.00
p95	5.00	16.65	5.00	5.00
p99	6.83	24.30	5.73	5.73
Max	8.00	27.00	6.00	6.00

*Only participants who contacted object included in calculation of median contact duration

Table F-29. Left Hand Median Contact Duration While Playing on the Turf (Second)

Distribution	Grass	Dietary	Non-Dietary	All objects
n*	42	28	56	56
Mean	3.52	7.57	3.28	3.29
SD	4.19	5.63	0.89	0.87
Min	0.50	1.00	2.00	2.00
p25	2.00	3.00	3.00	3.00
Median	2.00	6.50	3.00	3.00
p75	3.00	10.38	4.00	4.00
p95	12.33	18.13	5.00	4.63
p99	19.48	20.46	5.23	5.23
Max	20.50	21.00	5.50	5.50

*Only participants who contacted object included in calculation of median contact duration



Table F-30. Both Hands Median Contact Duration While Playing on the Turf (Second)

Distribution	Grass	Dietary	Non - Dietary	All objects
n*	42	28	56	56
Mean	2.72	7.47	3.28	3.31
SD	2.05	5.25	0.92	0.95
Min	0.50	1.00	1.00	1.00
p25	2.00	4.00	3.00	3.00
Median	2.00	6.75	3.00	3.00
p75	3.00	10.00	4.00	4.00
p95	5.00	18.00	5.00	5.00
p99	13.00	23.00	5.00	5.00
Max	13.00	23.00	5.00	5.00
p-value*	0.815	0.367	0.611	0.245

*p-value from Wilcoxon Signed Rank test comparison of left- and right-hand contact frequency

Hand Contact Activity by Age While Playing on Turf

Frequency, hourly duration and median duration for both hands contact activities while playing on turf are presented in Table F-31 to Table F-33 for the three age categories. Although there were more children in the 2 <9 years of age group, most of the contact with grass seem to have occurred in the older group (9 <16 years old). There were no significant differences in contact frequency, hourly contact duration, or median contact duration across all three age groups (Kruskal-Wallis Test). Also, there were no correlations between age groups and both hand activities (Table F-34).

Table F-31. Both Hands Contact Frequency (Event/Hour) While Playing on Turf by Age Groups

Age Group	Distribution	Grass	Dietary	Non-Dietary	All objects
<2 (n=8)	Mean	11.06	1.42	285.68	287.10
<2 (n=8)	SD	11.83	3.56	75.83	76.10
<2 (n=8)	Min	0.34	0.00	187.91	187.91
<2 (n=8)	p25	2.92	0.00	238.21	238.21
<2 (n=8)	Median	8.52	0.00	282.76	287.87
<2 (n=8)	p75	14.01	0.59	308.16	308.37



Age Group	Distribution	Grass	Dietary	Non-Dietary	All objects
<2 (n=8)	p95	37.24	10.22	439.28	440.02
<2 (n=8)	p99	37.24	10.22	439.28	440.02
<2 (n=8)	Max	37.24	10.22	439.28	440.02
2 to <9 (n=36)	Mean	8.00	6.76	271.32	278.08
2 to <9 (n=36)	SD	13.24	10.28	143.03	140.67
2 to <9 (n=36)	Min	0.00	0.00	107.69	109.66
2 to <9 (n=36)	p25	0.00	0.00	191.80	200.72
2 to <9 (n=36)	Median	2.61	1.47	218.08	228.40
2 to <9 (n=36)	p75	11.59	9.12	290.15	290.15
2 to <9 (n=36)	p95	29.39	27.91	680.29	680.29
2 to <9 (n=36)	p99	70.00	40.28	765.84	768.32
2 to <9 (n=36)	Max	70.00	40.28	765.84	768.32
9 to <16 (n=12)	Mean	36.41	7.13	287.30	294.43
9 to <16 (n=12)	SD	70.56	13.00	106.31	107.87
9 to <16 (n=12)	Min	0.00	0.00	138.75	138.75
9 to <16 (n=12)	p25	3.39	0.00	210.38	212.37
9 to <16 (n=12)	Median	7.51	0.59	276.31	296.81
9 to <16 (n=12)	p75	40.13	7.36	351.09	353.82
9 to <16 (n=12)	p95	251.87	40.33	494.87	494.87
9 to <16 (n=12)	p99	251.87	40.33	494.87	494.87
9 to <16 (n=12)	Max	251.87	40.33	494.87	494.87
p-value*	Not applicable	0.081	0.281	0.398	0.368

*p-value from Kruskal-Wallis Test

Table F-32. Both Hands Hourly Contact Duration (Min/Hour) While Playing on Turf by Age Groups

Age Group	Distribution	Grass	Dietary	Non-Dietary	All objects
<2 (n=8)	Mean	0.90	0.71	29.58	36.38
<2 (n=8)	SD	1.59	1.72	14.02	15.59



Age Group	Distribution	Grass	Dietary	Non-Dietary	All objects
<2 (n=8)	Min	0.01	0.00	15.63	26.78
<2 (n=8)	p25	0.10	0.00	22.66	28.18
<2 (n=8)	Median	0.49	0.01	26.90	29.61
<2 (n=8)	p75	0.61	0.39	29.87	38.14
<2 (n=8)	p95	4.79	4.91	62.19	72.41
<2 (n=8)	p99	4.79	4.91	62.19	72.41
<2 (n=8)	Max	4.79	4.91	62.19	72.41
2 to <9 (n=36)	Mean	0.63	2.19	32.79	37.84
2 to <9 (n=36)	SD	1.21	3.17	11.64	11.10
2 to <9 (n=36)	Min	0.00	0.00	18.85	22.36
2 to <9 (n=36)	p25	0.03	0.00	25.07	33.26
2 to <9 (n=36)	Median	0.20	0.16	30.97	35.40
2 to <9 (n=36)	p75	0.49	3.51	34.87	36.99
2 to <9 (n=36)	p95	4.63	8.16	62.51	67.25
2 to <9 (n=36)	p99	4.63	11.71	65.28	68.70
2 to <9 (n=36)	Max	4.63	11.71	65.28	68.70
9 to <16 (n=12)	Mean	1.87	1.39	34.82	41.35
9 to <16 (n=12)	SD	3.44	3.46	22.90	22.49
9 to <16 (n=12)	Min	0.00	0.00	14.72	22.89
9 to <16 (n=12)	p25	0.06	0.00	20.87	30.87
9 to <16 (n=12)	Median	0.51	0.01	30.87	34.02
9 to <16 (n=12)	p75	2.00	1.04	35.41	37.19
9 to <16 (n=12)	p95	11.96	12.14	96.03	105.10
9 to <16 (n=12)	p99	11.96	12.14	96.03	105.10
9 to <16 (n=12)	Max	11.96	12.14	96.03	105.10
p-value*	Not applicable	0.295	0.201	0.474	0.218

*p-value from Kruskal-Wallis Test



Table F-33. Both Hands Median Contact Duration (Second) While Playing on Turf by Age Groups

Age Group	Distribution	Grass	Dietary	Non-Dietary	All objects
<2	n	8	3	8	8
<2	Mean	2.50	4.00	3.63	3.63
<2	SD	0.76	4.77	0.52	0.52
<2	Min	2.00	1.00	3.00	3.00
<2	p25	2.00	1.00	3.00	3.00
<2	Median	2.00	1.50	4.00	4.00
<2	p75	3.00	9.50	4.00	4.00
<2	p95	4.00	9.50	4.00	4.00
<2	p99	4.00	9.50	4.00	4.00
<2	Max	4.00	9.50	4.00	4.00
2 to <9	n	24	22	36	36
2 to <9	Mean	2.54	9.09	3.38	3.43
2 to <9	SD	1.62	5.35	0.83	0.87
2 to <9	Min	0.50	1.00	2.00	2.00
2 to <9	p25	1.75	6.00	3.00	3.00
2 to <9	Median	2.00	7.75	3.25	3.75
2 to <9	p75	3.00	12.00	4.00	4.00
2 to <9	p95	5.00	18.00	5.00	5.00
2 to <9	p99	8.00	23.00	5.00	5.00
2 to <9	Max	8.00	23.00	5.00	5.00
9 to <16	n	11	7	12	12
9 to <16	Mean	3.27	3.86	2.75	2.75
9 to <16	SD	3.29	1.93	1.22	1.22
9 to <16	Min	1.00	1.00	1.00	1.00
9 to <16	p25	2.00	2.50	2.00	2.00
9 to <16	Median	2.00	4.00	2.50	2.50



Age Group	Distribution	Grass	Dietary	Non-Dietary	All objects
9 to <16	p75	3.00	6.00	3.00	3.00
9 to <16	p95	13.00	6.50	5.00	5.00
9 to <16	p99	13.00	6.50	5.00	5.00
9 to <16	Max	13.00	6.50	5.00	5.00
p-value*	Not applicable	0.744	0.134	0.054	0.054

*p-value from Kruskal-Wallis Test

Table F-34. Spearman Rank Correlation for Age (Years) and Hand Contact Events

Activity Variables	Contact	r	p-value
Hand contact frequency (event/hour)	Grass (n=56)	0.1371	0.314
Hand contact frequency (event/hour)	Non-Dietary (n=56)	0.0839	0.538
Hand contact frequency (event/hour)	Dietary (n=56)	-0.003	0.983
Hand contact frequency (event/hour)	All objects (n=56)	0.087	0.523
Hourly hand contact duration (min/hour)	Grass (n=56)	0.113	0.406
Hourly hand contact duration (min/hour)	Non-Dietary (n=56)	0.098	0.471
Hourly hand contact duration (min/hour)	Dietary (n=56)	-0.083	0.542
Hourly hand contact duration (min/hour)	All objects (n=56)	0.139	0.306
Hand contact median duration (second)	Grass (n=42)	0.137	0.314
Hand contact median duration (second)	Non-Dietary (n=56)	0.084	0.538
Hand contact median duration (second)	Dietary (n=28)	-0.003	0.983
Hand contact median duration (second)	All objects (n=56)	0.087	0.523

The hand contact frequency, hourly duration and median duration summary of the EPA recommended age categories (U.S. EPA, 2005) are presented in Supplemental Material Section S2.1. There were no significant differences in contact frequency, hourly contact duration, or median contact duration across the EPA recommended age groups (Kruskal-Wallis Test). Also, in Supplemental Material Section S2.1, the contact frequency, hourly duration and median duration for both hands contact activities while playing on turf are presented in Table S2-5 to Table S2-8 for children aged 1-6 years and 7-12. There were no significant difference between these two age groups and contact activities (Table S2-9).

Hand Contact Activity by Gender While Playing on Turf



As presented in Table F-35 to Table F-37 by gender (i.e., male, female), there were no significant differences in contact frequency, hourly contact duration, or median contact duration by gender while playing on turf.

Table F-35. Both Hands Contact Frequency (Event/Hour) While Playing on Turf by Gender

Gender	Distribution	Grass	Dietary	Non-Dietary	All Objects
Male (n=27)	Mean	12.54	5.11	267.70	272.80
Male (n=27)	SD	15.15	9.45	137.05	137.48
Male (n=27)	Min	0.00	0.00	107.69	109.66
Male (n=27)	Median	1.08	0.00	189.18	190.04
Male (n=27)	p25	7.34	0.86	227.39	228.25
Male (n=27)	p75	17.43	4.36	300.83	320.07
Male (n=27)	p95	45.03	26.61	528.37	532.73
Male (n=27)	p99	59.17	40.33	765.84	768.32
Male (n=27)	Max	59.17	40.33	765.84	768.32
Female (n=29)	Mean	16.38	6.98	285.27	292.25
Female (n=29)	SD	47.47	11.14	118.16	114.49
Female (n=29)	Min	0.00	0.00	163.11	173.66
Female (n=29)	Median	0.34	0.00	211.05	222.22
Female (n=29)	p25	3.78	0.00	253.24	253.24
Female (n=29)	p75	9.70	8.86	325.19	349.05
Female (n=29)	p95	70.00	27.91	494.87	494.87
Female (n=29)	p99	251.87	40.28	680.29	680.29
Female (n=29)	Max	251.87	40.28	680.29	680.29
p-value*	Not applicable	0.242	0.621	0.321	0.216

*p-value from Wilcoxon sum rank

Table F-36. Both Hands Hourly Contact Duration (Min/Hour) While Playing on Turf by Gender

Gender	Distribution	Grass	Dietary	Non-Dietary	All Objects
Male (n=27)	Mean	0.94	1.86	33.78	39.52



Gender	Distribution	Grass	Dietary	Non-Dietary	All Objects
Male (n=27)	SD	1.40	3.52	17.40	18.16
Male (n=27)	Min	0.00	0.00	14.72	22.36
Male (n=27)	Median	0.09	0.00	23.44	30.02
Male (n=27)	p25	0.40	0.13	29.74	34.01
Male (n=27)	p75	0.96	2.36	34.97	37.26
Male (n=27)	p95	4.63	11.71	62.19	72.41
Male (n=27)	p99	4.79	12.14	96.03	105.10
Male (n=27)	Max	4.79	12.14	96.03	105.10
Female (n=29)	Mean	0.92	1.76	31.82	37.32
Female (n=29)	SD	2.41	2.66	12.10	10.48
Female (n=29)	Min	0.00	0.00	16.75	23.24
Female (n=29)	Median	0.01	0.00	23.47	31.58
Female (n=29)	p25	0.11	0.11	29.83	35.19
Female (n=29)	p75	0.38	2.80	34.18	36.95
Female (n=29)	p95	4.63	8.08	62.51	65.20
Female (n=29)	p99	11.96	8.16	65.28	67.25
Female (n=29)	Max	11.96	8.16	65.28	67.25
p-value*	Not applicable	0.059	0.679	0.974	0.486

*p-value from Wilcoxon sum rank

Table F-37. Both Hands Contact Median Duration (Second) While Playing on Turf by Gender

Gender	Distribution	Grass	Dietary	Non-Dietary	All Objects
Male	n	22	27	27	27
Male	Mean	2.70	16.67	62.50	79.17
Male	SD	1.67	45.85	30.09	52.22
Male	Min	0.50	0.00	22.75	34.00
Male	Median	2.00	0.00	39.00	50.50
Male	p25	2.00	1.50	57.00	59.00



Gender	Distribution	Grass	Dietary	Non-Dietary	All Objects
Male	p75	3.00	19.00	80.00	87.00
Male	p95	5.00	31.50	114.50	173.00
Male	p99	8.00	241.00	159.00	282.50
Male	Max	8.00	241.00	159.00	282.50
Female	n	21	29	29	29
Female	Mean	2.54	9.69	77.80	87.49
Female	SD	2.12	33.31	54.47	65.71
Female	Min	0.50	0.00	22.50	22.50
Female	Median	1.50	0.00	52.25	52.25
Female	p25	2.00	0.00	61.00	74.50
Female	p75	2.50	16.00	85.25	90.00
Female	p95	4.00	42.50	135.50	191.50
Female	p99	13.00	56.00	415.50	445.00
Female	Max	13.00	56.00	415.50	445.00
p-value	Not applicable	0.324	0.567	0.358	0.566

*p-value from Wilcoxon sum rank

F.2.3.1.3. Mouthing Activity While Playing on Turf

The mouthing contact frequency for children playing on turf is summarized in Table F-38. Only 7% of the children (4/56) contacted grass with their mouth. Thus, the median mouth contact frequency for grass was 0.00 events/hour, and the maximum was 2.49 events/hour. The median mouthing frequencies were 7.56 events/hour and 10.87 events/hour for hands and non-dietary objects, respectively. These contacts could indicate additional exposure to crumb rubber if there have been transfers to the child's hands (Section F.2.3.1.2) or the objects that they are placing in their mouths.

Table F-38. Mouth Contact Frequency (Event/Hour) While Playing on the Turf (n=56)

Distribution	Grass	Hand	Dietary	Non-Dietary	All Objects
Mean	0.11	11.65	14.83	22.00	36.83
SD	0.44	14.16	28.40	29.87	44.66
Min	0.00	0.00	0.00	0.00	0.00
p25	0.00	3.38	0.00	6.61	7.28



Distribution	Grass	Hand	Dietary	Non-Dietary	All Objects
Median	0.00	7.56	0.84	10.87	20.39
p75	0.00	15.74	19.82	26.36	56.17
p95	1.17	41.04	74.00	76.19	160.77
p99	2.49	80.11	159.12	185.14	205.09
Max	2.49	80.11	159.12	185.14	205.09

The mouthing hourly contact duration for children playing on turf is summarized in Table F-39. The median mouthing hourly contact duration for grass was 0.00 min/hour, and the maximum was 0.10 min/hour. The median mouthing hourly durations were 0.21 min/hour and 0.30 min/hour for hands and non-dietary objects, respectively. These contacts could indicate additional exposure to crumb rubber if there have been transfers to the child's hands (Section F.2.3.1.2) or the objects that they are placing in their mouths.

Table F-39. Mouthing Hourly Contact Duration (Min/Hour) While Playing on the Turf (n=56)

Distribution	Grass	Hand	Dietary	Non-Dietary	All Objects
Mean	0.00	0.46	1.80	1.39	3.19
SD	0.01	0.76	5.30	4.65	7.35
Min	0.00	0.00	0.00	0.00	0.00
p25	0.00	0.10	0.00	0.14	0.24
Median	0.00	0.21	0.04	0.30	0.66
p75	0.00	0.42	1.17	1.31	3.06
p95	0.02	2.25	9.47	3.92	12.53
p99	0.10	3.54	36.52	34.71	39.65
Max	0.10	3.54	36.52	34.71	39.65

The mouthing median contact duration for children playing on turf is summarized in Table F-40. Median duration is only calculated from those children that contacted the object. The median mouthing contact duration for grass was 1.00 second, and the maximum was 2.5 seconds. The median mouthing contact durations were 1.00 second and 1.00 second for hands and non-dietary objects, respectively. These contacts could indicate additional exposure to crumb rubber if there have been transfers to the child's hands (Section F.2.3.1.2) or the objects that they are placing in their mouths.



Table F-40. Mouthing Median Contact Duration While Playing on Turf (Second)

Distribution	Grass	Hand	Dietary	Non-Dietary	All Objects
n*	4	49	30	52	53
Mean	1.38	1.86	7.40	1.88	2.89
SD	0.75	1.90	22.05	1.87	8.09
Min	1.00	0.50	1.00	0.00	0.00
p25	1.00	1.00	2.00	1.00	1.00
Median	1.00	1.00	2.00	1.00	1.00
p75	1.75	2.00	4.00	2.00	2.00
p95	2.50	5.00	14.00	5.00	5.50
p99	2.50	12.00	123.00	12.00	60.00
Max	2.50	12.00	123.00	12.00	60.00

*Only participants who contacted object included in calculation of median contact duration

Mouthing Contact Activity by Age While Playing on Turf

Frequency, hourly duration and median duration for mouthing contact activities while playing on turf are presented in Table F-41 to Table F-43 for three age categories (<2, 2 to <9, 9 to <16 years). There were no differences in mouthing contact frequency, hourly duration or median duration across these three age groups. Age (by year, 1-12 years) was negatively correlated ($p < 0.05$) with the hourly contact duration of mouthing activities for non-dietary objects and all objects (Table F-44). Age was also negatively correlated ($p < 0.05$) with the median contact duration of mouthing activities for hands and non-dietary objects (Table F-44).

Table F-41. Mouthing Contact Frequency (Event/Hour) While Playing on Turf by Age Groups

Age Group	Distribution	Grass	Hands	Dietary	Non-Dietary	All Objects
<2 (n=8)	Mean	0.34	12.74	2.79	25.69	28.48
<2 (n=8)	SD	0.65	11.69	4.39	22.61	23.62
<2 (n=8)	Min	0.00	0.66	0.00	1.48	2.96
<2 (n=8)	Median	0.00	3.70	0.00	9.44	9.44



Age Group	Distribution	Grass	Hands	Dietary	Non-Dietary	All Objects
<2 (n=8)	p25	0.00	11.99	0.74	16.30	19.68
<2 (n=8)	p75	0.51	16.56	4.41	44.37	51.41
<2 (n=8)	p95	1.69	36.78	12.05	63.82	63.82
<2 (n=8)	p99	1.69	36.78	12.05	63.82	63.82
<2 (n=8)	Max	1.69	36.78	12.05	63.82	63.82
2 to <9 (n=36)	Mean	0.10	8.95	19.03	20.81	39.84
2 to <9 (n=36)	SD	0.45	8.98	32.70	32.37	47.91
2 to <9 (n=36)	Min	0.00	0.00	0.00	0.00	0.00
2 to <9 (n=36)	Median	0.00	2.91	0.00	6.85	7.70
2 to <9 (n=36)	p25	0.00	7.12	1.21	9.85	23.53
2 to <9 (n=36)	p75	0.00	11.03	31.66	24.80	60.66
2 to <9 (n=36)	p95	1.17	33.69	80.19	76.19	189.59
2 to <9 (n=36)	p99	2.49	41.04	159.12	185.14	205.09
2 to <9 (n=36)	Max	2.49	41.04	159.12	185.14	205.09
9 to <16 (n=12)	Mean	ND	19.06	10.26	23.09	33.36
9 to <16 (n=12)	SD	ND	23.96	20.79	28.00	47.11
9 to <16 (n=12)	Min	ND	0.00	0.00	0.00	0.00
9 to <16 (n=12)	Median	ND	3.76	0.00	4.94	6.50
9 to <16 (n=12)	p25	ND	8.07	0.75	10.43	11.29
9 to <16 (n=12)	p75	ND	27.12	9.02	38.14	42.53
9 to <16 (n=12)	p95	ND	80.11	72.43	88.34	160.77
9 to <16 (n=12)	p99	ND	80.11	72.43	88.34	160.77
9 to <6 (n=12)	Max	ND	80.11	72.43	88.34	160.77
p-value*	Not applicable	0.091	0.444	0.643	0.690	0.726

*p-value from Kruskal-Wallis Test

ND: not determined



Table F-42. Mouthing Hourly Contact Duration (Min/Hour) While Playing on Turf by Age Groups

Age Group	Distribution	Grass	Hands	Dietary	Non-Dietary	All Objects
<2 (n=8)	Mean	0.01	1.13	0.65	1.54	2.18
<2 (n=8)	SD	0.01	1.50	1.39	1.52	1.84
<2 (n=8)	Min	0.00	0.03	0.00	0.16	0.16
<2 (n=8)	Median	0.00	0.24	0.00	0.30	0.31
<2 (n=8)	p25	0.00	0.33	0.01	0.96	2.09
<2 (n=8)	p75	0.01	2.17	0.58	2.84	3.98
<2 (n=8)	p95	0.03	3.54	4.02	3.95	4.57
<2 (n=8)	p99	0.03	3.54	4.02	3.95	4.57
<2 (n=8)	Max	0.03	3.54	4.02	3.95	4.57
2 to <9 (n=36)	Mean	0.00	0.27	2.45	1.61	4.05
2 to <9 (n=36)	SD	0.02	0.38	6.47	5.74	8.97
2 to <9 (n=36)	Min	0.00	0.00	0.00	0.00	0.00
2 to <9 (n=36)	Median	0.00	0.08	0.00	0.13	0.18
2 to <9 (n=36)	p25	0.00	0.16	0.07	0.28	0.78
2 to <9 (n=36)	p75	0.00	0.29	1.58	0.93	3.12
2 to <9 (n=36)	p95	0.02	1.59	10.53	3.92	36.66
2 to <9 (n=36)	p99	0.10	1.75	36.52	34.71	39.65
2 to <9 (n=36)	Max	0.10	1.75	36.52	34.71	39.65
9 to <16 (n=12)	Mean	ND	0.56	0.63	0.66	1.28
9 to <16 (n=12)	SD	ND	0.73	1.42	0.89	2.09
9 to <16 (n=12)	Min	ND	0.00	0.00	0.00	0.00
9 to <16 (n=12)	Median	ND	0.10	0.00	0.12	0.20
9 to <16 (n=12)	p25	ND	0.25	0.09	0.26	0.41
9 to <16 (n=12)	p75	ND	0.79	0.54	0.92	1.18
9 to <16 (n=12)	p95	ND	2.25	4.99	2.71	7.06
9 to <16 (n=12)	p99	ND	2.25	4.99	2.71	7.06



Age Group	Distribution	Grass	Hands	Dietary	Non-Dietary	All Objects
9 to <16 (n=12)	Max	ND	2.25	4.99	2.71	7.06
p-value*	Not applicable	0.094	0.075	0.797	0.145	0.415

*p-value from Kruskal-Wallis Test

ND: not determined

Table F-43. Mouthing Contact Median Duration (Second) While Playing on Turf by Age Groups

Age Group	Distribution	Grass	Hands	Dietary	Non-Dietary	All Objects
<2	n	2	8	4	8	8
<2	Mean	1.00	3.69	6.88	3.44	2.63
<2	SD	0.00	3.83	5.81	3.90	2.15
<2	Min	1.00	1.00	1.00	1.00	1.00
<2	Median	1.00	1.00	2.25	1.00	1.00
<2	p25	1.00	2.50	6.25	2.00	2.00
<2	p75	1.00	4.75	11.50	4.25	3.75
<2	p95	1.00	12.00	14.00	12.00	6.50
<2	p99	1.00	12.00	14.00	12.00	6.50
<2	Max	1.00	12.00	14.00	12.00	6.50
2 to <9	n	2	30	19	33	34
2 to <9	Mean	1.75	1.58	9.11	1.71	3.46
2 to <9	SD	1.06	1.11	27.65	1.19	10.05
2 to <9	Min	1.00	0.50	1.00	0.00	0.00
2 to <9	Median	1.00	1.00	1.00	1.00	1.00
2 to <9	p25	1.75	1.00	2.00	1.00	1.50
2 to <9	p75	2.50	2.00	4.00	2.00	2.00
2 to <9	p95	2.50	5.00	123.00	5.00	5.00
2 to <9	p99	2.50	5.00	123.00	5.00	60.00
2 to <9	Max	2.50	5.00	123.00	5.00	60.00
9 to <16	n	0	11	7	11	11



Age Group	Distribution	Grass	Hands	Dietary	Non-Dietary	All Objects
9 to <16	Mean	ND	1.27	3.07	1.23	1.32
9 to <16	SD	ND	0.47	2.65	0.41	0.46
9 to <16	Min	ND	1.00	1.50	1.00	1.00
9 to <16	Median	ND	1.00	2.00	1.00	1.00
9 to <16	p25	ND	1.00	2.00	1.00	1.00
9 to <16	p75	ND	2.00	3.00	1.50	2.00
9 to <16	p95	ND	2.00	9.00	2.00	2.00
9 to <16	p99	ND	2.00	9.00	2.00	2.00
9 to <16	Max	ND	2.00	9.00	2.00	2.00
p-value*	Not applicable	0.317	0.097	0.461	0.187	0.306

*p-value from Kruskal-Wallis Test

ND: not determined

Table F-44. Spearman Rank Correlation (r) for Age (1 - 12 Years) and Mouthing Events

Activity Variables	Contact	r	p-value
Mouth contact frequency (event/hour)	Grass (n=56)	-0.221	0.101
Mouth contact frequency (event/hour)	Hands (n=56)	-0.033	0.811
Mouth contact frequency (event/hour)	Non-Dietary (n=56)	-0.124	0.364
Mouth contact frequency (event/hour)	Dietary (n=56)	-0.058	0.672
Mouth contact frequency (event/hour)	All Objects (n=56)	-0.160	0.241
Mouth contact duration (min/hour)	Grass (n=56)	-0.221	0.102
Mouth contact duration (min/hour)	Hands (n=56)	-0.194	0.151
Mouth contact duration (min/hour)	Non-Dietary (n=56)*	-0.308	0.021
Mouth contact duration (min/hour)	Dietary (n=56)	-0.067	0.625
Mouth contact duration (min/hour)	All Objects (n=56)*	-0.266	0.048
Mouth median duration (second)	Grass (n=4)	0.272	0.728



Activity Variables	Contact	r	p-value
Mouth median duration (second)	Hands (n=49)*	-0.357	0.012
Mouth median duration (second)	Non-Dietary (n=52)*	-0.353	0.010
Mouth median duration (second)	Dietary (n=30)	-0.072	0.707
Mouth median duration (second)	All Objects (n=53)	-0.254	0.067

* Significant ($p < 0.05$) correlation with age (Spearman's rank correlation)

As presented in Table S2-10 to Table S2-12 in Supplemental Material Section S2.1, there were significant differences across the EPA recommended age groups (U.S. EPA, 2005) in mouthing hourly contact duration with hands and non-dietary objects (Table S2-11; Figure S2-1 and Figure S2-2). There were no significant differences in mouthing contact frequency and contact median duration. Also, for children aged 1-6 years and 7-12 years, the frequency, hourly duration and median duration for mouthing contact activities while playing on turf are presented in Table S2-13 to Table S2-15. There were significant differences between younger and older children in mouthing median duration with non-dietary objects and all objects (Table S2-15; Figure S2-3).

Mouthing Contact Activity by Gender While Playing on Turf

Frequency, hourly duration and median duration for mouthing contact activities while playing on turf are presented in Table F-45 to Table F-47 by gender (i.e., male, female). There were no significant differences in contact frequency, hourly contact duration, or median contact duration by gender.

Table F-45. Mouthing Contact Frequency (Event/Hour) While Playing on Turf by Gender

Gender	Distribution	Grass	Hands	Dietary	Non-Dietary	All Objects
Male (n=27)	Mean	0.10	12.76	14.30	19.97	34.28
Male (n=27)	SD	0.37	16.79	22.93	22.12	37.66
Male (n=27)	Min	0.00	0.00	0.00	0.00	0.00
Male (n=27)	p25	0.00	3.25	0.00	4.01	6.35
Male (n=27)	Median	0.00	8.14	1.48	13.31	22.77
Male (n=27)	p75	0.00	16.61	30.91	26.47	59.22
Male (n=27)	p95	1.02	41.04	72.43	63.82	106.17
Male (n=27)	p99	1.69	80.11	74.00	88.34	160.77
Male (n=27)	Max	1.69	80.11	74.00	88.34	160.77
Female (n=29)	Mean	0.13	10.63	15.33	23.88	39.20



Gender	Distribution	Grass	Hands	Dietary	Non-Dietary	All Objects
Female (n=29)	SD	0.50	11.38	33.10	35.92	50.87
Female (n=29)	Min	0.00	0.00	0.00	0.00	0.00
Female (n=29)	p25	0.00	4.51	0.00	7.14	7.42
Female (n=29)	Median	0.00	7.14	0.00	10.85	18.28
Female (n=29)	p75	0.00	12.63	19.69	26.24	53.12
Female (n=29)	p95	1.17	36.78	80.19	76.19	189.59
Female (n=29)	p99	2.49	47.70	159.12	185.14	205.09
Female (n=29)	Max	2.49	47.70	159.12	185.14	205.09
p-value*	Not applicable	0.971	0.825	0.412	0.737	0.941

*p-value from Wilcoxon sum rank

Table F-46. Mouthing Contact Duration (Min/Hour) While Playing on Turf by Gender

Gender	Distribution	Grass	Hands	Dietary	Non-Dietary	All Objects
Male (n=27)	Mean	0.00	0.47	1.00	0.70	1.70
Male (n=27)	SD	0.01	0.75	2.02	0.87	2.44
Male (n=27)	Min	0.00	0.00	0.00	0.00	0.00
Male (n=27)	p25	0.00	0.10	0.00	0.14	0.27
Male (n=27)	Median	0.00	0.20	0.08	0.30	0.81
Male (n=27)	p75	0.00	0.51	1.26	0.99	2.04
Male (n=27)	p95	0.02	1.75	4.99	2.14	7.06
Male (n=27)	p99	0.03	3.53	8.94	3.53	10.91
Male (n=27)	Max	0.03	3.53	8.94	3.53	10.91
Female (n=29)	Mean	0.00	0.45	2.55	2.03	4.59
Female (n=29)	SD	0.02	0.78	7.08	6.39	9.82
Female (n=29)	Min	0.00	0.00	0.00	0.00	0.00
Female (n=29)	p25	0.00	0.10	0.00	0.12	0.16
Female (n=29)	Median	0.00	0.22	0.00	0.29	0.50
Female (n=29)	p75	0.00	0.30	1.06	1.67	3.88



Female (n=29)	p95	0.02	2.25	10.53	3.95	36.66
Female (n=29)	p99	0.10	3.54	36.52	34.71	39.65
Female (n=29)	Max	0.10	3.54	36.52	34.71	39.65
p-value*	Not applicable	0.956	0.594	0.598	0.902	0.961

*p-value from Wilcoxon sum rank

Table F-47. Mouthing Median Duration (Second) by Gender

Gender	Distribution	Grass	Hands	Dietary	Non-Dietary	All Objects
Male	n ^a	2	25	17	25	26
Male	Mean	1.00	2.18	3.12	2.24	1.98
Male	SD	0.00	2.52	2.69	2.50	1.56
Male	Min	1.00	1.00	1.00	1.00	1.00
Male	p25	1.00	1.00	1.00	1.00	1.00
Male	Median	1.00	1.00	2.00	1.00	1.00
Male	p75	1.00	2.00	4.00	2.00	2.00
Male	p95	1.00	6.50	9.00	6.50	5.50
Male	p99	1.00	12.00	9.00	12.00	6.50
Male	Max	1.00	12.00	9.00	12.00	6.50
Female	n ^a	2	24	13	27	27
Female	Mean	1.75	1.52	13.00	1.54	3.76
Female	SD	1.06	0.81	33.24	0.91	11.27
Female	Min	1.00	0.50	1.00	0.00	0.00
Female	p25	1.00	1.00	2.00	1.00	1.00
Female	Median	1.75	1.00	3.00	1.00	2.00
Female	p75	2.50	2.00	3.50	2.00	2.00
Female	p95	2.50	3.00	123.00	3.00	4.00
Female	p99	2.50	3.50	123.00	4.00	60.00
Female	Max	2.50	3.50	123.00	4.00	60.00
p-value ^b	Not applicable	0.317	0.718	0.241	0.512	0.962

^aOnly participants who contacted object included in calculation of median contact duration.



^bp-value from Wilcoxon sum rank

F.2.3.2. Children Playing on Playgrounds:

F.2.3.2.1. Children's Description and Footage Length

From the analyzed total footage, we found that out of the total footage (2,548 min), children spent about 21% of the time playing on a playground. The median time per child spent playing on a playground was 21 min (See Table F-48 for details). 24 children (11 males and 13 females) played on playground structures at some point during the video recording and there was only one child in the oldest age group (9 to <16 years old) that played on playgrounds. See Table F-49 for details.

Table F-48. Time Spent on Playground (n=24)

Time Spent	Time in View	Time Not in View
Total playground time (minutes)	531.0	38.2
Median playground time per child (min)	21.0	0.3
Percentage of time spent on playground (%)	20.8	1.5

Table F-49. Number of Children Playing on Turf Grouped by Age Group and Gender

Gender	0<2 Years	2<9 Years	9<16 Years	Total
Male	3	8	0	11
Female	2	10	1	13
Total	5	18	1	24

F.2.3.2.2. Hands Object Activity While Playing on Playground Structures

The right- and left-hand frequencies are presented in Table F-50 to Table F-52. Since there were no significant differences in the contact frequency with object/surfaces between right hand and left hand, both hands were combined and summarized (Table F-52). This means that contact frequency for the left and right hands were summed, however, it does not mean that both hands were in contact with the object at the same time. The median contact frequency with floors for *both hands* combined was 12.12 events/hour.

Table F-50. Right Hand Frequency (Event/Hour) While Playing on the Playground (n=24)

Distribution	Floors	Dietary	Non-Dietary	All Objects
Mean	15.69	0.56	148.11	148.68
SD	32.22	1.66	86.26	86.11



Distribution	Floors	Dietary	Non-Dietary	All Objects
Min	0.00	0.00	15.88	15.88
p25	0.00	0.00	97.53	99.39
Median	4.81	0.00	128.13	128.13
p75	13.49	0.00	193.30	193.30
p95	69.23	3.71	327.27	327.27
p99	146.70	7.02	342.86	342.86
Max	146.70	7.02	342.86	342.86

Table F-51. Left Hand Frequency (Event/Hour) While Playing on Playground (n=24)

Distribution	Floors	Dietary	Non-Dietary	All Objects
Mean	13.95	1.02	144.55	145.57
SD	20.58	2.83	79.16	78.79
Min	0.00	0.00	10.71	10.71
p25	0.00	0.00	95.60	95.60
Median	5.69	0.00	143.56	145.79
p75	15.55	0.00	174.49	175.07
p95	62.43	8.86	291.34	291.34
p99	72.00	10.77	337.96	337.96
Max	72.00	10.77	337.96	337.96

Table F-52. Both Hands Object/Surface Contact Frequency (Event/Hour) on Playground (n=24)

Distribution	Floors	Dietary	Non-Dietary	All Objects
Mean	29.64	1.58	292.67	294.25
SD	48.01	3.87	160.75	160.35
Min	0.00	0.00	30.60	30.60
p25	0.00	0.00	198.42	204.18
Median	12.12	0.00	261.38	262.28
p75	36.31	0.59	377.91	378.50



Distribution	Floors	Dietary	Non-Dietary	All Objects
p95	141.23	10.77	602.49	602.49
p99	196.37	15.88	634.20	634.20
Max	196.37	15.88	634.20	634.20
p-value*	0.592	0.663	0.821	0.820

*p-value from Wilcoxon Signed Rank test comparison of left- and right-hand contact

The hourly contact with the objects for right- and left-hand duration are presented in Table F-53 to Table F-55. Since there were no significant differences in the contact duration with object/surfaces between right hand and left hand, both hands were combined and summarized (Table F-55). This means that contact hourly duration for the left and right hands were summed, however it does not mean that both hands were in contact with the object at the same time. If both hands were in contact with object for the entire time, the maximum value would be 120 min/hour. The median hourly object contact duration with floors was 0.58 min/hour for both hands, however, the 99th percentile was about ten times higher (10 min/hour).

Table F-53. Hourly Object/Surface Contact Duration (Min/Hour) cor Right Hand While Playing on Playground (n=24)

Distribution	Floors	Dietary	Non-Dietary	All Objects
Mean	1.04	0.25	16.50	16.74
SD	1.83	1.03	3.28	3.26
Min	0.00	0.00	3.64	3.64
p25	0.00	0.00	15.52	15.93
Median	0.23	0.00	17.21	17.52
p75	1.17	0.00	18.62	18.64
p95	6.13	0.75	19.23	19.23
p99	6.26	5.04	19.85	19.85
Max	6.26	5.04	19.85	19.85

Table F-54. Hourly Object/Surface Contact Duration (Min/Hour) for Left Hand While Playing on Playground (n=24)

Distribution	Floors	Dietary	Non-Dietary	All Objects
Mean	0.85	0.19	16.05	16.23
SD	1.21	0.80	3.29	3.33



Distribution	Floors	Dietary	Non-Dietary	All Objects
Min	0.00	0.00	4.38	4.38
p25	0.00	0.00	14.86	15.16
Median	0.19	0.00	16.97	17.15
p75	1.25	0.00	18.09	18.24
p95	3.54	0.43	19.82	19.82
p99	3.81	3.93	20.00	20.00
Max	3.81	3.93	20.00	20.00

Table F-55. Hourly Object/Surface Contact Duration (Min/Hour) for Both Hands While Playing on Playground (n=24)

Distribution	Floors	Dietary	Non-Dietary	All Objects
Mean	1.96	0.43	32.44	32.87
SD	2.98	1.28	4.43	4.63
Min	0.00	0.00	22.98	22.98
p25	0.00	0.00	31.61	31.66
Median	0.58	0.00	32.76	33.66
p75	2.32	0.01	35.42	36.65
p95	9.67	3.93	38.59	38.59
p99	10.07	5.04	39.67	39.67
Max	10.07	5.04	39.67	39.67
p-value	0.753	0.737	0.275	0.216

* p value from Wilcoxon Signed Rank test comparison of left- and right-hand contact

The median durations for right and left contact with objects are presented in Table F-56 to Table F-58. Since there were no significant differences in the contact median duration with object/surfaces between right hand and left hand, both hands were combined and summarized (Table F-58). This means that for each object category the durations of all the discrete contacts with an object for both the right and left hands were grouped and reranked to determine the median value of contact duration across all the “hand” contacts with that object. The median duration for contact of hands with floors is 2.00 seconds as presented in Table F-58. However, only 17 out of the 24 children touched the floors with their hands.



Table F-56. Right Hand Median Contact Duration (Second) While Playing on the Playground

Distribution	Floors	Dietary	Non-Dietary	All Objects
n	15	3	24	24
Mean	4.10	18.67	4.15	4.13
SD	2.35	26.31	2.31	2.31
Min	1.00	2.00	1.00	1.00
p25	2.50	3.50	3.00	3.00
Median	4.00	5.00	4.00	4.00
p75	5.25	27.00	5.00	5.00
p95	7.90	44.60	6.85	6.85
p99	9.58	48.12	11.24	11.24
Max	10.00	49.00	12.50	12.50

Table F-57. Left Hand Median Contact Duration (Second) While Playing on the Playground

Distribution	Floors	Dietary	Non-Dietary	All Objects
n	17	4	24	24
Mean	3.26	4.75	10.33	10.29
SD	2.89	6.18	23.87	23.89
Min	1.00	1.00	2.00	2.00
p25	1.50	1.75	3.00	3.00
Median	3.00	2.00	4.00	3.75
p75	4.00	5.00	5.00	5.00
p95	7.40	12.20	50.05	50.20
p99	11.88	13.64	97.66	97.66
Max	13.00	14.00	109.50	109.50

Table F-58. Both Hands Object/Surface Median Duration (Second)

Distribution	Floors	Dietary	Non-Dietary	All Objects
n	17	6	24	24



Distribution	Floors	Dietary	Non-Dietary	All Objects
Mean	3.32	11.75	3.90	3.94
SD	2.32	18.89	2.24	2.25
Min	1.00	1.00	1.00	1.00
p25	2.00	2.00	3.00	3.00
Median	2.00	2.25	4.00	4.00
p75	4.00	11.13	4.00	4.13
p95	6.90	40.25	6.00	6.00
p99	9.78	47.25	11.01	11.01
Max	10.50	49.00	12.50	12.50
p-value	0.136	0.279	0.942	0.883

*p-value from Wilcoxon Signed Rank test comparison of left- and right-hand contact

Hand Contact Activity by Age While Playing on Playground

Both hands contact frequency, hourly duration and median duration are summarized by age groups (<2, 2 to <9, 9 to <16 years) in Table F-59 to Table F-61. There were no significant differences between contact frequency, hourly contact duration, or median contact duration across all three age groups. Also, there were no correlations between age groups and both hand activities (Table F-62).

Table F-59. Both Hands Contact Frequency (Event/Hour) on Playground by Age Groups

Age Group	Distribution	Floors	Dietary	Non-Dietary	All Objects
<2 (n=5)	Mean	39.97	0.23	426.60	426.84
<2 (n=5)	SD	27.85	0.52	184.50	184.39
<2 (n=5)	Min	0.71	0.00	205.86	205.86
<2 (n=5)	Median	20.26	0.00	336.18	336.18
<2 (n=5)	p25	58.06	0.00	354.28	355.45
<2 (n=5)	p75	58.41	0.00	602.49	602.49
<2 (n=5)	p95	62.43	1.17	634.20	634.20
<2 (n=5)	p99	62.43	1.17	634.20	634.20
<2 (n=5)	Max	62.43	1.17	634.20	634.20
2 to <9 (n=18)	Mean	28.41	2.04	257.28	259.32



Age Group	Distribution	Floors	Dietary	Non-Dietary	All Objects
2 to <9 (n=18)	SD	53.40	4.39	142.43	142.33
2 to <9 (n=18)	Min	0.00	0.00	30.60	30.60
2 to <9 (n=18)	Median	0.00	0.00	182.39	182.39
2 to <9 (n=18)	p25	9.87	0.00	256.63	259.11
2 to <9 (n=18)	p75	19.48	2.80	288.28	304.16
2 to <9 (n=18)	p95	196.37	15.88	567.27	567.27
2 to <9 (n=18)	p99	196.37	15.88	567.27	567.27
2 to <9 (n=18)	Max	196.37	15.88	567.27	567.27
9 to <16 (n=1)	Mean	ND	ND	259.97	259.97
9 to <16 (n=1)	SD	ND	ND	ND	ND
9 to <16 (n=1)	Min	ND	ND	ND	ND
9 to <16 (n=1)	Median	ND	ND	ND	ND
9 to <16 (n=1)	p25	ND	ND	ND	ND
9 to <16 (n=1)	p75	ND	ND	ND	ND
9 to <16 (n=1)	p95	ND	ND	ND	ND
9 to <16 (n=1)	p99	ND	ND	ND	ND
9 to <16 (n=1)	Max	ND	ND	ND	ND
p-value*	NA	0.245	0.719	0.147	0.169

*p-value from Kruskal-Wallis Test

NA: not applicable; and ND: not determined

Table F-60. Both Hands Hourly Contact Duration (Min/Hour) on Playground by Age Groups

Age Group	Distribution	Floors	Dietary	Non-Dietary	All Objects
<2 (n=5)	Mean	3.16	0.00	29.74	29.74
<2 (n=5)	SD	4.05	0.01	3.66	3.67
<2 (n=5)	Min	0.01	0.00	24.90	24.90
<2 (n=5)	Median	0.78	0.00	26.87	26.87
<2 (n=5)	p25	1.65	0.00	31.56	31.56
<2 (n=5)	p75	3.28	0.00	31.86	31.88



Age Group	Distribution	Floors	Dietary	Non-Dietary	All Objects
<2 (n=5)	p95	10.07	0.02	33.51	33.51
<2 (n=5)	p99	10.07	0.02	33.51	33.51
<2 (n=5)	Max	10.07	0.02	33.51	33.51
2 to <9 (n=18)	Mean	1.73	0.58	33.23	33.80
2 to <9 (n=18)	SD	2.73	1.46	4.53	4.69
2 to <9 (n=18)	Min	0.00	0.00	22.98	22.98
2 to <9 (n=18)	Median	0.00	0.00	31.76	31.89
2 to <9 (n=18)	p25	0.36	0.00	34.04	34.98
2 to <9 (n=18)	p75	1.76	0.09	36.59	36.76
2 to <9 (n=18)	p95	9.67	5.04	39.67	39.67
2 to <9 (n=18)	p99	9.67	5.04	39.67	39.67
2 to <9 (n=18)	Max	9.67	5.04	39.67	39.67
9 to <16 (n=1)	Mean	ND	ND	31.83	31.83
9 to <16 (n=1)	SD	ND	ND	ND	ND
9 to <16 (n=1)	Min	ND	ND	ND	ND
9 to <16 (n=1)	Median	ND	ND	ND	ND
9 to <16 (n=1)	p25	ND	ND	ND	ND
9 to <16 (n=1)	p75	ND	ND	ND	ND
9 to <16 (n=1)	p95	ND	ND	ND	ND
9 to <16 (n=1)	p99	ND	ND	ND	ND
9 to <16 (n=1)	Max	ND	ND	ND	ND
p-value*	NA	0.233	0.719	0.151	0.088

*p-value from Kruskal-Wallis Test

NA: not applicable; and ND: not determined

Table F-61. Both Hands Contact Median Duration (Second) on Playground by Age Groups

Age Group	Distribution	Floors	Dietary	Non-Dietary	All Objects
<2	n	5	2	5	5



Age Group	Distribution	Floors	Dietary	Non-Dietary	All Objects
<2	Mean	2.60	1.00	3.50	3.50
<2	SD	1.95	ND	1.00	1.00
<2	Min	1.00	ND	2.00	2.00
<2	Median	2.00	ND	3.00	3.00
<2	p25	2.00	ND	4.00	4.00
<2	p75	2.00	ND	4.00	4.00
<2	p95	6.00	ND	4.50	4.50
<2	p99	6.00	ND	4.50	4.50
<2	Max	6.00	ND	4.50	4.50
2 to <9	n	12	5	18	18
2 to <9	Mean	3.63	13.90	4.00	4.06
2 to <9	SD	2.48	20.28	2.55	2.56
2 to <9	Min	1.00	2.00	1.00	1.00
2 to <9	Median	2.00	2.00	3.00	3.00
2 to <9	p25	3.50	2.50	4.00	4.00
2 to <9	p75	4.00	14.00	4.00	5.00
2 to <9	p95	10.50	49.00	12.50	12.50
2 to <9	p99	10.50	49.00	12.50	12.50
2 to <9	Max	10.50	49.00	12.50	12.50
9 to <16	n	0	0	1	1
9 to <16	Mean	ND	ND	4.00	4.00
9 to <16	SD	ND	ND	ND	ND
9 to <16	Min	ND	ND	ND	ND
9 to <16	Median	ND	ND	ND	ND
9 to <16	p25	ND	ND	ND	ND
9 to <16	p75	ND	ND	ND	ND
9 to <16	p95	ND	ND	ND	ND



Age Group	Distribution	Floors	Dietary	Non-Dietary	All Objects
9 to <16	p99	ND	ND	ND	ND
9 to <16	Max	ND	ND	ND	ND
p-value*	NA	0.296	0.138	0.933	0.941

*p-value from Kruskal-Wallis Test

NA: not applicable; and ND: not determined

Table F-62. Spearman Rank Correlation (r) for Both Hand Events and Age (Years)
While Playing on Playgrounds

Activity Variables	Contact	r	p-value
Both hands contact frequency (event/hour)	Floors (n=24)	-0.343	0.101
Both hands contact frequency (event/hour)	Non-Dietary (n=24)	0.235	0.270
Both hands contact frequency (event/hour)	Dietary (n=24)	0.054	0.801
Both hands contact frequency (event/hour)	All Objects (n=24)	-0.234	0.272
Both hands contact duration (min/hour)	Floors (n=24)	-0.163	0.448
Both hands contact duration (min/hour)	Non-Dietary (n=24)	0.360	0.084
Both hands contact duration (min/hour)	Dietary (n=24)	0.061	0.776
Both hands contact duration (min/hour)	All Objects (n=24)	0.391	0.059
Both hands median contact duration (second)	Floors (n=17)	0.274	0.287
Both hands median contact duration (second)	Non-Dietary* (n=24)	-0.072	0.737
Both hands median contact duration (second)	Dietary (n=7)	0.537	0.272
Both hands median contact duration (second)	All Objects* (n=24)	-0.023	0.916

* Significant (p < 0.05) correlation with age (Spearman's rank correlation).

As presented in Supplemental Material Section S2.2, contact frequency, hourly contact duration and median contact duration divided by the EPA recommended age categories (U.S. EPA, 2005) for both hands are presented in Table S2-17 to Table S2-19. There were no statistically significant differences between EPA age groups (Kruskal Wallis Test). Contact frequency, hourly duration and median duration for hand contact activities while playing on playgrounds are presented in Supplemental Material Section S2.2 in Table S2-20 to Table S2-22 for children aged 1-6 years and 7-12 years. There are no significant differences between younger and older children in hand contact activities Wilcoxon Rank Test.



Hand Contact Activity by Gender While Playing on Playground

Frequency, hourly duration and median duration for both hands' contact activities while playing on playgrounds are presented in Table F-63 to Table F-65 by gender (i.e., male, female). There were no significant differences in contact frequency, hourly contact duration, or median contact duration by gender.

Table F-63. Both Hands Contact Frequency While Children Play on Playgrounds by Gender

Gender	Distribution	Floors	Dietary	Non-Dietary	All Objects
Male (n=11)	Mean	42.16	0.36	272.99	273.35
Male (n=11)	SD	66.00	0.88	180.55	180.58
Male (n=11)	Min	0.00	0.00	30.60	30.60
Male (n=11)	p25	0.00	0.00	80.59	80.59
Male (n=11)	Median	16.60	0.00	275.85	275.85
Male (n=11)	p75	58.06	0.00	401.54	401.54
Male (n=11)	p95	196.37	2.80	602.49	602.49
Male (n=11)	p99	196.37	2.80	602.49	602.49
Male (n=11)	Max	196.37	2.80	602.49	602.49
Female (n=13)	Mean	19.04	2.61	309.32	311.93
Female (n=13)	SD	22.85	5.05	147.34	146.19
Female (n=13)	Min	0.00	0.00	171.54	171.54
Female (n=13)	p25	0.00	0.00	205.86	208.83
Female (n=13)	Median	11.01	0.00	259.97	261.77
Female (n=13)	p75	19.48	3.62	288.28	304.16
Female (n=13)	p95	62.43	15.88	634.20	634.20
Female (n=13)	p99	62.43	15.88	634.20	634.20
Female (n=13)	Max	62.43	15.88	634.20	634.20
p-value*	Not applicable	0.224	0.599	0.257	0.251

*p-value from Wilcoxon Sum Rank test comparison of male and female



Table F-64. Both Hands Hourly Contact Duration (Min/Hour) While Children Play on Playgrounds by Gender

Gender	Distribution	Floors	Dietary	Non-Dietary	All Objects
Male (n=11)	Mean	2.51	0.01	32.71	32.72
Male (n=11)	SD	3.31	0.03	4.43	4.43
Male (n=11)	Min	0.00	0.00	22.98	22.98
Male (n=11)	p25	0.00	0.00	31.56	31.56
Male (n=11)	Median	0.78	0.00	32.20	32.20
Male (n=11)	p75	5.60	0.00	36.34	36.34
Male (n=11)	p95	9.67	0.09	39.67	39.67
Male (n=11)	p99	9.67	0.09	39.67	39.67
Male (n=11)	Max	9.67	0.09	39.67	39.67
Female (n=13)	Mean	1.49	0.79	32.22	33.01
Female (n=13)	SD	2.72	1.69	4.59	4.97
Female (n=13)	Min	0.00	0.00	23.64	23.64
Female (n=13)	Median	0.00	0.00	31.66	31.76
Female (n=13)	p25	0.38	0.00	33.31	34.50
Female (n=13)	p75	1.65	0.12	34.50	36.70
Female (n=13)	p95	10.07	5.04	38.59	38.59
Female (n=13)	p99	10.07	5.04	38.59	38.59
Female (n=13)	Max	10.07	5.04	38.59	38.59
p-value*	Not applicable	0.630	0.224	0.599	0.257

*p-value from Wilcoxon Sum Rank test comparison of male and female

Table F-65. Both Hands Median Contact Duration (Second) While Children Play on Playgrounds by Gender

Gender	Distribution	Floors	Dietary	Non-Dietary	All Objects
Male	n	8	2	11	11
Male	Mean	3.44	1.50	4.59	4.59
Male	SD	3.18	0.71	3.01	3.01



Gender	Distribution	Floors	Dietary	Non-Dietary	All Objects
Male	Min	1.00	1.00	1.00	1.00
Male	Median	1.50	1.00	3.00	3.00
Male	p25	2.00	1.50	4.00	4.00
Male	p75	4.50	2.00	6.00	6.00
Male	p95	10.50	2.00	12.50	12.50
Male	p99	10.50	2.00	12.50	12.50
Male	Max	10.50	2.00	12.50	12.50
Female	n	9	4	13	13
Female	Mean	3.22	16.88	3.31	3.38
Female	SD	1.39	22.12	1.13	1.21
Female	Min	2.00	2.00	1.00	1.00
Female	Median	2.00	2.25	3.00	3.00
Female	p25	3.00	8.25	3.50	3.50
Female	p75	4.00	31.50	4.00	4.00
Female	p95	6.00	49.00	5.00	5.00
Female	p99	6.00	49.00	5.00	5.00
Female	Max	6.00	49.00	5.00	5.00
p-value*	Not applicable	0.753	0.638	0.675	0.573

*p-value from Wilcoxon Sum Rank test comparison of male and female

F.2.3.2.3. Mouthing Activity While Playing on Playground

The mouthing contact frequency for children playing on playground is summarized in Table F-66. Only one child (1/24) contacted “floors” with their mouth, resulting in a contact frequency of 2.30 events/hour. The median mouthing frequencies were 9.80 events/hour and 10.20 events/hour for hands and non-dietary objects, respectively. These contacts could indicate additional exposure to playmat chemicals that have transferred to their child’s hands (Section F.2.3.2.2) or the objects they are placing in their mouths.



Table F-66. Mouth Contact Frequency (Event/Hour) While Playing on the Playground (n=24)

Distribution	Floors	Hands	Dietary	Non-Dietary	All Objects
Mean	2.30	19.60	43.10	28.20	71.30
SD	ND	20.30	112.30	46.80	114.30
Min	ND	1.40	0.00	0.00	0.00
p25	ND	4.30	0.00	2.90	4.00
Median	ND	9.80	0.00	10.20	20.40
p75	ND	25.40	3.20	30.30	66.00
p95	ND	67.50	313.40	82.50	335.00
p99	ND	67.50	378.95	218.18	378.95
Max	ND	67.50	378.95	218.18	378.95

ND: not determined

The mouthing hourly contact duration for children playing on playgrounds is summarized in Table F-67. The hourly contact duration for the one child who contacted “floors” with their mouth was 0.04 min/hour. The median hourly mouthing duration was 0.10 and 0.00 min/hour for hands and non-dietary objects, respectively. These contacts could indicate additional exposure to playmat chemicals that have transferred to their child’s hands (Section F.2.3.2.2) or the objects they are placing in their mouths.

Table F-67. Hourly Mouthing Contact Duration (Min/Hour) While Playing on Playground (n=24)

Distribution	Floors	Hands	Dietary	Non-Dietary	All Objects
Mean	0.04	1.00	1.50	1.70	3.20
SD	ND	1.60	4.10	4.10	5.50
Min	ND	0.00	0.00	0.00	0.00
Median	ND	0.10	0.00	0.00	0.00
p25	ND	0.30	0.00	0.30	0.80
p75	ND	0.70	0.20	1.30	3.80
p95	ND	5.00	11.40	5.00	16.60
p99	ND	5.00	16.50	20.00	20.00
Max	ND	5.00	16.50	20.00	20.00

ND: not determined



The mouthing median contact duration for children playing on playgrounds is summarized in Table F-68. Median duration is only contacted from those children that contacted the object. The median mouthing contact duration for the one child that contacted “floors” with their mouth is 1.00 second. Median mouthing contact duration was 1.00 second and 1.50 seconds for hands and non-dietary objects, respectively.

Table F-68. Mouth Median Contact Duration (Second) While Playing on Playground

Distribution	Floors	Hands	Dietary	Non-Dietary	All Objects
n	1	14	6	13	20
Mean	1.00	2.07	25.58	2.21	7.83
SD	ND	1.87	59.00	1.91	26.11
Min	ND	1.00	0.50	0.00	0.00
p25	ND	1.00	1.00	1.00	1.00
Median	ND	1.00	1.50	1.00	1.00
p75	ND	2.00	2.75	3.00	2.25
p95	ND	6.18	110.25	6.10	12.10
p99	ND	6.44	138.85	6.42	97.22
Max	ND	6.50	146.00	6.50	118.50

ND: not determined

Mouthing Activity While Playing on Playground by Age

Frequency, hourly duration and median duration for mouthing contact activities while playing on playgrounds are presented in Table F-69 to Table F-71 for three age categories (<2, 2 to <9, 9 to <16 years). There were no significant differences in mouthing contact frequency across these age groups. Out of the 24 children, only one child (less than 2 years of age) made a mouthing contact with any type of floor while playing on playground structures. Age (by year, 1-12 years) was negatively correlated ($p < 0.05$) with the contact frequency, hourly contact duration and median contact duration of mouthing activities with total objects (Table F-72). Age was also negatively correlated ($p < 0.05$) with hourly mouthing duration and frequency for hands. Finally, the non-dietary objects were negatively correlated with mouthing duration and median duration.

Table F-69. Mouthing Contact Frequency (Event/Hour) While on Playground (n=24) by Age Groups

Age Group	Distribution	Floors	Hands	Dietary	Non-Dietary	All Objects
<2 (n=5)	Mean	2.30	26.60	6.39	27.23	33.63



Age Group	Distribution	Floors	Hands	Dietary	Non-Dietary	All Objects
<2 (n=5)	SD	ND	18.56	9.31	14.84	18.91
<2 (n=5)	Min	ND	9.22	0.00	4.96	4.96
<2 (n=5)	p25	ND	9.22	0.00	24.43	24.43
<2 (n=5)	Median	ND	24.43	0.00	29.95	41.47
<2 (n=5)	p75	ND	46.15	11.52	30.68	46.15
<2 (n=5)	p95	ND	46.15	20.45	46.15	51.13
<2 (n=5)	p99	ND	46.15	20.45	46.15	51.13
<2 (n=5)	Max	ND	46.15	20.45	46.15	51.13
2 to <9 (n=18)	Mean	ND	17.74	55.69	29.98	85.67
2 to <9 (n=18)	SD	ND	21.18	127.94	53.51	129.04
2 to <9 (n=18)	Min	ND	1.39	0.00	0.00	0.00
2 to <9 (n=18)	p25	ND	3.85	0.00	2.78	2.94
2 to <9 (n=18)	Median	ND	9.12	0.00	9.54	13.42
2 to <9 (n=18)	p75	ND	25.42	0.00	25.42	82.50
2 to <9 (n=18)	p95	ND	67.50	378.95	218.18	378.95
2 to <9 (n=18)	p99	ND	67.50	378.95	218.18	378.95
2 to <9 (n=18)	Max	ND	67.50	378.95	218.18	378.95
9 to <16 (n=0)	Mean	ND	ND	ND	ND	ND
9 to <16 (n=0)	SD	ND	ND	ND	ND	ND
9 to <16 (n=0)	Min	ND	ND	ND	ND	ND
9 to <16 (n=0)	p25	ND	ND	ND	ND	ND
9 to <16 (n=0)	Median	ND	ND	ND	ND	ND
9 to <16 (n=0)	p75	ND	ND	ND	ND	ND
9 to <16 (n=0)	p95	ND	ND	ND	ND	ND
9 to <16 (n=0)	p99	ND	ND	ND	ND	ND
9 to <16 (n=0)	Max	ND	ND	ND	ND	ND
p-value*	NA	0.150	0.312	0.711	0.161	0.341

*p-value from Kruskal-Wallis Test



NA: not applicable; and ND: not determined

Table F-70. Mouthing Contact Duration (Min/Hour) on Playground (n=24) by Age Groups

Age Group	Distribution	Floors	Hands	Dietary	Non-Dietary	All Objects
<2 (n=5)	Mean	0.04	3.09	0.26	2.66	2.92
<2 (n=5)	SD	ND	2.60	0.42	2.05	1.88
<2 (n=5)	Min	ND	0.12	0.00	0.33	0.33
<2 (n=5)	p25	ND	0.12	0.00	0.78	1.74
<2 (n=5)	Median	ND	4.14	0.00	3.07	3.41
<2 (n=5)	p75	ND	5.00	0.34	4.14	4.14
<2 (n=5)	p95	ND	5.00	0.96	5.00	5.00
<2 (n=5)	p99	ND	5.00	0.96	5.00	5.00
<2 (n=5)	Max	ND	5.00	0.96	5.00	5.00
2 to <9 (n=18)	Mean	ND	0.41	1.98	1.53	3.52
2 to <9 (n=18)	SD	ND	0.49	4.62	4.64	6.23
2 to <9 (n=18)	Min	ND	0.02	0.00	0.00	0.00
2 to <9 (n=18)	p25	ND	0.06	0.00	0.00	0.04
2 to <9 (n=18)	Median	ND	0.27	0.00	0.24	0.55
2 to <9 (n=18)	p75	ND	0.51	0.00	1.00	3.16
2 to <9 (n=18)	p95	ND	1.75	16.52	20.00	20.00
2 to <9 (n=18)	p99	ND	1.75	16.52	20.00	20.00
2 to <9 (n=18)	Max	ND	1.75	16.52	20.00	20.00
9 to <16 (n=0)	Mean	ND	ND	ND	ND	ND
9 to <16 (n=0)	SD	ND	ND	ND	ND	ND
9 to <16 (n=0)	Min	ND	ND	ND	ND	ND
9 to <16 (n=0)	p25	ND	ND	ND	ND	ND
9 to <16 (n=0)	Median	ND	ND	ND	ND	ND
9 to <16 (n=0)	p75	ND	ND	ND	ND	ND
9 to <16 (n=0)	p95	ND	ND	ND	ND	ND



Age Group	Distribution	Floors	Hands	Dietary	Non-Dietary	All Objects
9 to <16 (n=0)	p99	ND	ND	ND	ND	ND
9 to <16 (n=0)	Max	ND	ND	ND	ND	ND
p-value*	NA	0.149	0.187	0.782	0.043	0.176

*p-value from Kruskal-Wallis Test

NA: not applicable; and ND: not determined

Table F-71. Mouthing Contact Median Duration (Second) on Playground (n=24) by Age Groups

Age Group	Distribution	Floors	Hands	Dietary	Non-Dietary	All Objects
<2	n	1	3	2		5
<2	Mean	1.00	4.50	2.00	4.13	3.50
<2	SD	ND	3.04	1.41	2.59	2.65
<2	Min	ND	1.00	1.00	1.00	1.00
<2	p25	ND	1.00	1.00	2.00	1.00
<2	Median	ND	6.00	2.00	4.50	3.00
<2	p75	ND	6.50	3.00	6.25	6.00
<2	p95	ND	6.50	3.00	6.50	6.50
<2	p99	ND	6.50	3.00	6.50	6.50
<2	Max	ND	6.50	3.00	6.50	6.50
2 to <9	n	0	11	4	13	15
2 to <9	Mean	ND	1.41	37.38	1.62	9.27
2 to <9	SD	ND	0.66	72.42	1.26	30.24
2 to <9	Min	ND	1.00	0.50	0.00	0.00
2 to <9	p25	ND	1.00	0.75	1.00	1.00
2 to <9	Median	ND	1.00	1.50	1.00	1.00
2 to <9	p75	ND	2.00	74.00	2.00	2.00
2 to <9	p95	ND	3.00	146.00	5.00	118.50
2 to <9	p99	ND	3.00	146.00	5.00	118.50
2 to <9	Max	ND	3.00	146.00	5.00	118.50



Age Group	Distribution	Floors	Hands	Dietary	Non-Dietary	All Objects
9 to <16	n	0	0	0	0	0
9 to <16	Mean	ND	ND	ND	ND	ND
9 to <16	SD	ND	ND	ND	ND	ND
9 to <16	Min	ND	ND	ND	ND	ND
9 to <16	p25	ND	ND	ND	ND	ND
9 to <16	Median	ND	ND	ND	ND	ND
9 to <16	p75	ND	ND	ND	ND	ND
9 to <16	p95	ND	ND	ND	ND	ND
9 to <16	p99	ND	ND	ND	ND	ND
9 to <16	Max	ND	ND	ND	ND	ND
p-value*	NA	ND	0.121	0.814	0.055	0.183

*p-value from Kruskal-Wallis Test

NA: not applicable; and ND: not determined

Table F-72. Spearman Rank Correlation for Age (Years) and Mouthing Events While Playing on Playgrounds

Activity Variables	Contact	r	p-value
Mouth contact frequency (event/hour)	Floors (n=24)	-0.289	0.170
Mouth contact frequency (event/hour)	Hands* (n=24)	-0.631	0.016*
Mouth contact frequency (event/hour)	Non-Dietary (n=24)	0.378	0.705
Mouth contact frequency (event/hour)	Dietary	-0.163	0.448
Mouth contact frequency (event/hour)	All objects* (n=24)	-0.459	0.024*
Mouth contact duration (min/hour)	Floors (n=24)	-0.289	0.170
Mouth contact duration (min/hour)	Hands* (n=24)	-0.593	0.025*
Mouth contact duration (min/hour)	Non-Dietary* (n=24)	-0.624	0.011*
Mouth contact duration (min/hour)	Dietary	-0.095	0.657
Mouth contact duration (min/hour)	All objects* (n=24)	-0.025	0.025*
Mouth median contact duration (second)	Floors (n=1)	-0.289	0.169
Mouth median contact duration (second)	Hands (n=14)	-0.322	0.261



Activity Variables	Contact	r	p-value
Mouth median contact duration (second)	Non-Dietary* (n=13)	-0.555	0.005*
Mouth median contact duration (second)	Dietary (n=6)	-0.122	0.570
Mouth median contact duration (second)	All objects* (n=20)	-0.418	0.042*

*Significant ($p < 0.05$) correlation with age (Spearman's rank correlation).

In Supplemental Material Section S2.2, the mouthing events are summarized by the EPA recommended age categories (U.S. EPA, 2005) in Table S2-23 to Table S2-25. There were significant differences between the EPA age categories with respect to mouthing frequency for non-dietary objects and all objects. Similarly, a significant difference was observed for the mouthing duration for non-dietary objects.

Also, in Supplemental Material Section S2.2, the frequency, hourly duration and median duration for mouthing contact activities while playing on playgrounds are presented in Table S2-26 to Table S2-28 for children aged 1-6 and 7-12 years. There were significant differences between younger (1-6 years old) and older children (7-12 years old) groups for the mouthing frequency and duration with all objects (Table S2-26 and Table S2-27) and for the mouthing duration with non-dietary objects (Table S2-27).

Mouthing Activity While Playing on Playground by Gender

Frequency, hourly duration and median duration for mouthing contact activities while playing on playgrounds are presented in Table F-73 to Table F-75 by gender (i.e., male, female). There were no significant differences in contact frequency or hourly contact duration by gender. However, female children had significantly longer contact durations with "All objects".

Table F-73. Mouthing Frequency (Event/Hour) While Children Play on Playgrounds by Gender

Gender	Distribution	Floors	Hands	Dietary	Non-Dietary	All objects
Male (n=11)	Mean	2.30	27.87	64.96	27.91	92.87
Male (n=11)	SD	ND	24.68	137.82	30.70	128.10
Male (n=11)	Min	ND	2.94	0.00	0.00	0.00
Male (n=11)	p25	ND	9.22	0.00	0.00	2.94
Male (n=11)	Median	ND	20.36	0.00	24.43	51.13
Male (n=11)	p75	ND	46.84	20.45	65.16	82.50
Male (n=11)	p95	ND	67.50	378.95	82.50	378.95
Male (n=11)	p99	ND	67.50	378.95	82.50	378.95



Gender	Distribution	Floors	Hands	Dietary	Non-Dietary	All objects
Male (n=11)	Max	ND	67.50	378.95	82.50	378.95
Female (n=13)	Mean	ND	13.46	24.60	28.37	52.97
Female (n=13)	SD	ND	15.13	86.80	58.44	102.80
Female (n=13)	Min	ND	1.39	0.00	0.00	0.00
Female (n=13)	p25	ND	4.08	0.00	3.85	4.96
Female (n=13)	Median	ND	8.09	0.00	10.03	10.27
Female (n=13)	p75	ND	17.90	0.00	21.56	25.42
Female (n=13)	p95	ND	46.15	313.43	218.18	334.99
Female (n=13)	p99	ND	46.15	313.43	218.18	334.99
Female (n=13)	Max	ND	46.15	313.43	218.18	334.99
p-value*	NA	ND	0.197	0.223	0.705	0.212

* p-value from Wilcoxon Sum Rank test comparison of male and female

NA: not applicable; and ND: not determined

Table F-74. Mouthing Duration (Min/Hour) While Children Play on Playgrounds by Gender

Gender	Distribution	Floors	Hands	Dietary	Non-Dietary	All Objects
Male (n=11)	Mean	0.04	1.17	0.83	1.10	1.93
Male (n=11)	SD	ND	1.59	1.58	1.40	1.71
Male (n=11)	Min	ND	0.05	0.00	0.00	0.00
Male (n=11)	p25	ND	0.12	0.00	0.00	0.05
Male (n=11)	Median	ND	0.47	0.00	0.78	1.74
Male (n=11)	p75	ND	1.75	0.96	1.87	3.41
Male (n=11)	p95	ND	4.14	4.62	4.14	4.77
Male (n=11)	p99	ND	4.14	4.62	4.14	4.77
Male (n=11)	Max	ND	4.14	4.62	4.14	4.77
Female (n=13)	Mean	ND	0.84	2.15	2.21	4.36
Female (n=13)	SD	ND	1.69	5.35	5.51	7.20
Female (n=13)	Min	ND	0.02	0.00	0.00	0.00



Gender	Distribution	Floors	Hands	Dietary	Non-Dietary	All Objects
Female (n=13)	p25	ND	0.17	0.00	0.04	0.04
Female (n=13)	Median	ND	0.28	0.00	0.33	0.33
Female (n=13)	p75	ND	0.42	0.00	0.58	5.00
Female (n=13)	p95	ND	5.00	16.52	20.00	20.00
Female (n=13)	p99	ND	5.00	16.52	20.00	20.00
Female (n=13)	Max	ND	5.00	16.52	20.00	20.00
p-value*	NA	ND	0.697	0.402	0.770	0.662

*p-value from Wilcoxon Sum Rank test comparison of male and female

NA: not applicable; and ND: not determined

Table F-75. Mouthing Median Duration (Second) While Children Play on Playgrounds by Gender

Gender	Distribution	Floors	Hands	Dietary	Non-Dietary	All Objects
Male	n	1	6	4	6	9
Male	Mean	1.00	1.83	1.38	1.83	1.50
Male	SD	ND	2.04	1.11	2.04	1.70
Male	Min	ND	1.00	0.50	1.00	0.50
Male	p25	ND	1.00	0.75	1.00	1.00
Male	Median	ND	1.00	1.00	1.00	1.00
Male	p75	ND	1.00	2.00	1.00	1.00
Male	p95	ND	6.00	3.00	6.00	6.00
Male	p99	ND	6.00	3.00	6.00	6.00
Male	Max	ND	6.00	3.00	6.00	6.00
Female	n	0	8	2	11	11
Female	Mean	ND	2.25	74.00	2.41	13.00
Female	SD	ND	1.85	101.82	1.91	35.04
Female	Min	ND	1.00	2.00	0.00	0.00
Female	p25	ND	1.00	2.00	1.00	1.00
Female	Median	ND	1.75	74.00	2.00	2.00



Gender	Distribution	Floors	Hands	Dietary	Non-Dietary	All Objects
Female	p75	ND	2.50	146.00	3.00	5.00
Female	p95	ND	6.50	146.00	6.50	118.50
Female	p99	ND	6.50	146.00	6.50	118.50
Female	Max	ND	6.50	146.00	6.50	118.50
p-value*	NA	ND	0.174	0.156	0.264	0.041*

*p-value from Wilcoxon Sum Rank test comparison of male and female
NA: not applicable; and ND: not determined



F.2.4. References

- AuYeung, W., Canales, R.A., Beamer, P., Ferguson, A.C., and J.O. Leckie. (2004). "Young Children's Mouthing Behavior: An Observational Study via Videotaping in a Primarily Outdoor Residential Setting." *J Children's Health*, 2 (3-4), pg. 271-295
- AuYeung, W., Canales, R.A., Beamer, P., Ferguson, A.C., and J.O. Leckie. (2006). "Young children's hand contact activities: An observational study via videotaping in primarily outdoor residential settings." *J Exp Sci Environ Epidemiol*, 16: 434-446.
- Beamer, P., Key, M.E., Ferguson, A.C., Canales, R.A., AuYeung, W., and J.O. Leckie. (2008) "Quantified Activity Pattern Data from 6-to-27-Month-Old Farmworker Children for Use in Exposure Assessment." *Environ Res*, 108: 239-246. PMID: 18723168.
- Beamer, P., Luik, C.E., Canales, R.A., Leckie, J.O. "Quantified Outdoor Micro-Activity Data for Children Aged 7-12 Years Old." *J Exp Sci Environ Epidemiol*, 22:82-92. PMID: 21989500.
- Ferguson, A.C., Canales, R.A., Beamer, P., AuYeung, W., Key, M., Munninghoff, A., Lee, K.T., Robertson, A., and J.O. Leckie. (2006) "Video methods in the quantification of children's exposures." *J Exp Anal Environ Epidemiol*, 16: 287-298.
- Hubal, E., Sheldon, L., Burke, J., McCurdy, T., Berry, M., Rigas, M., Zartarian, V. and Freeman, N. (2000). Children's Exposure Assessment: A Review of Factors Influencing Children's Exposure, and the Data Available to Characterize and Assess That Exposure. *Environmental Health Perspectives*, 108(6), p.475.
- Juberg D, Alfano K, Coughlin R, Thompson K. (2001). An Observational Study of Object Mouthing Behavior by Young Children. *Pediatrics*. 2001;107:135–142.
- RStudio Team (2015). RStudio: Integrated Development for R. RStudio, Inc., Boston, MA URL <http://www.rstudio.com/>.
- Zartarian V, Ferguson A, Ong C, Leckie J. (1997) Quantifying Videotaped Activity Patterns: Video Translation Software and Training Methodologies. *J Expo Anal Environ Epidemiol*: 7:535–542.



F.3. Quantification of Micro-level Activities from Recorded Data of Children Playing Soccer on Artificial Turf - by The University of Arizona



**Part II: Quantification of Micro-level Activities from Recorded Data of Children
Playing Soccer on Artificial Turf**

Prepared for the California Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Principal Investigator: Paloma Beamer

Prepared by: Nicolas Lopez-Galvez

University of Arizona, College of Public Health

August 20, 2018



Table of Contents

F.1.2.3.1. Survey Development.....	F-13
F.1.2.3.2. Recruitment.....	F-13
F.1.2.3.3. Data Collection	F-14
F.1.2.3.4. Data Management and Analysis.....	F-14
F.1.2.4.1. Questionnaire Development.....	F-14
F.1.2.4.2. Recruitment.....	F-14
F.1.2.4.3. Data Collection	F-15
F.1.2.4.4. Data Management and Analysis.....	F-15
F.1.3.4.1. Demographic Information.....	F-19
F.1.3.4.2. Player History	F-22
F.1.3.4.3. Longest Amount of Time Spent on Synthetic Turf Field in Past Year ..F-25	
F.1.3.4.4. Reported Exertion During Practices and Games.....	F-26
F.1.3.4.5. Contact with Crumb Rubber During Practices and Games	F-27
F.1.3.4.6. Food and Water Consumption Before and During Practices and Games F-36	
F.1.3.4.7. Observation of Crumb Rubber in Home	F-37
F.2.2.2.1. Children Playing on Turf.....	F-49
F.2.2.2.2. Children Playing on Playgrounds	F-49
F.2.3.1.1. Amount of Footage with Turf and Demographics of Children Who Played on Turf.....	F-51
F.2.3.1.2. Hand Contact Activity While Playing on Turf.....	F-52
F.2.3.1.3. Mouthing Activity While Playing on Turf.....	F-64
F.2.3.2.1. Children’s Description and Footage Length	F-74
F.2.3.2.2. Hands Object Activity While Playing on Playground Structures	F-74
F.2.3.2.3. Mouthing Activity While Playing on Playground	F-86
F.3.2.2.1. Training	F-105
F.3.2.2.2. Quality Control and Quality Assurance.....	F-105
F.3.2.3.1. Analysis for Both Hands and Mouthing Events.....	F-106
F.3.2.3.2. Analysis for Macro-Activities, Actions and Intensity	F-107



F.3.2.3.3. Statistical Analysis	F-108
F.3.3.6.1. Event Schedule (Moment/Time)	F-119
F.3.3.6.2. Location (On or Off Field) (Supplemental Material Sections S3.2.2.1 and S3.2.2.2) F-120	
F.3.3.6.3. Use Of Gloves (On or Off) (Supplemental Material Sections S3.2.3.1 and S3.2.3.2) F-121	
F.3.3.6.4. Player Age (Supplemental Material Section S3.3)	F-121
F.3.3.6.5. Player Gender	F-126
F.3.3.6.6. Player Position (Goalie or Others)	F-131
F.3.3.6.7. Event Type (Game or Practice).....	F-136



F.3.1. Introduction

There is a lack of information on the exposure to crumb rubber infill for artificial soccer fields. The dermal and ingestion pathways of exposure have not been fully assessed and may pose a particular risk for children. Risk assessment models are used to evaluate the non-dietary ingestion and dermal pathway of exposure, which require the use of precise activity data on hand and mouthing contacts. However, there is limited information on the behaviors of soccer players while playing on artificial turf. Direct observation (including videotaping) is considered the most accurate way to record a child's dermal contact and mouthing behaviors (Cohen Hubal et al., 2000a). While eliminating recall bias, videotaping preserves real-time activities that can later be carefully analyzed for detailed contact frequency and duration, and for inter- and intra-observer reliability, as well as any other needs should new concerns arise (Zartarian et al., 1997a). This approach may be the only viable approach for collecting activity patterns to estimate non-dietary ingestion since it allows quantification of frequent mouthing events of short duration (Cohen Hubal et al., 2000b; Juberg et al., 2001). Therefore, this study focused on the collection and analysis of micro-level activity time series (MLATS) from video data of participants playing soccer in the San Francisco Bay Area and the Sacramento area. The objective of this report is to provide quantified activity data to support estimating potential health risks of Californians that play soccer on synthetic turf fields.

F.3.2. Methods

F.3.2.1. Data Collection

In order to collect the MLATS, a total of 40 participants were video recorded while playing soccer by a team of researchers from the University of California, Berkeley (UCB) and The Office of Environmental Health Hazard Assessment (OEHHA). The videotaping occurred during scheduled games or practices in Northern California. A total of 10 events (an average of 4 players per event) were recorded. Research staff completed an "Object Palette Log Sheet", where they recorded descriptions of the objects contacted by participants. This was later used to confirm objects in the video recording. The "Taper's Log Time" sheet was completed to record all of the start and stop times, and to describe any particular change during the recording.

The collected video data were saved to an encrypted computer at UCB and then the data were received by the University of Arizona (UA) via a secure platform. Once at UA, the data was saved to an encrypted protected external hard drive, which was stored in a locked drawer. The UA Institutional Review Board (IRB) approved to cede the project to the UC Berkeley Committee for the Protection of Human Subjects (CPHS) and the California State IRB. The UC Berkeley CPHS approval was received on November 13, 2017 and the California State IRB approved on December 4th, 2017.



F.3.2.2. Video Translation and Data Processing

The video data was translated for body part (hands and mouth) to object/surface contact data behavior, macro-activities, specific actions and intensity of players during a soccer event on artificial turf. Before transcription, each video was converted from an MTS file type to an MP4 file type using Windows Movie Maker V2013 to be analyzed. The Video Translator -TE software developed by Dr. Robert Canales in the College of Public Health at the University of Arizona was adapted from the Virtual Timing Device previously developed by Stanford's Exposure Research Group (Ferguson et al., 2006). This computer software was used to translate the video footage into computer text files. Video Translator -TE software is a tool designed to quantify real-time, sequential micro-activity pattern data collected via videos. The software couples grid activation with a computer clock that records (to fractions of a second), the total duration of each contact/activity event in the sequence that the activities occurred. A computer text file of the time series is produced from the translation process (Figure F-19).




Figure F-19. Example of Computer Text File Produced from Video Translator -TE Interface and Processing of Repetitive and Continuous Contacts (Adapted from AuYeung et al., 2006).

The Video Translator -TE interface is an on-screen window containing multiple grids. This interface is referred to as a "palette" and can be customized for a specific study. For this study, two palettes were designed. The first consists of four grids, each comprising cells labeled with names of: (a) locations, (b) moments/time of event, (c) glove use, and (d) object/surfaces relevant to people playing soccer (Figure F-20). While monitoring a contact boundary (e.g., right hand, left hand, mouth) on the video, the video translator collects data by activating (i.e., positioning the cursor over a cell and



clicking the pointing device) the appropriate cell in each grid.

The screenshot shows a software interface titled "Video Translator" with a menu bar containing "Options", "Settings", and "About". The main area is divided into three primary sections:

- Location:** A vertical list of four buttons: "Sideline", "Soccer_field", "Other_outdoor", and "Vehicle".
- Moment / Time of event:** A vertical list of six buttons: "Before_game/practice", "Player_warming_up", "During_game/practice", "Snack_time/break", and "After_game/practice".
- Object/Surfaces:** A 5x6 grid of buttons. The first two rows are grouped under a bracket labeled "Goggles".

Location	Moment / Time of event	Object/Surfaces
Sideline	Before_game/practice	Artificial_turf
		Wood_structure
Soccer_field	Player_warming_up	People
		Clothes_towel
Other_outdoor	During_game/practice	Other_food
		Head_face
Vehicle	Snack_time/break	Black_crumb
		Plast_structure
	After_game/practice	Wood_tool
		Footwear
		Sticky_food
		Body_skin
		Grass
		Metal_structure
		Plast_tool_appl
		Shin_guards
		Tap_water
		Other_hand_s
		Dirt
		Fabric_structure
		Metal_tool_appl
		Gloves
		Puddle_water
		Mouth
		Plastic_rubber_mat
		Porous_plast_struct
		Paper_wrapper
		Mth_guard_pacifier
		Other_beverages
		Hair
		Asphalt
		Fabric_toy
		Vegetation
		Bag_backpack
		Water_bottle
		Not_in_view
		Conc_rock_floor
		Wood_toy
		Animal
		Soccer_ball
		Food_cont
		Nothing

Figure F-20. Palette Used to Collect Data on Contact Frequency and Duration of Body Parts with Objects, Location of Participant, Glove Use, and The Moment/Time of The Event

As presented in Figure F-21, to collect data on macro-activities, specific actions and player intensity, the Video Translator -TE palette was modified to have four grids (palette) with: (a) locations, (b) moments/time of event, (c) macro-activities/actions and (d) player intensity during the soccer event.



		Video Translator Options	Settings	About	
Location		Sideline	Stretching	Running	Highly
		Soccer_field	Push_ups	Walking	Moderately
		Other_outdoor	Sit_ups	Resting_standing	Lightly
		Vehicle	Jumping	Rest_sit_chair	Resting
Moment / Time of event		Before_game/practice	Heading	Rest_sit_on_ground	
		Player_warming_up	Diving	Fall_ground	
		During_game/practice	Tackling_sliding	Not_in_view	
		Snack_time/break			
		After_game/practice			

Macro-activities/Actions

Figure F-21. Palette used to collect data on macro-activities, specific actions, and intensity during the soccer event.

F.3.2.2.1. Training

All of the videos were translated by six members of the research team who passed a rigorous training. The training consisted of analyzing four videos of increasing difficulty. The training videos were transcribed using the two palettes (Figure F-20 and Figure F-21). The transcribed data was then compared to a “gold standard” (previously transcribed by an experienced transcriber). To acquire a high inter-observer reliability, the trainee’s transcription had to be less than 10% different compared to the gold standard before the trainee could move on to the next training video and eventually to collect data once all 4 training videos were completed adequately. For additional information on the training process, see Supplemental Material Section S3.1.1. Code was written in Rstudio V1.1 (RStudio Team, 2017) to compare the files and calculate the percent difference between them (Supplemental Material Section S3.1).

F.3.2.2.2. Quality Control and Quality Assurance

In addition to the translation training requirements, in order to maintain consistency and quality of translations, 10% of the translated video files were “spot-checked.” These files were re-translated by experienced researchers. To pass the “spot-check,” a 90% agreement between translator and experienced translator was required. For every 12 hours of translated video data, a re-translation was randomly conducted for the body parts (mouth and hands) and macro-activities. In case there was a discrepancy of more than 10% between the translator file and the experienced translator file, both worked together to resolve the discrepancy. Some of the errors may have occurred due to mis-



designations of objects, lack of attention, or the participant not being in view. In order to check to correct and minimize the possible errors, a portion of videos were randomly reviewed following strict protocols attached in Supplemental Material Section S3.1. For intra-observer reliability, some of the videos were translated twice by the same translator. All these files had 90% agreement. All of the QC/QA checks were completed using code developed in Rstudio V1.1 (Supplemental Material Section S3.1.4).

F.3.2.3. Data Analysis

F.3.2.3.1. Analysis for Both Hands and Mouthing Events

The translated plain text files were exported into Microsoft Excel 2013 files before analyzing them using Rstudio V1.1. (Rstudio team, 2017). Contact frequency, hourly contact duration and median contact duration were calculated for the mouth and each hand separately.

- Contact frequency (events/hour) is the total number of contacts of a body part with a specific object divided by the total time that the child was in view (hour).
- Hourly contact duration (min/hour) is the total time that the body part was in contact with the object (min) divided by the total time that the child was in view (hour).
- Median contact duration (second) for each child was calculated from the duration (second) of all the contacts a body part had with an object during the time in view. Median contact duration is only calculated for participants that contacted the object.

To represent the overall distribution for the population, children who did not contact an object/surface were still included in the calculation of contact frequency and hourly contact duration. However, median duration of contact was only calculated among those participants who contacted the object. The object/surfaces were selected and grouped as presented in Table F-76. More specific objects were assigned to the palette categories, and examples are available in Supplemental Material Section S3.1.

All calculations were made for individual categories and also for “non-dietary” and “all objects.” While for contact frequency and hourly contact duration, the values for individual object categories were summed, median contact duration for these aggregate categories was calculated across all of the contacts for that aggregate category. For example, median duration for “all objects” was calculated as the median duration of contact with any object.

Table F-76. Selected Object/Surfaces Summarized in This Study

Object “Super” Categories	Specific Categories from Video Translator -TE Interface
Shin guards	Shin guards
Footwear	Footwear



Object "Super" Categories	Specific Categories from Video Translator -TE Interface
Water bottle	Water bottle
Soccer ball	Soccer ball
Artificial turf	Artificial turf, Black crumb
Body	Body skin, People, Other hand
Clothes	Clothes and towels, Bags and backpacks, Fabric toys
Head	Head face, Hair
Field structures	Plastic tools, Plastic structures, Fabric structures, Metal structure
Dietary objects	Tap water, Other beverages, Water bottle, Other food, Food container
Non-Dietary objects	Everything, but dietary categories
Hands*	Hands
All objects/surfaces	Shin guards, Artificial turf, Black crumb, Head and face, Hair, Clothes and towels, Bag and backpack, Fabric toy, Plastic tools, Plastic structures, Fabric structures, Metal structures, Vegetation, Grass, Body skin, People, Concrete or rock floor, Paper wrapper, Water bottle, Soccer ball, Gloves, Footwear, Tap water, Food container, Wood toy, Other food, Dirt, Puddle water, Plastic or rubber mat, Asphalt, Wood structure, Porous plastic structure, Wood tool, Metal tools, Animal, Mouth guard - pacifier, Sticky food, Other beverages.

* Only used for mouthing events

F.3.2.3.2. Analysis for Macro-Activities, Actions and Intensity

The macro-activities, actions and intensity categories used in this study are presented in Table F-105. Macro-activities are discrete major behavior or activities, such as walking, running, or resting. Using RStudio V1.1, activity patterns were also summarized for macro-activities, specific actions, and player intensity as follows:

- Frequency of specific actions (event/hour) is the total number times a participant did a specific action (event) divided by participant's time in view (hour)
- Hourly duration of macro-activities (min/hour) is the total time a participant spent doing the macro-activity (min) over their total time in view (hour)
- Hourly duration of player intensity (min/hour) is the total time a player spent at each intensity (min) divided by their total time in view (hour)



Table F-77. Macro-Activities, Specific Actions and Player Intensity Categories

Activity Variables	Specific Categories from Video Translator -TE Interface
Macro-activity	Resting standing, Running, Stretching, Walking, Rest, Sit on the ground, Resting sitting on a chair, Push-ups, Sit ups.
Specific Actions	Jumping, Diving, Fall or slip on the ground, Tackling or sliding, Heading
Intensity	Highly, Moderately, Lightly, Resting

All of the previous activity variables were also calculated according to the following categories: event schedule (moment/time), player location, glove use, player age, player gender, player position, and event type. (Table F-106).

Table F-78. Description of Analysis Categories

Category	Description
Event Schedule	During game or practice, warming up, after/before/breaks, other combined time (after/before/breaks, and warming up with exception of during the game).
Player Location	On field, Off field (Sideline, Vehicle, Other)
Glove Use	Gloves on, Gloves off
Player Age	2 - <9 years, 9-16 years, >16 years
Player Gender	Male, Female
Player Position	Goalie, Others
Type of Event	Game, Practice

F.3.2.3.3. Statistical Analysis

Using Rstudio V1.1 (RStudio Team, 2017), the activity pattern variables (i.e., hand and mouth contact frequency, hourly contact duration, median contact duration, duration of macro-activities, frequency of specific actions, and duration of player intensity) were evaluated for significant differences as a function of each of the analysis categories (Table F-106). The Wilcoxon Signed Rank Test for paired data was used to assess if there is a difference in hand contact activity between the left and right hands. The Kruskal-Wallis Test was used to assess for differences across the four age groups. To determine if there are differences between gender, position, event type, we used the Wilcoxon Rank Sum Test.

F.3.3. Results

40 participants were recruited and video recorded at 10 events for a total of 74 hours of footage (Table F-79). Unless specified (i.e., by event schedule), activity patterns are



summarized for the entire time each player was in view on the footage.

Table F-79. Footage and View Time for All 10 Soccer Events (n=40)

Unit	Total Footage	Time not in view	Time in view
Minutes	4410.30	272.30	4138.00
Hours	73.50	4.50	69.00
Percentage	100.00	6.20	93.80

F.3.3.1. Activities with Hands

The left and right hand contact activities with objects are presented in Table F-80 to Table F-85, respectively. The contact frequency, duration and median duration seem to be similar for the left and right hand. There was no significant difference for right and left hand contact activities (Table F-86). Consequently, the activities for left and right hands were combined into both hands.

Table F-80. Left Hand Contact Frequency (Event/Hour) (n=40)

Object	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Artificial turf	4.34	6.06	0.00	0.73	1.75	4.83	16.60	23.96	28.43
Body	12.21	11.52	0.73	3.48	10.42	16.37	30.22	49.55	52.62
Clothes	76.33	39.32	17.15	53.55	65.18	94.63	150.49	194.80	212.21
Field structures	7.33	5.90	0.50	2.18	5.88	10.20	16.83	23.50	26.41
Footwear	5.77	5.33	0.00	1.88	4.33	8.71	12.90	22.34	28.28
Head/face/hair	29.46	26.47	1.80	12.36	20.30	38.68	94.33	105.76	111.06
Shin guards	1.23	2.96	0.00	0.00	0.00	0.91	5.99	12.45	16.44
Soccer ball	20.73	25.79	0.43	6.25	10.35	23.63	63.62	107.16	108.32
Water bottle	6.15	5.23	0.00	1.86	5.25	9.28	13.65	19.73	22.80
Dietary objects	6.21	5.26	0.00	1.86	5.25	9.28	13.65	19.73	22.80
Non-dietary objects	200.97	79.63	87.56	152.38	187.48	217.86	354.94	421.31	429.36
All objects	207.18	80.85	91.21	158.70	193.75	226.28	363.74	427.43	429.85



Table F-81. Left Hand Contact Duration (Min/Hour) (n=40)

Object	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Artificial turf	0.24	0.52	0.00	0.02	0.09	0.27	0.58	2.29	3.21
Body	1.26	1.87	0.01	0.12	0.37	1.79	4.75	7.27	8.41
Clothes	13.94	12.14	1.71	5.51	9.39	17.83	36.88	47.70	52.61
Field structures	0.77	0.83	0.03	0.13	0.46	1.35	2.56	2.97	3.15
Footwear	0.54	0.59	0.00	0.10	0.32	0.88	1.87	2.14	2.23
Head/face/hair	1.02	1.35	0.05	0.33	0.67	1.27	2.45	6.05	8.27
Shin guards	0.12	0.34	0.00	0.00	0.00	0.08	0.67	1.53	1.81
Soccer ball	0.78	0.89	0.01	0.16	0.45	1.12	3.17	3.57	3.81
Water bottle	1.00	1.17	0.00	0.25	0.78	1.52	2.04	5.13	6.73
Dietary objects	1.01	1.17	0.00	0.25	0.80	1.53	2.04	5.13	6.73
Non-dietary objects	21.31	12.68	5.93	12.96	17.40	26.87	45.67	54.05	58.09
All objects	22.32	12.88	6.13	13.11	18.59	28.23	46.10	55.61	58.80

Table F-82. Left Hand Contact Median Duration (second)

Object	N	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Artificial turf	35	2.55	3.72	0.19	0.87	1.22	2.55	9.59	16.73	18.23
Body	40	2.01	1.90	0.57	1.11	1.39	2.30	4.19	9.09	12.17
Clothes	40	4.21	2.55	1.68	2.62	3.38	4.53	9.35	12.44	13.65
Field structures	40	3.47	2.27	0.38	1.85	2.90	4.71	7.71	9.87	10.22
Footwear	40	4.11	3.52	0.93	2.05	2.88	5.24	9.00	15.82	19.75
Head/face/hair	40	1.34	0.51	0.55	1.08	1.20	1.56	2.25	2.87	3.16
Shin guards	17	4.62	4.26	1.41	1.82	3.31	5.38	12.01	15.76	16.70
Soccer ball	40	1.69	0.77	0.64	1.20	1.55	2.03	3.16	3.93	4.39
Water bottle	38	7.88	5.83	0.73	4.66	6.39	10.18	16.14	27.90	31.77
Dietary objects	38	7.86	5.84	0.73	4.71	6.39	10.18	16.14	27.90	31.77



Object	N	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Non-dietary objects	40	2.33	1.14	0.54	1.61	2.15	2.79	3.65	6.21	7.19
All objects	40	2.41	1.12	0.54	1.65	2.22	2.98	3.75	6.09	6.95

Table F-83. Right Hand Contact Frequency (Event/Hour) (n=40)

Object	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Artificial turf	4.46	5.72	0.00	1.26	2.55	4.89	12.58	25.24	26.94
Body	14.77	11.99	1.85	6.33	10.90	18.56	38.45	50.19	56.20
Clothes	72.63	34.94	18.11	47.21	68.24	87.23	151.98	154.52	156.13
Field structures	7.86	6.59	1.34	3.43	5.77	10.97	19.72	27.47	30.60
Footwear	6.75	5.89	0.73	2.76	5.74	8.36	14.37	25.84	33.06
Head/face/hair	27.32	24.59	1.39	9.49	19.78	37.98	70.45	103.02	116.46
Shin guards	1.47	3.69	0.00	0.00	0.00	1.23	5.53	16.32	21.16
Soccer ball	20.90	25.07	0.51	4.92	12.00	31.55	68.54	101.57	104.03
Water bottle	7.08	6.03	0.00	3.28	5.72	8.83	18.70	25.89	29.73
Dietary objects	7.12	6.04	0.00	3.28	5.72	8.83	18.70	25.89	29.73
Non-dietary objects	199.52	70.80	95.10	145.07	184.38	230.37	308.10	378.36	418.10
All objects	206.64	71.95	101.04	149.90	195.47	249.85	318.85	382.38	419.59

Table F-84. Right Hand Contact Frequency (Min/Hour) (n=40)

Object	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Artificial Turf	0.27	0.53	0.00	0.02	0.11	0.30	0.81	2.42	2.90
Body	1.25	1.65	0.04	0.16	0.54	1.62	4.73	5.67	5.73
Clothes	12.21	11.16	1.53	5.72	8.35	14.32	34.98	47.52	52.84
Field Structures	0.84	0.99	0.02	0.20	0.43	1.05	3.40	3.51	3.55
Footwear	0.59	0.56	0.02	0.15	0.31	0.86	1.73	1.89	1.93
Head/face/hair	0.99	1.42	0.01	0.31	0.61	1.20	2.15	6.32	8.87
Shin guards	0.12	0.34	0.00	0.00	0.00	0.09	0.65	1.50	1.89



Object	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Soccer ball	0.95	1.06	0.01	0.25	0.67	1.19	3.85	4.22	4.38
Water bottle	1.42	1.44	0.00	0.53	1.01	1.84	3.77	6.10	6.80
Dietary Objects	1.43	1.44	0.00	0.53	1.01	1.84	3.77	6.10	6.80
Non-dietary Objects	19.70	11.82	5.88	12.39	15.35	25.67	44.15	53.47	57.80
All Objects	21.13	12.06	6.53	12.87	18.15	26.10	44.64	55.62	59.75

Table F-85. Right Hand Contact Median Duration (Second) (n=40)

Object	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Artificial Turf	1.37	0.84	0.18	0.74	1.18	1.74	3.18	3.25	3.27
Body	1.59	1.39	0.43	0.91	1.18	1.85	3.15	7.11	7.70
Clothes	4.31	2.66	1.51	2.61	3.49	4.99	8.66	12.62	12.88
Field Structures	3.52	3.62	0.72	1.46	2.66	4.23	7.74	16.61	22.10
Footwear	3.55	3.18	0.83	1.55	2.30	4.99	9.91	13.84	15.40
Head/face/hair	1.34	0.47	0.54	1.05	1.24	1.48	2.13	2.94	3.42
Shin guards	4.96	5.07	0.90	2.28	3.02	5.39	16.97	18.00	18.26
Soccer ball	2.54	1.79	0.46	1.53	1.97	3.24	6.76	8.02	8.40
Water bottle	8.18	5.07	2.01	4.78	6.57	10.25	15.96	23.88	24.74
Dietary Objects	8.17	5.07	2.01	4.78	6.57	10.25	15.96	23.88	24.74
Non-dietary Objects	2.27	1.09	0.33	1.49	2.02	2.89	3.79	5.62	6.34
All Objects	2.36	1.12	0.33	1.55	2.26	2.98	3.94	5.79	6.52

Table F-86. p-Values from Comparing the Left and Right Hand Activities Using Wilcoxon Signed Rank Test

Object	Frequency* (n=40)	Duration* (n=40)	Median Duration*
Artificial Turf	0.519	0.703	0.401 (n=70)
Body	0.154	0.413	0.062 (n=80)
Clothes	0.765	0.557	0.946 (n=80)
Field Structures	0.707	0.686	0.447 (n=80)



Object	Frequency* (n=40)	Duration* (n=40)	Median Duration*
Footwear	0.303	0.651	0.168 (n=78)
Head/face/hair	0.630	0.729	0.871 (n=80)
Shin guards	0.499	0.472	0.917 (n=30)
Soccer ball	0.950	0.356	0.153 (n=80)
Water bottle	0.567	0.164	0.567 (n=74)
Dietary Objects	0.564	0.167	0.566 (n=74)
Non-dietary Objects	0.954	0.535	0.825 (n=80)
All Objects	0.908	0.744	0.817 (n=80)

*There were no significant differences between left and right hand for frequency, duration and median duration.

Since there was no significant difference between left and right hand contact activities, the data sets were combined to calculate activity patterns for both hands (Table F-86). The both hands contact frequency for participants playing soccer on turf is summarized in Table F-87. The median hand contact frequency with artificial turf is 2.39 events/hour, with a maximum of 27.68 events/hour. Aside from clothes, body and their head, participants contacted soccer balls the most frequently with a median contact frequency of 10.83 event/hour. The median hand contact frequency was 160.99 events/hour for non-dietary objects.

Table F-87. Table 3.1.8 Both Hands Contact Frequency (Event/Hour) (n=40)

Objects	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Artificial turf	4.49	5.90	0.00	1.09	2.39	4.83	17.44	23.79	27.68
Body	13.54	11.52	1.52	5.10	9.78	16.48	35.48	49.38	54.40
Clothes	74.72	35.55	17.63	54.03	65.91	91.02	151.33	174.31	182.29
Field structures	7.62	6.01	1.46	3.04	6.03	10.64	19.65	23.88	24.75
Footwear	6.32	5.52	0.72	2.26	5.05	9.15	12.68	23.74	30.66
Head/face/hair	28.74	25.09	2.28	12.35	20.21	33.70	78.41	103.75	113.79
Shin guards	1.36	3.30	0.00	0.00	0.00	1.10	5.76	14.38	18.79
Soccer ball	21.40	27.34	0.55	5.60	10.83	29.33	65.67	116.98	123.89
Water bottle	6.67	5.37	0.00	3.23	5.15	9.04	17.57	20.82	20.89



Objects	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Dietary Objects	6.72	5.39	0.00	3.27	5.15	9.04	17.57	20.82	20.89
Non-dietary objects	169.80	66.85	73.89	122.44	160.99	188.22	303.96	352.05	361.74
All objects	176.52	68.47	80.69	131.96	165.08	197.66	308.60	364.63	371.23

The hourly hand contact duration for participants playing soccer on artificial turf is summarized in Table F-88. The median hourly hand contact duration for artificial turf was 0.11 min/hour, and the maximum is 3.05 min/hour. The median hand hourly duration with non-dietary objects was 15.02 min/hour.

Table F-88. Both Hands Hourly Contact Duration (Min/Hour) (n=40)

Objects	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Artificial turf	0.26	0.51	0.00	0.03	0.11	0.31	0.79	2.24	3.05
Body	1.26	1.72	0.03	0.16	0.43	2.03	4.56	6.25	6.99
Clothes	12.75	11.81	1.60	5.74	9.02	16.84	35.19	47.61	52.73
Field structures	0.81	0.87	0.03	0.18	0.44	1.29	2.43	3.20	3.30
Footwear	0.57	0.57	0.01	0.16	0.32	0.86	1.85	1.98	1.98
Head/face/hair	1.01	1.37	0.03	0.35	0.65	1.44	2.15	6.09	8.57
Shin guards	0.12	0.34	0.00	0.00	0.00	0.08	0.66	1.51	1.85
Soccer ball	0.89	1.03	0.01	0.22	0.57	1.19	3.61	4.20	4.27
Water bottle	1.23	1.26	0.00	0.42	0.89	1.77	2.74	5.48	6.77
Dietary objects	1.23	1.26	0.00	0.42	0.91	1.77	2.74	5.48	6.77
Non-dietary objects	18.44	12.01	1.27	9.83	15.02	22.65	42.63	52.64	56.91
All objects	19.67	12.11	3.44	11.78	15.92	24.87	43.15	54.50	58.83

The median hand contact duration for participants while playing on artificial turf is summarized in Table F-89. Median duration is only calculated from those participants that contacted the object. The median hand contact duration for artificial turf was 1.20 second, and the maximum was 5.14 seconds. The median hand contact duration was 2.10 for non-dietary objects. Note that for the all objects (Table F-76), median duration was calculated across all of the contacts with all of the object categories included. Thus, sometimes the values will be lower than for some of the individual object categories,



particularly if there was proportionally a lower number of contacts with that object.

Table F-89. Both Hands Contact Median Duration (Second)

Objects	n*	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Artificial turf	35	1.53	1.11	0.46	0.84	1.20	1.65	3.85	4.95	5.14
Body	40	1.70	1.62	0.53	0.96	1.26	1.92	3.13	8.43	9.78
Clothes	40	4.21	2.54	1.85	2.50	3.43	4.77	9.26	12.23	13.16
Field structures	40	3.47	3.27	0.41	1.78	2.56	4.40	6.93	15.19	19.96
Footwear	40	3.74	3.38	0.89	1.66	2.47	4.83	9.91	15.18	15.34
Head/face/hair	40	1.35	0.46	0.60	1.11	1.21	1.59	2.08	2.79	3.09
Shin guards	17	4.28	4.26	0.90	1.61	2.64	4.68	13.55	16.07	16.70
Soccer ball	40	1.86	0.87	0.64	1.20	1.64	2.09	3.78	3.90	3.95
Water bottle	38	6.87	2.98	2.09	5.19	6.64	8.41	13.40	13.76	13.79
Dietary objects	38	6.84	2.97	2.09	5.16	6.64	8.30	13.40	13.76	13.79
Non-dietary objects	40	2.48	1.20	1.23	1.70	2.10	2.93	4.22	6.50	6.93
All objects	40	2.57	1.20	1.25	1.79	2.18	3.01	4.28	6.58	6.94

*Only calculated from participants who contacted the object.

F.3.3.2. Mouthing activities

The mouthing contact frequency for participants playing soccer on turf is summarized in Table F-90. None of the participants observed contacted artificial turf (including black crumb) with their mouths during the recordings. There were also no mouthing contacts with shin guards or footwear observed. The median mouth contact frequency was 7.55 and 14.48 events/hour for hands and non-dietary objects, respectively.

Table F-90. Mouthing Contact Frequency (Event/Hour) (n=40)

Objects	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Artificial turf*	NA	NA	NA	NA	NA	NA	NA	NA	NA
Body skin	0.55	1.71	0.00	0.00	0.00	0.12	2.1	7.71	9.93



Objects	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Clothes	9.13	16.98	0.00	1.33	4.05	11.1	24.3	74.82	104.27
Field structures	0.01	0.06	0.00	0.00	0.00	0.00	0.00	0.24	0.4
Footwear*	NA	NA	NA	NA	NA	NA	NA	NA	NA
Hands	8.91	5.87	0.71	5.64	7.55	11.31	18.41	25.8	26.03
Head/face/hair	0.03	0.15	0.00	0.00	0.00	0.00	0.02	0.67	0.83
Shin guards*	NA	NA	NA	NA	NA	NA	NA	NA	NA
Soccer ball	0.01	0.09	0.00	0.00	0.00	0.00	0.00	0.34	0.55
Water bottle	4.96	4.65	0.00	1.7	3.75	6.43	15.58	17.7	17.81
Dietary objects	5.17	4.56	0.00	2.04	4.03	6.85	15.58	17.7	17.81
Non-dietary objects	19.81	18.76	2.94	9.78	14.48	23.68	42.97	91	111.92
All objects	24.97	19.94	3.57	12.18	18.87	30.93	50.75	96.76	118.87

*No participants contacted artificial turf, shin guards or footwear with their mouths

NA: not applicable

The hourly mouthing contact duration for participants playing soccer on artificial turf is summarized in Table F-91. There were no mouthing contacts observed with artificial turf, shin guards, or footwear. The median mouthing hourly duration was 0.21 and 0.41 min/hour with hands and non-dietary objects, respectively.

Table F-91. Mouthing Hourly Contact Duration (Min/Hour) (n=40)

Objects	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Artificial turf*	NA	NA	NA	NA	NA	NA	NA	NA	NA
Body skin	0.01	0.04	0.00	0.00	0.00	0.00	0.04	0.2	0.25
Clothes	0.31	0.52	0.00	0.03	0.12	0.37	1.00	2.32	2.76
Field structures	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.03
Footwear*	NA	NA	NA	NA	NA	NA	NA	NA	NA
Hands	0.41	0.76	0.01	0.10	0.21	0.37	0.92	3.32	4.74
Head/face/hair	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.01
Shin guards*	NA	NA	NA	NA	NA	NA	NA	NA	NA
Soccer ball	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00



Water bottle	0.36	0.55	0.00	0.10	0.24	0.41	1.12	2.5	3.17
Dietary objects	0.37	0.54	0.00	0.10	0.25	0.42	1.12	2.5	3.17
Non-dietary objects	0.76	0.92	0.05	0.20	0.41	0.98	1.8	4.21	4.99
All objects	1.13	1.18	0.06	0.40	0.82	1.42	3.06	5.34	6.45

*No participants contacted these objects with their mouth.

NA: not applicable

The median mouthing contact duration for participants while playing on artificial turf is summarized in Table F-92. Median duration is only contacted from those children that contacted the object with their mouth. No participants were observed contacting artificial turf, shin guards or footwear with their mouths. The median mouthing contact duration was 1.10 second for both hands and 1.14 second non-dietary objects. Thus, sometimes the values will be lower than for some of the individual object categories, particularly if there was proportionally a lower number of contacts with that object.

Table F-92. Mouthing Median Contact Duration (Second)

Objects	n	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Artificial turf	0	NA	NA	NA	NA	NA	NA	NA	NA	NA
Body skin	10	1.00	0.41	0.54	0.61	0.98	1.25	1.59	1.60	1.60
Clothes	36	1.72	1.7	0.62	0.91	1.36	1.78	3.62	8.35	10.61
Field structures	1	5.18	ND	ND	ND	ND	ND	ND	ND	ND
Footwear	0	NA	NA	NA	NA	NA	NA	NA	NA	NA
Hands	40	1.35	0.79	0.66	0.91	1.10	1.45	2.56	4.15	4.76
Head/face/hair	2	1.15	0.5	0.79	0.97	1.15	1.33	1.47	1.50	1.50
Shin guards	0	NA	NA	NA	NA	NA	NA	NA	NA	NA
Soccer ball	1	0.13	ND	ND	ND	ND	ND	ND	ND	ND
Water bottle	35	3.60	1.33	1.71	2.59	3.37	4.40	5.81	6.5	6.83
Dietary objects	37	3.48	1.31	1.71	2.48	3.15	4.34	5.73	6.46	6.83
Non-dietary objects	37	1.33	0.73	0.74	0.87	1.14	1.40	2.55	3.87	4.45
All objects	39	1.64	0.86	0.79	1.08	1.34	2.06	3.07	4.5	5.36

NA: not applicable; and ND: not determined

F.3.3.3. Macro-Activities (e.g., Walking, Running)

The hourly duration for participants engaging in macro-activities on artificial turf is



summarized in Table F-93. The macro-activities that participants engaged in for the largest proportion of their time were walking, standing, and running with median durations of 24.51, 16.75, 13.85 min/hour respectively. All 40 participants were observed doing those activities. 39 participants were observed sitting on the ground, while only 17 were observed sitting in a chair. 31 participants were observed stretching and only 2 participants were observed doing push-ups.

Table F-93. Hourly Duration (Min/Hour) of Macro-Activities (n=40)

Macro-Activities	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Push ups	0.0	0.02	0.0	0.0	0.0	0.0	0.0	0.06	0.1
Running	13.69	6.13	2.47	9.13	13.85	18.21	23.61	24.49	24.99
Sit ups*	NA	NA	NA	NA	NA	NA	NA	NA	NA
Sitting on chair	2.26	4.5	0.00	0.00	0.00	2.73	11.61	17.65	17.67
Sitting on ground	1.18	1.73	0.00	0.11	0.55	1.57	3.21	7.69	9.75
Standing	17.31	5.54	6.98	14.99	16.75	19.89	27.23	31.06	32.5
Stretching	0.75	1.27	0.00	0.09	0.37	0.81	3.24	5.6	5.86
Walking	24.49	6.08	10.5	21.18	24.51	28.52	33.63	35.87	37.28

*No participants were observed doing sit ups.

NA: not applicable

F.3.3.4. Specific Actions (e.g., Tackling, Falling)

The event frequency for participants engaging in specific actions on artificial turf is summarized in Table F-94. The specific actions that participants engaged in the most frequently was jumping with a median frequency of 3.65 events/hour, followed by slipping/falling with a median frequency of 0.78 event/hour. 34 participants were observed jumping. 27 participants were observed slipping or falling on the artificial turf. Fewer participants were observed diving, heading, and tackling, with only 8, 12, and 13 participants observed conducting these actions, respectively.

Table F-94. Frequency (Event/Hour) of Actions During the Whole Event (n=40)

Specific Actions	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Diving	0.53	1.69	0.00	0.00	0.00	0.00	2.44	7.40	9.89
Heading	0.33	0.61	0.00	0.00	0.00	0.48	1.91	2.07	2.10
Jumping	9.46	17.72	0.00	0.68	3.65	9.30	65.16	71.43	75.30
Slip/fall	2.19	3.45	0.00	0.00	0.78	2.31	11.30	13.06	13.19



Specific Actions	Mean	SD	Min	p25	Median	p75	p95	p99	Max
Tackling or sliding	0.42	0.78	0.00	0.00	0.00	0.51	2.20	2.81	2.81

F.3.3.5. Player Intensity

The hourly duration for player intensity while on artificial turf is summarized in Table F-95. Players spent the majority of their time at “low” intensity, with a median duration of 24.00 min/hour. Low intensity corresponds to players who were moving but walking. Moderate and high intensity correspond to jogging and running, respectively. All 40 participants were observed at least for some time at each of the 4 intensities.

Table F-95. Hourly Duration (Min/Hour) of Player Intensity (n=40)

Intensity	Mean	SD	Min	p25	Median	p75	p95	p99	Max
High	2.00	0.99	0.09	1.15	1.99	2.68	3.64	3.84	3.88
Moderate	12.52	5.09	3.75	8.65	12.98	15.88	20.80	22.05	22.82
Low	24.04	5.31	10.65	21.58	24.00	27.47	31.05	33.75	34.29
Resting	21.44	7.81	10.86	16.50	18.68	26.98	33.83	40.91	41.81

F.3.3.6. Analyzed by Event Schedule, Location, Gloves Usage, Age, Gender, Players' Position, and Event Type

F.3.3.6.1. Event Schedule (Moment/Time)

The data was summarized based on the event schedule which included: the whole event (Section F.3.3.1 to Section F.3.3.5), during the game/practice, warming up, after/before/break time, and other combined time with the exception of during the game (i.e., warming up and after/before/break time combined). The hand to object, mouthing events, actions, macro-activities, and intensity according to different event times are presented in Supplemental Material Section S3.2. As presented in Supplemental Material Section S3.2, the data was summarized for each specific time of the event such as *during the game, game/practice, warming up, after/before/break time, and other combined time*.

During the Game (Supplemental Material Section S3.2.1.1)

Data during the game/practice were very similar to the whole event. There were more contacts of hands to artificial turf (median frequency: 2.16 event/hour), followed by contacts of hands to footwear (median frequency: 3.44 event/hour), and hands to soccer ball (median frequency: 9.97 event/hour) than contacts with dietary objects (median frequency: 1.30 event/hour). However, the median contact duration was longer for dietary objects than all of the other objects, which may indicate that there was a longer holding time of these dietary objects than the rest of the objects. The mouth to hand median contact frequency was 7.49 event/hour. Also, the most common action event per hour was jumping (median: 2.32 event/hour) followed by slipping/falling on the



artificial turf (median 0.81 event/hour). With respect to macro-activities, players spent the majority of the time walking (median: 25.8 min/hour) and running (median: 15.10 min/hour) during the game and only about 0.37 min/hour sitting on the ground, which corresponds to their intensity, as they spent only 1.84 min/hour doing high intensity activities. (See Supplemental Material S3.2.1.1 of activity patterns during games/practice for details).

Warming Up (Supplemental Material Section S3.2.1.2)

The median frequency of players hands to artificial turf (0.00 event/hour) contact was less during warm up than during the game as only four players made hand to artificial turf contact while warming up. Also, the median mouthing frequency event for mouth to hand (0.00 event/hour) contact was lower than during the game. Similar to during the game, players spent the majority of the time walking (median: 21.12 min/hour) and running (median: 15.21 min/hour). However, while warming up, they spent less time sitting on the ground (0.00 min/hour) than during the game. Additionally, with respect to intensity, players spent most of the time resting (median: 5.30 min/hour) while warming up (See Supplemental material Section S3.2.2.2 for details). *Please note that the intensity corresponds to actions and macro-activities together.*

After/Before/Break Time (Supplemental Material Section S3.2.1.3)

Similar to warming up, the median frequency of players hands to artificial turf contacts was 0.00 event/hour. For the mouthing with artificial turf, regardless of the time of the event, there was not any mouthing events with artificial turf and only one person had a mouth to soccer ball contact. The median mouthing frequency event for mouth to hand was 0.00 event/hour. For macro-activities, "Resting standing" was the macro-activity that players did most of the time (median: 30.40 min/hour) on after/before/or during break time, which coincides with the intensity where they spent most of the time resting (37.00 min/hour). (See Supplemental Material Section S3.2.1.3 for details).

Other Combined Time (Supplemental Material Section S3.2.1.4)

The median frequency of players hands to artificial turf contacts was 0.00 event/hour. The hand to non-dietary objects median contact frequency was 206.41. However, there were no mouthing events with artificial turf. Similarly to when players were on before/after/break time, dietary objects were contacted most frequently, for longer hourly durations and for longer periods of time (8.50 event/hour, 1.10 min/hour, and 3.70 seconds) than non-dietary objects. Also, "Resting standing" was the macro-activity that players did most of the time (median: 26.39 min/hour) on other combined time, while the intensity that they spent most of the time was resting (34.95 min/hour).

F.3.3.6.2. Location (On or Off Field) (Supplemental Material Sections S3.2.2.1 and S3.2.2.2)

There seems to not be some differences between hand to mouth contact when players were on or off the field. The hands to artificial turf median frequency contact for players



recorded on field (2.61 event/hour) was higher than players recorded off field (0.00 event/hour). The mouthing median frequency with hands was higher for players on the field (6.99 event/hour) than off the field (5.90 event/hour). However, mouthing median frequency contact with clothes was higher off field than on field. While jumping was the most common event per hour for players on the field and off the field, players spent more time walking (median: 27.19 min/hour) on the field and more than resting (median: 25.99 min/hour) when off the field. (See Supplemental Material Sections S3.2.2.1 and S3.2.2.2 for details).

F.3.3.6.3. Use Of Gloves (On or Off) (Supplemental Material Sections S3.2.3.1 and S3.2.3.2)

There seem not be a large difference between the median frequency of players hands to artificial turf contacts when players wore gloves or not (2.30 event/hour and 0.0 event/hour, respectively). However, more mouthing events with hands were recorded for players who did not wear any gloves during the than the players who wore gloves at some point during the whole event (1.88 event/hour and 7.18 event/hour for people wearing gloves and not wearing gloves, respectively). (See Supplemental Material Sections S3.2.3.1 and S3.2.3.2 for details).

F.3.3.6.4. Player Age (Supplemental Material Section S3.3)

Activity statistics by age and results from statistical tests (Wilcoxon sum rank test) are presented in Supplemental Material Section S3.3.

During the Whole Event (Supplemental Material Section S3.3.1):

Hand to Object Contacts. There were significant differences ($p < 0.05$) in contact frequency (events/hour) between the three age groups with artificial turf, clothes, soccer ball, and field structures. The younger age group (2 to <9 years) had more contacts per hour with artificial turf, soccer ball, and field structure than the other two older age groups. There were significant differences in hourly contact duration between all three age groups with artificial turf and field structures. While the youngest children had the longer hourly contact duration with artificial turf, the oldest age group had longer hourly contact duration with field structures. However, for median duration, contact with artificial turf was not significantly different between the three age groups, but contact with non-dietary objects, clothes, and soccer ball were significantly different between the three age groups. The middle age group had the longest durations of contact with clothes. The oldest age group had the longest durations of contact with the soccer ball and the shortest with non-dietary objects.

Mouthing Events: There were significant differences ($p < 0.05$) in mouthing contact frequency (event/hour) between age groups with water bottle and non-dietary objects. The younger age group (2 to <9 years) had more mouthing contacts per hour with water bottle and non-dietary objects than the other two age groups. There were also significant differences ($p < 0.05$) in mouthing duration (min/hour) between age groups with water bottle, non-dietary objects and all objects. For the mouthing median duration,



there was a significant difference ($p < 0.05$) between age groups with hands. The younger age group (2 to <9 years) had longer periods of hand to mouth contacts than the other age groups.

Actions: There were significant differences in frequency (event/hour) between the three age groups while players were jumping and slipping/falling. Older soccer players (16+ years) were jumping and slipping/falling on the ground more frequently per hour than the younger age groups.

Macro-Activities. There was a significant difference in hourly duration (min/hour) between the three age groups while players were stretching. The older group (16 yrs +) spent more minutes per hour stretching than the other two age groups.

Intensity. There were no significant differences between the three age groups in hourly duration of player intensity.

During the Game/Practice (Supplemental Material Section S3.3.2.1):

Hands to Object Contacts. There were significant differences ($p < 0.05$) in contact frequency (event/hour) between the three age groups with artificial turf, water bottle and dietary objects during the game/practice. The younger age group (2 to <9 years) had more contacts per hour with artificial turf, water bottle and dietary objects than the other two age groups. There was only one significant difference in hourly duration between three age groups with soccer ball. The middle age group of players (9 to <16 years) contacted soccer ball more minutes per hour than the other age groups. For median duration (second), there were significant differences between the three age groups with artificial turf, body, head/face/hair, water bottle, and dietary objects.

Mouthing Events. There were no significant differences ($p < 0.05$) in mouthing contact frequency (event/hour) between age groups during the game/practice. There was a significant difference in mouthing duration (min/hour) of hands between the age group. The younger age group (2 to <9 years) had longer mouthing of hands per hour than the other age groups. Similarly, for the median duration (second), there was a significant difference in mouthing between age groups with hands. The younger age group (2 to <9 years) had longer durations of hand to mouth contacts than other two age groups.

Actions. There were significant differences in frequency (event/hour) between the three age groups for players jumping. Older soccer players (16+ years) jumped more frequently per hour than the younger age groups.

Macro-Activities. There were no significant differences between the three age groups in hourly duration of macro-activities.

Intensity. There were no significant differences between the three age groups in hourly duration of player intensity.

Warming up (Supplemental Material Section S3.3.2.2):

Hands to Object Contacts. There were no significant differences in contact frequency



(event/hour), duration (min/hour), or median duration (second) between the three age groups with any of the observed objects during warm up.

Mouthing Events. Similarly, as hand to object activities, there were no significant differences in mouthing contact frequency (event/hour), duration (min/hour), or median duration (second) between the three age groups with any of the observed objects during warm up.

Actions. There were no significant differences between the three age groups for any of the specific actions while players were warming up.

Macro-Activities. There was a significant difference in hourly duration between age groups for time spent walking during warm up. The older groups (9 to <16 years and 16 years +) spent more time walking than the younger age groups.

Intensity. There were no significant differences between the three age groups in hourly duration of player intensity.

After/Before/Break Time (Supplemental Material Section S3.3.2.3):

Hands to Object Contacts. There were significant differences ($p < 0.05$) in contact frequency (event/hour) between age groups with body, soccer ball, non-dietary objects, and all objects during after or before the event and during break times. The older age group (16+ years) had fewer contacts per hour with body, soccer ball, non-dietary objects, and all objects than the other two age groups. There was a significant difference in contact duration (min/hour) between age groups with non-dietary objects. The middle age group of players (9 to <16 years) contacted non-dietary objects for more minutes per hour than the other age groups. For the median duration, there were no significant differences between age groups.

Mouthing Events. There were no significant differences in mouthing contact frequency (event/hour), duration (min/hour), or median duration (second) between the three age groups with any of the observed objects after or before the event and during break times.

Actions. There were no significant differences in frequency (event/hour) of specific actions between age groups.

Macro-Activities. There were no significant differences in contact frequency and median duration between age groups. However, there were significant differences in hourly duration (min/hour) between age groups while players were sitting on a chair or stretching. The older groups (9 to <16 years and 16 years +) had less minutes per hour stretching or sitting on a chair than the younger age group. Also, there were significant difference in hourly duration between age groups for time players spent walking and standing. The older groups (9 to <16 years and 16 years +) had more minutes per hour walking or standing than the younger age group.

Intensity. There were significant differences in hourly duration (min/hour) of intensity



between age groups. Older groups (9 to <16 years and 16 years +) had longer periods of light intensity, but shorter periods of moderate and high intensity than the younger age group.

Other Combined Time (After/Before/Break/Warming Up Time) But During the Game (Supplemental Material Section S3.3.2.4):

Hands to Object Contacts. There was a significant difference in contact frequency (event/hour) with non-dietary objects between age groups during the other combined times that did not include the actual game or practice. The younger age group (2 to <9 years) had less contacts per hour with non-dietary objects than the other two age groups. There were significant differences in hourly duration (min/hour) between age groups with soccer ball and non-dietary objects. The middle age group (9 to <16 years) contacted the ball and non-dietary objects for more minutes per hour than the other age groups. There were no significant differences in median duration (second) between age groups with any of the observed objects during this other combined time.

Mouthing events. There were significant differences ($p < 0.05$) in mouthing frequency (event/hour) between age groups with water bottle, non-dietary objects, and all objects. The older age group (16+ years) had fewer mouthing contacts with these objects than the other two groups. Also, there were significant differences ($p < 0.05$) in hourly contact duration (min/hour) between age groups with water bottle, non-dietary objects, and all objects. The older age group (16+ years) had a shorter contact duration (min/hour) with these objects than the other two groups. For the median duration (second), there were no significant differences between the three age groups.

Actions. There were no significant differences in frequency of any specific action between age groups not during the game or practice.

Macro-activities. Not during the game or practice, there was a significant difference in duration (min/hour) between age groups while players were resting standing. The older group (16 years +) had shorter periods of standing per hour than the younger age groups. The younger age group (2 to <9 years) also had longer durations for running than other age groups. The older age group (16+ years) spent more time stretching than the other two groups.

Intensity. There were no significant differences between the three age groups in hourly duration of player intensity not during the game or practice (note: intensity includes activities and actions)

On Field (Supplemental Material Section S3.3.3.1):

Hands to Object Contacts. There was a significant difference in contact frequency (event/hour) between age groups with body while participants were on the field. The older age group (16+ years) had more contacts per hour with body than the other two age groups. Another significant difference in contact frequency (event/hour) between age groups was observed with soccer ball. There were significant differences in hourly



duration (min/hour) between age groups with soccer ball. The middle age group of players (9 to <16 years) contacted the ball for more minutes per hour than the other age groups. For the median contact duration, there were no significant differences between age groups with any of the observed objects.

Mouthing Events. There were significant differences ($p < 0.05$) in mouthing hourly duration between age groups with hands. The middle age group (2 to <9 years) had a greater contact duration (min/hour) with hands than the other two age groups while on the field. The mouthing frequency and median duration were not significantly different between age groups while players were on the field.

Actions. There were significant differences in frequency (event/hour) of specific actions between age groups when players were on the field. The older age group (16+ years) jumped and slipped/fell on the ground more times per hour than the younger age groups.

Macro-Activities. There were no significant difference between age groups in hourly duration of player macro-activities while on the field.

Intensity. There were no significant differences between age groups in hourly duration of player intensity while on the field.

Off Field (Supplemental Material Section S3.3.3.2):

Hands to Object Contacts. There was a significant difference in contact frequency (event/hour) between age groups with artificial turf while players were off the field. The younger age group (2 to <9 years) had more contacts per hour with artificial turf than the other two age groups. There were significant differences in contact hourly duration (min/hour) between age groups with water bottles and artificial turf while off the field. The younger age group (2 to <9 years) contacted the turf and water bottles for more minutes per hour than the other age groups. For median contact duration, there were no significant differences between age groups with any of the observed objects (note that artificial turf can be present on sideline and not just on soccer field).

Mouthing Events. There was a significant difference ($p < 0.05$) in mouthing frequency (event/hour) between age groups with “all objects” while players were off the field. The youngest group (2 to <9 years) had more mouthing events with all objects per hour than the other two groups. There were no statistical differences in mouthing hourly duration between age groups. Also, there were no significant difference in median duration between age groups with contact with any object, while players were off the field.

Actions. There were no significant differences in frequency of specific actions between age groups while players were off the field.

Macro-Activities. There was a significant difference in hourly duration between age groups when players were walking off the field. The younger age group (2 to <9 years) had less minutes walking per hour than the older age groups.



Intensity. There were no significant differences between age groups in hourly duration of player intensity.

Gloves On (Supplemental Material Section S3.3.4.1):

Hands to Object Contacts. There were no significant differences ($p < 0.05$) in contact frequency (event/hour) between age groups. There were no significant differences in contact hourly duration (min/hour) between age groups with field structures and artificial turf. There were significant differences in median contact duration (second) between age groups with water bottle, and dietary objects. The middle age group (9 to <16 years) had longer contacts with clothes and non-dietary than the other age groups. The younger age group (2 to <9 years) contacted water bottle and dietary objects for longer periods than the other age groups.

Mouthing Events. There were no significant difference ($p < 0.05$) in mouthing contact frequency (event/hour) between age groups. There were no significant difference ($p < 0.05$) in hourly contact duration (min/hour) between the age groups. There was a significant difference ($p < 0.05$) in mouthing median duration between the age groups with water bottle. The younger group (2 to <9 years) had shorter mouthing periods with water bottle than the other age groups, even while wearing gloves.

Gloves Off (Supplemental Material Section S3.3.4.2):

Hands to Object Contacts. There were significant differences ($p < 0.05$) in hand contact frequency (event/hour) between age groups with clothes, water bottle, dietary objects, non-dietary objects, and all objects when players were not wearing gloves. The older age group (16+ years) had more contacts per hour with water bottle, dietary objects, non-dietary objects, and all objects than the other two age groups. There was a significant difference in hourly contact duration (min/hour) between age groups with soccer ball, water bottle and dietary objects. The older age group (16+ years) contacted turf for less minutes per hour than the other age groups, while not wearing gloves. There were no significant differences in hand contact median duration (second) between age groups.

Mouthing Events. There were significant differences ($p < 0.05$) in mouthing frequency (event/hour) between age groups with water bottle and dietary objects. The younger group (2 to <9 years) had more frequent mouthing with these two objects than the other two groups. There was a significant difference ($p < 0.05$) in hourly mouthing duration (min/hour) between age groups with water bottle, dietary, and all objects. The younger group (2 to <9 years) had more mouthing minutes per hour with these objects than the other two groups. There were no significant difference ($p < 0.05$) in mouthing median duration (second) between age groups.

F.3.3.6.5. Player Gender

Activity statistics by gender and results from statistical tests (Wilcoxon sum rank test) are presented in **Supplemental Material Section S3.4.**



During the Whole Event (Supplemental Material Section S3.4.1):

Hands to Object Contacts. There were significant differences ($p < 0.05$) in contact frequency (events/hour) between males and females with field structures, head/face/hair, water bottle, dietary, non-dietary, and all objects. Females had more contacts per hour with field structures, head/face/hair, water bottle, dietary, non-dietary, and all objects than males. There were significant differences ($p < 0.05$) in hourly contact duration between males and females for head/face/hair, shin guards, water bottle, and dietary objects. Females' hands contacted these objects for more minutes per hour than males. For median contact duration, there were no significant differences between males and females.

Mouthing Events. There were significant differences ($p < 0.05$) in mouthing contact frequency (events/hour) between males and females with water bottle and dietary objects. Females had more mouthing contacts with these objects than males. Similarly, there were significant differences ($p < 0.05$) in mouthing duration (min/hour) between males and females with water bottle, and dietary objects. Females had longer mouthing of water bottle and dietary objects per hour than males. For the mouthing median contact duration, there were significant differences ($p < 0.05$) between males and females with dietary and all objects. Females touched these objects with their mouths for longer periods of times than males.

Actions. There were no significant differences between males and females in frequency of specific actions.

Macro-Activities. There were no significant differences between males and females in hourly duration of macro-activities.

Intensity. There were no significant differences between males and females in hourly duration of player intensity.

During the Game/Practice (Supplemental Material Section S3.4.2.1):

Hands to Object Contacts. There were significant differences ($p < 0.05$) in hand contact frequency (event/hour) between males and females with head/face/hair, non-dietary objects, and all objects while during the game or practice. Females had more contacts per hour with these objects than males. There was a significant difference in contact hourly duration (min/hour) between males and females with head/face/hair. Females' hands contacted head/face/hair for more minutes per hour than males. For the median duration, there were no significant differences between males and females.

Mouthing Events. There were no significant differences in contact frequency, duration or median duration between males and females while during the game or practice.

Actions. There were no significant differences between males and females in frequency of specific actions while during the game or practice.

Macro-Activities. There was a significant difference in hourly duration (min/hour)



between males and females with duration of time spent stretching. Females stretched more minutes per hour than males during the game.

Intensity. There were no significant differences between males and females in hourly duration of player intensity while during the game.

Warming Up (Supplemental Material Section S3.4.2.2):

Hands to Object Contacts. There were no significant differences in hand contact frequency, hourly contact duration, or median contact duration between males and females while warming up.

Mouthing Events. There were no significant differences by gender in mouthing frequency, hourly contact duration and median contact duration while warming up.

Actions. There were no significant differences between males and females in frequency of specific actions while warming up.

Macro-Activities. There were no significant differences between males and females in hourly contact duration of macro-activities while warming up.

Intensity. There were no significant differences between males and females in hourly duration of player intensity while warming up.

Before/After/Break Time (Supplemental Material Section S3.4.2.3):

Hands to Object Contacts. There were significant differences ($p < 0.05$) in hand contact frequency (event/hour) between males and females with head/face/hair, water bottle and dietary objects. Females had more contacts per hour with these objects than males. There were significant differences in contact hourly duration (min/hour) between males and females with head/face/hair, water bottle, shin guards and dietary objects. Females' hands contacted head/face/hair, water bottle and dietary objects for more minutes per hour than males. However, males contacted shin guards for more minutes per hour than females. For the median duration, there were no significant differences between males and females.

Mouthing Events. There were significant differences in mouthing frequency (event/hour) between the males and females in water bottle, dietary and all objects. Females had more mouthing events per hour with these objects than males. Similarly, there were significant differences ($p < 0.05$) in mouthing hourly duration (min/hour) between males and females with water bottle, dietary and all objects. Females contacted these objects with their hands for more minutes per hour than males. Finally, there was a significant difference in mouthing median duration (second) between males and females for all objects. Females contacted all objects for longer period of time than males.

Actions. There were no significant differences between males and females in frequency of specific actions.



Macro-Activities. There were no significant differences between males and females in hourly duration of macro-activities.

Intensity. There were no significant differences between males and females in hourly duration of player intensity.

Other Combined Time (After/Before/Break/Warming Up Time) But Not During the Game (Supplemental Material Section S3.4.2.4):

Hands to Object Contacts. There were significant differences ($p < 0.05$) in hand contact frequency (event/hour) between males and females with head/face/hair, water bottle and dietary objects. Females had more contacts per hour with these objects than males. There were significant differences in contact hourly duration (min/hour) between males and females with head/face/hair, water bottle and dietary objects. Females' hands contacted head/face/hair, water bottle and dietary objects for more minutes per hour than males. For the contact median duration, there were no significant differences between males and females.

Mouthing Events. There were significant differences in mouthing frequency (event/hour) between males and females with water bottle, dietary and all objects. Females had more mouthing events per hour with these objects than males. There were significant differences ($p < 0.05$) in mouthing hourly duration (min/hour) between males and females with water bottle, dietary and all objects. Females mouthing contacts with these objects was more minutes per hour than males. There was a significant difference in mouthing median duration between males and females with all objects. Females had longer mouthing contacts with all objects than males.

Actions. There were no significant differences between males and females in frequency of specific actions.

Macro-Activities. There were no significant differences between males and females in hourly duration of macro-activities.

Intensity. There were no significant differences between males and females in hourly duration of player intensity.

On Field (Supplemental Material Section S3.4.3.1):

Hands to Object Contacts. While on the field, there were significant differences ($p < 0.05$) in contact frequency (event/hour) between males and females with head/face/hair, non-dietary objects, and all objects. Females had more contacts per hour with these objects than males. There were significant differences in contact hourly duration (min/hour) between males and females with head/face/hair, and shin guards. Females' hands contacted head/face/hair for more minutes per hour than males, while males' hands contacted shin guards for more minutes per hour than females. For the median contact duration, there were no significant differences between males and females.

Mouthing Events. There were significant differences in mouthing frequency (event/hour)



between males and females with water bottle, dietary and all objects while on the field. Females had more mouthing events per hour with these objects than males. There were significant differences ($p < 0.05$) in mouthing hourly duration (min/hour) between males and females with water bottle, dietary and all objects. Females contacted these objects with their mouth for more minutes per hour than males. There was a significant difference in mouthing median duration (second) between males and females with all objects. Females contacted all objects with their mouth for longer period of time than males while on the field.

Actions. Males tackled or slid significantly ($p < 0.05$) more times than females (event/hour) while on the field.

Macro-Activities. There was a significant difference in hourly duration (min/hour) between males and females with time spent stretching while on the field. Females stretched more minutes per hour than males on the field.

Intensity. There were no significant differences between males and females in hourly duration of player intensity while on the field.

Off field (Supplemental Material Section S3.4.3.2):

Hands to Object Contacts. There were significant differences ($p < 0.05$) in contact frequency (event/hour) between males and females with field structures, head/face/hair, water bottle, and dietary objects while off the field. Females had more contacts per hour with these objects than males. There were significant differences ($p < 0.05$) in hourly contact duration (min/hour) between males and females with field structures, head/face/hair, water bottle, and dietary objects. Females' hands contacted these objects for more minutes per hour than males. There was a significant difference in contact median duration (second) between males and females with soccer ball. Females contacted soccer ball with their hands for longer periods than males.

Mouthing Events. There were significant differences ($p < 0.05$) in mouthing frequency (event/hour) between males and females with water bottle, and dietary objects while off the field. Females had more mouthing events per hour with these objects than males. There were significant differences ($p < 0.05$) in mouthing hourly duration (min/hour) between males and females with water bottle, and dietary objects. Females contacted these objects with their mouth for more minutes per hour than males. There was a significant difference in mouthing median duration (second) between males and females with all objects. Females contacted all objects with the mouth for longer periods of time than males.

Actions. There were no significant differences between males and females in frequency of specific actions while off the field.

Macro-Activities. There were no significant differences between males and females in hourly duration of macro-activities while off the field.

Intensity. There were no significant differences between males and females in hourly



duration of player intensity while off the field.

Gloves On (Supplemental Material Section S3.4.4.1):

Hands to Object Contacts. There were significant differences ($p < 0.05$) in contact frequency (event/hour) between males and females with non-dietary objects, and all objects when participants were wearing gloves. Females had more contacts per hour with these objects than males. There were significant differences in contact hourly duration (min/hour) between males and females with all objects. Females' hands contacted these objects for more minutes per hour than males, while wearing gloves. For the median duration, there were no significant differences between males and females while wearing gloves.

Mouthing Events. There was a significant difference in mouthing frequency (event/hour) between males and females with hands. Males had more mouthing contacts per hour with their hands than females. There was significant differences ($p < 0.05$) in mouthing hourly duration (min/hour) between males and females with hands. Males contacted their mouths with their mouths for more minutes per hour than females. There were no significant differences in mouthing median duration (second) between males and females.

Gloves Off (Supplemental Material Section S3.4.4.2):

Hands to Object Contacts. There were significant differences ($p < 0.05$) in contact frequency (event/hour) between males and females with field structures, and head/face/hair when they were not wearing gloves. Females had more contacts per hour with these objects than males. There was a significant difference in contact hourly duration (min/hour) between males and females with head head/face/hair. Females' hands contacted these objects for more minutes per hour than males. For median hand contact duration, there were no significant differences between males and females while wearing gloves.

Mouthing Events. There were significant differences in mouthing frequency (event/hour) between males and females with hands. Females had more mouthing contacts per hour with hands than males while not wearing gloves. There were significant differences ($p < 0.05$) in mouthing hourly duration (min/hour) between males and females with water bottle, and dietary objects. Females contacted these objects with their mouth for more minutes per hour than males. There were no significant differences in mouthing median duration (second) between males and females.

F.3.3.6.6. Player Position (Goalie or Others)

Activity statistics by player position and results from statistical tests (Wilcoxon sum rank test) are presented in Supplemental Material Section S3.5.

Whole Event (Supplemental Material Section S3.5.1):

Hands to Object Contacts. There were no significant differences ($p < 0.05$) in contact



frequency and hourly contact duration between goalies and other positions. There was a significant difference in contact median duration (second) between goalies and other positions with head/face/hair. Goalies contacted their face/hair/head contact for longer periods of time than other players' positions.

Mouthing Events. There were no significant differences in mouthing frequency and hourly contact duration between goalies and other positions. There was a significant difference ($p < 0.05$) in mouthing median duration (second) between goalies and other positions with clothes. Goalies mouthing of clothes was for longer periods of times than other players' positions.

Actions. There were no significant differences between goalies and other positions in frequency of specific actions.

Macro-Activities. There were significant differences in hourly duration (min/hour) between goalies and other positions for the amount of time players spent stretching, sitting on the ground, or running. Goalies spent more time sitting on the ground than other players. Goalies ran and stretched for less minutes per hour than other players positions.

Intensity. There were significant differences on intensity between goalies and other positions for high, light, and resting. Goalies played more time at light and resting intensities in comparison to other players.

During the Game/Practice (Supplemental Material Section S3.5.2.1):

Hands to Object Contacts. There were significant differences ($p < 0.05$) in contact frequency (event/hour) between goalies and other positions with contact with the body while during the game/practice. Goalies had less contacts per hour with body parts than other positions. There were not significant differences in contact hourly duration (min/hour) between goalies and other positions. There was a significant difference in contact median duration (second) between goalies and other positions with face/head/hair. Goalies had longer contact with head/face/hair than other players' positions.

Mouthing Events. There were no significant differences in mouthing frequency, and hourly duration between goalies and other positions while during the game/practice. There was a significant difference in mouthing median duration (second) between goalies and other positions with clothes. Goalies had longer mouthing contact with clothes than other the players' positions.

Actions. There were no significant differences between goalies and other positions in frequency of specific actions.

Macro-Activities. There were significant differences in hourly duration (min/hour) between goalies and other positions for players in the time spent running and stretching while during the game. Goalies were running and stretching less minutes per hour than other players during the game.



Intensity. There were significant differences in hourly duration of player intensity between goalies and other positions for high, median and resting intensity. Goalies had less minutes per hour of high and median intensity than the other players' positions. However, goalies had more minutes per hour of resting intensity than other positions.

Warming Up (Supplemental Material Section S3.5.2.2):

Hands to Object Contacts. There were no significant differences in frequency, duration, or median duration between goalies and other positions while warming up.

Mouthing Events. For the mouthing frequency, duration and median duration, there were no significant differences between goalies and other positions while warming up.

Actions. There were no significant differences between goalies and other positions in frequency of specific actions while warming up.

Macro-Activities. There were no significant differences between goalies and other positions in hourly duration of macro-activities while warming up.

Intensity. There was a significant difference between goalies and other positions in hourly duration of player intensity. Goalies rested for more minutes per hour than other positions while warming up.

Before/After/Break Time (Supplemental Material Section S3.5.2.3):

Hands to Object Contacts. There were significant differences ($p < 0.05$) in contact frequency (event/hour) between goalies and other positions with field structures and dietary objects. Goalies had less contacts per hour with field structures than other positions. Also, goalies had more contacts per hour with dietary objects than other positions. There was a significant difference in contact hourly duration (min/hour) between goalies and other positions with field structures. Goalies contacted field structured in less minutes per hour than other players' positions. There were significant differences in contact median duration (second) between goalies and other positions with water bottle and dietary objects. Goalies had shorter contact with these objects than other players' positions.

Mouthing events. There were no significant differences in mouthing frequency, hourly duration, and median duration between goalies and other positions.

Actions. There were no significant differences between goalies and other positions in frequency of specific actions.

Macro-Activities. There were no significant differences between goalies and other positions in hourly duration of macro-activities.

Intensity. There were no significant differences between goalies and other positions in hourly duration of player intensity.

Other Combined Time (After/Before/Break/Warming Up Time) But During the Game (Supplemental Material Section S3.5.2.4):



Hands to Object Contacts. There were significant differences ($p < 0.05$) in contact frequency (event/hour) between goalies and other positions with field structures. Goalies had less contacts per hour with field structures than other positions. There was a significant difference in contact hourly duration (min/hour) between goalies and other positions with field structures. Goalies contacted field structured for less minutes per hour than other players' positions. There were significant differences in contact median duration (second) between goalies and other positions with field structures, head/face/hair, and dietary objects. Goalies had shorter contact with field structures and dietary objects than other players' positions. However, goalies had longer contacts with head/face/hair than other players' positions.

Mouthing Events. There were no significant differences in mouthing frequency, and hourly duration, and median duration between goalies and other positions.

Actions. There were no significant differences between goalies and other positions in frequency of specific actions.

Macro-Activities. There was a significant difference between goalies and other positions in hourly duration of players walking. Goalies had less minutes per hour walking than other players positions.

Intensity. There were no significant differences between goalies and other positions in hourly duration of player intensity.

On Field (Supplemental Material Section S3.5.3.1):

Hands to Object Contacts. There was significant difference ($p < 0.05$) in contact frequency (event/hour) between goalies and other positions with field structures while on the field. Goalies had more contacts per hour with field structures. There was a significant difference in contact hourly duration (min/hour) between goalies and other positions with field structures. Goalies contacted field structured for less minutes per hour than other players' positions. There was a significant difference in contact median duration (second) between goalies and other positions dietary objects. Goalies had shorter contacts with these objects than other players' positions.

Mouthing Events. There were no significant differences in mouthing frequency, and hourly duration between goalies and other positions while on the field. There was significant difference in mouthing median duration (second) between goalies and other positions with clothes. Goalies had longer mouthing contacts with clothes than other players' positions.

Actions. There were no significant differences between goalies and other positions in frequency of specific actions while on the field.

Macro-Activities. There were significant differences between goalies and other positions in hourly duration for time spent by players running and sitting on the ground. Goalies ran less minutes per hour than other players' positions, but goalies sat on the ground more minutes per hour than other positions.



Intensity. There were significant differences between goalies and other positions in hourly duration of player intensity while players were at high, moderate, and resting intensities. Goalies rested for more minutes per hour than other positions. However, goalies spent less time playing at high and moderate intensities than other positions.

Off field (Supplemental Material Section S3.5.3.2):

Hands to Object Contacts. There were significant differences ($p < 0.05$) in contact frequency (event/hour) between goalies and other positions with field structures and all objects while off the field. Goalies had less contacts per hour with field structures and all objects compared to other players. There was a significant difference in contact hourly duration (min/hour) between goalies and other positions with field structures. Goalies contacted field structures for less minutes per hour than other players' positions. There were no significant differences in contact median duration (second) between goalies and other positions.

Mouthing Events. There were no significant differences in mouthing frequency, hourly duration, and median duration between goalies and other positions while off the field.

Actions. There were no significant differences between goalies and other positions in frequency of specific actions while off the field.

Macro-Activities. There were no significant differences between goalies and other positions in hourly duration for each micro-activity while off the field.

Intensity. There was a significant difference between goalies and other positions in hourly duration of player intensity for time that players spent resting while off the field. Goalies rested for more minutes per hour than other positions.

Gloves On (Supplemental Material Section S3.5.4.1):

Hands to Object Contacts. There was significant differences ($p < 0.05$) in contact frequency between goalies with head/face/hair while wearing gloves. Goalies contacted hand/face/hair less times per hour than players in other positions. There were no significant differences ($p < 0.05$) in contact hourly duration between goalies and other positions while players were wearing gloves. There was a significant difference in contact median duration (second) between goalies and other positions with body. Goalies contacted body for shorter durations than other players while wearing gloves.

Mouthing Events. There were no significant differences in mouthing frequency, hourly duration, and median duration between goalies and other positions while wearing gloves.

Gloves Off (Supplemental Material Section S3.5.4.2):

Hands to Object Contacts. There was significant differences ($p < 0.05$) in contact frequency between goalies with footwear while not wearing gloves. Goalies contacted footwear less times per hour than players in other positions. There was a significant difference in contact hourly (min/hour) between goalies and other positions with non-



dietary objects and all objects. Goalies contacted these objects more minutes per hour than other positions when not wearing gloves. There were significant differences in contact median duration (second) between goalies and other positions with head/face/hair, soccer ball, non-dietary objects, and all objects. Goalies contacted these objects for longer durations than other players without gloves.

Mouthing Events. There were no significant differences in mouthing frequency, hourly duration and median duration between goalies and other positions when not wearing gloves.

F.3.3.6.7. Event Type (Game or Practice)

Activity statistics by event type and results from statistical tests (Wilcoxon sum rank test) are presented in the Supplemental Material S3.6.

Whole Event (Supplemental Material Section S3.6.1):

Hands to Object Contacts. Players had significantly more frequent contacts between their hands and clothes during a practice compared to a game. There were no significant differences ($p < 0.05$) in contact hourly duration and median duration between players playing a game or practice.

Mouthing Events. There were no significant differences in mouthing frequency and median duration between players playing a game and practice. There was a significant difference ($p < 0.05$) in mouthing hourly duration (min/hour) between game and practice with dietary objects. Players who were playing a game had more mouthing minutes per hour with dietary objects than players who played in practices.

Actions. There were no significant differences in frequency of specific actions between players who played soccer in a practice and players who played in a game.

Macro-Activities. There was a significant difference in hourly duration (min/hour) between players who played in practice and game when stretching. During practices players stretched for more minutes per hour than players who played in games.

Intensity. There were no significant differences in hourly duration of player intensity between players who played in practices and players who played in games.

During the game/practice (Supplemental Material Section S3.6.2.1):

Hands to Object Contacts. There was a significant difference ($p < 0.05$) in contact frequency (event/hour) between players playing games and practices with field structures while during the game/practice. Players who played in games had less contacts per hour with field structures than other players who played in practices. There was a significant difference in contact hourly duration (min/hour) between players in games or practices with field structures. Players who played in games had less minutes per hour with field structures than players in practices. There was a significant difference in hand contact median duration (second) between players playing games and practices with water bottle and dietary objects. Players who played in games had



longer contact with water bottle and dietary objects than players who played in practice while during the game/practice.

Mouthing Events. There was a significant difference in mouthing frequency (event/hour) between players playing in games and practices with body, while during the game and practice. Players who played in games had less mouthing events per hour with body than players who played in practices. There was a significant difference in mouthing hourly duration (min/hour) between players playing games and practices with body while during the game or practice. Players who played in games had less mouthing minutes per hour with body than players who played in practices. There were no significant differences in mouthing median duration between players playing a game and practice, while during the game or practice.

Actions. There were no significant differences in frequency of specific actions between players who played in games and players who played in practices while during the game/practice.

Macro-Activities. There were no significant differences in hourly duration of macro-activities between players who played in games and players who played in practices while during the game/practice.

Intensity. There were no significant differences in hourly duration of intensity between players who played in games and players who played in practices while during the game/practice.

Warming up (Supplemental Material Section S3.6.2.2):

Hands to Object Contacts. There were no significant differences in hourly hand contact duration, or median hand contact duration between players who played in games and players who played in practices while warming up. There was a significant difference in contact frequency (event/hour) between players who played in games and players who played in practices with body, non-dietary and all objects. Players who played in games had less hand contact events per hour with body, non-dietary and all objects than players who played in practices while warming up.

Mouthing Events. There were no significant differences in mouthing frequency, duration and median duration between players who played in games and players who played in practices while warming up.

Actions. There were no significant differences between players who played in games and players who played in practices while warming up in frequency of specific actions.

Macro-Activities. There was a significant difference between players who played in games and players who played in practices for time spent running. Players who played in games spent less minutes per hour running than players who played in practices.

Intensity. There were no significant differences in hourly duration of intensity between players who played in games and players who played in practices during warm up.



Before/after/break time (Supplemental Material Section S3.6.2.3):

Hands to Object Contacts. There was a significant difference ($p < 0.05$) in hand contact frequency (event/hour) between players playing games and practices with body and clothes. Players who played in games had less hand contacts per hour with clothes than other players who played in practices. Players who played in games had more contacts per hour with body than other players who played in practices. There was a significant difference in contact hourly duration (min/hour) between players playing games or practices with body. Players who played in games had more hand hourly contact duration with body than players who played in practices. There was a significant difference in contact median duration (second) between players playing games and practices with artificial turf, field structure, and non-dietary objects. Players who played in games had shorter contact with artificial turf, field structure, and non-dietary objects than players who played in practice.

Mouthing events. There were significant differences in mouthing frequency (event/hour) between players during games and practices with water bottle, dietary objects and all objects. Players who played in games had less mouthing events per hour with water bottle, dietary objects and all objects than players who played in practices. There were significant differences in hourly duration between players playing games and practices with water bottle, dietary objects and all objects. Players who played in games had less mouthing hourly duration with water bottle, dietary objects and all objects than other players who played in practices. There were no significant differences in mouthing median duration between players playing games and practices.

Actions. There were no significant differences in frequency of specific actions between players who played in games and players who played in practices.

Macro-Activities. There were no significant differences in hourly duration of macro-activities between players who played in games and players who played in practices.

Intensity. There was a significant difference in hourly duration between players who played in games and players who played in practices. Players who played in games had more minutes per hour of high intensity activities than players during practices.

Other Combined Time (After/Before/Break/Warming Up Time) But During the Game (Supplemental Material Section S3.6.2.4):

Hands to Object Contacts. There was a significant difference ($p < 0.05$) in hand contact frequency (event/hour) between players playing in games and practices with body, clothes, non-dietary and all objects. Players who played in games had less hand contacts per hour with clothes, non-dietary and all objects than other players during practices. Players who played in games had more contacts per hour with body than other players during practices. There was a significant difference in contact hourly duration (min/hour) between players playing games and practices with body. Players who played in games had longer hourly duration of contact with body than players during practice. There was a significant difference in contact median duration (second)



between players playing games and practices with artificial turf. Players who played in games had shorter contacts with artificial turf than players during practice.

Mouthing Events. There were significant differences in mouthing frequency (event/hour) between players playing in games and practices with water bottle, dietary objects and all objects. Players who played in games had less mouthing events per hour with water bottle, dietary objects and all objects than other players who played in practices. There were significant differences in hourly duration between players playing games and practices with water bottle, dietary objects and all objects. Players who played in games had less mouthing hourly duration with water bottle, dietary objects and all objects than other players who played in practices. There were no significant differences in mouthing median duration between players playing games and practices.

Actions. There were no significant differences in frequency of specific actions between players who played in games and players who played in practices.

Macro-Activities. There were no significant differences in hourly contact duration of macro-activities between players who played in games and players who played in practices.

Intensity. There were no significant differences in hourly duration of intensity between players who played in games and players who played in practices.

On Field (Supplemental Material Section S3.6.3.1):

Hands to Object Contacts. There was a significant difference ($p < 0.05$) in contact frequency (event/hour) between players playing games and practices with clothes, field structure, water bottle, dietary, non-dietary and all objects while on the field. Players who played in games had less contacts per hour with clothes, field structure, water bottle, non-dietary and all objects than players during practices. There were significant differences in contact hourly duration (min/hour) between players playing games and practices with field structure, water bottle, non-dietary and all objects. Players who played in games had shorter hourly hand contact duration with field structure, water bottle, non-dietary and all objects than other players who played in practices. There were no significant differences in contact median duration (second) between players playing games and practices.

Mouthing Events. There were significant differences in mouthing frequency (event/hour) between players playing games and practices with water bottle, dietary objects and all objects. Players who played in games had less mouthing events per hour with water bottle, dietary objects and all objects than other players who played in practices. There were significant differences in hourly contact duration between players during games and practices with water bottle, dietary objects, non-dietary and all objects. Players who played in games had shorter hourly mouthing duration with water bottle, dietary objects, non-dietary and all objects than other players who played in practices. Players had significantly longer median mouthing duration with all objects during games than during practices.



Actions. There were no significant differences in frequency of specific actions between players who played in games and players who played in practices while on the field.

Macro-Activities. There was a significant difference in hourly duration between players who played in games and players who played in practices. Players who played in games spent more minutes per hour stretching than players during practices while on the field.

Intensity. There were no significant differences in hourly duration of intensity between players who played in games and players who played in practices while on the field.

Off Field (Supplemental Material Section S3.6.3.2):

Hands to Object Contacts. There was a significant difference ($p < 0.05$) in contact frequency (event/hour) between players playing games and practices with body. Players who played in games had more contacts per hour with body than those during practice. There were significant differences in hand contact hourly duration (min/hour) between players playing games and practices for contacts with body and head/face/hair. Players who played in games had longer hourly duration with body and head/face/hair than players during practices. There were no significant differences in contact median duration (second) between players playing games and practices.

Mouthing Events. There were no significant differences in mouthing frequency and median duration between players during games and practices. Players during games had significantly less hourly mouthing duration with water bottle.

Actions. There were no significant differences in frequency of specific actions between players who played in games and players who played in practices while off the field.

Macro-Activities. There was a significant difference in hourly duration between players who played in games and players who played in practices. Players who played in games spent more minutes per hour stretching than players who played in practices.

Intensity. There were no significant differences in hourly duration of intensity between players who played in games and players who played in practices.

Gloves On (Supplemental Material Section S3.6.4.1):

Hands to Object Contacts. There were no significant differences in contact frequency, hourly duration and median duration between players playing games and practices with gloves on.

Mouthing Events. There were no significant differences in mouthing frequency and hourly duration between players playing games and practices with glove on. There was a significant difference in median duration between players playing games and practices with water bottle and dietary objects. Players who played in games had less mouthing time with water bottles and dietary objects than players who played in practices.

Gloves Off (Supplemental Material Section S3.6.4.2):



Hands to Object Contacts. There was a significant difference ($p < 0.05$) in contact frequency (event/hour) between players playing games and practices with non-dietary objects and all objects. Players who played in games had less contacts per hour with these objects. There were no significant differences in contact hourly duration and median duration between players playing games and practices.

Mouthing Events. There were no significant differences in mouthing frequency and median mouthing duration between players during games and practices. There was a significant difference in hourly mouthing duration between players playing games and practices with dietary objects while gloves were off. Players who played in games had less mouthing minutes per hour with dietary objects than players during practice while gloves were off.



F.3.4. References

- AuYeung, W., Canales, R.A., Beamer, P., Ferguson, A.C., and J.O. Leckie. (2004). "Young Children's Mouthing Behavior: An Observational Study via Videotaping in a Primarily Outdoor Residential Setting." *J Children's Health*, 2 (3-4), pg. 271-295
- AuYeung, W., Canales, R.A., Beamer, P., Ferguson, A.C., and J.O. Leckie. (2006). "Young children's hand contact activities: An observational study via videotaping in primarily outdoor residential settings." *J Exp Sci Environ Epidemiol*, 16: 434-446.
- Beamer, P., Key, M.E., Ferguson, A.C., Canales, R.A., AuYeung, W., and J.O. Leckie. (2008) "Quantified Activity Pattern Data from 6-to-27-Month-Old Farmworker Children for Use in Exposure Assessment." *Environ Res*, 108: 239-246. PMID: 18723168.
- Beamer, P., Luik, C.E., Canales, R.A., Leckie, J.O. "Quantified Outdoor Micro-Activity Data for Children Aged 7-12 Years Old." *J Exp Sci Environ Epidemiol*, 22:82-92. PMID: 21989500.
- Ferguson, A.C., Canales, R.A., Beamer, P., AuYeung, W., Key, M., Munninghoff, A., Lee, K.T., Robertson, A., and J.O. Leckie. (2006) "Video methods in the quantification of children's exposures." *J Exp Anal Environ Epidemiol*, 16: 287-298.
- Hubal, E., Sheldon, L., Burke, J., McCurdy, T., Berry, M., Rigas, M., Zartarian, V. and Freeman, N. (2000). Children's Exposure Assessment: A Review of Factors Influencing Children's Exposure, and the Data Available to Characterize and Assess That Exposure. *Environmental Health Perspectives*, 108(6), p.475.
- Juberg D, Alfano K, Coughlin R, Thompson K. (2001). An Observational Study of Object Mouthing Behavior by Young Children. *Pediatrics*. 2001;107:135–142.
- RStudio Team (2015). RStudio: Integrated Development for R. RStudio, Inc., Boston, MA URL <http://www.rstudio.com/>.
- Zartarian V, Ferguson A, Ong C, Leckie J. (1997) Quantifying Videotaped Activity Patterns: Video Translation Software and Training Methodologies. *J Expo Anal Environ Epidemiol*: 7:535–542.



F.4. Adjustment Factors for Inhalation Exposure (AF_{inh}) and Inhalation Exposure Concentration (C_{inh})

This section presents:

- the acute inhalation exposure concentration (Acute C_{inh}) for exposure to chemicals with available acute inhalation toxicity criteria (Acute TC_{inh}).
- the adjustment factors of chronic inhalation exposure to general chemicals (AF_{inh}) and one-day inhalation exposure to developmental and reproductive toxicants (DARTs, $AF_{inh-DART}$).
- the one-day inhalation exposure concentration of DARTs ($C_{inh-DART}$), the chronic inhalation exposure concentration of general chemicals (Chronic C_{inh}) and sensory irritants (Chronic $C_{inh-sensory}$) for chemicals with available inhalation toxicity criteria (DART TC_{inh} , TC_{inh} , and Sensory TC_{inh}).

Acute C_{inh} values are used to assess acute non-cancer hazard, while $C_{inh-DART}$, Chronic C_{inh} , and Chronic $C_{inh-sensory}$ values are used to assess one-day or chronic non-cancer hazards from inhalation exposure to the three groups of chemicals on synthetic turf fields. The receptor- and age-specific $AF_{inh-DART}$ addresses a less than 24 hours per day exposure of DARTs at the fields, whereas the receptor- and age-specific AF_{inh} addresses a less than 365 days per year and a less than 24 hours per day exposure of general chemicals at synthetic turf fields. These adjustment factors are used to derive the values of $C_{inh-DART}$ and Chronic C_{inh} , respectively, by modifying the field-specific average concentrations ($C_{air-field}$) of DARTs detected in air on each synthetic turf field and the mean of the 35 individual field average concentrations ($C_{air-avg}$) of general chemicals. No adjustment factor is applied to modify the $C_{air-avg}$ in the calculation of Chronic $C_{inh-sensory}$. Details of how to calculate the $AF_{inh-DART}$, AF_{inh} , $C_{inh-DART}$, Chronic C_{inh} , and Chronic $C_{inh-sensory}$ are presented in Main Report Section 5.4 and Appendix B Section B.3.4. An example calculation is presented in Appendix Section G.1.

F.4.1. Acute Inhalation Exposure Concentration (Acute C_{inh}) for Non-Cancer Hazard Assessment of Chemicals with Available Acute Inhalation Toxicity Criteria

Table F-96. Acute Inhalation Exposure Concentration^a (Acute C_{inh} , nanograms per cubic meter) for Exposure to Chemicals^b with Available Acute Inhalation Toxicity Criteria—Combined Gender and **All Receptor Categories**

Chemical ^a	CASRN	Acute C _{inh} ^b	
		On Field	Off Field
Field-Related Chemical			
Styrene	100-42-5	1600	1400
Non-Field-Related Chemicals			



Chemical ^a	CASRN	Acute C _{inh} ^b	
		On Field	Off Field
Acetaldehyde	75-07-0	9700	not sampled
Benzene	71-43-2	3700	3500
2-Butanone	78-93-3	1900	not sampled
2-Butoxyethanol	111-76-2	1000	480
Formaldehyde	50-00-0	17000	not sampled
Phenol	108-95-2	210	210
Tetrachloroethylene	127-18-4	420	420
Toluene	108-88-3	12000	10000
m/p-Xylene	106-42-3	5200	5200
o-Xylene	95-47-6	2100	2000

^a Only chemicals with available acute toxicity criteria for inhalation are shown in the table.

^b On-field Acute C_{inh} equals the maximum concentration of a chemical detected in air on the 35 synthetic turf fields. Off-field Acute C_{inh} equals the maximum concentration of a chemical detected in air off the 35 synthetic turf fields.

^c Selected carbonyls, collected using 2,4-dinitrophenylhydrazine-cartiridges, were not sampled at off the field locations during the statewide sample collections.

CASRN: Chemical Abstracts Service Registry Number. Values are rounded to two significant figures.

F.4.2. Adjustment Factor for One-Day Inhalation Exposure for Non-Cancer Hazard Assessment of DARTs (AF_{inh-DART}).

Table F-97. Values of Adjustment Factor of One-Day Inhalation Exposure to DARTs (AF_{inh-DART}, unitless)—Combined Gender and **All Receptor Categories**

Age Group, years	AF _{inh-DART}			
	Athletes	Coaches	Referees	Spectators
Third Trimester Fetus	Not applicable	Not applicable	Not applicable	0.049
0<2	Not applicable	Not applicable	Not applicable	0.40
2<6	0.73	Not applicable	Not applicable	0.21
6<11	0.55	Not applicable	Not applicable	0.20
11<16	0.46	Not applicable	Not applicable	0.086
16<30	0.68	0.29	0.27	0.057
30<40	0.45	0.25	0.23	0.050
40<50	0.45	0.25	0.24	0.050
50<70	0.49	0.25	0.24	0.050

Values are rounded to two significant figures.



Table F-98. Values of Exposure Factors Used to Calculate Adjustment Factors of One-Day Inhalation Exposure to DARTs ($AF_{inh-DART}$, unitless)—Combined Gender **Athletes**

Age Group, years	Practice BR_{TW} , cubic meters per hour	Practice ET, hours per event	Game BR_{TW} , cubic meters per hour	Game ET, hours per event	V_{event} , cubic meters per event	BW, kilograms	$AF_{inh-DART}$
2<6	1.4	1.5	1.5	4.3	4.3	20.5	0.73
6<11	2.3	2.1	2.3	2.3	5.1	32.0	0.55
11<16	2.8	2.3	2.9	2.2	6.4	48.7	0.46
16<30	4.2	2.8	4.4	3.1	12.7	65.8	0.68
30<40	3.6	3.1	4.0	2.1	9.8	75.3	0.45
40<50	3.9	2.5	4.0	2.4	9.7	74.5	0.45
50<70	4.1	2.3	4.4	2.6	10.4	74.3	0.49

$AF_{inh-DART}$: adjustment factor of chronic exposure to a reproductive or developmental toxicant for a receptor category in an age group, values are rounded to two significant figures; BR_{TW} : time-weighted one-hour breathing rate of a receptor category in an age group for an event type, values are rounded to one decimal place; ET: mean of the maximum time spent on field per event of a receptor category in an age group for an event type, values are rounded to one decimal place; V_{event} : average volume of air a receptor inhales during a soccer event, values are rounded to one decimal place; and BW: receptor category and age-specific bodyweight, values are rounded to one decimal place.

Table F-99. Values of Exposure Factors Used to Calculate Adjustment Factors of One-Day Inhalation Exposure to DARTs ($AF_{inh-DART}$, unitless)—Combined Gender **Coaches**

Age Group, years	Practice BR_{TW} , cubic meters per hour	Practice ET, hours per event	Game BR_{TW} , cubic meters per hour	Game ET, hours per event	V_{event} , cubic meters per event	BW, kilograms	$AF_{inh-DART}$
16<30	1.9	3.1	1.6	3.1	5.4	65.8	0.29
30<40	1.9	3.1	1.6	3.1	5.4	75.3	0.25
40<50	1.9	3.1	1.6	3.1	5.4	74.5	0.25
50<70	1.9	3.1	1.6	3.1	5.4	74.3	0.25

$AF_{inh-DART}$: adjustment factor of chronic exposure to a reproductive or developmental toxicant for a receptor category in an age group, values are rounded to two significant figures; BR_{TW} : time-weighted one-hour breathing rate of a receptor category in an age group for an event type, values are rounded to one decimal place; ET: mean of the maximum time spent on field per event of a receptor category in an age group for an event type, values are rounded to one decimal place; V_{event} : average volume of air a receptor inhales during a soccer event, values are rounded to one decimal place; and BW: receptor category and age-specific bodyweight, values are rounded to one decimal place.

Table F-100. Values of Exposure Factors and Adjustment Factor of One-Day Inhalation Exposure to DARTs ($AF_{inh-DART}$, unitless)—Combined Gender **Referees**

Age Group, years	Practice BR_{TW} , cubic meters per hour	Practice ET, hours per event	Game BR_{TW} , cubic meters per hour	Game ET, hours per event	V_{event} , cubic meters per event	BW, kilograms	$AF_{inh-DART}$
16<30	Not applicable	Not applicable	1.6	3.1	5.0	65.8	0.27
30<40	Not applicable	Not applicable	1.6	3.1	5.0	75.3	0.23
40<50	Not applicable	Not applicable	1.6	3.1	5.0	74.5	0.24
50<70	Not applicable	Not applicable	1.6	3.1	5.0	74.3	0.24

$AF_{inh-DART}$: adjustment factor of chronic exposure to a reproductive or developmental toxicant for a



receptor category in an age group, values are rounded to two significant figures; BR_{TW} : time-weighted one-hour breathing rate of a receptor category in an age group for an event type, values are rounded to one decimal place; ET: mean of the maximum time spent on field per event of a receptor category in an age group for an event type, values are rounded to one decimal place; V_{event} : average volume of air a receptor inhales during a soccer event, values are rounded to one decimal place; and BW: receptor category and age-specific bodyweight, values are rounded to one decimal place.

Table F-101. Values of Exposure Factors and Adjustment Factor of One-Day Inhalation Exposure to DARTs ($AF_{inh-DART}$, unitless)—Combined Gender **Spectators**

Age Group, years	Practice BR_{TW} , cubic meters per hour	Practice ET, hours per event	Game BR_{TW} , cubic meters per hour	Game ET, hours per event	V_{event} , cubic meters per event	BW, kilograms	$AF_{inh-DART}$
Third Trimester Fetus	0.4	2.1	0.4	3.1	1.1	75.6	0.049
0<2	0.4	2.1	0.4	3.1	1.1	9.4	0.40
2<6	0.5	2.1	0.5	3.1	1.2	20.5	0.21
6<11	0.7	2.1	0.7	3.1	1.8	32.0	0.20
11<16	0.5	2.1	0.5	3.1	1.2	48.7	0.086
16<30	0.4	2.1	0.4	3.1	1.1	65.8	0.057
30<40	0.4	2.1	0.4	3.1	1.1	75.3	0.050
40<50	0.4	2.1	0.4	3.1	1.1	74.5	0.050
50<70	0.4	2.1	0.4	3.1	1.1	74.3	0.050

$AF_{inh-DART}$: adjustment factor of chronic exposure to a reproductive or developmental toxicant for a receptor category in an age group, values are rounded to two significant figures; BR_{TW} : time-weighted one-hour breathing rate of a receptor category in an age group for an event type, values are rounded to one decimal place; ET: mean of the maximum time spent on field per event of a receptor category in an age group for an event type, values are rounded to one decimal place; V_{event} : average volume of air a receptor inhales during a soccer event, values are rounded to one decimal place; and BW: receptor category and age-specific bodyweight, values are rounded to one decimal place.

F.4.3. Adjustment Factor for Chronic Inhalation Exposure for Non-Cancer Hazard Assessment of General Chemicals (AF_{inh})

Table F-102. Values of Adjustment Factor of Chronic Inhalation Exposure to General Chemicals (AF_{inh} , unitless) for Each Receptor Category—Combined Gender and **All Receptor Categories**

Age Group, years	AF_{inh}			
	Athletes	Coaches	Referees	Spectators
Third Trimester Fetus	Not applicable	Not applicable	Not applicable	0.021
0<2	Not applicable	Not applicable	Not applicable	0.17
2<6	0.16	Not applicable	Not applicable	0.087
6<11	0.20	Not applicable	Not applicable	0.083
11<16	0.20	Not applicable	Not applicable	0.036
16<30	0.37	0.15	0.057	0.024
30<40	0.24	0.13	0.050	0.021
40<50	0.24	0.14	0.050	0.021



Age Group, years	AF _{inh}			
	Athletes	Coaches	Referees	Spectators
50<70	0.23	0.14	0.050	0.021

Values are rounded to two significant figures.

Table F-103. Values of Mean Exposure Parameters and Calculated Adjustment Factor of Chronic Inhalation Exposure to General Chemicals (AF_{inh}, unitless)—Combined Gender **Athletes**

Age Group, years	Practice BR _{TW} , cubic meters per hour	Practice ET, hours per event	Game BR _{TW} , cubic meters per hour	Game ET, hours per event	V _{event} , cubic meters per event	BW, kilograms
2<6	1.4	126	1.5	104	20.5	0.16
6<11	2.3	156	2.3	137	32.0	0.20
11<16	2.8	223	2.9	129	48.7	0.20
16<30	4.2	354	4.4	241	65.8	0.37
30<40	3.6	343	4.0	167	75.3	0.24
40<50	3.9	256	4.0	214	74.5	0.24
50<70	4.1	187	4.4	239	74.3	0.23

AET: mean annual event time spent on the field of a receptor category in an age group for an event type, values are rounded to the nearest whole integers; AF_{inh}: adjustment factor of chronic exposure to a chemical for a receptor category in an age group, values are rounded to two significant figures; BR_{TW}: mean time-weighted one-hour breathing rate of a receptor category in an age group for an event type, values are rounded to one decimal place; and BW: mean study-specific bodyweight of an age group, values are rounded to two decimal places.

Table F-104. Values of Mean Exposure Factors and Calculated Adjustment Factor of Chronic Inhalation Exposure to General Chemicals (AF_{inh}, unitless)—Combined Gender **Coaches**

Age Group, years	Practice BR _{TW} , cubic meters per hour	Practice AET, hours per year	Game BR _{TW} , cubic meters per hour	Game AET, hours per year	BW, kilograms	AF _{inh}
16<30	1.9	354	1.6	241	65.8	0.15
30<40	1.9	354	1.6	241	75.3	0.13
40<50	1.9	354	1.6	241	74.5	0.14
50<70	1.9	354	1.6	241	74.3	0.14

AET: mean annual event time spent on the field of a receptor category in an age group for an event type, values are rounded to the nearest whole integers; AF_{inh}: adjustment factor of chronic exposure to a chemical for a receptor category in an age group, values are rounded to two significant figures; BR_{TW}: mean time-weighted one-hour breathing rate of a receptor category in an age group for an event type, values are rounded to one decimal place; and BW: mean study-specific bodyweight of an age group, values are rounded to one decimal place.



Table F-105. Values of Mean Exposure Parameters and Calculated Adjustment Factor of Chronic Inhalation Exposure to General Chemicals (AF_{inh} , unitless)—Combined Gender **Referees**

Age Group, years	Practice BR_{TW} , cubic meters per hour	Practice AET, hours per year	Game BR_{TW} , cubic meters per hour	Game AET, hours per year	BW, kilograms	AF_{inh}
16<30	Not applicable	Not applicable	1.6	241	65.8	0.057
30<40	Not applicable	Not applicable	1.6	241	75.3	0.050
40<50	Not applicable	Not applicable	1.6	241	74.5	0.050
50<70	Not applicable	Not applicable	1.6	241	74.3	0.050

AET: mean annual event time spent on the field of a receptor category in an age group for an event type, values are rounded to the nearest whole integers; AF_{inh} : adjustment factor of chronic exposure to a chemical for a receptor category in an age group, values are rounded to two significant figures; BR_{TW} : mean time-weighted one-hour breathing rate of a receptor category in an age group for an event type, values are rounded to one decimal place; and BW: mean study-specific bodyweight of an age group, values are rounded to one decimal place.

Table F-106. Values of Mean Exposure Factors and Calculated Adjustment Factor of Chronic Inhalation Exposure to General Chemicals (AF_{inh} , unitless)—Combined Gender **Spectators**

Age Group, years	Practice BR_{TW} , cubic meters per hour	Practice AET, hours per year	Game BR_{TW} , cubic meters per hour	Game AET, hours per year	BW, kilograms	AF_{inh}
3 rd Trimester Fetus	0.4	156	0.4	241	75.6	0.021
0<2	0.4	156	0.4	241	9.4	0.17
2<6	0.5	156	0.5	241	20.5	0.087
6<11	0.7	156	0.7	241	32.0	0.083
11<16	0.5	156	0.5	241	48.7	0.036
16<30	0.4	156	0.4	241	65.8	0.024
30<40	0.4	156	0.4	241	75.3	0.021
40<50	0.4	156	0.4	241	74.5	0.021
50<70	0.4	156	0.4	241	74.3	0.021

AET: mean annual event time spent on the field of a receptor category in an age group for an event type, values are rounded to the nearest whole integers; AF_{inh} : adjustment factor of chronic exposure to a chemical for a receptor category in an age group, values are rounded to two significant figures; BR_{TW} : mean time-weighted one-hour breathing rate of a receptor category in an age group for an event type, values are rounded to one decimal places; and BW: mean study-specific bodyweight of an age group, values are rounded to one decimal place.

F.4.4. One-Day Field-Specific Inhalation Exposure Concentration for DARTs ($C_{inh-DART-field}$)

INDIVIDUAL FIELD ASSESSMENT (Table F-107 to Table F-139)



Table F-107. **On-Field One-Day Field-Specific^a Inhalation Exposure Concentration^b**
($C_{inh-DART-field}$, nanograms per cubic meter) for **DARTs—Combined Gender Athletes 2<6 Years**

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	420	300	330	890	1000
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.83	0.88	0.45	2.2	3.4
Benzo[e]pyrene	192-97-2	6	0	0.014	0.031	0	0.093	0.093
Benzo[g,h,i]perylene	191-24-2	19	0	0.095	0.14	0.036	0.38	0.55
n-Caproic acid vinyl ester	3050-69-9	1	0	4.9	29	0	0	170
Chrysene	218-01-9	13	0	0.15	0.23	0	0.56	0.96
Coronene	191-07-1	10	0	0.062	0.12	0	0.22	0.53
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.25	0.42	0.17	0.75	2.2
Cyclohexylamine	108-91-8	4	0	1.7	4.9	0	11	23
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.053	0.064	0.02	0.15	0.3
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.24	0.34	0.18	1	1.5
Dimethyl phthalate	131-11-3	4	0	2.6	8.7	0	18	45
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	16	92	0	0	540
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.078	0.26	0	0.89	0.89
Methyl stearate	112-61-8	10	0	4	8.5	0	27	32
4-tert-Octylphenol	140-66-9	10	0	1.3	2.8	0	7.5	10
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.013	0.053	0	0.077	0.22
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.32	0.87	0	1.5	4.6

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-108. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender **Athletes**
6<11 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	320	230	250	670	780
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.63	0.66	0.34	1.7	2.6
Benzo[e]pyrene	192-97-2	6	0	0.01	0.024	0	0.07	0.07
Benzo[g,h,i]perylene	191-24-2	19	0	0.072	0.11	0.028	0.29	0.42
n-Caproic acid vinyl ester	3050-69-9	1	0	3.8	22	0	0	130
Chrysene	218-01-9	13	0	0.11	0.18	0	0.42	0.73
Coronene	191-07-1	10	0	0.047	0.089	0	0.17	0.4
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.19	0.32	0.13	0.57	1.7
Cyclohexylamine	108-91-8	4	0	1.3	3.7	0	8.5	17
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.04	0.049	0.015	0.12	0.23
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.18	0.26	0.14	0.77	1.1
Dimethyl phthalate	131-11-3	4	0	2.0	6.6	0	13	34
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	12	70	0	0	410
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.059	0.19	0	0.67	0.67
Methyl stearate	112-61-8	10	0	3.1	6.5	0	21	25
4-tert-Octylphenol	140-66-9	10	0	1.0	2.1	0	5.7	7.7
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.0098	0.04	0	0.058	0.17
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.24	0.66	0	1.1	3.5

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-109. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender **Athletes**
11<16 Years

Chemical	CASRN	Detection	C _{inh} -DART					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	260	190	210	560	650
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.52	0.55	0.28	1.4	2.2
Benzo[e]pyrene	192-97-2	6	0	0.0086	0.02	0	0.059	0.059
Benzo[g,h,i]perylene	191-24-2	19	0	0.06	0.089	0.023	0.24	0.35
n-Caproic acid vinyl ester	3050-69-9	1	0	3.1	18	0	0	110
Chrysene	218-01-9	13	0	0.094	0.15	0	0.35	0.61
Coronene	191-07-1	10	0	0.039	0.074	0	0.14	0.34
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.16	0.26	0.11	0.47	1.4
Cyclohexylamine	108-91-8	4	0	1	3.1	0	7.1	14
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.033	0.041	0.012	0.097	0.19
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.15	0.22	0.12	0.64	0.92
Dimethyl phthalate	131-11-3	4	0	1.7	5.5	0	11	29
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	10	58	0	0	340
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.05	0.16	0	0.56	0.56
Methyl stearate	112-61-8	10	0	2.6	5.4	0	17	20
4-tert-Octylphenol	140-66-9	10	0	0.8	1.7	0	4.7	6.4
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.0082	0.033	0	0.049	0.14
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.2	0.55	0	0.95	2.9

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-110. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender **Athletes**
16<30 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	390	280	310	820	950
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.77	0.81	0.41	2.1	3.2
Benzo[e]pyrene	192-97-2	6	0	0.013	0.029	0	0.086	0.086
Benzo[g,h,i]perylene	191-24-2	19	0	0.088	0.13	0.034	0.35	0.51
n-Caproic acid vinyl ester	3050-69-9	1	0	4.6	27	0	0	160
Chrysene	218-01-9	13	0	0.14	0.21	0	0.52	0.89
Coronene	191-07-1	10	0	0.058	0.11	0	0.2	0.49
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.23	0.38	0.16	0.69	2.1
Cyclohexylamine	108-91-8	4	0	1.5	4.5	0	10	21
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.049	0.06	0.018	0.14	0.28
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.22	0.32	0.17	0.93	1.3
Dimethyl phthalate	131-11-3	4	0	2.4	8.1	0	16	42
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	15	85	0	0	500
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.073	0.24	0	0.82	0.82
Methyl stearate	112-61-8	10	0	3.7	7.9	0	25	30
4-tert-Octylphenol	140-66-9	10	0	1.2	2.6	0	7	9.5
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.012	0.049	0	0.071	0.2
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.3	0.8	0	1.4	4.2

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-field-DART}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-111. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender **Athletes**
30<40 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	260	190	210	550	640
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.52	0.55	0.28	1.4	2.1
Benzo[e]pyrene	192-97-2	6	0	0.0085	0.02	0	0.058	0.058
Benzo[g,h,i]perylene	191-24-2	19	0	0.059	0.088	0.023	0.23	0.34
n-Caproic acid vinyl ester	3050-69-9	1	0	3.1	18	0	0	100
Chrysene	218-01-9	13	0	0.093	0.14	0	0.35	0.6
Coronene	191-07-1	10	0	0.039	0.073	0	0.14	0.33
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.15	0.26	0.1	0.47	1.4
Cyclohexylamine	108-91-8	4	0	1	3.1	0	7	14
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.033	0.04	0.012	0.095	0.19
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.15	0.21	0.11	0.63	0.9
Dimethyl phthalate	131-11-3	4	0	1.6	5.4	0	11	28
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	9.8	57	0	0	330
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.049	0.16	0	0.55	0.55
Methyl stearate	112-61-8	10	0	2.5	5.3	0	17	20
4-tert-Octylphenol	140-66-9	10	0	0.8	1.7	0	4.7	6.4
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.0081	0.033	0	0.048	0.14
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.2	0.54	0	0.94	2.9

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-112. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender **Athletes**
40<50 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	260	190	210	550	640
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.52	0.55	0.28	1.4	2.1
Benzo[e]pyrene	192-97-2	6	0	0.0085	0.02	0	0.058	0.058
Benzo[g,h,i]perylene	191-24-2	19	0	0.059	0.088	0.023	0.23	0.34
n-Caproic acid vinyl ester	3050-69-9	1	0	3.1	18	0	0	100
Chrysene	218-01-9	13	0	0.093	0.14	0	0.35	0.6
Coronene	191-07-1	10	0	0.039	0.073	0	0.14	0.33
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.15	0.26	0.1	0.47	1.4
Cyclohexylamine	108-91-8	4	0	1	3.1	0	7	14
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.033	0.04	0.012	0.095	0.19
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.15	0.21	0.11	0.63	0.9
Dimethyl phthalate	131-11-3	4	0	1.6	5.4	0	11	28
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	9.8	57	0	0	330
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.049	0.16	0	0.55	0.55
Methyl stearate	112-61-8	10	0	2.5	5.3	0	17	20
4-tert-Octylphenol	140-66-9	10	0	0.8	1.7	0	4.7	6.4
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.0081	0.033	0	0.048	0.14
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.2	0.54	0	0.94	2.9

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-113. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender **Athletes**
50<70 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	280	200	220	600	690
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.56	0.59	0.3	1.5	2.3
Benzo[e]pyrene	192-97-2	6	0	0.0092	0.021	0	0.062	0.062
Benzo[g,h,i]perylene	191-24-2	19	0	0.064	0.095	0.024	0.25	0.37
n-Caproic acid vinyl ester	3050-69-9	1	0	3.3	19	0	0	110
Chrysene	218-01-9	13	0	0.1	0.16	0	0.38	0.65
Coronene	191-07-1	10	0	0.042	0.079	0	0.15	0.36
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.17	0.28	0.11	0.5	1.5
Cyclohexylamine	108-91-8	4	0	1.1	3.3	0	7.6	15
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.036	0.043	0.013	0.1	0.2
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.16	0.23	0.12	0.68	0.98
Dimethyl phthalate	131-11-3	4	0	1.8	5.9	0	12	30
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	11	62	0	0	360
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.053	0.17	0	0.6	0.6
Methyl stearate	112-61-8	10	0	2.7	5.7	0	18	22
4-tert-Octylphenol	140-66-9	10	0	0.9	1.9	0	5.1	6.9
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.0087	0.035	0	0.052	0.15
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.22	0.58	0	1	3.1

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-114. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender **Coaches**
16<30 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	170	120	130	350	410
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.33	0.35	0.18	0.88	1.4
Benzo[e]pyrene	192-97-2	6	0	0.0054	0.012	0	0.037	0.037
Benzo[g,h,i]perylene	191-24-2	19	0	0.037	0.055	0.014	0.15	0.22
n-Caproic acid vinyl ester	3050-69-9	1	0	1.9	11	0	0	66
Chrysene	218-01-9	13	0	0.059	0.091	0	0.22	0.38
Coronene	191-07-1	10	0	0.025	0.046	0	0.087	0.21
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.097	0.16	0.066	0.29	0.88
Cyclohexylamine	108-91-8	4	0	0.65	1.9	0	4.4	8.9
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.021	0.025	0.0078	0.06	0.12
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.096	0.13	0.072	0.4	0.57
Dimethyl phthalate	131-11-3	4	0	1.0	3.4	0	7	18
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	6.2	36	0	0	210
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.031	0.1	0	0.35	0.35
Methyl stearate	112-61-8	10	0	1.6	3.4	0	11	13
4-tert-Octylphenol	140-66-9	10	0	0.5	1.1	0	3	4
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.0051	0.021	0	0.03	0.087
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.13	0.34	0	0.59	1.8

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-115. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender **Coaches**
30<40 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	140	100	110	310	350
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.29	0.3	0.15	0.77	1.2
Benzo[e]pyrene	192-97-2	6	0	0.0047	0.011	0	0.032	0.032
Benzo[g,h,i]perylene	191-24-2	19	0	0.033	0.048	0.013	0.13	0.19
n-Caproic acid vinyl ester	3050-69-9	1	0	1.7	9.9	0	0	58
Chrysene	218-01-9	13	0	0.051	0.08	0	0.19	0.33
Coronene	191-07-1	10	0	0.021	0.04	0	0.076	0.18
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.085	0.14	0.058	0.26	0.77
Cyclohexylamine	108-91-8	4	0	0.57	1.7	0	3.9	7.8
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.018	0.022	0.0068	0.053	0.1
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.084	0.12	0.063	0.35	0.5
Dimethyl phthalate	131-11-3	4	0	0.9	3	0	6.1	16
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	5.4	32	0	0	180
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.027	0.088	0	0.31	0.31
Methyl stearate	112-61-8	10	0	1.4	2.9	0	9.4	11
4-tert-Octylphenol	140-66-9	10	0	0.5	0.95	0	2.6	3.5
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.0045	0.018	0	0.027	0.076
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.11	0.3	0	0.52	1.6

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-116. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender **Coaches**
40<50 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	150	100	120	310	360
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.29	0.3	0.15	0.78	1.2
Benzo[e]pyrene	192-97-2	6	0	0.0048	0.011	0	0.032	0.032
Benzo[g,h,i]perylene	191-24-2	19	0	0.033	0.049	0.013	0.13	0.19
n-Caproic acid vinyl ester	3050-69-9	1	0	1.7	10	0	0	59
Chrysene	218-01-9	13	0	0.052	0.081	0	0.2	0.34
Coronene	191-07-1	10	0	0.022	0.041	0	0.076	0.19
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.086	0.14	0.059	0.26	0.78
Cyclohexylamine	108-91-8	4	0	0.58	1.7	0	3.9	7.8
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.018	0.022	0.0069	0.053	0.1
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.085	0.12	0.064	0.35	0.51
Dimethyl phthalate	131-11-3	4	0	0.9	3	0	6.2	16
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	5.5	32	0	0	190
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.027	0.089	0	0.31	0.31
Methyl stearate	112-61-8	10	0	1.4	3	0	9.5	11
4-tert-Octylphenol	140-66-9	10	0	0.5	0.96	0	2.6	3.6
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.0045	0.018	0	0.027	0.077
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.11	0.3	0	0.52	1.6

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-117. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender **Coaches**
50<70 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	150	110	120	310	360
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.29	0.31	0.16	0.78	1.2
Benzo[e]pyrene	192-97-2	6	0	0.0048	0.011	0	0.032	0.032
Benzo[g,h,i]perylene	191-24-2	19	0	0.033	0.049	0.013	0.13	0.19
n-Caproic acid vinyl ester	3050-69-9	1	0	1.7	10	0	0	59
Chrysene	218-01-9	13	0	0.052	0.081	0	0.2	0.34
Coronene	191-07-1	10	0	0.022	0.041	0	0.077	0.19
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.086	0.15	0.059	0.26	0.78
Cyclohexylamine	108-91-8	4	0	0.58	1.7	0	3.9	7.9
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.018	0.022	0.0069	0.054	0.11
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.085	0.12	0.064	0.35	0.51
Dimethyl phthalate	131-11-3	4	0	0.9	3	0	6.2	16
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	5.5	32	0	0	190
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.027	0.089	0	0.31	0.31
Methyl stearate	112-61-8	10	0	1.4	3	0	9.5	11
4-tert-Octylphenol	140-66-9	10	0	0.5	0.97	0	2.6	3.6
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.0045	0.018	0	0.027	0.077
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.11	0.3	0	0.52	1.6

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-118. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender **Referees**
16<30 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	150	110	120	320	380
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.3	0.32	0.16	0.82	1.3
Benzo[e]pyrene	192-97-2	6	0	0.005	0.011	0	0.034	0.034
Benzo[g,h,i]perylene	191-24-2	19	0	0.035	0.052	0.013	0.14	0.2
n-Caproic acid vinyl ester	3050-69-9	1	0	1.8	11	0	0	62
Chrysene	218-01-9	13	0	0.055	0.085	0	0.21	0.35
Coronene	191-07-1	10	0	0.023	0.043	0	0.08	0.19
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.09	0.15	0.062	0.27	0.82
Cyclohexylamine	108-91-8	4	0	0.61	1.8	0	4.1	8.3
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.019	0.024	0.0072	0.056	0.11
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.089	0.13	0.067	0.37	0.53
Dimethyl phthalate	131-11-3	4	0	1.0	3.2	0	6.5	17
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	5.8	34	0	0	200
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.029	0.094	0	0.33	0.33
Methyl stearate	112-61-8	10	0	1.5	3.1	0	9.9	12
4-tert-Octylphenol	140-66-9	10	0	0.5	1	0	2.7	3.7
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.0047	0.019	0	0.028	0.081
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.12	0.32	0	0.55	1.7

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-119. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender **Referees**
30<40 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	130	96	110	280	330
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.27	0.28	0.14	0.72	1.1
Benzo[e]pyrene	192-97-2	6	0	0.0044	0.01	0	0.03	0.03
Benzo[g,h,i]perylene	191-24-2	19	0	0.03	0.045	0.012	0.12	0.18
n-Caproic acid vinyl ester	3050-69-9	1	0	1.6	9.2	0	0	54
Chrysene	218-01-9	13	0	0.048	0.074	0	0.18	0.31
Coronene	191-07-1	10	0	0.02	0.038	0	0.07	0.17
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.079	0.13	0.054	0.24	0.72
Cyclohexylamine	108-91-8	4	0	0.53	1.6	0	3.6	7.2
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.017	0.021	0.0063	0.049	0.096
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.078	0.11	0.059	0.32	0.46
Dimethyl phthalate	131-11-3	4	0	0.8	2.8	0	5.7	14
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	5	29	0	0	170
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.025	0.082	0	0.28	0.28
Methyl stearate	112-61-8	10	0	1.3	2.7	0	8.7	10
4-tert-Octylphenol	140-66-9	10	0	0.4	0.89	0	2.4	3.3
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.0041	0.017	0	0.025	0.07
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.1	0.28	0	0.48	1.5

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-120. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender **Referees**
40<50 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	140	97	110	290	330
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.27	0.28	0.14	0.72	1.1
Benzo[e]pyrene	192-97-2	6	0	0.0044	0.01	0	0.03	0.03
Benzo[g,h,i]perylene	191-24-2	19	0	0.031	0.045	0.012	0.12	0.18
n-Caproic acid vinyl ester	3050-69-9	1	0	1.6	9.3	0	0	54
Chrysene	218-01-9	13	0	0.048	0.075	0	0.18	0.31
Coronene	191-07-1	10	0	0.02	0.038	0	0.071	0.17
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.08	0.13	0.054	0.24	0.73
Cyclohexylamine	108-91-8	4	0	0.54	1.6	0	3.6	7.3
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.017	0.021	0.0064	0.05	0.097
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.078	0.11	0.059	0.33	0.47
Dimethyl phthalate	131-11-3	4	0	0.9	2.8	0	5.7	15
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	5.1	30	0	0	170
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.025	0.083	0	0.29	0.29
Methyl stearate	112-61-8	10	0	1.3	2.8	0	8.8	10
4-tert-Octylphenol	140-66-9	10	0	0.4	0.9	0	2.4	3.3
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.0042	0.017	0	0.025	0.071
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.1	0.28	0	0.49	1.5

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-121. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender **Referees**
50<70 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	140	98	110	290	330
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.27	0.28	0.14	0.73	1.1
Benzo[e]pyrene	192-97-2	6	0	0.0044	0.01	0	0.03	0.03
Benzo[g,h,i]perylene	191-24-2	19	0	0.031	0.046	0.012	0.12	0.18
n-Caproic acid vinyl ester	3050-69-9	1	0	1.6	9.3	0	0	55
Chrysene	218-01-9	13	0	0.048	0.075	0	0.18	0.31
Coronene	191-07-1	10	0	0.02	0.038	0	0.071	0.17
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.08	0.13	0.055	0.24	0.73
Cyclohexylamine	108-91-8	4	0	0.54	1.6	0	3.7	7.3
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.017	0.021	0.0064	0.05	0.097
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.079	0.11	0.059	0.33	0.47
Dimethyl phthalate	131-11-3	4	0	0.9	2.8	0	5.8	15
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	5.1	30	0	0	170
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.025	0.083	0	0.29	0.29
Methyl stearate	112-61-8	10	0	1.3	2.8	0	8.8	10
4-tert-Octylphenol	140-66-9	10	0	0.4	0.9	0	2.4	3.3
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.0042	0.017	0	0.025	0.071
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.1	0.28	0	0.49	1.5

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-122. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender
Spectators Third Trimester Fetus<0 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	28	20	22	60	70
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.056	0.059	0.03	0.15	0.23
Benzo[e]pyrene	192-97-2	6	0	0.00092	0.0021	0	0.0063	0.0063
Benzo[g,h,i]perylene	191-24-2	19	0	0.0064	0.0095	0.0025	0.025	0.037
n-Caproic acid vinyl ester	3050-69-9	1	0	0.33	2	0	0	11
Chrysene	218-01-9	13	0	0.01	0.016	0	0.038	0.065
Coronene	191-07-1	10	0	0.0042	0.0079	0	0.015	0.036
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.017	0.028	0.011	0.051	0.15
Cyclohexylamine	108-91-8	4	0	0.11	0.33	0	0.76	1.5
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.0036	0.0044	0.0013	0.01	0.02
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.016	0.023	0.012	0.068	0.098
Dimethyl phthalate	131-11-3	4	0	0.2	0.59	0	1.2	3.1
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	1.1	6.2	0	0	36
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.0053	0.017	0	0.06	0.06
Methyl stearate	112-61-8	10	0	0.27	0.58	0	1.8	2.2
4-tert-Octylphenol	140-66-9	10	0	0.1	0.19	0	0.51	0.69
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.00088	0.0036	0	0.0052	0.015
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.022	0.059	0	0.1	0.31

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-123. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender
Spectators 0<2 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	230	160	180	480	560
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.45	0.48	0.24	1.2	1.9
Benzo[e]pyrene	192-97-2	6	0	0.0074	0.017	0	0.05	0.05
Benzo[g,h,i]perylene	191-24-2	19	0	0.052	0.077	0.02	0.21	0.3
n-Caproic acid vinyl ester	3050-69-9	1	0	2.7	16	0	0	91
Chrysene	218-01-9	13	0	0.081	0.13	0	0.3	0.52
Coronene	191-07-1	10	0	0.034	0.064	0	0.12	0.29
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.13	0.23	0.092	0.41	1.2
Cyclohexylamine	108-91-8	4	0	0.9	2.7	0	6.1	12
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.029	0.035	0.011	0.083	0.16
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.13	0.19	0.1	0.55	0.79
Dimethyl phthalate	131-11-3	4	0	1.4	4.7	0	9.7	25
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	8.6	50	0	0	290
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.043	0.14	0	0.48	0.48
Methyl stearate	112-61-8	10	0	2.2	4.6	0	15	18
4-tert-Octylphenol	140-66-9	10	0	0.7	1.5	0	4.1	5.6
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.007	0.029	0	0.042	0.12
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.18	0.47	0	0.82	2.5

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-124. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender
Spectators 2<6 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	120	86	95	250	290
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.24	0.25	0.13	0.64	0.98
Benzo[e]pyrene	192-97-2	6	0	0.0039	0.009	0	0.027	0.027
Benzo[g,h,i]perylene	191-24-2	19	0	0.027	0.04	0.01	0.11	0.16
n-Caproic acid vinyl ester	3050-69-9	1	0	1.4	8.2	0	0	48
Chrysene	218-01-9	13	0	0.043	0.066	0	0.16	0.28
Coronene	191-07-1	10	0	0.018	0.034	0	0.063	0.15
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.071	0.12	0.048	0.21	0.64
Cyclohexylamine	108-91-8	4	0	0.47	1.4	0	3.2	6.4
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.015	0.018	0.0056	0.044	0.086
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.069	0.098	0.052	0.29	0.42
Dimethyl phthalate	131-11-3	4	0	0.8	2.5	0	5.1	13
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	4.5	26	0	0	150
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.022	0.073	0	0.25	0.25
Methyl stearate	112-61-8	10	0	1.2	2.4	0	7.8	9.3
4-tert-Octylphenol	140-66-9	10	0	0.4	0.79	0	2.1	2.9
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.0037	0.015	0	0.022	0.063
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.092	0.25	0	0.43	1.3

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-125. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender
Spectators 6<11 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	110	82	91	240	280
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.23	0.24	0.12	0.61	0.94
Benzo[e]pyrene	192-97-2	6	0	0.0037	0.0085	0	0.025	0.025
Benzo[g,h,i]perylene	191-24-2	19	0	0.026	0.038	0.0099	0.1	0.15
n-Caproic acid vinyl ester	3050-69-9	1	0	1.3	7.9	0	0	46
Chrysene	218-01-9	13	0	0.041	0.063	0	0.15	0.26
Coronene	191-07-1	10	0	0.017	0.032	0	0.06	0.15
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.067	0.11	0.046	0.2	0.61
Cyclohexylamine	108-91-8	4	0	0.45	1.3	0	3.1	6.1
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.014	0.018	0.0054	0.042	0.082
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.066	0.093	0.05	0.28	0.4
Dimethyl phthalate	131-11-3	4	0	0.7	2.4	0	4.8	12
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	4.3	25	0	0	150
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.021	0.07	0	0.24	0.24
Methyl stearate	112-61-8	10	0	1.1	2.3	0	7.4	8.8
4-tert-Octylphenol	140-66-9	10	0	0.4	0.76	0	2	2.8
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.0035	0.014	0	0.021	0.06
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.088	0.24	0	0.41	1.2

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-126. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender
Spectators 11<16 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	49	35	39	100	120
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.098	0.1	0.052	0.26	0.41
Benzo[e]pyrene	192-97-2	6	0	0.0016	0.0037	0	0.011	0.011
Benzo[g,h,i]perylene	191-24-2	19	0	0.011	0.017	0.0043	0.044	0.065
n-Caproic acid vinyl ester	3050-69-9	1	0	0.58	3.4	0	0	20
Chrysene	218-01-9	13	0	0.018	0.027	0	0.066	0.11
Coronene	191-07-1	10	0	0.0073	0.014	0	0.026	0.063
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.029	0.049	0.02	0.088	0.26
Cyclohexylamine	108-91-8	4	0	0.2	0.58	0	1.3	2.7
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.0062	0.0076	0.0023	0.018	0.035
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.029	0.04	0.022	0.12	0.17
Dimethyl phthalate	131-11-3	4	0	0.3	1	0	2.1	5.3
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	1.9	11	0	0	63
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.0092	0.03	0	0.1	0.1
Methyl stearate	112-61-8	10	0	0.48	1	0	3.2	3.8
4-tert-Octylphenol	140-66-9	10	0	0.2	0.33	0	0.88	1.2
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.0015	0.0062	0	0.0091	0.026
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.038	0.1	0	0.18	0.54

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-127. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender
Spectators 16<30 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	33	23	26	69	80
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.065	0.068	0.035	0.17	0.27
Benzo[e]pyrene	192-97-2	6	0	0.0011	0.0024	0	0.0072	0.0072
Benzo[g,h,i]perylene	191-24-2	19	0	0.0074	0.011	0.0028	0.029	0.043
n-Caproic acid vinyl ester	3050-69-9	1	0	0.38	2.2	0	0	13
Chrysene	218-01-9	13	0	0.012	0.018	0	0.044	0.075
Coronene	191-07-1	10	0	0.0048	0.0091	0	0.017	0.041
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.019	0.032	0.013	0.058	0.17
Cyclohexylamine	108-91-8	4	0	0.13	0.38	0	0.88	1.8
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.0041	0.005	0.0015	0.012	0.023
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.019	0.027	0.014	0.078	0.11
Dimethyl phthalate	131-11-3	4	0	0.2	0.68	0	1.4	3.5
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	1.2	7.1	0	0	42
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.0061	0.02	0	0.069	0.069
Methyl stearate	112-61-8	10	0	0.31	0.66	0	2.1	2.5
4-tert-Octylphenol	140-66-9	10	0	0.1	0.22	0	0.58	0.79
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.001	0.0041	0	0.006	0.017
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.025	0.067	0	0.12	0.36

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-128. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender
Spectators 30<40 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	28	20	23	60	70
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.056	0.059	0.03	0.15	0.23
Benzo[e]pyrene	192-97-2	6	0	0.00093	0.0021	0	0.0063	0.0063
Benzo[g,h,i]perylene	191-24-2	19	0	0.0065	0.0096	0.0025	0.026	0.038
n-Caproic acid vinyl ester	3050-69-9	1	0	0.34	2	0	0	11
Chrysene	218-01-9	13	0	0.01	0.016	0	0.038	0.065
Coronene	191-07-1	10	0	0.0042	0.008	0	0.015	0.036
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.017	0.028	0.011	0.051	0.15
Cyclohexylamine	108-91-8	4	0	0.11	0.33	0	0.77	1.5
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.0036	0.0044	0.0013	0.01	0.02
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.016	0.023	0.012	0.069	0.099
Dimethyl phthalate	131-11-3	4	0	0.2	0.59	0	1.2	3.1
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	1.1	6.2	0	0	36
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.0053	0.017	0	0.06	0.06
Methyl stearate	112-61-8	10	0	0.27	0.58	0	1.8	2.2
4-tert-Octylphenol	140-66-9	10	0	0.1	0.19	0	0.51	0.69
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.00088	0.0036	0	0.0052	0.015
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.022	0.059	0	0.1	0.31

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-129. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender
Spectators 40<50 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	29	21	23	61	71
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.057	0.06	0.031	0.15	0.24
Benzo[e]pyrene	192-97-2	6	0	0.00094	0.0022	0	0.0064	0.0064
Benzo[g,h,i]perylene	191-24-2	19	0	0.0065	0.0097	0.0025	0.026	0.038
n-Caproic acid vinyl ester	3050-69-9	1	0	0.34	2	0	0	12
Chrysene	218-01-9	13	0	0.01	0.016	0	0.038	0.066
Coronene	191-07-1	10	0	0.0043	0.0081	0	0.015	0.037
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.017	0.029	0.012	0.051	0.15
Cyclohexylamine	108-91-8	4	0	0.11	0.34	0	0.77	1.5
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.0036	0.0044	0.0014	0.011	0.021
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.017	0.023	0.013	0.069	0.1
Dimethyl phthalate	131-11-3	4	0	0.2	0.6	0	1.2	3.1
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	1.1	6.3	0	0	37
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.0054	0.018	0	0.061	0.061
Methyl stearate	112-61-8	10	0	0.28	0.58	0	1.9	2.2
4-tert-Octylphenol	140-66-9	10	0	0.1	0.19	0	0.52	0.7
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.00089	0.0036	0	0.0053	0.015
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.022	0.06	0	0.1	0.31

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-130. **On-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **DARTs**—Combined Gender
Spectators 50<70 Years

Chemical	CASRN	Detection	C _{inh} -DART-field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field Related Chemical								
2-Butanone	78-93-3	30	0	29	21	23	61	71
Field-Related Chemicals								
Benzo[a]pyrene	50-32-8	28	0	0.057	0.06	0.031	0.15	0.24
Benzo[e]pyrene	192-97-2	6	0	0.00094	0.0022	0	0.0064	0.0064
Benzo[g,h,i]perylene	191-24-2	19	0	0.0065	0.0097	0.0025	0.026	0.038
n-Caproic acid vinyl ester	3050-69-9	1	0	0.34	2	0	0	12
Chrysene	218-01-9	13	0	0.01	0.016	0	0.039	0.066
Coronene	191-07-1	10	0	0.0043	0.0081	0	0.015	0.037
Cyclohexanamine, N-cyclohexyl-	101-83-7	19	0	0.017	0.029	0.012	0.051	0.15
Cyclohexylamine	108-91-8	4	0	0.11	0.34	0	0.78	1.6
Cyclopenta[cd]pyrene	27208-37-3	28	0	0.0036	0.0044	0.0014	0.011	0.021
N,N-Dicyclohexylmethylamine	7560-83-0	18	0	0.017	0.024	0.013	0.069	0.1
Dimethyl phthalate	131-11-3	4	0	0.2	0.6	0	1.2	3.1
Bis(2-Ethylhexyl)adipate	103-23-1	1	0	1.1	6.3	0	0	37
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	0.0054	0.018	0	0.061	0.061
Methyl stearate	112-61-8	10	0	0.28	0.59	0	1.9	2.2
4-tert-Octylphenol	140-66-9	10	0	0.1	0.19	0	0.52	0.7
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	2	0	0.00089	0.0036	0	0.0053	0.015
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.022	0.06	0	0.1	0.32

^a 34 field-specific $C_{inh-DART-field}$ values are included in the table.

^b On-field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-131. **Off-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b ($C_{inh-DART-field}$, nanograms per cubic meter air) for **Field-Related^c DARTs**—Combined Gender **Spectators Third Trimester Fetus<0 Years**

Chemical	CASRN	Detection	$C_{inh-DART-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	24	0	0.069	0.11	0.029	0.25	0.49
Benzo[e]pyrene	192-97-2	6	0	0.0013	0.003	0	0.0063	0.013
Benzo[g,h,i]perylene	191-24-2	16	0	0.0066	0.011	0	0.024	0.049
n-Caproic acid vinyl ester	3050-69-9	3	0	0.65	2.6	0	3.4	15
Chrysene	218-01-9	7	0	0.0052	0.011	0	0.021	0.046
Coronene	191-07-1	7	0	0.0033	0.0075	0	0.013	0.035
Cyclohexanamine, N-cyclohexyl-	101-83-7	10	0	0.0099	0.017	0	0.052	0.061
Cyclohexylamine	108-91-8	1	0	0.046	0.27	0	0	1.5
Cyclopenta[cd]pyrene	27208-37-3	22	0	0.0031	0.004	0.0013	0.011	0.013
N,N-Dicyclohexylmethylamine	7560-83-0	12	0	0.017	0.026	0	0.066	0.089
Dimethyl phthalate	131-11-3	3	0	0.2	0.58	0	1.2	2.9
Bis(2-Ethylhexyl)adipate	103-23-1	2	0	0.84	3.4	0	5.6	14
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	0.0036	0.015	0	0.024	0.06
Methyl stearate	112-61-8	8	0	0.27	0.59	0	1.7	2.2
4-tert-Octylphenol	140-66-9	7	0	0.0	0.063	0	0.15	0.28
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	0	0	0	0	0	0	0
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.032	0.07	0	0.13	0.34

^a 33 field-specific $C_{inh-DART-field}$ values are included in the table.

^b Off-Field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

^c Non-field DART, 2-butanone, was not sampled off-field. Selected carbonyls, collected using 2,4-dinitrophenylhydrazine-cartridges, were not sampled at off the field locations during the statewide sample collections.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures



Table F-132. **Off-Field Field-Specific^a One-Day Inhalation Exposure Concentration^b**
($C_{inh-DART-field}$, nanograms per cubic meter air) for **Field-Related^c DARTs**—Combined
Gender **Spectators 0<2 Years**

Chemical	CASRN	Detection	$C_{inh-DART-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	24	0	0.56	0.86	0.23	2	3.9
Benzo[e]pyrene	192-97-2	6	0	0.011	0.025	0	0.05	0.1
Benzo[g,h,i]perylene	191-24-2	16	0	0.053	0.086	0	0.19	0.39
n-Caproic acid vinyl ester	3050-69-9	3	0	5.2	21	0	28	120
Chrysene	218-01-9	7	0	0.042	0.088	0	0.17	0.37
Coronene	191-07-1	7	0	0.027	0.06	0	0.1	0.28
Cyclohexanamine, N-cyclohexyl-	101-83-7	10	0	0.08	0.14	0	0.42	0.49
Cyclohexylamine	108-91-8	1	0	0.37	2.1	0	0	12
Cyclopenta[cd]pyrene	27208-37-3	22	0	0.025	0.032	0.011	0.089	0.1
N,N-Dicyclohexylmethylamine	7560-83-0	12	0	0.14	0.21	0	0.53	0.72
Dimethyl phthalate	131-11-3	3	0	1.3	4.7	0	9.7	24
Bis(2-Ethylhexyl)adipate	103-23-1	2	0	6.8	27	0	45	110
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	0.029	0.12	0	0.19	0.48
Methyl stearate	112-61-8	8	0	2.2	4.7	0	14	17
4-tert-Octylphenol	140-66-9	7	0	0.2	0.5	0	1.2	2.3
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	0	0	0	0	0	0	0
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.26	0.56	0	1.1	2.8

^a 33 field-specific $C_{inh-DART-field}$ values are included in the table.

^b Off-Field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

^c Non-field DART, 2-butanone, was not sampled off-field. Selected carbonyls, collected using 2,4-dinitrophenylhydrazine-cartridges, were not sampled at off the field locations during the statewide sample collections.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures



Table F-133. **Off-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **Field-Related^c DARTs**—Combined
Gender **Spectators 2<6 Years**

Chemical	CASRN	Detection	$C_{inh-DART-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	24	0	0.29	0.45	0.12	1	2.1
Benzo[e]pyrene	192-97-2	6	0	0.0056	0.013	0	0.027	0.053
Benzo[g,h,i]perylene	191-24-2	16	0	0.028	0.045	0	0.1	0.21
n-Caproic acid vinyl ester	3050-69-9	3	0	2.8	11	0	15	62
Chrysene	218-01-9	7	0	0.022	0.046	0	0.088	0.19
Coronene	191-07-1	7	0	0.014	0.032	0	0.053	0.15
Cyclohexanamine, N-cyclohexyl-	101-83-7	10	0	0.042	0.074	0	0.22	0.26
Cyclohexylamine	108-91-8	1	0	0.2	1.1	0	0	6.4
Cyclopenta[cd]pyrene	27208-37-3	22	0	0.013	0.017	0.0056	0.047	0.054
N,N-Dicyclohexylmethylamine	7560-83-0	12	0	0.073	0.11	0	0.28	0.38
Dimethyl phthalate	131-11-3	3	0	0.7	2.4	0	5.1	12
Bis(2-Ethylhexyl)adipate	103-23-1	2	0	3.6	14	0	23	59
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	0.015	0.062	0	0.1	0.25
Methyl stearate	112-61-8	8	0	1.1	2.5	0	7.4	9.1
4-tert-Octylphenol	140-66-9	7	0	0.1	0.26	0	0.64	1.2
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	0	0	0	0	0	0	0
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.13	0.3	0	0.55	1.5

^a 33 field-specific $C_{inh-DART-field}$ values are included in the table.

^b Off-Field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

^c Non-field DART, 2-butanone, was not sampled off-field. Selected carbonyls, collected using 2,4-dinitrophenylhydrazine-cartridges, were not sampled at off the field locations during the statewide sample collections.

CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures



Table F-134. **Off-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **Field-Related^c DARTs**—Combined
Gender **Spectators 6<11 Years**

Chemical	CASRN	Detection	$C_{inh-DART-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	24	0	0.28	0.43	0.12	1	2
Benzo[e]pyrene	192-97-2	6	0	0.0054	0.012	0	0.025	0.051
Benzo[g,h,i]perylene	191-24-2	16	0	0.026	0.043	0	0.096	0.2
n-Caproic acid vinyl ester	3050-69-9	3	0	2.6	11	0	14	59
Chrysene	218-01-9	7	0	0.021	0.044	0	0.084	0.18
Coronene	191-07-1	7	0	0.013	0.03	0	0.051	0.14
Cyclohexanamine, N-cyclohexyl-	101-83-7	10	0	0.04	0.071	0	0.21	0.25
Cyclohexylamine	108-91-8	1	0	0.19	1.1	0	0	6.1
Cyclopenta[cd]pyrene	27208-37-3	22	0	0.013	0.016	0.0054	0.045	0.052
N,N-Dicyclohexylmethylamine	7560-83-0	12	0	0.069	0.11	0	0.27	0.36
Dimethyl phthalate	131-11-3	3	0	0.7	2.3	0	4.8	12
Bis(2-Ethylhexyl)adipate	103-23-1	2	0	3.4	14	0	22	56
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	0.015	0.059	0	0.097	0.24
Methyl stearate	112-61-8	8	0	1.1	2.4	0	7	8.7
4-tert-Octylphenol	140-66-9	7	0	0.1	0.25	0	0.61	1.1
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	0	0	0	0	0	0	0
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.13	0.28	0	0.53	1.4

^a 33 field-specific $C_{inh-DART-field}$ values are included in the table.

^b Off-Field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

^c Non-field DART, 2-butanone, was not sampled off-field. Selected carbonyls, collected using 2,4-dinitrophenylhydrazine-cartridges, were not sampled at off the field locations during the statewide sample collections.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures



Table F-135. **Off-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b ($C_{inh-DART-field}$, nanograms per cubic meter air) for **Field-Related^c DARTs**—Combined Gender **Spectators 11<16 Years**

Chemical	CASRN	Detection	$C_{inh-DART-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	24	0	0.12	0.19	0.05	0.43	0.85
Benzo[e]pyrene	192-97-2	6	0	0.0023	0.0053	0	0.011	0.022
Benzo[g,h,i]perylene	191-24-2	16	0	0.011	0.019	0	0.042	0.085
n-Caproic acid vinyl ester	3050-69-9	3	0	1.1	4.6	0	6	26
Chrysene	218-01-9	7	0	0.009	0.019	0	0.036	0.08
Coronene	191-07-1	7	0	0.0058	0.013	0	0.022	0.061
Cyclohexanamine, N-cyclohexyl-	101-83-7	10	0	0.017	0.03	0	0.091	0.11
Cyclohexylamine	108-91-8	1	0	0.08	0.46	0	0	2.7
Cyclopenta[cd]pyrene	27208-37-3	22	0	0.0055	0.007	0.0023	0.019	0.022
N,N-Dicyclohexylmethylamine	7560-83-0	12	0	0.03	0.046	0	0.12	0.15
Dimethyl phthalate	131-11-3	3	0	0.3	1	0	2.1	5.1
Bis(2-Ethylhexyl)adipate	103-23-1	2	0	1.5	5.9	0	9.7	24
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	0.0063	0.025	0	0.042	0.1
Methyl stearate	112-61-8	8	0	0.47	1	0	3	3.8
4-tert-Octylphenol	140-66-9	7	0	0.0	0.11	0	0.26	0.5
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	0	0	0	0	0	0	0
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.055	0.12	0	0.23	0.6

^a 33 field-specific $C_{inh-DART-field}$ values are included in the table.

^b Off-Field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

^c Non-field DART, 2-butanone, was not sampled off-field. Selected carbonyls, collected using 2,4-dinitrophenylhydrazine-cartridges, were not sampled at off the field locations during the statewide sample collections.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures



Table F-136. **Off-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b ($C_{inh-DART-field}$, nanograms per cubic meter air) for **Field-Related^c DARTs**—Combined Gender **Spectators 16<30 Years**

Chemical	CASRN	Detection	$C_{inh-DART-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	24	0	0.079	0.12	0.033	0.28	0.56
Benzo[e]pyrene	192-97-2	6	0	0.0015	0.0035	0	0.0072	0.015
Benzo[g,h,i]perylene	191-24-2	16	0	0.0075	0.012	0	0.027	0.056
n-Caproic acid vinyl ester	3050-69-9	3	0	0.75	3	0	4	17
Chrysene	218-01-9	7	0	0.006	0.013	0	0.024	0.053
Coronene	191-07-1	7	0	0.0038	0.0086	0	0.014	0.04
Cyclohexanamine, N-cyclohexyl-	101-83-7	10	0	0.011	0.02	0	0.06	0.071
Cyclohexylamine	108-91-8	1	0	0.053	0.3	0	0	1.8
Cyclopenta[cd]pyrene	27208-37-3	22	0	0.0036	0.0046	0.0015	0.013	0.015
N,N-Dicyclohexylmethylamine	7560-83-0	12	0	0.02	0.03	0	0.076	0.1
Dimethyl phthalate	131-11-3	3	0	0.2	0.66	0	1.4	3.4
Bis(2-Ethylhexyl)adipate	103-23-1	2	0	0.97	3.9	0	6.4	16
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	0.0042	0.017	0	0.028	0.069
Methyl stearate	112-61-8	8	0	0.31	0.67	0	2	2.5
4-tert-Octylphenol	140-66-9	7	0	0.0	0.072	0	0.17	0.33
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	0	0	0	0	0	0	0
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.037	0.081	0	0.15	0.39

^a 33 field-specific $C_{inh-DART-field}$ values are included in the table.

^b Off-Field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

^c Non-field DART, 2-butanone, was not sampled off-field. Selected carbonyls, collected using 2,4-dinitrophenylhydrazine-cartridges, were not sampled at off the field locations during the statewide sample collections.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures



Table F-137. **Off-Field Field-Specific^a One-Day Inhalation Exposure Concentration^b**
($C_{inh-DART-field}$, nanograms per cubic meter air) for **Field-Related^c DARTs—Combined**
Gender Spectators 30<40 Years

Chemical	CASRN	Detection	$C_{inh-DART-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	24	0	0.069	0.11	0.029	0.25	0.49
Benzo[e]pyrene	192-97-2	6	0	0.0013	0.0031	0	0.0063	0.013
Benzo[g,h,i]perylene	191-24-2	16	0	0.0066	0.011	0	0.024	0.049
n-Caproic acid vinyl ester	3050-69-9	3	0	0.65	2.7	0	3.5	15
Chrysene	218-01-9	7	0	0.0052	0.011	0	0.021	0.046
Coronene	191-07-1	7	0	0.0034	0.0075	0	0.013	0.035
Cyclohexanamine, N-cyclohexyl-	101-83-7	10	0	0.0099	0.018	0	0.052	0.062
Cyclohexylamine	108-91-8	1	0	0.046	0.27	0	0	1.5
Cyclopenta[cd]pyrene	27208-37-3	22	0	0.0031	0.004	0.0013	0.011	0.013
N,N-Dicyclohexylmethylamine	7560-83-0	12	0	0.017	0.027	0	0.067	0.089
Dimethyl phthalate	131-11-3	3	0	0.2	0.58	0	1.2	3
Bis(2-Ethylhexyl)adipate	103-23-1	2	0	0.85	3.4	0	5.6	14
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	0.0037	0.015	0	0.024	0.06
Methyl stearate	112-61-8	8	0	0.27	0.59	0	1.7	2.2
4-tert-Octylphenol	140-66-9	7	0	0.0	0.063	0	0.15	0.29
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	0	0	0	0	0	0	0
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.032	0.07	0	0.13	0.35

^a 33 field-specific $C_{inh-DART-field}$ values are included in the table.

^b Off-Field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

^c Non-field DART, 2-butanone, was not sampled off-field. Selected carbonyls, collected using 2,4-dinitrophenylhydrazine-cartridges, were not sampled at off the field locations during the statewide sample collections.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures



Table F-138. **Off-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **Field-Related^c DARTs**—Combined
Gender **Spectators 40<50 Years**

Chemical	CASRN	Detection	$C_{inh-DART-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	24	0	0.07	0.11	0.029	0.25	0.5
Benzo[e]pyrene	192-97-2	6	0	0.0014	0.0031	0	0.0064	0.013
Benzo[g,h,i]perylene	191-24-2	16	0	0.0067	0.011	0	0.024	0.05
n-Caproic acid vinyl ester	3050-69-9	3	0	0.66	2.7	0	3.5	15
Chrysene	218-01-9	7	0	0.0053	0.011	0	0.021	0.047
Coronene	191-07-1	7	0	0.0034	0.0076	0	0.013	0.035
Cyclohexanamine, N-cyclohexyl-	101-83-7	10	0	0.01	0.018	0	0.053	0.062
Cyclohexylamine	108-91-8	1	0	0.047	0.27	0	0	1.5
Cyclopenta[cd]pyrene	27208-37-3	22	0	0.0032	0.0041	0.0014	0.011	0.013
N,N-Dicyclohexylmethylamine	7560-83-0	12	0	0.017	0.027	0	0.067	0.09
Dimethyl phthalate	131-11-3	3	0	0.2	0.59	0	1.2	3
Bis(2-Ethylhexyl)adipate	103-23-1	2	0	0.85	3.4	0	5.6	14
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	0.0037	0.015	0	0.024	0.061
Methyl stearate	112-61-8	8	0	0.27	0.59	0	1.8	2.2
4-tert-Octylphenol	140-66-9	7	0	0.0	0.063	0	0.15	0.29
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	0	0	0	0	0	0	0
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.032	0.071	0	0.13	0.35

^a 33 field-specific $C_{inh-DART-field}$ values are included in the table.

^b Off-Field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

^c Non-field DART, 2-butanone, was not sampled off-field. Selected carbonyls, collected using 2,4-dinitrophenylhydrazine-cartridges, were not sampled at off the field locations during the statewide sample collections.

CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures



Table F-139. **Off-Field** Field-Specific^a One-Day Inhalation Exposure Concentration^b
($C_{inh-DART-field}$, nanograms per cubic meter air) for **Field-Related^c DARTs**—Combined
Gender **Spectators 50<70 Years**

Chemical	CASRN	Detection	$C_{inh-DART-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	24	0	0.07	0.11	0.029	0.25	0.5
Benzo[e]pyrene	192-97-2	6	0	0.0014	0.0031	0	0.0064	0.013
Benzo[g,h,i]perylene	191-24-2	16	0	0.0067	0.011	0	0.024	0.05
n-Caproic acid vinyl ester	3050-69-9	3	0	0.66	2.7	0	3.5	15
Chrysene	218-01-9	7	0	0.0053	0.011	0	0.021	0.047
Coronene	191-07-1	7	0	0.0034	0.0076	0	0.013	0.035
Cyclohexanamine, N-cyclohexyl-	101-83-7	10	0	0.01	0.018	0	0.053	0.063
Cyclohexylamine	108-91-8	1	0	0.047	0.27	0	0	1.6
Cyclopenta[cd]pyrene	27208-37-3	22	0	0.0032	0.0041	0.0014	0.011	0.013
N,N-Dicyclohexylmethylamine	7560-83-0	12	0	0.017	0.027	0	0.068	0.09
Dimethyl phthalate	131-11-3	3	0	0.2	0.59	0	1.2	3
Bis(2-Ethylhexyl)adipate	103-23-1	2	0	0.86	3.4	0	5.7	14
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	0.0037	0.015	0	0.024	0.061
Methyl stearate	112-61-8	8	0	0.27	0.6	0	1.8	2.2
4-tert-Octylphenol	140-66-9	7	0	0.0	0.064	0	0.15	0.29
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	0	0	0	0	0	0	0
Phenol, 4-(1-phenylethyl)-	1988-89-2	9	0	0.032	0.071	0	0.13	0.35

^a 33 field-specific $C_{inh-DART-field}$ values are included in the table.

^b Off-Field field-specific $C_{inh-DART-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh-DART}$) of a DART (see Appendix Section F.4.2).

^c Non-field DART, 2-butanone, was not sampled off-field. Selected carbonyls, collected using 2,4-dinitrophenylhydrazine-cartridges, were not sampled at off the field locations during the statewide sample collections.

CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures



F.4.5. Chronic Inhalation Exposure Concentration for Sensory Irritants (Chronic $C_{inh-sensory}$)

Table F-140. Chronic Inhalation Exposure Concentration^a (Chronic $C_{inh-sensory}$, nanograms per cubic meter air) for **Sensory Irritants—All Receptor Categories**

Chemical	CASRN	Chronic C _{inh-sensory}	
		On-Field	Off-Field
Field-Related Chemical			
Styrene	100-42-5	59	60
Non-Field-Related Chemicals			
Acetaldehyde	75-07-0	2500	Not measured
Formaldehyde	50-00-0	3800	Not measured

^a On-Field Chronic $C_{inh-sensory}$ equals to mean of the 35 individual field average concentrations ($C_{air-avg}$) of a chemical detected in air on the fields. Off-Field Chronic $C_{inh-sensory}$ equals to mean of the 35 individual field average concentrations ($C_{air-avg}$) of a chemical detected in air off the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

INDIVIDUAL FIELD ASSESSMENT (Table F-141)

Table F-141. Field-Specific Chronic Inhalation Exposure Concentration (Chronic $C_{inh-sensory-field}$, nanograms per cubic meter air) for **Sensory Irritants—All Receptor Categories**

Location	Chemical	CASRN	Detection	Chronic C _{inh-sensory-field}					
				Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
On Field ^a	Field Related Chemical								
	Styrene	100-42-5	17	0	59	120	0	200	660
	Non-Field-Related Chemicals								
	Acetaldehyde	75-07-0	34	260	2500	1900	1900	4800	9600
	Formaldehyde	50-00-0	34	810	3800	2900	3200	6400	16000
Off Field ^b	Field Related Chemical								
	Styrene	100-42-5	13	0	60	130	0	220	670

^a 35 field-specific on-field $C_{inh-sensory-field}$ values are included in the table.

^b 34 field-specific off-field $C_{inh-sensory-field}$ values are included in the table. Selected carbonyls, collected using 2,4-dinitrophenylhydrazine-cartiridges, were not sampled at off the field locations during the statewide sample collections.



F.4.6. Chronic Inhalation Exposure Concentration for General Chemicals (Chronic C_{inh})

Table F-142. **On-Field** Chronic Inhalation Exposure Concentration^a for **All General Chemicals** (Chronic C_{inh}, nanograms per cubic meter)—Combined Gender **Athletes**

Chemical	CASRN	Chronic C _{inh}						
		2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Field-Related Chemicals								
Acenaphthylene	208-96-8	0.16	0.2	0.2	0.37	0.24	0.24	0.24
Acetone	67-64-1	3000	3900	3800	7300	4700	4700	4600
Aniline	62-53-3	1	1.3	1.3	2.4	1.6	1.5	1.5
Anthracene	120-12-7	0.051	0.066	0.064	0.12	0.079	0.078	0.077
Anthracene, 2-methyl-	613-12-7	0.013	0.017	0.017	0.032	0.021	0.02	0.02
Anthracene, 9,10-dimethyl	781-43-1	0.015	0.02	0.019	0.036	0.024	0.023	0.023
Anthracene, 9-phenyl	602-55-1	0.0014	0.0018	0.0018	0.0033	0.0022	0.0022	0.0021
Benz[a]anthracene	56-55-3	0.00076	0.0009 9	0.00097	0.0018	0.0012	0.0012	0.0012
Benzaldehyde	100-52-7	14	18	18	33	22	21	21
Benzene, 1,2,3-trimethyl-	526-73-8	2	2.7	2.6	4.9	3.2	3.1	3.1
Benzene, 1,2,4,5-tetramethyl-	95-93-2	0.39	0.5	0.49	0.93	0.61	0.6	0.59
Benzene, 1,2,4-trimethyl-	95-63-6	18	24	23	44	28	28	28
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	0.82	1.1	1	1.9	1.3	1.3	1.2
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	1.2	1.5	1.5	2.8	1.8	1.8	1.8
Benzene, butyl-	104-51-8	0.41	0.53	0.52	0.97	0.63	0.63	0.61
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	0.58	0.75	0.73	1.4	0.9	0.89	0.87
Benzo[b]fluoranthene	205-99-2	0.0038	0.005	0.0048	0.0091	0.006	0.0059	0.0058
7H-Benzo[c]fluorene	205-12-9	0.0084	0.011	0.011	0.02	0.013	0.013	0.013
Benzo[k]fluoranthene	207-08-9	0.0048	0.0062	0.006	0.011	0.0074	0.0073	0.0072
Benzothiazole	95-16-9	5.7	7.4	7.3	14	9	8.8	8.7
Benzothiazole, 2-phenyl-	883-93-2	0.46	0.59	0.58	1.1	0.71	0.7	0.69
2-Benzothiazolone	934-34-9	0.72	0.93	0.91	1.7	1.1	1.1	1.1
Benzyl butyl phthalate	85-68-7	0.53	0.69	0.67	1.3	0.83	0.82	0.8
Butanal	123-72-8	47	62	60	110	74	73	72
Cyclopentasiloxane, decamethyl-	541-02-6	25	32	32	60	39	38	38
Cyclotetrasiloxane, octamethyl-	556-67-2	9	12	11	21	14	14	14
p-Cymene	99-87-6	3.9	5	4.9	9.3	6.1	6	5.9
Decane	124-18-5	8.4	11	11	20	13	13	13
Dibenz[a,h]anthracene	53-70-3	0.022	0.029	0.028	0.053	0.035	0.034	0.034
Dibenzothiophene	132-65-0	0.19	0.25	0.24	0.46	0.3	0.29	0.29
Dibutyl phthalate	84-74-2	58	76	74	140	91	90	88
Diethyl phthalate	84-66-2	0.41	0.53	0.52	0.97	0.64	0.63	0.62
Diisobutyl phthalate	84-69-5	2.2	2.9	2.8	5.3	3.4	3.4	3.3



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Chronic C _{inh}						
		2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Diisooctylphthalate	27554-26-3	7.7	9.9	9.7	18	12	12	12
Di-n-octyl phthalate	117-84-0	0.015	0.019	0.019	0.036	0.023	0.023	0.023
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	5	6.5	6.4	12	7.8	7.7	7.6
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	2.6	3.4	3.3	6.3	4.1	4	4
Dodecane	112-40-3	1	1.4	1.3	2.5	1.6	1.6	1.6
Fluoranthene	206-44-0	0.59	0.76	0.74	1.4	0.91	0.9	0.88
Fluorene	86-73-7	0.93	1.2	1.2	2.2	1.4	1.4	1.4
Furan, 2-methyl	534-22-5	17	21	21	39	26	25	25
Heptanal	111-71-7	2.3	3	2.9	5.5	3.6	3.5	3.4
Hexadecane	544-76-3	4.9	6.4	6.2	12	7.7	7.6	7.4
2,5-Hexanedione	110-13-4	4.2	5.5	5.3	10	6.6	6.5	6.4
Indan	496-11-7	2.1	2.7	2.7	5	3.3	3.3	3.2
Mesitylene	108-67-8	4.3	5.6	5.4	10	6.7	6.6	6.5
Methacrolein	78-85-3	12	15	15	28	18	18	18
Methyl Isobutyl Ketone	108-10-1	2.4	3.1	3.1	5.8	3.8	3.7	3.7
Naphthalene	91-20-3	4.1	5.4	5.2	9.9	6.4	6.3	6.2
Naphthalene, 1,2-dimethyl-	573-98-8	0.058	0.075	0.073	0.14	0.09	0.089	0.087
Naphthalene, 1,6-dimethyl-	575-43-9	0.41	0.54	0.52	0.99	0.64	0.63	0.62
Naphthalene, 1-methyl-	90-12-0	3.4	4.4	4.3	8	5.2	5.2	5.1
Naphthalene, 2-(bromomethyl)-	939-26-4	0.11	0.15	0.15	0.27	0.18	0.18	0.17
Naphthalene, 2,3-dimethyl-	581-40-8	0.31	0.4	0.39	0.74	0.48	0.48	0.47
Naphthalene, 2-methyl-	91-57-6	5.1	6.7	6.5	12	8	7.9	7.8
1-Octadecene	112-88-9	0.68	0.89	0.87	1.6	1.1	1.1	1
Octanal	124-13-0	6.9	9	8.8	17	11	11	10
Octane	111-65-9	9.4	12	12	22	15	14	14
17-Pentatriacontene	6971-40-0	0.11	0.15	0.14	0.27	0.18	0.17	0.17
N-Phenylbenzamide	93-98-1	1.4	1.8	1.8	3.3	2.2	2.1	2.1
Phenanthrene	85-01-8	2	2.6	2.6	4.8	3.1	3.1	3
Phenanthrene, 1-methyl	832-69-9	0.13	0.17	0.16	0.3	0.2	0.2	0.19
Phenanthrene, 2-methyl-	2531-84-2	0.23	0.3	0.29	0.55	0.36	0.36	0.35
Phenanthrene, 3-methyl	832-71-3	0.28	0.36	0.35	0.66	0.43	0.42	0.42
Propionaldehyde	123-38-6	29	37	36	68	45	44	43
Pyrene	129-00-0	0.5	0.64	0.63	1.2	0.77	0.76	0.75
Pyridine, 2-(4-methylphenyl)-	4467-06-5	0.0055	0.0071	0.007	0.013	0.0086	0.0084	0.0083
Resorcinol	108-46-3	2.9	3.8	3.7	7	4.5	4.5	4.4
m-Tolualdehyde	620-23-5	42	54	53	100	65	64	63
TXIB "Kodaflex"	6846-50-0	0.31	0.4	0.39	0.73	0.48	0.47	0.46
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	0.59	0.76	0.74	1.4	0.92	0.9	0.89



Chemical	CASRN	Chronic C _{inh}						
		2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Undecane	1120-21-4	2	2.6	2.6	4.9	3.2	3.1	3.1
Valeraldehyde	110-62-3	140	190	180	350	230	220	220
Non-Field-Related Chemicals								
Benzene	71-43-2	94	120	120	220	150	140	140
Benzene, 1,4-dichloro	106-46-7	3	3.9	3.8	7.1	4.6	4.6	4.5
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	89	120	110	210	140	140	130
2-Butoxyethanol	111-76-2	0.77	1	0.97	1.8	1.2	1.2	1.2
Cyclotrisiloxane, hexamethyl-	541-05-9	11	15	14	27	18	17	17
Decanal	112-31-2	4.4	5.6	5.5	10	6.8	6.7	6.6
Ethylbenzene	100-41-4	27	35	34	64	42	41	40
Heptane	142-82-5	36	46	45	85	56	55	54
Hexanal	66-25-1	120	160	150	290	190	190	180
Hexane	110-54-3	100	130	130	250	160	160	160
1-Hexanol, 2-ethyl-	104-76-7	1.2	1.6	1.5	2.9	1.9	1.9	1.8
Nonanal	124-19-6	1.2	1.6	1.5	2.9	1.9	1.9	1.8
Phenol	108-95-2	9.1	12	11	22	14	14	14
Tetrachloroethylene	127-18-4	7.4	9.6	9.4	18	12	11	11
Tetradecane	629-59-4	1.3	1.6	1.6	3	2	1.9	1.9
Texanol, TXIB (mono-isomer)	25265-77-4	16	21	20	39	25	25	24
Toluene	108-88-3	210	270	270	500	330	320	320
Trichloroethylene	79-01-6	1.5	2	1.9	3.6	2.4	2.3	2.3
Trichloromethane	67-66-3	5.9	7.7	7.5	14	9.2	9.1	8.9
m/p-Xylene	106-42-3	90	120	110	210	140	140	140
o-Xylene	95-47-6	30	39	38	72	47	46	45

^a On-Field Chronic C_{inh} equals the modified mean of the 35 individual field average concentrations (C_{air-avg} x AF_{inh}) of a chemical detected in air on the fields (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-143. On-Field Chronic Inhalation Exposure Concentration^a for All General Chemicals (Chronic C_{inh}, nanograms per cubic meter)—Combined Gender Coaches

Chemical	CASRN	Chronic C _{inh}			
		16<30 years	30<40 years	40<50 years	50<70 years
Field-Related Chemicals					
Acenaphthylene	208-96-8	0.15	0.13	0.14	0.14
Acetone	67-64-1	3000	2600	2600	2700
Aniline	62-53-3	0.99	0.87	0.88	0.88
Anthracene	120-12-7	0.05	0.044	0.044	0.044
Anthracene, 2-methyl-	613-12-7	0.013	0.011	0.012	0.012



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Chronic C _{inh}			
		16<30 years	30<40 years	40<50 years	50<70 years
Anthracene, 9,10-dimethyl	781-43-1	0.015	0.013	0.013	0.013
Anthracene, 9-phenyl	602-55-1	0.0014	0.0012	0.0012	0.0012
Benz[a]anthracene	56-55-3	0.00075	0.00066	0.00067	0.00067
Benzaldehyde	100-52-7	14	12	12	12
Benzene, 1,2,3-trimethyl-	526-73-8	2	1.8	1.8	1.8
Benzene, 1,2,4,5-tetramethyl-	95-93-2	0.38	0.33	0.34	0.34
Benzene, 1,2,4-trimethyl-	95-63-6	18	16	16	16
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	0.81	0.7	0.71	0.71
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	1.1	1	1	1
Benzene, butyl-	104-51-8	0.4	0.35	0.35	0.36
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	0.57	0.5	0.5	0.51
Benzo[b]fluoranthene	205-99-2	0.0038	0.0033	0.0033	0.0033
7H-Benzo[c]fluorene	205-12-9	0.0083	0.0072	0.0073	0.0073
Benzo[k]fluoranthene	207-08-9	0.0047	0.0041	0.0041	0.0042
Benzothiazole	95-16-9	5.7	4.9	5	5
Benzothiazole, 2-phenyl-	883-93-2	0.45	0.39	0.4	0.4
2-Benzothiazolone	934-34-9	0.71	0.62	0.62	0.63
Benzyl butyl phthalate	85-68-7	0.53	0.46	0.46	0.46
Butanal	123-72-8	47	41	41	41
Cyclopentasiloxane, decamethyl-	541-02-6	25	22	22	22
Cyclotetrasiloxane, octamethyl-	556-67-2	8.9	7.8	7.8	7.9
p-Cymene	99-87-6	3.8	3.3	3.4	3.4
Decane	124-18-5	8.3	7.2	7.3	7.3
Dibenz[a,h]anthracene	53-70-3	0.022	0.019	0.019	0.019
Dibenzothiophene	132-65-0	0.19	0.17	0.17	0.17
Dibutyl phthalate	84-74-2	57	50	51	51
Diethyl phthalate	84-66-2	0.4	0.35	0.36	0.36
Diisobutyl phthalate	84-69-5	2.2	1.9	1.9	1.9
Diisooctylphthalate	27554-26-3	7.6	6.6	6.7	6.7
Di-n-octyl phthalate	117-84-0	0.015	0.013	0.013	0.013
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	5	4.3	4.4	4.4
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	2.6	2.3	2.3	2.3
Dodecane	112-40-3	1	0.9	0.91	0.92
Fluoranthene	206-44-0	0.58	0.5	0.51	0.51
Fluorene	86-73-7	0.91	0.8	0.81	0.81
Furan, 2-methyl	534-22-5	16	14	14	14
Heptanal	111-71-7	2.3	2	2	2
Hexadecane	544-76-3	4.9	4.2	4.3	4.3
2,5-Hexanedione	110-13-4	4.2	3.6	3.7	3.7
Indan	496-11-7	2.1	1.8	1.8	1.8



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Chronic C _{inh}			
		16<30 years	30<40 years	40<50 years	50<70 years
Mesitylene	108-67-8	4.2	3.7	3.7	3.7
Methacrolein	78-85-3	12	10	10	10
Methyl Isobutyl Ketone	108-10-1	2.4	2.1	2.1	2.1
Naphthalene	91-20-3	4.1	3.6	3.6	3.6
Naphthalene, 1,2-dimethyl-	573-98-8	0.057	0.05	0.05	0.05
Naphthalene, 1,6-dimethyl-	575-43-9	0.41	0.36	0.36	0.36
Naphthalene, 1-methyl-	90-12-0	3.3	2.9	2.9	2.9
Naphthalene, 2-(bromomethyl)-	939-26-4	0.11	0.099	0.1	0.1
Naphthalene, 2,3-dimethyl-	581-40-8	0.31	0.27	0.27	0.27
Naphthalene, 2-methyl-	91-57-6	5.1	4.4	4.5	4.5
1-Octadecene	112-88-9	0.68	0.59	0.6	0.6
Octanal	124-13-0	6.9	6	6.1	6.1
Octane	111-65-9	9.3	8.1	8.2	8.2
17-Pentatriacontene	6971-40-0	0.11	0.098	0.099	0.099
N-Phenylbenzamide	93-98-1	1.4	1.2	1.2	1.2
Phenanthrene	85-01-8	2	1.7	1.8	1.8
Phenanthrene, 1-methyl	832-69-9	0.13	0.11	0.11	0.11
Phenanthrene, 2-methyl-	2531-84-2	0.23	0.2	0.2	0.2
Phenanthrene, 3-methyl	832-71-3	0.27	0.24	0.24	0.24
Propionaldehyde	123-38-6	28	25	25	25
Pyrene	129-00-0	0.49	0.43	0.43	0.43
Pyridine, 2-(4-methylphenyl)-	4467-06-5	0.0054	0.0047	0.0048	0.0048
Resorcinol	108-46-3	2.9	2.5	2.5	2.5
m-Tolualdehyde	620-23-5	41	36	36	37
TXIB "Kodaflex"	6846-50-0	0.3	0.26	0.27	0.27
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	0.58	0.51	0.51	0.51
Undecane	1120-21-4	2	1.8	1.8	1.8
Valeraldehyde	110-62-3	140	120	130	130
Non-Field-Related Chemicals					
Benzene	71-43-2	93	81	82	82
Benzene, 1,4-dichloro	106-46-7	2.9	2.6	2.6	2.6
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	88	77	78	78
2-Butoxyethanol	111-76-2	0.76	0.66	0.67	0.67
Cyclotrisiloxane, hexamethyl-	541-05-9	11	9.8	9.9	9.9
Decanal	112-31-2	4.3	3.8	3.8	3.8
Ethylbenzene	100-41-4	26	23	23	23
Heptane	142-82-5	35	31	31	31
Hexanal	66-25-1	120	110	110	110
Hexane	110-54-3	100	90	91	91
1-Hexanol, 2-ethyl-	104-76-7	1.2	1	1.1	1.1



Chemical	CASRN	Chronic C _{inh}			
		16<30 years	30<40 years	40<50 years	50<70 years
Nonanal	124-19-6	1.2	1	1.1	1.1
Phenol	108-95-2	8.9	7.8	7.9	7.9
Tetrachloroethylene	127-18-4	7.3	6.4	6.5	6.5
Tetradecane	629-59-4	1.2	1.1	1.1	1.1
Texanol, TXIB (mono-isomer)	25265-77-4	16	14	14	14
Toluene	108-88-3	210	180	180	180
Trichloroethylene	79-01-6	1.5	1.3	1.3	1.3
Trichloromethane	67-66-3	5.8	5.1	5.1	5.2
m/p-Xylene	106-42-3	88	77	78	78
o-Xylene	95-47-6	30	26	26	26

^a On-Field Chronic C_{inh} equals the modified mean of the 35 individual field average concentrations (C_{air-avg} x AF_{inh}) of a chemical detected in air on the fields (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-144. On-Field Chronic Inhalation Exposure Concentration^a for All General Chemicals (Chronic C_{inh}, nanograms per cubic meter)—Combined Gender Referees

Chemical	CASRN	Chronic C _{inh}			
		16<30 years	30<40 years	40<50 years	50<70 years
Field-Related Chemicals					
Acenaphthylene	208-96-8	0.057	0.05	0.05	0.05
Acetone	67-64-1	1100	970	980	990
Aniline	62-53-3	0.37	0.32	0.33	0.33
Anthracene	120-12-7	0.019	0.016	0.016	0.016
Anthracene, 2-methyl-	613-12-7	0.0049	0.0042	0.0043	0.0043
Anthracene, 9,10-dimethyl	781-43-1	0.0055	0.0048	0.0049	0.0049
Anthracene, 9-phenyl	602-55-1	0.00051	0.00045	0.00045	0.00045
Benz[a]anthracene	56-55-3	0.00028	0.00024	0.00025	0.00025
Benzaldehyde	100-52-7	5.1	4.4	4.5	4.5
Benzene, 1,2,3-trimethyl-	526-73-8	0.75	0.65	0.66	0.66
Benzene, 1,2,4,5-tetramethyl-	95-93-2	0.14	0.12	0.13	0.13
Benzene, 1,2,4-trimethyl-	95-63-6	6.7	5.8	5.9	5.9
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	0.3	0.26	0.26	0.26
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	0.42	0.37	0.37	0.38
Benzene, butyl-	104-51-8	0.15	0.13	0.13	0.13
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	0.21	0.19	0.19	0.19
Benzo[b]fluoranthene	205-99-2	0.0014	0.0012	0.0012	0.0012
7H-Benzo[c]fluorene	205-12-9	0.0031	0.0027	0.0027	0.0027
Benzo[k]fluoranthene	207-08-9	0.0017	0.0015	0.0015	0.0015
Benzothiazole	95-16-9	2.1	1.8	1.9	1.9



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Chronic C _{inh}			
		16<30 years	30<40 years	40<50 years	50<70 years
Benzothiazole, 2-phenyl-	883-93-2	0.17	0.15	0.15	0.15
2-Benzothiazolone	934-34-9	0.26	0.23	0.23	0.23
Benzyl butyl phthalate	85-68-7	0.19	0.17	0.17	0.17
Butanal	123-72-8	17	15	15	15
Cyclopentasiloxane, decamethyl-	541-02-6	9.1	8	8.1	8.1
Cyclotetrasiloxane, octamethyl-	556-67-2	3.3	2.9	2.9	2.9
p-Cymene	99-87-6	1.4	1.2	1.3	1.3
Decane	124-18-5	3.1	2.7	2.7	2.7
Dibenz[a,h]anthracene	53-70-3	0.0081	0.0071	0.0072	0.0072
Dibenzothiophene	132-65-0	0.07	0.061	0.062	0.062
Dibutyl phthalate	84-74-2	21	19	19	19
Diethyl phthalate	84-66-2	0.15	0.13	0.13	0.13
Diisobutyl phthalate	84-69-5	0.81	0.71	0.72	0.72
Diisooctylphthalate	27554-26-3	2.8	2.5	2.5	2.5
Di-n-octyl phthalate	117-84-0	0.0055	0.0048	0.0048	0.0049
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	1.8	1.6	1.6	1.6
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	0.96	0.84	0.85	0.85
Dodecane	112-40-3	0.38	0.34	0.34	0.34
Fluoranthene	206-44-0	0.21	0.19	0.19	0.19
Fluorene	86-73-7	0.34	0.3	0.3	0.3
Furan, 2-methyl	534-22-5	6.1	5.3	5.3	5.4
Heptanal	111-71-7	0.84	0.73	0.74	0.74
Hexadecane	544-76-3	1.8	1.6	1.6	1.6
2,5-Hexanedione	110-13-4	1.5	1.3	1.4	1.4
Indan	496-11-7	0.77	0.68	0.68	0.69
Mesitylene	108-67-8	1.6	1.4	1.4	1.4
Methacrolein	78-85-3	4.3	3.8	3.8	3.8
Methyl Isobutyl Ketone	108-10-1	0.89	0.78	0.79	0.79
Naphthalene	91-20-3	1.5	1.3	1.3	1.3
Naphthalene, 1,2-dimethyl-	573-98-8	0.021	0.018	0.019	0.019
Naphthalene, 1,6-dimethyl-	575-43-9	0.15	0.13	0.13	0.13
Naphthalene, 1-methyl-	90-12-0	1.2	1.1	1.1	1.1
Naphthalene, 2-(bromomethyl)-	939-26-4	0.042	0.037	0.037	0.037
Naphthalene, 2,3-dimethyl-	581-40-8	0.11	0.099	0.1	0.1
Naphthalene, 2-methyl-	91-57-6	1.9	1.6	1.7	1.7
1-Octadecene	112-88-9	0.25	0.22	0.22	0.22
Octanal	124-13-0	2.5	2.2	2.2	2.3
Octane	111-65-9	3.4	3	3	3
17-Pentatriacontene	6971-40-0	0.042	0.036	0.037	0.037
N-Phenylbenzamide	93-98-1	0.51	0.45	0.45	0.45



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Chronic C _{inh}			
		16<30 years	30<40 years	40<50 years	50<70 years
Phenanthrene	85-01-8	0.74	0.64	0.65	0.65
Phenanthrene, 1-methyl	832-69-9	0.047	0.041	0.041	0.041
Phenanthrene, 2-methyl-	2531-84-2	0.085	0.074	0.075	0.075
Phenanthrene, 3-methyl	832-71-3	0.1	0.088	0.089	0.09
Propionaldehyde	123-38-6	10	9.1	9.2	9.3
Pyrene	129-00-0	0.18	0.16	0.16	0.16
Pyridine, 2-(4-methylphenyl)-	4467-06-5	0.002	0.0018	0.0018	0.0018
Resorcinol	108-46-3	1.1	0.93	0.94	0.94
m-Tolualdehyde	620-23-5	15	13	14	14
TXIB "Kodaflex"	6846-50-0	0.11	0.098	0.099	0.099
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	0.21	0.19	0.19	0.19
Undecane	1120-21-4	0.75	0.65	0.66	0.66
Valeraldehyde	110-62-3	53	46	47	47
Non-Field-Related Chemicals					
Benzene	71-43-2	34	30	30	30
Benzene, 1,4-dichloro	106-46-7	1.1	0.95	0.96	0.97
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	33	29	29	29
2-Butoxyethanol	111-76-2	0.28	0.25	0.25	0.25
Cyclotrisiloxane, hexamethyl-	541-05-9	4.2	3.6	3.7	3.7
Decanal	112-31-2	1.6	1.4	1.4	1.4
Ethylbenzene	100-41-4	9.8	8.5	8.6	8.6
Heptane	142-82-5	13	11	12	12
Hexanal	66-25-1	45	39	40	40
Hexane	110-54-3	38	33	34	34
1-Hexanol, 2-ethyl-	104-76-7	0.44	0.39	0.39	0.39
Nonanal	124-19-6	0.44	0.39	0.39	0.39
Phenol	108-95-2	3.3	2.9	2.9	2.9
Tetrachloroethylene	127-18-4	2.7	2.4	2.4	2.4
Tetradecane	629-59-4	0.46	0.4	0.41	0.41
Texanol, TXIB (mono-isomer)	25265-77-4	5.9	5.2	5.2	5.2
Toluene	108-88-3	77	67	68	68
Trichloroethylene	79-01-6	0.55	0.48	0.49	0.49
Trichloromethane	67-66-3	2.2	1.9	1.9	1.9
m/p-Xylene	106-42-3	33	29	29	29
o-Xylene	95-47-6	11	9.6	9.7	9.7

^a On-Field Chronic C_{inh} equals the modified mean of the 35 individual field average concentrations (C_{air-avg} x AF_{inh}) of a chemical detected in air on the fields (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-145. **On-Field** Chronic Inhalation Exposure Concentration^a for **All General Chemicals** (Chronic C_{inh}, nanograms per cubic meter)—Combined Gender **Spectators**

Chemical	CASRN	Chronic C _{inh}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Field-Related Chemicals										
Acenaphthylene	208-96-8	0.021	0.17	0.087	0.083	0.036	0.024	0.021	0.021	0.021
Acetone	67-64-1	400	3200	1700	1600	700	460	400	410	410
Aniline	62-53-3	0.13	1.1	0.56	0.54	0.23	0.15	0.13	0.14	0.14
Anthracene	120-12-7	0.0067	0.054	0.028	0.027	0.012	0.0077	0.0068	0.0068	0.0069
Anthracene, 2-methyl-	613-12-7	0.0018	0.014	0.0074	0.0071	0.0031	0.002	0.0018	0.0018	0.0018
Anthracene, 9,10-dimethyl	781-43-1	0.002	0.016	0.0085	0.0081	0.0035	0.0023	0.002	0.002	0.002
Anthracene, 9-phenyl	602-55-1	0.00019	0.0015	0.00079	0.00075	0.00032	0.00021	0.00019	0.00019	0.00019
Benz[a]anthracene	56-55-3	0.0001	0.00082	0.00043	0.00041	0.00018	0.00012	0.0001	0.0001	0.0001
Benzaldehyde	100-52-7	1.8	15	7.8	7.4	3.2	2.1	1.8	1.9	1.9
Benzene, 1,2,3-trimethyl-	526-73-8	0.27	2.2	1.1	1.1	0.47	0.31	0.27	0.28	0.28
Benzene, 1,2,4,5-tetramethyl-	95-93-2	0.052	0.41	0.22	0.21	0.09	0.059	0.052	0.052	0.052
Benzene, 1,2,4-trimethyl-	95-63-6	2.4	19	10	9.8	4.2	2.8	2.4	2.5	2.5
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	0.11	0.87	0.46	0.44	0.19	0.12	0.11	0.11	0.11
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	0.15	1.2	0.65	0.62	0.27	0.18	0.15	0.16	0.16
Benzene, butyl-	104-51-8	0.054	0.43	0.23	0.22	0.094	0.062	0.054	0.055	0.055
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	0.077	0.62	0.33	0.31	0.13	0.088	0.077	0.078	0.078
Benzo[b]fluoranthene	205-99-2	0.00051	0.0041	0.0021	0.002	0.00088	0.00058	0.00051	0.00052	0.00052
7H-Benzo[c]fluorene	205-12-9	0.0011	0.0089	0.0047	0.0045	0.0019	0.0013	0.0011	0.0011	0.0011
Benzo[k]fluoranthene	207-08-9	0.00063	0.0051	0.0027	0.0025	0.0011	0.00073	0.00063	0.00064	0.00064
Benzothiazole	95-16-9	0.76	6.1	3.2	3.1	1.3	0.88	0.77	0.77	0.78
Benzothiazole, 2-phenyl-	883-93-2	0.061	0.49	0.26	0.24	0.11	0.07	0.061	0.062	0.062
2-Benzothiazolone	934-34-9	0.095	0.76	0.4	0.38	0.17	0.11	0.095	0.096	0.097
Benzyl butyl phthalate	85-68-7	0.071	0.57	0.3	0.29	0.12	0.081	0.071	0.072	0.072



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Chronic C _{inh}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Butanal	123-72-8	6.3	51	27	25	11	7.2	6.3	6.4	6.4
Cyclopentasiloxane, decamethyl-	541-02-6	3.3	27	14	13	5.8	3.8	3.3	3.4	3.4
Cyclotetrasiloxane, octamethyl-	556-67-2	1.2	9.6	5	4.8	2.1	1.4	1.2	1.2	1.2
p-Cymene	99-87-6	0.52	4.2	2.2	2.1	0.9	0.59	0.52	0.52	0.53
Decane	124-18-5	1.1	9	4.7	4.5	1.9	1.3	1.1	1.1	1.1
Dibenz[a,h]anthracene	53-70-3	0.003	0.024	0.012	0.012	0.0051	0.0034	0.003	0.003	0.003
Dibenzothiophene	132-65-0	0.025	0.2	0.11	0.1	0.044	0.029	0.026	0.026	0.026
Dibutyl phthalate	84-74-2	7.7	62	33	31	13	8.9	7.8	7.9	7.9
Diethyl phthalate	84-66-2	0.054	0.44	0.23	0.22	0.094	0.062	0.054	0.055	0.055
Diisobutyl phthalate	84-69-5	0.29	2.4	1.2	1.2	0.51	0.34	0.29	0.3	0.3
Diisooctylphthalate	27554-26-3	1	8.2	4.3	4.1	1.8	1.2	1	1	1
Di-n-octyl phthalate	117-84-0	0.002	0.016	0.0084	0.008	0.0035	0.0023	0.002	0.002	0.002
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	0.67	5.4	2.8	2.7	1.2	0.77	0.67	0.68	0.68
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	0.35	2.8	1.5	1.4	0.61	0.4	0.35	0.35	0.36
Dodecane	112-40-3	0.14	1.1	0.59	0.56	0.24	0.16	0.14	0.14	0.14
Fluoranthene	206-44-0	0.078	0.63	0.33	0.31	0.14	0.089	0.078	0.079	0.079
Fluorene	86-73-7	0.12	0.99	0.52	0.5	0.21	0.14	0.12	0.12	0.13
Furan, 2-methyl	534-22-5	2.2	18	9.3	8.9	3.8	2.5	2.2	2.2	2.2
Heptanal	111-71-7	0.3	2.4	1.3	1.2	0.53	0.35	0.3	0.31	0.31
Hexadecane	544-76-3	0.65	5.3	2.8	2.6	1.1	0.75	0.66	0.66	0.67
2,5-Hexanedione	110-13-4	0.56	4.5	2.4	2.3	0.97	0.64	0.56	0.57	0.57
Indan	496-11-7	0.28	2.3	1.2	1.1	0.49	0.32	0.28	0.28	0.29
Mesitylene	108-67-8	0.57	4.6	2.4	2.3	0.99	0.65	0.57	0.58	0.58
Methacrolein	78-85-3	1.6	13	6.6	6.3	2.7	1.8	1.6	1.6	1.6
Methyl Isobutyl Ketone	108-10-1	0.32	2.6	1.4	1.3	0.56	0.37	0.32	0.33	0.33
Naphthalene	91-20-3	0.55	4.4	2.3	2.2	0.95	0.63	0.55	0.56	0.56
Naphthalene, 1,2-dimethyl-	573-98-8	0.0077	0.062	0.032	0.031	0.013	0.0088	0.0077	0.0078	0.0078



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Chronic C _{inh}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Naphthalene, 1,6-dimethyl-	575-43-9	0.055	0.44	0.23	0.22	0.095	0.063	0.055	0.056	0.056
Naphthalene, 1-methyl-	90-12-0	0.45	3.6	1.9	1.8	0.78	0.51	0.45	0.45	0.45
Naphthalene, 2-(bromomethyl)-	939-26-4	0.015	0.12	0.064	0.061	0.026	0.017	0.015	0.015	0.015
Naphthalene, 2,3-dimethyl-	581-40-8	0.041	0.33	0.17	0.17	0.072	0.047	0.041	0.042	0.042
Naphthalene, 2-methyl-	91-57-6	0.68	5.5	2.9	2.8	1.2	0.78	0.68	0.69	0.69
1-Octadecene	112-88-9	0.091	0.73	0.38	0.37	0.16	0.1	0.091	0.092	0.093
Octanal	124-13-0	0.92	7.4	3.9	3.7	1.6	1.1	0.93	0.94	0.94
Octane	111-65-9	1.2	10	5.3	5	2.2	1.4	1.3	1.3	1.3
17-Pentatriacontene	6971-40-0	0.015	0.12	0.064	0.061	0.026	0.017	0.015	0.015	0.015
N-Phenylbenzamide	93-98-1	0.18	1.5	0.78	0.75	0.32	0.21	0.19	0.19	0.19
Phenanthrene	85-01-8	0.27	2.2	1.1	1.1	0.47	0.31	0.27	0.27	0.27
Phenanthrene, 1-methyl	832-69-9	0.017	0.14	0.072	0.068	0.029	0.019	0.017	0.017	0.017
Phenanthrene, 2-methyl-	2531-84-2	0.031	0.25	0.13	0.12	0.054	0.035	0.031	0.031	0.031
Phenanthrene, 3-methyl	832-71-3	0.037	0.3	0.16	0.15	0.064	0.042	0.037	0.037	0.037
Propionaldehyde	123-38-6	3.8	31	16	15	6.6	4.4	3.8	3.9	3.9
Pyrene	129-00-0	0.066	0.53	0.28	0.27	0.11	0.076	0.066	0.067	0.067
Pyridine, 2-(4-methylphenyl)-	4467-06-5	0.00073	0.0059	0.0031	0.0029	0.0013	0.00084	0.00073	0.00074	0.00074
Resorcinol	108-46-3	0.39	3.1	1.6	1.6	0.67	0.44	0.39	0.39	0.39
m-Tolualdehyde	620-23-5	5.6	45	23	22	9.7	6.4	5.6	5.6	5.7
TXIB "Kodaflex"	6846-50-0	0.041	0.33	0.17	0.16	0.071	0.047	0.041	0.041	0.041
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	0.078	0.63	0.33	0.31	0.14	0.09	0.078	0.079	0.079
Undecane	1120-21-4	0.27	2.2	1.1	1.1	0.47	0.31	0.27	0.27	0.28
Valeraldehyde	110-62-3	19	150	81	78	33	22	19	20	20
Non-Field-Related Chemicals										
Benzene	71-43-2	12	100	53	50	22	14	13	13	13
Benzene, 1,4-dichloro	106-46-7	0.4	3.2	1.7	1.6	0.69	0.45	0.4	0.4	0.4



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Chronic C _{inh}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	12	95	50	48	21	14	12	12	12
2-Butoxyethanol	111-76-2	0.1	0.82	0.43	0.41	0.18	0.12	0.1	0.1	0.1
Cyclotrisiloxane, hexamethyl-	541-05-9	1.5	12	6.4	6.1	2.6	1.7	1.5	1.5	1.5
Decanal	112-31-2	0.58	4.6	2.4	2.3	1	0.66	0.58	0.59	0.59
Ethylbenzene	100-41-4	3.5	28	15	14	6.2	4.1	3.6	3.6	3.6
Heptane	142-82-5	4.8	38	20	19	8.3	5.5	4.8	4.8	4.8
Hexanal	66-25-1	16	130	69	65	28	19	16	16	17
Hexane	110-54-3	14	110	58	56	24	16	14	14	14
1-Hexanol, 2-ethyl-	104-76-7	0.16	1.3	0.68	0.65	0.28	0.18	0.16	0.16	0.16
Nonanal	124-19-6	0.16	1.3	0.68	0.65	0.28	0.18	0.16	0.16	0.16
Phenol	108-95-2	1.2	9.7	5.1	4.9	2.1	1.4	1.2	1.2	1.2
Tetrachloroethylene	127-18-4	0.98	7.9	4.2	4	1.7	1.1	0.99	1	1
Tetradecane	629-59-4	0.17	1.3	0.71	0.67	0.29	0.19	0.17	0.17	0.17
Texanol, TXIB (mono-isomer)	25265-77-4	2.1	17	9.1	8.7	3.7	2.5	2.2	2.2	2.2
Toluene	108-88-3	28	220	120	110	49	32	28	28	28
Trichloroethylene	79-01-6	0.2	1.6	0.85	0.81	0.35	0.23	0.2	0.2	0.2
Trichloromethane	67-66-3	0.78	6.3	3.3	3.2	1.4	0.9	0.79	0.8	0.8
m/p-Xylene	106-42-3	12	96	50	48	21	14	12	12	12
o-Xylene	95-47-6	4	32	17	16	6.9	4.6	4	4	4.1

^a On-Field Chronic C_{inh} equals the modified mean of the 35 individual field average concentrations (C_{air-avg} x AF_{inh}) of a chemical detected in air on the fields (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-146. **Off-Field** Chronic Inhalation Exposure Concentration^a for **All General Chemicals** (Chronic C_{inh}, nanograms per cubic meter)—Combined Gender **Spectators**

Chemical	CASRN	Chronic C _{inh}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Field-Related Chemicals										



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Chronic C _{inh}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Acenaphthylene	208-96-8	0.022	0.18	0.092	0.088	0.038	0.025	0.022	0.022	0.022
Aniline	62-53-3	0.15	1.2	0.62	0.6	0.26	0.17	0.15	0.15	0.15
Anthracene	120-12-7	0.0041	0.033	0.018	0.017	0.0072	0.0048	0.0042	0.0042	0.0042
Anthracene, 2-methyl-	613-12-7	0.0011	0.0087	0.0046	0.0044	0.0019	0.0012	0.0011	0.0011	0.0011
Anthracene, 9,10-dimethyl	781-43-1	0.0017	0.014	0.0073	0.007	0.003	0.002	0.0017	0.0017	0.0018
Anthracene, 9-phenyl	602-55-1	0.00077	0.0062	0.0032	0.0031	0.0013	0.00088	0.00077	0.00078	0.00078
Benz[a]anthracene	56-55-3	0	0	0	0	0	0	0	0	0
Benzaldehyde	100-52-7	2	16	8.5	8.1	3.5	2.3	2	2	2
Benzene, 1,2,3-trimethyl-	526-73-8	0.34	2.8	1.5	1.4	0.6	0.4	0.35	0.35	0.35
Benzene, 1,2,4,5-tetramethyl-	95-93-2	0.066	0.53	0.28	0.26	0.11	0.075	0.066	0.067	0.067
Benzene, 1,2,4-trimethyl-	95-63-6	2.6	21	11	11	4.6	3	2.6	2.7	2.7
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	0.12	0.99	0.52	0.5	0.21	0.14	0.12	0.12	0.12
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	0.17	1.4	0.73	0.7	0.3	0.2	0.17	0.18	0.18
Benzene, butyl-	104-51-8	0.089	0.72	0.38	0.36	0.16	0.1	0.089	0.09	0.091
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	0.087	0.7	0.37	0.35	0.15	0.099	0.087	0.088	0.088
Benzo[b]fluoranthene	205-99-2	0.00079	0.0063	0.0033	0.0032	0.0014	0.0009	0.00079	0.0008	0.0008
7H-Benzo[c]fluorene	205-12-9	0.00065	0.0052	0.0027	0.0026	0.0011	0.00074	0.00065	0.00066	0.00066
Benzo[k]fluoranthene	207-08-9	0.00065	0.0052	0.0028	0.0026	0.0011	0.00075	0.00065	0.00066	0.00066
Benzothiazole	95-16-9	0.1	0.8	0.42	0.4	0.17	0.11	0.1	0.1	0.1
Benzothiazole, 2-phenyl-	883-93-2	0.0089	0.071	0.037	0.036	0.015	0.01	0.0089	0.009	0.009
2-Benzothiazolone	934-34-9	0.084	0.67	0.35	0.34	0.15	0.096	0.084	0.085	0.085
Benzyl butyl phthalate	85-68-7	0.09	0.73	0.38	0.36	0.16	0.1	0.091	0.092	0.092
Butanal	123-72-8	5.9	47	25	24	10	6.7	5.9	6	6
Cyclopentasiloxane, decamethyl-	541-02-6	3.1	25	13	12	5.3	3.5	3.1	3.1	3.1
Cyclotetrasiloxane, octamethyl-	556-67-2	1.1	8.6	4.5	4.3	1.9	1.2	1.1	1.1	1.1
p-Cymene	99-87-6	0.54	4.3	2.3	2.2	0.94	0.62	0.54	0.55	0.55



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Chronic C _{inh}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Decane	124-18-5	1.2	10	5.2	5	2.2	1.4	1.2	1.3	1.3
Dibenz[a,h]anthracene	53-70-3	0.0019	0.015	0.0078	0.0075	0.0032	0.0021	0.0019	0.0019	0.0019
Dibenzothiophene	132-65-0	0.024	0.19	0.1	0.098	0.042	0.028	0.024	0.025	0.025
Dibutyl phthalate	84-74-2	6.2	50	26	25	11	7.2	6.3	6.3	6.3
Diethyl phthalate	84-66-2	0.22	1.8	0.94	0.9	0.39	0.26	0.22	0.23	0.23
Diisobutyl phthalate	84-69-5	0.32	2.5	1.3	1.3	0.55	0.36	0.32	0.32	0.32
Diisooctylphthalate	27554-26-3	0.3	2.4	1.3	1.2	0.53	0.35	0.31	0.31	0.31
Di-n-octyl phthalate	117-84-0	0.002	0.016	0.0087	0.0083	0.0036	0.0024	0.0021	0.0021	0.0021
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	0.81	6.5	3.4	3.3	1.4	0.93	0.81	0.82	0.83
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	0.23	1.8	0.96	0.91	0.4	0.26	0.23	0.23	0.23
Dodecane	112-40-3	0.18	1.4	0.76	0.72	0.31	0.21	0.18	0.18	0.18
Fluoranthene	206-44-0	0.046	0.37	0.19	0.19	0.08	0.053	0.046	0.047	0.047
Fluorene	86-73-7	0.1	0.83	0.43	0.41	0.18	0.12	0.1	0.1	0.1
Furan, 2-methyl	534-22-5	0.78	6.3	3.3	3.2	1.4	0.9	0.79	0.79	0.8
Heptanal	111-71-7	0.31	2.5	1.3	1.3	0.54	0.36	0.31	0.32	0.32
Hexadecane	544-76-3	0.94	7.6	4	3.8	1.6	1.1	0.94	0.95	0.96
2,5-Hexanedione	110-13-4	0.73	5.9	3.1	2.9	1.3	0.84	0.73	0.74	0.74
Indan	496-11-7	0.28	2.2	1.2	1.1	0.48	0.32	0.28	0.28	0.28
Mesitylene	108-67-8	0.59	4.7	2.5	2.4	1	0.67	0.59	0.59	0.6
Methacrolein	78-85-3	1.4	12	6.1	5.8	2.5	1.6	1.4	1.5	1.5
Methyl Isobutyl Ketone	108-10-1	0.074	0.59	0.31	0.3	0.13	0.085	0.074	0.075	0.075
Naphthalene	91-20-3	0.6	4.8	2.5	2.4	1	0.69	0.6	0.61	0.61
Naphthalene, 1,2-dimethyl-	573-98-8	0.016	0.13	0.068	0.065	0.028	0.018	0.016	0.016	0.016
Naphthalene, 1,6-dimethyl-	575-43-9	0.057	0.46	0.24	0.23	0.099	0.066	0.057	0.058	0.058
Naphthalene, 1-methyl-	90-12-0	0.44	3.6	1.9	1.8	0.77	0.51	0.45	0.45	0.45
Naphthalene, 2-(bromomethyl)-	939-26-4	0.016	0.13	0.067	0.064	0.028	0.018	0.016	0.016	0.016
Naphthalene, 2,3-dimethyl-	581-40-8	0.043	0.34	0.18	0.17	0.075	0.049	0.043	0.043	0.044
Naphthalene, 2-methyl-	91-57-6	0.73	5.8	3.1	2.9	1.3	0.83	0.73	0.74	0.74
1-Octadecene	112-88-9	0.084	0.67	0.35	0.34	0.15	0.096	0.084	0.085	0.085



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Chronic C _{inh}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Octanal	124-13-0	0.9	7.2	3.8	3.6	1.6	1	0.9	0.91	0.91
Octane	111-65-9	1.5	12	6.3	6	2.6	1.7	1.5	1.5	1.5
17-Pentatriacontene	6971-40-0	0.01	0.083	0.044	0.042	0.018	0.012	0.01	0.011	0.011
N-Phenylbenzamide	93-98-1	0.18	1.5	0.77	0.74	0.32	0.21	0.18	0.19	0.19
Phenanthrene	85-01-8	0.22	1.8	0.94	0.9	0.39	0.26	0.22	0.23	0.23
Phenanthrene, 1-methyl	832-69-9	0.01	0.082	0.043	0.041	0.018	0.012	0.01	0.01	0.01
Phenanthrene, 2-methyl-	2531-84-2	0.02	0.16	0.083	0.08	0.034	0.023	0.02	0.02	0.02
Phenanthrene, 3-methyl	832-71-3	0.023	0.18	0.095	0.091	0.039	0.026	0.023	0.023	0.023
Pyrene	129-00-0	0.033	0.26	0.14	0.13	0.057	0.038	0.033	0.033	0.033
Pyridine, 2-(4-methylphenyl)-	4467-06-5	0.00011	0.00089	0.00047	0.00045	0.00019	0.00013	0.00011	0.00011	0.00011
Resorcinol	108-46-3	0.66	5.3	2.8	2.7	1.2	0.76	0.66	0.67	0.67
TXIB "Kodaflex"	6846-50-0	0.084	0.67	0.35	0.34	0.15	0.096	0.084	0.085	0.085
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	0.056	0.45	0.24	0.22	0.097	0.064	0.056	0.056	0.057
Undecane	1120-21-4	0.37	3	1.6	1.5	0.65	0.43	0.37	0.38	0.38
Non-Field-Related Chemicals										
Benzene	71-43-2	13	110	55	53	23	15	13	13	13
Benzene, 1,4-dichloro	106-46-7	0.36	2.9	1.5	1.4	0.63	0.41	0.36	0.36	0.37
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	12	96	50	48	21	14	12	12	12
2-Butoxyethanol	111-76-2	0.098	0.79	0.41	0.4	0.17	0.11	0.098	0.099	0.1
Cyclotrisiloxane, hexamethyl-	541-05-9	1.6	13	6.8	6.4	2.8	1.8	1.6	1.6	1.6
Decanal	112-31-2	0.35	2.8	1.5	1.4	0.62	0.41	0.35	0.36	0.36
Ethylbenzene	100-41-4	3.7	30	16	15	6.5	4.3	3.8	3.8	3.8
Heptane	142-82-5	4.8	38	20	19	8.3	5.5	4.8	4.8	4.8
Hexanal	66-25-1	18	140	75	72	31	21	18	18	18
Hexane	110-54-3	9.4	76	40	38	16	11	9.4	9.5	9.6
1-Hexanol, 2-ethyl-	104-76-7	0.36	2.9	1.5	1.5	0.63	0.42	0.36	0.37	0.37
Nonanal	124-19-6	0	0	0	0	0	0	0	0	0
Phenol	108-95-2	1.1	8.8	4.6	4.4	1.9	1.3	1.1	1.1	1.1
Tetrachloroethylene	127-18-4	1	8.2	4.3	4.1	1.8	1.2	1	1	1



Chemical	CASRN	Chronic C _{inh}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Tetradecane	629-59-4	0.13	1	0.55	0.52	0.23	0.15	0.13	0.13	0.13
Texanol, TXIB (mono-isomer)	25265-77-4	2.1	17	8.8	8.4	3.6	2.4	2.1	2.1	2.1
Toluene	108-88-3	29	230	120	120	50	33	29	29	29
Trichloroethylene	79-01-6	0.2	1.6	0.86	0.82	0.35	0.23	0.2	0.21	0.21
Trichloromethane	67-66-3	0.76	6.1	3.2	3.1	1.3	0.87	0.76	0.77	0.77
m/p-Xylene	106-42-3	13	100	54	51	22	15	13	13	13
o-Xylene	95-47-6	3.9	32	17	16	6.8	4.5	3.9	4	4

^a Off-Field Chronic C_{inh} equals the modified mean of the 35 individual field average concentrations (C_{air-avg} x AF_{inh}) of a chemical detected in air off the fields (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

INDIVIDUAL FIELD ASSESSMENT (Table F-147 to Table F-179)

Table F-147. **On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic C_{inh-field}, nanograms per cubic meter) —Combined Gender Athletes 2<6 Years**

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.16	0.31	0	0.66	1.4
Acetone	67-64-1	34	240	3000	3400	1600	9900	11000
Aniline	62-53-3	10	0	1	1.8	0	3.9	6.8
Anthracene	120-12-7	12	0	0.051	0.095	0	0.22	0.37
Anthracene, 2-methyl-	613-12-7	15	0	0.013	0.019	0	0.051	0.065
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.015	0.027	0	0.042	0.15
Anthracene, 9-phenyl	602-55-1	1	0	0.0014	0.0082	0	0	0.048
Benz[a]anthracene	56-55-3	1	0	0.00076	0.0045	0	0	0.026
Benzaldehyde	100-52-7	15	0	14	21	0	50	84
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	2	6	0	9	32
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.39	1	0	2.4	4.2
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	18	32	0	65	150
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.82	2.1	0	4.2	11
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	1.2	2.3	0	5.1	11
Benzene, butyl-	104-51-8	2	0	0.41	2	0	0.85	11



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.58	0.68	0.35	1.9	2.3
Benzo[b]fluoranthene	205-99-2	2	0	0.0038	0.016	0	0.023	0.065
7H-Benzo[c]fluorene	205-12-9	13	0	0.0084	0.013	0	0.022	0.055
Benzo[k]fluoranthene	207-08-9	4	0	0.0048	0.014	0	0.036	0.054
Benzothiazole	95-16-9	19	0	5.7	6.6	3.2	19	19
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.46	0.6	0.23	1.6	2.7
2-Benzothiazolone	934-34-9	5	0	0.72	1.9	0	4.7	6.9
Benzyl butyl phthalate	85-68-7	7	0	0.53	1.1	0	3.6	3.6
Butanal	123-72-8	13	0	47	120	0	280	590
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	25	42	7.7	89	200
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	9	14	0	35	51
p-Cymene	99-87-6	21	0	3.9	4.7	2.5	13	18
Decane	124-18-5	11	0	8.4	15	0	46	47
Dibenz[a,h]anthracene	53-70-3	14	0	0.022	0.044	0	0.11	0.19
Dibenzothiophene	132-65-0	14	0	0.19	0.29	0	0.82	1.1
Dibutyl phthalate	84-74-2	7	0	58	150	0	260	760
Diethyl phthalate	84-66-2	1	0	0.41	2.4	0	0	14
Diisobutyl phthalate	84-69-5	15	0	2.2	4.3	0	7	23
Diisooctylphthalate	27554-26-3	9	0	7.7	26	0	20	150
Di-n-octyl phthalate	117-84-0	5	0	0.015	0.039	0	0.085	0.17
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	5	6.8	2.6	20	22
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	2.6	5.2	0	13	17
Dodecane	112-40-3	4	0	1	3.5	0	6	18
Fluoranthene	206-44-0	21	0	0.59	0.75	0.21	2	2.6
Fluorene	86-73-7	15	0	0.93	1.8	0	3.3	9.5
Furan, 2-methyl	534-22-5	34	0	17	14	12	39	63
Heptanal	111-71-7	7	0	2.3	5.7	0	20	20
Hexadecane	544-76-3	11	0	4.9	9	0	26	35
2,5-Hexanedione	110-13-4	5	0	4.2	11	0	28	41
Indan	496-11-7	12	0	2.1	3.9	0	9.1	17
Mesitylene	108-67-8	15	0	4.3	7.6	0	17	35
Methacrolein	78-85-3	20	0	12	15	6.7	35	67
Methyl Isobutyl Ketone	108-10-1	5	0	2.4	6.6	0	19	28
Naphthalene	91-20-3	9	0	4.1	8.6	0	18	40
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.058	0.17	0	0.56	0.56



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.41	0.64	0	2	2.3
Naphthalene, 1-methyl-	90-12-0	15	0	3.4	5.8	0	15	23
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.11	0.17	0	0.39	0.39
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.31	0.46	0	1.4	1.6
Naphthalene, 2-methyl-	91-57-6	10	0	5.1	11	0	25	49
1-Octadecene	112-88-9	16	0	0.68	0.88	0	2.4	2.8
Octanal	124-13-0	24	0	6.9	8	4.3	21	32
Octane	111-65-9	10	0	9.4	17	0	39	65
17-Pentatriacontene	6971-40-0	2	0	0.11	0.49	0	0.45	2.6
N-Phenylbenzamide	93-98-1	6	0	1.4	3.1	0	8.6	8.6
Phenanthrene	85-01-8	17	0	2	3.1	1.1	8.8	13
Phenanthrene, 1-methyl	832-69-9	17	0	0.13	0.17	0.057	0.46	0.52
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.23	0.3	0.16	0.85	0.97
Phenanthrene, 3-methyl	832-71-3	18	0	0.28	0.36	0.15	1	1.2
Propionaldehyde	123-38-6	12	0	29	58	0	110	280
Pyrene	129-00-0	20	0	0.5	0.63	0.2	1.8	2.1
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.0055	0.028	0	0.0096	0.16
Resorcinol	108-46-3	18	0	2.9	4.2	1.2	9.5	18
m-Tolualdehyde	620-23-5	19	0	42	44	34	110	140
TXIB "Kodaflex"	6846-50-0	1	0	0.31	1.8	0	0	11
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.59	0.82	0	2.1	3
Undecane	1120-21-4	5	0	2	5.3	0	16	20
Valeraldehyde	110-62-3	11	0	140	250	0	650	710
Non-Field-Related Chemicals								
Benzene	71-43-2	35	14	94	76	66	230	380
Benzene, 1,4-dichloro	106-46-7	7	0	3	6.5	0	18	18
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	89	93	59	270	330
2-Butoxyethanol	111-76-2	1	0	0.77	4.5	0	0	27
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	11	67	0	0	400
Decanal	112-31-2	7	0	4.4	15	0	16	85
Ethylbenzene	100-41-4	13	0	27	45	0	110	180
Heptane	142-82-5	20	0	36	53	6.9	150	230
Hexanal	66-25-1	30	0	120	160	57	470	620
Hexane	110-54-3	26	0	100	230	46	280	1400
1-Hexanol, 2-ethyl-	104-76-7	6	0	1.2	3.1	0	6.1	14
Nonanal	124-19-6	2	0	1.2	5	0	6.3	21



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Phenol	108-95-2	18	0	9.1	11	5.4	32	32
Tetrachloroethylene	127-18-4	7	0	7.4	18	0	50	65
Tetradecane	629-59-4	3	0	1.3	4.4	0	11	22
m/p-Xylene	106-42-3	25	0	90	130	36	310	540
o-Xylene	95-47-6	13	0	30	50	0	120	200

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-148. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter) —Combined Gender Athletes 6<11 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.2	0.41	0	0.85	1.8
Acetone	67-64-1	34	320	3900	4400	2100	13000	14000
Aniline	62-53-3	10	0	1.3	2.3	0	5.1	8.8
Anthracene	120-12-7	12	0	0.066	0.12	0	0.29	0.49
Anthracene, 2-methyl-	613-12-7	15	0	0.017	0.024	0	0.066	0.084
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.02	0.035	0	0.055	0.19
Anthracene, 9-phenyl	602-55-1	1	0	0.0018	0.011	0	0	0.062
Benz[a]anthracene	56-55-3	1	0	0.00099	0.0058	0	0	0.034
Benzaldehyde	100-52-7	15	0	18	28	0	65	110
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	2.7	7.8	0	12	42
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.5	1.3	0	3.1	5.4
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	24	42	0	84	190
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	1.1	2.8	0	5.4	14
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	1.5	3	0	6.6	14
Benzene, butyl-	104-51-8	2	0	0.53	2.6	0	1.1	15
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.75	0.88	0.45	2.5	3
Benzo[b]fluoranthene	205-99-2	2	0	0.005	0.02	0	0.03	0.084
7H-Benzo[c]fluorene	205-12-9	13	0	0.011	0.016	0	0.028	0.072
Benzo[k]fluoranthene	207-08-9	4	0	0.0062	0.018	0	0.047	0.07
Benzothiazole	95-16-9	19	0	7.4	8.6	4.1	25	25
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.59	0.78	0.3	2	3.5
2-Benzothiazolone	934-34-9	5	0	0.93	2.4	0	6.1	9



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh} -field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzyl butyl phthalate	85-68-7	7	0	0.69	1.5	0	4.7	4.7
Butanal	123-72-8	13	0	62	150	0	360	760
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	32	55	10	120	260
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	12	19	0	46	66
p-Cymene	99-87-6	21	0	5	6.1	3.2	17	24
Decane	124-18-5	11	0	11	19	0	60	62
Dibenz[a,h]anthracene	53-70-3	14	0	0.029	0.058	0	0.14	0.24
Dibenzothiophene	132-65-0	14	0	0.25	0.38	0	1.1	1.4
Dibutyl phthalate	84-74-2	7	0	76	190	0	340	990
Diethyl phthalate	84-66-2	1	0	0.53	3.1	0	0	18
Diisobutyl phthalate	84-69-5	15	0	2.9	5.6	0	9	30
Diisooctylphthalate	27554-26-3	9	0	9.9	33	0	26	190
Di-n-octyl phthalate	117-84-0	5	0	0.019	0.05	0	0.11	0.22
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	6.5	8.8	3.4	26	29
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	3.4	6.7	0	17	22
Dodecane	112-40-3	4	0	1.4	4.6	0	7.7	24
Fluoranthene	206-44-0	21	0	0.76	0.97	0.27	2.6	3.4
Fluorene	86-73-7	15	0	1.2	2.3	0	4.2	12
Furan, 2-methyl	534-22-5	34	0	21	18	16	50	82
Heptanal	111-71-7	7	0	3	7.4	0	26	26
Hexadecane	544-76-3	11	0	6.4	12	0	34	46
2,5-Hexanedione	110-13-4	5	0	5.5	14	0	36	53
Indan	496-11-7	12	0	2.7	5.1	0	12	22
Mesitylene	108-67-8	15	0	5.6	9.9	0	22	46
Methacrolein	78-85-3	20	0	15	19	8.7	45	86
Methyl Isobutyl Ketone	108-10-1	5	0	3.1	8.6	0	24	37
Naphthalene	91-20-3	9	0	5.4	11	0	23	52
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.075	0.21	0	0.73	0.73
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.54	0.82	0	2.5	3
Naphthalene, 1-methyl-	90-12-0	15	0	4.4	7.5	0	20	30
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.15	0.22	0	0.5	0.5
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.4	0.6	0	1.8	2.1
Naphthalene, 2-methyl-	91-57-6	10	0	6.7	14	0	32	63
1-Octadecene	112-88-9	16	0	0.89	1.1	0	3.1	3.7
Octanal	124-13-0	24	0	9	10	5.6	27	42
Octane	111-65-9	10	0	12	22	0	51	85
17-Pentatriacontene	6971-40-0	2	0	0.15	0.63	0	0.58	3.3



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
N-Phenylbenzamide	93-98-1	6	0	1.8	4.1	0	11	11
Phenanthrene	85-01-8	17	0	2.6	4	1.4	11	17
Phenanthrene, 1-methyl	832-69-9	17	0	0.17	0.21	0.075	0.6	0.68
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.3	0.39	0.2	1.1	1.3
Phenanthrene, 3-methyl	832-71-3	18	0	0.36	0.46	0.19	1.3	1.5
Propionaldehyde	123-38-6	12	0	37	75	0	150	360
Pyrene	129-00-0	20	0	0.64	0.82	0.25	2.3	2.8
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.0071	0.036	0	0.013	0.21
Resorcinol	108-46-3	18	0	3.8	5.5	1.5	12	24
m-Tolualdehyde	620-23-5	19	0	54	57	44	150	180
TXIB "Kodaflex"	6846-50-0	1	0	0.4	2.4	0	0	14
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.76	1.1	0	2.7	3.9
Undecane	1120-21-4	5	0	2.6	6.9	0	20	26
Valeraldehyde	110-62-3	11	0	190	320	0	850	920
Non-Field-Related Chemicals								
Benzene	71-43-2	35	18	120	98	86	300	490
Benzene, 1,4-dichloro	106-46-7	7	0	3.9	8.5	0	24	24
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	120	120	77	340	430
2-Butoxyethanol	111-76-2	1	0	1	5.9	0	0	35
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	15	87	0	0	520
Decanal	112-31-2	7	0	5.6	19	0	21	110
Ethylbenzene	100-41-4	13	0	35	58	0	140	230
Heptane	142-82-5	20	0	46	69	8.9	190	300
Hexanal	66-25-1	30	0	160	200	74	610	800
Hexane	110-54-3	26	0	130	300	60	360	1800
1-Hexanol, 2-ethyl-	104-76-7	6	0	1.6	4	0	7.9	18
Nonanal	124-19-6	2	0	1.6	6.5	0	8.2	27
Phenol	108-95-2	18	0	12	15	6.9	42	42
Tetrachloroethylene	127-18-4	7	0	9.6	23	0	64	84
Tetradecane	629-59-4	3	0	1.6	5.8	0	14	29
m/p-Xylene	106-42-3	25	0	120	170	47	400	700
o-Xylene	95-47-6	13	0	39	65	0	150	260

^a 35 field-specific C_{inh-field} values are included in the table.

^b On-Field field-specific Chronic C_{inh-field} equals the modified field-specific average concentration of a field (C_{air-field} x AF_{inh}) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-149. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic C_{inh-field}, nanograms per cubic meter) —Combined Gender Athletes 11<16 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.2	0.4	0	0.83	1.7
Acetone	67-64-1	34	310	3800	4300	2100	13000	14000
Aniline	62-53-3	10	0	1.3	2.2	0	5	8.6
Anthracene	120-12-7	12	0	0.064	0.12	0	0.28	0.47
Anthracene, 2-methyl-	613-12-7	15	0	0.017	0.024	0	0.065	0.082
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.019	0.034	0	0.053	0.19
Anthracene, 9-phenyl	602-55-1	1	0	0.0018	0.01	0	0	0.06
Benz[a]anthracene	56-55-3	1	0	0.00097	0.0057	0	0	0.033
Benzaldehyde	100-52-7	15	0	18	27	0	63	110
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	2.6	7.7	0	11	41
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.49	1.3	0	3	5.3
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	23	41	0	82	190
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	1	2.7	0	5.3	14
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	1.5	2.9	0	6.5	13
Benzene, butyl-	104-51-8	2	0	0.52	2.5	0	1.1	14
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.73	0.86	0.44	2.4	2.9
Benzo[b]fluoranthene	205-99-2	2	0	0.0048	0.02	0	0.029	0.082
7H-Benzo[c]fluorene	205-12-9	13	0	0.011	0.016	0	0.028	0.07
Benzo[k]fluoranthene	207-08-9	4	0	0.006	0.018	0	0.046	0.068
Benzothiazole	95-16-9	19	0	7.3	8.4	4	24	24
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.58	0.76	0.29	2	3.4
2-Benzothiazolone	934-34-9	5	0	0.91	2.4	0	5.9	8.8
Benzyl butyl phthalate	85-68-7	7	0	0.67	1.4	0	4.6	4.6
Butanal	123-72-8	13	0	60	150	0	350	740
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	32	54	9.7	110	260
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	11	18	0	45	64
p-Cymene	99-87-6	21	0	4.9	6	3.1	16	23
Decane	124-18-5	11	0	11	19	0	58	60
Dibenz[a,h]anthracene	53-70-3	14	0	0.028	0.056	0	0.14	0.24
Dibenzothiophene	132-65-0	14	0	0.24	0.37	0	1	1.3
Dibutyl phthalate	84-74-2	7	0	74	190	0	330	970
Diethyl phthalate	84-66-2	1	0	0.52	3	0	0	18



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Diisobutyl phthalate	84-69-5	15	0	2.8	5.5	0	8.8	30
Diisooctylphthalate	27554-26-3	9	0	9.7	32	0	25	190
Di-n-octyl phthalate	117-84-0	5	0	0.019	0.049	0	0.11	0.21
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	6.4	8.6	3.3	25	28
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	3.3	6.5	0	17	22
Dodecane	112-40-3	4	0	1.3	4.5	0	7.5	23
Fluoranthene	206-44-0	21	0	0.74	0.95	0.27	2.6	3.4
Fluorene	86-73-7	15	0	1.2	2.3	0	4.1	12
Furan, 2-methyl	534-22-5	34	0	21	18	15	49	80
Heptanal	111-71-7	7	0	2.9	7.3	0	25	25
Hexadecane	544-76-3	11	0	6.2	11	0	33	45
2,5-Hexanedione	110-13-4	5	0	5.3	14	0	35	52
Indan	496-11-7	12	0	2.7	5	0	12	21
Mesitylene	108-67-8	15	0	5.4	9.7	0	22	45
Methacrolein	78-85-3	20	0	15	19	8.5	44	84
Methyl Isobutyl Ketone	108-10-1	5	0	3.1	8.4	0	24	36
Naphthalene	91-20-3	9	0	5.2	11	0	23	51
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.073	0.21	0	0.71	0.71
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.52	0.81	0	2.5	2.9
Naphthalene, 1-methyl-	90-12-0	15	0	4.3	7.3	0	19	29
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.15	0.22	0	0.49	0.49
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.39	0.58	0	1.8	2.1
Naphthalene, 2-methyl-	91-57-6	10	0	6.5	14	0	32	62
1-Octadecene	112-88-9	16	0	0.87	1.1	0	3	3.6
Octanal	124-13-0	24	0	8.8	10	5.4	27	41
Octane	111-65-9	10	0	12	22	0	49	83
17-Pentatriacontene	6971-40-0	2	0	0.14	0.62	0	0.57	3.3
N-Phenylbenzamide	93-98-1	6	0	1.8	4	0	11	11
Phenanthrene	85-01-8	17	0	2.6	4	1.4	11	16
Phenanthrene, 1-methyl	832-69-9	17	0	0.16	0.21	0.073	0.58	0.66
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.29	0.38	0.2	1.1	1.2
Phenanthrene, 3-methyl	832-71-3	18	0	0.35	0.45	0.19	1.3	1.5
Propionaldehyde	123-38-6	12	0	36	73	0	140	350
Pyrene	129-00-0	20	0	0.63	0.8	0.25	2.3	2.7
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.007	0.035	0	0.012	0.2
Resorcinol	108-46-3	18	0	3.7	5.3	1.5	12	23
m-Tolualdehyde	620-23-5	19	0	53	56	43	150	180



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
TXIB "Kodaflex"	6846-50-0	1	0	0.39	2.3	0	0	14
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.74	1	0	2.6	3.8
Undecane	1120-21-4	5	0	2.6	6.7	0	20	26
Valeraldehyde	110-62-3	11	0	180	320	0	830	900
Non-Field-Related Chemicals								
Benzene	71-43-2	35	18	120	96	84	300	480
Benzene, 1,4-dichloro	106-46-7	7	0	3.8	8.3	0	23	23
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	110	120	75	340	420
2-Butoxyethanol	111-76-2	1	0	0.97	5.8	0	0	34
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	14	85	0	0	500
Decanal	112-31-2	7	0	5.5	19	0	21	110
Ethylbenzene	100-41-4	13	0	34	57	0	140	230
Heptane	142-82-5	20	0	45	68	8.7	190	290
Hexanal	66-25-1	30	0	150	200	72	600	780
Hexane	110-54-3	26	0	130	290	58	350	1700
1-Hexanol, 2-ethyl-	104-76-7	6	0	1.5	3.9	0	7.8	18
Nonanal	124-19-6	2	0	1.5	6.3	0	8	27
Phenol	108-95-2	18	0	11	15	6.8	41	41
Tetrachloroethylene	127-18-4	7	0	9.4	22	0	63	82
Tetradecane	629-59-4	3	0	1.6	5.6	0	14	28
m/p-Xylene	106-42-3	25	0	110	160	46	390	680
o-Xylene	95-47-6	13	0	38	64	0	150	250

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-150. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter) —Combined Gender Athletes 16<30 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.37	0.75	0	1.6	3.2
Acetone	67-64-1	34	580	7300	8100	3900	24000	26000
Aniline	62-53-3	10	0	2.4	4.2	0	9.3	16
Anthracene	120-12-7	12	0	0.12	0.23	0	0.53	0.89



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh} -field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Anthracene, 2-methyl-	613-12-7	15	0	0.032	0.045	0	0.12	0.16
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.036	0.065	0	0.1	0.35
Anthracene, 9-phenyl	602-55-1	1	0	0.0033	0.02	0	0	0.11
Benz[a]anthracene	56-55-3	1	0	0.0018	0.011	0	0	0.062
Benzaldehyde	100-52-7	15	0	33	51	0	120	200
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	4.9	14	0	22	77
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.93	2.4	0	5.7	10
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	44	77	0	150	360
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	1.9	5.1	0	10	26
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	2.8	5.6	0	12	25
Benzene, butyl-	104-51-8	2	0	0.97	4.7	0	2	27
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	1.4	1.6	0.83	4.5	5.4
Benzo[b]fluoranthene	205-99-2	2	0	0.0091	0.037	0	0.054	0.16
7H-Benzo[c]fluorene	205-12-9	13	0	0.02	0.03	0	0.052	0.13
Benzo[k]fluoranthene	207-08-9	4	0	0.011	0.034	0	0.087	0.13
Benzothiazole	95-16-9	19	0	14	16	7.6	46	46
Benzothiazole, 2-phenyl-	883-93-2	27	0	1.1	1.4	0.55	3.8	6.5
2-Benzothiazolone	934-34-9	5	0	1.7	4.5	0	11	17
Benzyl butyl phthalate	85-68-7	7	0	1.3	2.7	0	8.6	8.6
Butanal	123-72-8	13	0	110	280	0	670	1400
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	60	100	18	210	480
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	21	34	0	84	120
p-Cymene	99-87-6	21	0	9.3	11	5.9	31	44
Decane	124-18-5	11	0	20	36	0	110	110
Dibenz[a,h]anthracene	53-70-3	14	0	0.053	0.11	0	0.27	0.45
Dibenzothiophene	132-65-0	14	0	0.46	0.7	0	1.9	2.5
Dibutyl phthalate	84-74-2	7	0	140	360	0	630	1800
Diethyl phthalate	84-66-2	1	0	0.97	5.7	0	0	33
Diisobutyl phthalate	84-69-5	15	0	5.3	10	0	17	56
Diisooctylphthalate	27554-26-3	9	0	18	61	0	48	350
Di-n-octyl phthalate	117-84-0	5	0	0.036	0.093	0	0.2	0.4
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	12	16	6.2	47	54
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	6.3	12	0	31	41
Dodecane	112-40-3	4	0	2.5	8.4	0	14	44



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh} -field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Fluoranthene	206-44-0	21	0	1.4	1.8	0.5	4.9	6.3
Fluorene	86-73-7	15	0	2.2	4.3	0	7.8	23
Furan, 2-methyl	534-22-5	34	0	39	33	29	93	150
Heptanal	111-71-7	7	0	5.5	14	0	48	48
Hexadecane	544-76-3	11	0	12	21	0	63	84
2,5-Hexanedione	110-13-4	5	0	10	26	0	66	98
Indan	496-11-7	12	0	5	9.4	0	22	40
Mesitylene	108-67-8	15	0	10	18	0	41	84
Methacrolein	78-85-3	20	0	28	36	16	83	160
Methyl Isobutyl Ketone	108-10-1	5	0	5.8	16	0	45	68
Naphthalene	91-20-3	9	0	9.9	21	0	43	96
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.14	0.4	0	1.3	1.3
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.99	1.5	0	4.7	5.5
Naphthalene, 1-methyl-	90-12-0	15	0	8	14	0	36	55
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.27	0.41	0	0.93	0.93
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.74	1.1	0	3.4	3.9
Naphthalene, 2-methyl-	91-57-6	10	0	12	26	0	60	120
1-Octadecene	112-88-9	16	0	1.6	2.1	0	5.7	6.7
Octanal	124-13-0	24	0	17	19	10	50	77
Octane	111-65-9	10	0	22	41	0	93	160
17-Pentatriacontene	6971-40-0	2	0	0.27	1.2	0	1.1	6.1
N-Phenylbenzamide	93-98-1	6	0	3.3	7.5	0	21	21
Phenanthrene	85-01-8	17	0	4.8	7.4	2.6	21	31
Phenanthrene, 1-methyl	832-69-9	17	0	0.3	0.39	0.14	1.1	1.3
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.55	0.72	0.37	2	2.3
Phenanthrene, 3-methyl	832-71-3	18	0	0.66	0.85	0.36	2.4	2.8
Propionaldehyde	123-38-6	12	0	68	140	0	270	660
Pyrene	129-00-0	20	0	1.2	1.5	0.47	4.3	5.1
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.013	0.066	0	0.023	0.38
Resorcinol	108-46-3	18	0	7	10	2.8	23	43
m-Tolualdehyde	620-23-5	19	0	100	110	81	270	330
TXIB "Kodaflex"	6846-50-0	1	0	0.73	4.3	0	0	26
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	1.4	2	0	4.9	7.2
Undecane	1120-21-4	5	0	4.9	13	0	38	48
Valeraldehyde	110-62-3	11	0	350	600	0	1600	1700
Non-Field-Related Chemicals								
Benzene	71-43-2	35	33	220	180	160	560	910
Benzene, 1,4-dichloro	106-46-7	7	0	7.1	16	0	44	44



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	210	220	140	630	780
2-Butoxyethanol	111-76-2	1	0	1.8	11	0	0	64
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	27	160	0	0	950
Decanal	112-31-2	7	0	10	35	0	39	200
Ethylbenzene	100-41-4	13	0	64	110	0	260	430
Heptane	142-82-5	20	0	85	130	16	350	550
Hexanal	66-25-1	30	0	290	370	140	1100	1500
Hexane	110-54-3	26	0	250	550	110	660	3200
1-Hexanol, 2-ethyl-	104-76-7	6	0	2.9	7.4	0	15	34
Nonanal	124-19-6	2	0	2.9	12	0	15	50
Phenol	108-95-2	18	0	22	27	13	77	77
Tetrachloroethylene	127-18-4	7	0	18	42	0	120	150
Tetradecane	629-59-4	3	0	3	11	0	26	53
m/p-Xylene	106-42-3	25	0	210	310	87	730	1300
o-Xylene	95-47-6	13	0	72	120	0	280	470

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-151. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter) —Combined Gender Athletes 30<40 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.24	0.49	0	1	2.1
Acetone	67-64-1	34	380	4700	5300	2600	15000	17000
Aniline	62-53-3	10	0	1.6	2.8	0	6.1	11
Anthracene	120-12-7	12	0	0.079	0.15	0	0.35	0.58
Anthracene, 2-methyl-	613-12-7	15	0	0.021	0.029	0	0.079	0.1
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.024	0.042	0	0.066	0.23
Anthracene, 9-phenyl	602-55-1	1	0	0.0022	0.013	0	0	0.074
Benz[a]anthracene	56-55-3	1	0	0.0012	0.007	0	0	0.041
Benzaldehyde	100-52-7	15	0	22	33	0	78	130
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	3.2	9.4	0	14	51
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.61	1.6	0	3.7	6.5
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	28	50	0	100	230



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	1.3	3.3	0	6.6	17
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	1.8	3.6	0	8	17
Benzene, butyl-	104-51-8	2	0	0.63	3.1	0	1.3	18
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.9	1.1	0.54	2.9	3.6
Benzo[b]fluoranthene	205-99-2	2	0	0.006	0.024	0	0.036	0.1
7H-Benzo[c]fluorene	205-12-9	13	0	0.013	0.02	0	0.034	0.086
Benzo[k]fluoranthene	207-08-9	4	0	0.0074	0.022	0	0.057	0.084
Benzothiazole	95-16-9	19	0	9	10	5	30	30
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.71	0.94	0.36	2.5	4.2
2-Benzothiazolone	934-34-9	5	0	1.1	2.9	0	7.3	11
Benzyl butyl phthalate	85-68-7	7	0	0.83	1.8	0	5.6	5.6
Butanal	123-72-8	13	0	74	180	0	440	910
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	39	66	12	140	320
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	14	22	0	55	79
p-Cymene	99-87-6	21	0	6.1	7.4	3.8	20	29
Decane	124-18-5	11	0	13	23	0	72	74
Dibenz[a,h]anthracene	53-70-3	14	0	0.035	0.069	0	0.17	0.29
Dibenzothiophene	132-65-0	14	0	0.3	0.46	0	1.3	1.7
Dibutyl phthalate	84-74-2	7	0	91	230	0	410	1200
Diethyl phthalate	84-66-2	1	0	0.64	3.7	0	0	22
Diisobutyl phthalate	84-69-5	15	0	3.4	6.8	0	11	36
Diisooctylphthalate	27554-26-3	9	0	12	40	0	31	230
Di-n-octyl phthalate	117-84-0	5	0	0.023	0.061	0	0.13	0.26
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	7.8	11	4.1	31	35
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	4.1	8.1	0	21	27
Dodecane	112-40-3	4	0	1.6	5.5	0	9.3	29
Fluoranthene	206-44-0	21	0	0.91	1.2	0.33	3.2	4.1
Fluorene	86-73-7	15	0	1.4	2.8	0	5.1	15
Furan, 2-methyl	534-22-5	34	0	26	22	19	61	99
Heptanal	111-71-7	7	0	3.6	8.9	0	31	31
Hexadecane	544-76-3	11	0	7.7	14	0	41	55
2,5-Hexanedione	110-13-4	5	0	6.6	17	0	43	64
Indan	496-11-7	12	0	3.3	6.2	0	14	26
Mesitylene	108-67-8	15	0	6.7	12	0	27	55
Methacrolein	78-85-3	20	0	18	23	10	54	100
Methyl Isobutyl Ketone	108-10-1	5	0	3.8	10	0	29	44



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene	91-20-3	9	0	6.4	13	0	28	63
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.09	0.26	0	0.87	0.87
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.64	0.99	0	3	3.6
Naphthalene, 1-methyl-	90-12-0	15	0	5.2	9	0	23	36
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.18	0.27	0	0.61	0.61
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.48	0.72	0	2.2	2.5
Naphthalene, 2-methyl-	91-57-6	10	0	8	17	0	39	76
1-Octadecene	112-88-9	16	0	1.1	1.4	0	3.7	4.4
Octanal	124-13-0	24	0	11	13	6.7	33	50
Octane	111-65-9	10	0	15	27	0	61	100
17-Pentatriacontene	6971-40-0	2	0	0.18	0.76	0	0.7	4
N-Phenylbenzamide	93-98-1	6	0	2.2	4.9	0	13	13
Phenanthrene	85-01-8	17	0	3.1	4.9	1.7	14	20
Phenanthrene, 1-methyl	832-69-9	17	0	0.2	0.26	0.09	0.72	0.82
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.36	0.47	0.24	1.3	1.5
Phenanthrene, 3-methyl	832-71-3	18	0	0.43	0.56	0.23	1.6	1.8
Propionaldehyde	123-38-6	12	0	45	90	0	180	430
Pyrene	129-00-0	20	0	0.77	0.99	0.3	2.8	3.3
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.0086	0.043	0	0.015	0.25
Resorcinol	108-46-3	18	0	4.5	6.6	1.8	15	28
m-Tolualdehyde	620-23-5	19	0	65	69	53	180	220
TXIB "Kodaflex"	6846-50-0	1	0	0.48	2.8	0	0	17
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.92	1.3	0	3.2	4.7
Undecane	1120-21-4	5	0	3.2	8.3	0	25	32
Valeraldehyde	110-62-3	11	0	230	390	0	1000	1100
Non-Field-Related Chemicals								
Benzene	71-43-2	35	22	150	120	100	370	590
Benzene, 1,4-dichloro	106-46-7	7	0	4.6	10	0	29	29
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	140	140	92	410	510
2-Butoxyethanol	111-76-2	1	0	1.2	7.1	0	0	42
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	18	100	0	0	620
Decanal	112-31-2	7	0	6.8	23	0	26	130
Ethylbenzene	100-41-4	13	0	42	70	0	170	280
Heptane	142-82-5	20	0	56	83	11	230	360
Hexanal	66-25-1	30	0	190	240	89	740	960
Hexane	110-54-3	26	0	160	360	72	430	2100
1-Hexanol, 2-ethyl-	104-76-7	6	0	1.9	4.8	0	9.5	22
Nonanal	124-19-6	2	0	1.9	7.8	0	9.9	33



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Phenol	108-95-2	18	0	14	18	8.4	50	50
Tetrachloroethylene	127-18-4	7	0	12	27	0	78	100
Tetradecane	629-59-4	3	0	2	6.9	0	17	34
m/p-Xylene	106-42-3	25	0	140	200	57	480	840
o-Xylene	95-47-6	13	0	47	79	0	190	310

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-152. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter) —Combined Gender Athletes 40<50 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.24	0.48	0	1	2.1
Acetone	67-64-1	34	370	4700	5200	2500	15000	16000
Aniline	62-53-3	10	0	1.5	2.7	0	6	10
Anthracene	120-12-7	12	0	0.078	0.15	0	0.34	0.58
Anthracene, 2-methyl-	613-12-7	15	0	0.02	0.029	0	0.078	0.1
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.023	0.042	0	0.065	0.23
Anthracene, 9-phenyl	602-55-1	1	0	0.0022	0.013	0	0	0.073
Benz[a]anthracene	56-55-3	1	0	0.0012	0.0069	0	0	0.04
Benzaldehyde	100-52-7	15	0	21	33	0	77	130
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	3.1	9.3	0	14	50
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.6	1.6	0	3.7	6.4
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	28	50	0	100	230
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	1.3	3.3	0	6.5	17
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	1.8	3.6	0	7.9	16
Benzene, butyl-	104-51-8	2	0	0.63	3	0	1.3	18
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.89	1	0.54	2.9	3.5
Benzo[b]fluoranthene	205-99-2	2	0	0.0059	0.024	0	0.035	0.1
7H-Benzo[c]fluorene	205-12-9	13	0	0.013	0.02	0	0.034	0.085
Benzo[k]fluoranthene	207-08-9	4	0	0.0073	0.022	0	0.056	0.083
Benzothiazole	95-16-9	19	0	8.8	10	4.9	29	29



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.7	0.93	0.36	2.4	4.2
2-Benzothiazolone	934-34-9	5	0	1.1	2.9	0	7.2	11
Benzyl butyl phthalate	85-68-7	7	0	0.82	1.8	0	5.6	5.6
Butanal	123-72-8	13	0	73	180	0	430	900
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	38	65	12	140	310
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	14	22	0	54	78
p-Cymene	99-87-6	21	0	6	7.3	3.8	20	28
Decane	124-18-5	11	0	13	23	0	71	73
Dibenz[a,h]anthracene	53-70-3	14	0	0.034	0.068	0	0.17	0.29
Dibenzothiophene	132-65-0	14	0	0.29	0.45	0	1.3	1.6
Dibutyl phthalate	84-74-2	7	0	90	230	0	400	1200
Diethyl phthalate	84-66-2	1	0	0.63	3.7	0	0	21
Diisobutyl phthalate	84-69-5	15	0	3.4	6.7	0	11	36
Diisooctylphthalate	27554-26-3	9	0	12	39	0	31	230
Di-n-octyl phthalate	117-84-0	5	0	0.023	0.06	0	0.13	0.26
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	7.7	10	4	30	35
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	4	7.9	0	20	26
Dodecane	112-40-3	4	0	1.6	5.4	0	9.2	28
Fluoranthene	206-44-0	21	0	0.9	1.1	0.32	3.1	4.1
Fluorene	86-73-7	15	0	1.4	2.8	0	5	15
Furan, 2-methyl	534-22-5	34	0	25	21	19	60	98
Heptanal	111-71-7	7	0	3.5	8.8	0	31	31
Hexadecane	544-76-3	11	0	7.6	14	0	40	54
2,5-Hexanedione	110-13-4	5	0	6.5	17	0	42	63
Indan	496-11-7	12	0	3.3	6.1	0	14	26
Mesitylene	108-67-8	15	0	6.6	12	0	26	54
Methacrolein	78-85-3	20	0	18	23	10	53	100
Methyl Isobutyl Ketone	108-10-1	5	0	3.7	10	0	29	44
Naphthalene	91-20-3	9	0	6.3	13	0	27	62
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.089	0.25	0	0.86	0.86
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.63	0.98	0	3	3.6
Naphthalene, 1-methyl-	90-12-0	15	0	5.2	8.9	0	23	36
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.18	0.27	0	0.6	0.6
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.48	0.71	0	2.2	2.5
Naphthalene, 2-methyl-	91-57-6	10	0	7.9	17	0	38	75



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
1-Octadecene	112-88-9	16	0	1.1	1.4	0	3.6	4.3
Octanal	124-13-0	24	0	11	12	6.6	32	49
Octane	111-65-9	10	0	14	26	0	60	100
17-Pentatriacontene	6971-40-0	2	0	0.17	0.75	0	0.69	4
N-Phenylbenzamide	93-98-1	6	0	2.1	4.8	0	13	13
Phenanthrene	85-01-8	17	0	3.1	4.8	1.7	14	20
Phenanthrene, 1-methyl	832-69-9	17	0	0.2	0.25	0.088	0.71	0.81
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.36	0.46	0.24	1.3	1.5
Phenanthrene, 3-methyl	832-71-3	18	0	0.42	0.55	0.23	1.6	1.8
Propionaldehyde	123-38-6	12	0	44	89	0	170	430
Pyrene	129-00-0	20	0	0.76	0.97	0.3	2.7	3.3
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.0084	0.042	0	0.015	0.24
Resorcinol	108-46-3	18	0	4.5	6.5	1.8	15	28
m-Tolualdehyde	620-23-5	19	0	64	68	52	180	220
TXIB "Kodaflex"	6846-50-0	1	0	0.47	2.8	0	0	17
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.9	1.3	0	3.2	4.6
Undecane	1120-21-4	5	0	3.1	8.2	0	24	31
Valeraldehyde	110-62-3	11	0	220	380	0	1000	1100
Non-Field-Related Chemicals								
Benzene	71-43-2	35	22	140	120	100	360	590
Benzene, 1,4-dichloro	106-46-7	7	0	4.6	10	0	28	28
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	140	140	91	410	510
2-Butoxyethanol	111-76-2	1	0	1.2	7	0	0	41
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	17	100	0	0	610
Decanal	112-31-2	7	0	6.7	23	0	25	130
Ethylbenzene	100-41-4	13	0	41	69	0	170	280
Heptane	142-82-5	20	0	55	82	11	230	360
Hexanal	66-25-1	30	0	190	240	88	730	950
Hexane	110-54-3	26	0	160	360	71	420	2100
1-Hexanol, 2-ethyl-	104-76-7	6	0	1.9	4.8	0	9.4	22
Nonanal	124-19-6	2	0	1.9	7.7	0	9.8	33
Phenol	108-95-2	18	0	14	18	8.2	49	49
Tetrachloroethylene	127-18-4	7	0	11	27	0	76	100
Tetradecane	629-59-4	3	0	1.9	6.8	0	17	34
m/p-Xylene	106-42-3	25	0	140	200	56	470	830
o-Xylene	95-47-6	13	0	46	77	0	180	300

^a 35 field-specific C_{inh-field} values are included in the table.

^b On-Field field-specific Chronic C_{inh-field} equals the modified field-specific average concentration of a field



($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-153. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter) —Combined Gender Athletes 50<70 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.24	0.48	0	0.99	2.1
Acetone	67-64-1	34	370	4600	5100	2500	15000	16000
Aniline	62-53-3	10	0	1.5	2.7	0	5.9	10
Anthracene	120-12-7	12	0	0.077	0.14	0	0.33	0.57
Anthracene, 2-methyl-	613-12-7	15	0	0.02	0.028	0	0.077	0.098
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.023	0.041	0	0.064	0.22
Anthracene, 9-phenyl	602-55-1	1	0	0.0021	0.012	0	0	0.072
Benz[a]anthracene	56-55-3	1	0	0.0012	0.0067	0	0	0.039
Benzaldehyde	100-52-7	15	0	21	32	0	76	130
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	3.1	9.1	0	14	49
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.59	1.5	0	3.6	6.3
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	28	49	0	98	230
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	1.2	3.2	0	6.3	17
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	1.8	3.5	0	7.7	16
Benzene, butyl-	104-51-8	2	0	0.61	3	0	1.3	17
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.87	1	0.53	2.9	3.4
Benzo[b]fluoranthene	205-99-2	2	0	0.0058	0.023	0	0.034	0.098
7H-Benzo[c]fluorene	205-12-9	13	0	0.013	0.019	0	0.033	0.084
Benzo[k]fluoranthene	207-08-9	4	0	0.0072	0.021	0	0.055	0.081
Benzothiazole	95-16-9	19	0	8.7	10	4.8	29	29
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.69	0.91	0.35	2.4	4.1
2-Benzothiazolone	934-34-9	5	0	1.1	2.8	0	7.1	10
Benzyl butyl phthalate	85-68-7	7	0	0.8	1.7	0	5.5	5.5
Butanal	123-72-8	13	0	72	180	0	420	880
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	38	64	12	130	310
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	14	22	0	53	77
p-Cymene	99-87-6	21	0	5.9	7.1	3.7	20	28
Decane	124-18-5	11	0	13	23	0	69	72
Dibenz[a,h]anthracene	53-70-3	14	0	0.034	0.067	0	0.17	0.28



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dibenzothiophene	132-65-0	14	0	0.29	0.44	0	1.2	1.6
Dibutyl phthalate	84-74-2	7	0	88	220	0	400	1200
Diethyl phthalate	84-66-2	1	0	0.62	3.6	0	0	21
Diisobutyl phthalate	84-69-5	15	0	3.3	6.6	0	11	35
Diisooctylphthalate	27554-26-3	9	0	12	39	0	30	220
Di-n-octyl phthalate	117-84-0	5	0	0.023	0.059	0	0.13	0.26
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	7.6	10	3.9	30	34
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	4	7.8	0	20	26
Dodecane	112-40-3	4	0	1.6	5.3	0	9	28
Fluoranthene	206-44-0	21	0	0.88	1.1	0.32	3.1	4
Fluorene	86-73-7	15	0	1.4	2.7	0	4.9	14
Furan, 2-methyl	534-22-5	34	0	25	21	18	59	96
Heptanal	111-71-7	7	0	3.4	8.7	0	30	30
Hexadecane	544-76-3	11	0	7.4	14	0	40	53
2,5-Hexanedione	110-13-4	5	0	6.4	17	0	42	62
Indan	496-11-7	12	0	3.2	6	0	14	25
Mesitylene	108-67-8	15	0	6.5	12	0	26	53
Methacrolein	78-85-3	20	0	18	23	10	52	100
Methyl Isobutyl Ketone	108-10-1	5	0	3.7	10	0	29	43
Naphthalene	91-20-3	9	0	6.2	13	0	27	61
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.087	0.25	0	0.85	0.85
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.62	0.96	0	2.9	3.5
Naphthalene, 1-methyl-	90-12-0	15	0	5.1	8.7	0	23	35
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.17	0.26	0	0.59	0.59
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.47	0.7	0	2.1	2.5
Naphthalene, 2-methyl-	91-57-6	10	0	7.8	16	0	38	74
1-Octadecene	112-88-9	16	0	1	1.3	0	3.6	4.3
Octanal	124-13-0	24	0	10	12	6.5	32	48
Octane	111-65-9	10	0	14	26	0	59	99
17-Pentatriacontene	6971-40-0	2	0	0.17	0.74	0	0.68	3.9
N-Phenylbenzamide	93-98-1	6	0	2.1	4.7	0	13	13
Phenanthrene	85-01-8	17	0	3	4.7	1.6	13	20
Phenanthrene, 1-methyl	832-69-9	17	0	0.19	0.25	0.087	0.69	0.79
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.35	0.46	0.24	1.3	1.5
Phenanthrene, 3-methyl	832-71-3	18	0	0.42	0.54	0.23	1.5	1.8
Propionaldehyde	123-38-6	12	0	43	87	0	170	420
Pyrene	129-00-0	20	0	0.75	0.95	0.3	2.7	3.2
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.0083	0.042	0	0.015	0.24



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Resorcinol	108-46-3	18	0	4.4	6.4	1.8	14	27
m-Tolualdehyde	620-23-5	19	0	63	67	51	170	210
TXIB "Kodaflex"	6846-50-0	1	0	0.46	2.7	0	0	16
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.89	1.2	0	3.1	4.5
Undecane	1120-21-4	5	0	3.1	8	0	24	31
Valeraldehyde	110-62-3	11	0	220	380	0	990	1100
Non-Field-Related Chemicals								
Benzene	71-43-2	35	21	140	110	100	350	580
Benzene, 1,4-dichloro	106-46-7	7	0	4.5	9.9	0	28	28
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	130	140	89	400	500
2-Butoxyethanol	111-76-2	1	0	1.2	6.9	0	0	41
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	17	100	0	0	600
Decanal	112-31-2	7	0	6.6	22	0	25	130
Ethylbenzene	100-41-4	13	0	40	68	0	160	270
Heptane	142-82-5	20	0	54	81	10	220	350
Hexanal	66-25-1	30	0	180	240	86	710	930
Hexane	110-54-3	26	0	160	350	70	420	2100
1-Hexanol, 2-ethyl-	104-76-7	6	0	1.8	4.7	0	9.2	21
Nonanal	124-19-6	2	0	1.8	7.5	0	9.6	32
Phenol	108-95-2	18	0	14	17	8.1	49	49
Tetrachloroethylene	127-18-4	7	0	11	27	0	75	98
Tetradecane	629-59-4	3	0	1.9	6.7	0	17	33
m/p-Xylene	106-42-3	25	0	140	190	55	460	810
o-Xylene	95-47-6	13	0	45	76	0	180	300

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-154. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter) —Combined Gender Coaches 16<30 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.15	0.31	0	0.65	1.3
Acetone	67-64-1	34	240	3000	3400	1600	9800	11000



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	10	0	0.99	1.7	0	3.9	6.7
Anthracene	120-12-7	12	0	0.05	0.094	0	0.22	0.37
Anthracene, 2-methyl-	613-12-7	15	0	0.013	0.019	0	0.05	0.064
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.015	0.027	0	0.042	0.14
Anthracene, 9-phenyl	602-55-1	1	0	0.0014	0.0081	0	0	0.047
Benz[a]anthracene	56-55-3	1	0	0.00075	0.0044	0	0	0.026
Benzaldehyde	100-52-7	15	0	14	21	0	49	83
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	2	6	0	8.9	32
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.38	1	0	2.4	4.1
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	18	32	0	64	150
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.81	2.1	0	4.1	11
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	1.1	2.3	0	5	10
Benzene, butyl-	104-51-8	2	0	0.4	1.9	0	0.84	11
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.57	0.67	0.34	1.9	2.3
Benzo[b]fluoranthene	205-99-2	2	0	0.0038	0.015	0	0.022	0.064
7H-Benzo[c]fluorene	205-12-9	13	0	0.0083	0.013	0	0.022	0.055
Benzo[k]fluoranthene	207-08-9	4	0	0.0047	0.014	0	0.036	0.053
Benzothiazole	95-16-9	19	0	5.7	6.5	3.1	19	19
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.45	0.59	0.23	1.6	2.7
2-Benzothiazolone	934-34-9	5	0	0.71	1.8	0	4.6	6.9
Benzyl butyl phthalate	85-68-7	7	0	0.53	1.1	0	3.6	3.6
Butanal	123-72-8	13	0	47	120	0	280	580
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	25	42	7.6	88	200
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	8.9	14	0	35	50
p-Cymene	99-87-6	21	0	3.8	4.7	2.4	13	18
Decane	124-18-5	11	0	8.3	15	0	45	47
Dibenz[a,h]anthracene	53-70-3	14	0	0.022	0.044	0	0.11	0.18
Dibenzothiophene	132-65-0	14	0	0.19	0.29	0	0.81	1
Dibutyl phthalate	84-74-2	7	0	57	150	0	260	750
Diethyl phthalate	84-66-2	1	0	0.4	2.3	0	0	14
Diisobutyl phthalate	84-69-5	15	0	2.2	4.3	0	6.9	23
Diisooctylphthalate	27554-26-3	9	0	7.6	25	0	20	150
Di-n-octyl phthalate	117-84-0	5	0	0.015	0.038	0	0.084	0.17
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	5	6.7	2.6	19	22
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	2.6	5.1	0	13	17



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh} -field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dodecane	112-40-3	4	0	1	3.5	0	5.9	18
Fluoranthene	206-44-0	21	0	0.58	0.74	0.21	2	2.6
Fluorene	86-73-7	15	0	0.91	1.8	0	3.2	9.4
Furan, 2-methyl	534-22-5	34	0	16	14	12	38	63
Heptanal	111-71-7	7	0	2.3	5.7	0	20	20
Hexadecane	544-76-3	11	0	4.9	8.9	0	26	35
2,5-Hexanedione	110-13-4	5	0	4.2	11	0	27	40
Indan	496-11-7	12	0	2.1	3.9	0	9	17
Mesitylene	108-67-8	15	0	4.2	7.5	0	17	35
Methacrolein	78-85-3	20	0	12	15	6.6	34	66
Methyl Isobutyl Ketone	108-10-1	5	0	2.4	6.5	0	19	28
Naphthalene	91-20-3	9	0	4.1	8.5	0	18	40
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.057	0.16	0	0.55	0.55
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.41	0.63	0	1.9	2.3
Naphthalene, 1-methyl-	90-12-0	15	0	3.3	5.7	0	15	23
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.11	0.17	0	0.38	0.38
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.31	0.45	0	1.4	1.6
Naphthalene, 2-methyl-	91-57-6	10	0	5.1	11	0	25	48
1-Octadecene	112-88-9	16	0	0.68	0.87	0	2.3	2.8
Octanal	124-13-0	24	0	6.9	7.9	4.2	21	32
Octane	111-65-9	10	0	9.3	17	0	39	64
17-Pentatriacontene	6971-40-0	2	0	0.11	0.48	0	0.44	2.5
N-Phenylbenzamide	93-98-1	6	0	1.4	3.1	0	8.5	8.5
Phenanthrene	85-01-8	17	0	2	3.1	1.1	8.7	13
Phenanthrene, 1-methyl	832-69-9	17	0	0.13	0.16	0.057	0.45	0.52
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.23	0.3	0.15	0.84	0.96
Phenanthrene, 3-methyl	832-71-3	18	0	0.27	0.35	0.15	1	1.1
Propionaldehyde	123-38-6	12	0	28	57	0	110	270
Pyrene	129-00-0	20	0	0.49	0.62	0.19	1.8	2.1
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.0054	0.027	0	0.0095	0.16
Resorcinol	108-46-3	18	0	2.9	4.2	1.2	9.4	18
m-Tolualdehyde	620-23-5	19	0	41	43	33	110	140
TXIB "Kodaflex"	6846-50-0	1	0	0.3	1.8	0	0	11
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.58	0.81	0	2	3
Undecane	1120-21-4	5	0	2	5.2	0	16	20
Valeraldehyde	110-62-3	11	0	140	250	0	640	700
Non-Field-Related Chemicals								
Benzene	71-43-2	35	14	93	75	65	230	380



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene, 1,4-dichloro	106-46-7	7	0	2.9	6.5	0	18	18
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	88	91	58	260	320
2-Butoxyethanol	111-76-2	1	0	0.76	4.5	0	0	27
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	11	66	0	0	390
Decanal	112-31-2	7	0	4.3	15	0	16	84
Ethylbenzene	100-41-4	13	0	26	44	0	110	180
Heptane	142-82-5	20	0	35	53	6.8	150	230
Hexanal	66-25-1	30	0	120	150	56	470	610
Hexane	110-54-3	26	0	100	230	45	270	1300
1-Hexanol, 2-ethyl-	104-76-7	6	0	1.2	3.1	0	6	14
Nonanal	124-19-6	2	0	1.2	4.9	0	6.3	21
Phenol	108-95-2	18	0	8.9	11	5.3	32	32
Tetrachloroethylene	127-18-4	7	0	7.3	17	0	49	64
Tetradecane	629-59-4	3	0	1.2	4.4	0	11	22
m/p-Xylene	106-42-3	25	0	88	130	36	300	530
o-Xylene	95-47-6	13	0	30	50	0	120	200

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-155. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter) —Combined Gender Coaches 30<40 Years

Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.13	0.27	0	0.57	1.2
Acetone	67-64-1	34	210	2600	2900	1400	8600	9200
Aniline	62-53-3	10	0	0.87	1.5	0	3.4	5.8
Anthracene	120-12-7	12	0	0.044	0.082	0	0.19	0.32
Anthracene, 2-methyl-	613-12-7	15	0	0.011	0.016	0	0.044	0.056
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.013	0.023	0	0.036	0.13
Anthracene, 9-phenyl	602-55-1	1	0	0.0012	0.007	0	0	0.041
Benz[a]anthracene	56-55-3	1	0	0.00066	0.0038	0	0	0.022
Benzaldehyde	100-52-7	15	0	12	18	0	43	73
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	1.8	5.2	0	7.8	28
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.33	0.88	0	2.1	3.6



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	16	28	0	56	130
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.7	1.9	0	3.6	9.5
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	1	2	0	4.4	9.1
Benzene, butyl-	104-51-8	2	0	0.35	1.7	0	0.74	9.8
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.5	0.58	0.3	1.6	2
Benzo[b]fluoranthene	205-99-2	2	0	0.0033	0.013	0	0.02	0.056
7H-Benzo[c]fluorene	205-12-9	13	0	0.0072	0.011	0	0.019	0.048
Benzo[k]fluoranthene	207-08-9	4	0	0.0041	0.012	0	0.031	0.046
Benzothiazole	95-16-9	19	0	4.9	5.7	2.7	16	16
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.39	0.52	0.2	1.4	2.3
2-Benzothiazolone	934-34-9	5	0	0.62	1.6	0	4	6
Benzyl butyl phthalate	85-68-7	7	0	0.46	0.98	0	3.1	3.1
Butanal	123-72-8	13	0	41	100	0	240	500
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	22	36	6.6	77	170
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	7.8	12	0	30	44
p-Cymene	99-87-6	21	0	3.3	4.1	2.1	11	16
Decane	124-18-5	11	0	7.2	13	0	40	41
Dibenz[a,h]anthracene	53-70-3	14	0	0.019	0.038	0	0.096	0.16
Dibenzothiophene	132-65-0	14	0	0.17	0.25	0	0.7	0.91
Dibutyl phthalate	84-74-2	7	0	50	130	0	230	660
Diethyl phthalate	84-66-2	1	0	0.35	2	0	0	12
Diisobutyl phthalate	84-69-5	15	0	1.9	3.7	0	6	20
Diisooctylphthalate	27554-26-3	9	0	6.6	22	0	17	130
Di-n-octyl phthalate	117-84-0	5	0	0.013	0.034	0	0.073	0.15
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	4.3	5.9	2.2	17	19
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	2.3	4.5	0	11	15
Dodecane	112-40-3	4	0	0.9	3	0	5.1	16
Fluoranthene	206-44-0	21	0	0.5	0.64	0.18	1.8	2.3
Fluorene	86-73-7	15	0	0.8	1.6	0	2.8	8.2
Furan, 2-methyl	534-22-5	34	0	14	12	11	33	55
Heptanal	111-71-7	7	0	2	4.9	0	17	17
Hexadecane	544-76-3	11	0	4.2	7.7	0	23	30
2,5-Hexanedione	110-13-4	5	0	3.6	9.5	0	24	35
Indan	496-11-7	12	0	1.8	3.4	0	7.9	14



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Mesitylene	108-67-8	15	0	3.7	6.6	0	15	30
Methacrolein	78-85-3	20	0	10	13	5.8	30	57
Methyl Isobutyl Ketone	108-10-1	5	0	2.1	5.7	0	16	24
Naphthalene	91-20-3	9	0	3.6	7.4	0	15	35
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.05	0.14	0	0.48	0.48
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.36	0.55	0	1.7	2
Naphthalene, 1-methyl-	90-12-0	15	0	2.9	5	0	13	20
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.099	0.15	0	0.34	0.34
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.27	0.4	0	1.2	1.4
Naphthalene, 2-methyl-	91-57-6	10	0	4.4	9.4	0	22	42
1-Octadecene	112-88-9	16	0	0.59	0.76	0	2	2.4
Octanal	124-13-0	24	0	6	6.9	3.7	18	28
Octane	111-65-9	10	0	8.1	15	0	34	56
17-Pentatriacontene	6971-40-0	2	0	0.098	0.42	0	0.39	2.2
N-Phenylbenzamide	93-98-1	6	0	1.2	2.7	0	7.4	7.4
Phenanthrene	85-01-8	17	0	1.7	2.7	0.93	7.6	11
Phenanthrene, 1-methyl	832-69-9	17	0	0.11	0.14	0.05	0.4	0.45
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.2	0.26	0.14	0.73	0.84
Phenanthrene, 3-methyl	832-71-3	18	0	0.24	0.31	0.13	0.88	1
Propionaldehyde	123-38-6	12	0	25	50	0	98	240
Pyrene	129-00-0	20	0	0.43	0.54	0.17	1.5	1.8
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.0047	0.024	0	0.0083	0.14
Resorcinol	108-46-3	18	0	2.5	3.6	1	8.2	16
m-Tolualdehyde	620-23-5	19	0	36	38	29	99	120
TXIB "Kodaflex"	6846-50-0	1	0	0.26	1.6	0	0	9.3
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.51	0.71	0	1.8	2.6
Undecane	1120-21-4	5	0	1.8	4.6	0	14	17
Valeraldehyde	110-62-3	11	0	120	220	0	560	610
Non-Field-Related Chemicals								
Benzene	71-43-2	35	12	81	65	57	200	330
Benzene, 1,4-dichloro	106-46-7	7	0	2.6	5.6	0	16	16
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	77	80	51	230	280
2-Butoxyethanol	111-76-2	1	0	0.66	3.9	0	0	23
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	9.8	58	0	0	340
Decanal	112-31-2	7	0	3.8	13	0	14	74
Ethylbenzene	100-41-4	13	0	23	39	0	94	150
Heptane	142-82-5	20	0	31	46	5.9	130	200



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Hexanal	66-25-1	30	0	110	140	49	410	530
Hexane	110-54-3	26	0	90	200	40	240	1200
1-Hexanol, 2-ethyl-	104-76-7	6	0	1	2.7	0	5.3	12
Nonanal	124-19-6	2	0	1	4.3	0	5.5	18
Phenol	108-95-2	18	0	7.8	9.9	4.6	28	28
Tetrachloroethylene	127-18-4	7	0	6.4	15	0	43	56
Tetradecane	629-59-4	3	0	1.1	3.8	0	9.5	19
m/p-Xylene	106-42-3	25	0	77	110	31	260	470
o-Xylene	95-47-6	13	0	26	43	0	100	170

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-156. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter) —Combined Gender Coaches 40<50 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.14	0.27	0	0.57	1.2
Acetone	67-64-1	34	210	2600	3000	1400	8600	9300
Aniline	62-53-3	10	0	0.88	1.5	0	3.4	5.9
Anthracene	120-12-7	12	0	0.044	0.083	0	0.19	0.33
Anthracene, 2-methyl-	613-12-7	15	0	0.012	0.016	0	0.044	0.057
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.013	0.024	0	0.037	0.13
Anthracene, 9-phenyl	602-55-1	1	0	0.0012	0.0071	0	0	0.042
Benz[a]anthracene	56-55-3	1	0	0.00067	0.0039	0	0	0.023
Benzaldehyde	100-52-7	15	0	12	19	0	44	73
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	1.8	5.3	0	7.9	28
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.34	0.89	0	2.1	3.6
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	16	28	0	56	130
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.71	1.9	0	3.7	9.6
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	1	2	0	4.5	9.2
Benzene, butyl-	104-51-8	2	0	0.35	1.7	0	0.74	9.9
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.5	0.59	0.3	1.6	2
Benzo[b]fluoranthene	205-99-2	2	0	0.0033	0.014	0	0.02	0.057



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
7H-Benzo[c]fluorene	205-12-9	13	0	0.0073	0.011	0	0.019	0.048
Benzo[k]fluoranthene	207-08-9	4	0	0.0041	0.012	0	0.032	0.047
Benzothiazole	95-16-9	19	0	5	5.8	2.8	17	17
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.4	0.53	0.2	1.4	2.4
2-Benzothiazolone	934-34-9	5	0	0.62	1.6	0	4.1	6.1
Benzyl butyl phthalate	85-68-7	7	0	0.46	0.99	0	3.2	3.2
Butanal	123-72-8	13	0	41	100	0	240	510
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	22	37	6.7	78	180
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	7.8	13	0	31	44
p-Cymene	99-87-6	21	0	3.4	4.1	2.1	11	16
Decane	124-18-5	11	0	7.3	13	0	40	41
Dibenz[a,h]anthracene	53-70-3	14	0	0.019	0.039	0	0.097	0.16
Dibenzothiophene	132-65-0	14	0	0.17	0.26	0	0.71	0.92
Dibutyl phthalate	84-74-2	7	0	51	130	0	230	660
Diethyl phthalate	84-66-2	1	0	0.36	2.1	0	0	12
Diisobutyl phthalate	84-69-5	15	0	1.9	3.8	0	6.1	20
Diisooctylphthalate	27554-26-3	9	0	6.7	22	0	17	130
Di-n-octyl phthalate	117-84-0	5	0	0.013	0.034	0	0.074	0.15
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	4.4	5.9	2.3	17	20
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	2.3	4.5	0	11	15
Dodecane	112-40-3	4	0	0.91	3.1	0	5.2	16
Fluoranthene	206-44-0	21	0	0.51	0.65	0.18	1.8	2.3
Fluorene	86-73-7	15	0	0.81	1.6	0	2.8	8.3
Furan, 2-methyl	534-22-5	34	0	14	12	11	34	55
Heptanal	111-71-7	7	0	2	5	0	17	17
Hexadecane	544-76-3	11	0	4.3	7.8	0	23	31
2,5-Hexanedione	110-13-4	5	0	3.7	9.6	0	24	36
Indan	496-11-7	12	0	1.8	3.4	0	8	15
Mesitylene	108-67-8	15	0	3.7	6.6	0	15	31
Methacrolein	78-85-3	20	0	10	13	5.8	30	58
Methyl Isobutyl Ketone	108-10-1	5	0	2.1	5.8	0	16	25
Naphthalene	91-20-3	9	0	3.6	7.5	0	16	35
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.05	0.14	0	0.49	0.49
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.36	0.55	0	1.7	2
Naphthalene, 1-methyl-	90-12-0	15	0	2.9	5	0	13	20
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.1	0.15	0	0.34	0.34



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.27	0.4	0	1.2	1.4
Naphthalene, 2-methyl-	91-57-6	10	0	4.5	9.5	0	22	43
1-Octadecene	112-88-9	16	0	0.6	0.77	0	2.1	2.5
Octanal	124-13-0	24	0	6.1	7	3.7	18	28
Octane	111-65-9	10	0	8.2	15	0	34	57
17-Pentatriacontene	6971-40-0	2	0	0.099	0.42	0	0.39	2.2
N-Phenylbenzamide	93-98-1	6	0	1.2	2.7	0	7.5	7.5
Phenanthrene	85-01-8	17	0	1.8	2.7	0.94	7.6	11
Phenanthrene, 1-methyl	832-69-9	17	0	0.11	0.14	0.05	0.4	0.46
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.2	0.26	0.14	0.74	0.85
Phenanthrene, 3-methyl	832-71-3	18	0	0.24	0.31	0.13	0.89	1
Propionaldehyde	123-38-6	12	0	25	50	0	99	240
Pyrene	129-00-0	20	0	0.43	0.55	0.17	1.6	1.9
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.0048	0.024	0	0.0084	0.14
Resorcinol	108-46-3	18	0	2.5	3.7	1	8.3	16
m-Tolualdehyde	620-23-5	19	0	36	38	30	100	120
TXIB "Kodaflex"	6846-50-0	1	0	0.27	1.6	0	0	9.4
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.51	0.72	0	1.8	2.6
Undecane	1120-21-4	5	0	1.8	4.6	0	14	18
Valeraldehyde	110-62-3	11	0	130	220	0	570	620
Non-Field-Related Chemicals								
Benzene	71-43-2	35	12	82	66	58	200	330
Benzene, 1,4-dichloro	106-46-7	7	0	2.6	5.7	0	16	16
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	78	81	52	230	290
2-Butoxyethanol	111-76-2	1	0	0.67	4	0	0	23
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	9.9	59	0	0	350
Decanal	112-31-2	7	0	3.8	13	0	14	74
Ethylbenzene	100-41-4	13	0	23	39	0	95	160
Heptane	142-82-5	20	0	31	46	6	130	200
Hexanal	66-25-1	30	0	110	140	50	410	540
Hexane	110-54-3	26	0	91	200	40	240	1200
1-Hexanol, 2-ethyl-	104-76-7	6	0	1.1	2.7	0	5.3	12
Nonanal	124-19-6	2	0	1.1	4.3	0	5.5	18
Phenol	108-95-2	18	0	7.9	10	4.7	28	28
Tetrachloroethylene	127-18-4	7	0	6.5	15	0	43	57
Tetradecane	629-59-4	3	0	1.1	3.9	0	9.6	19



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
m/p-Xylene	106-42-3	25	0	78	110	32	270	470
o-Xylene	95-47-6	13	0	26	44	0	100	170

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-157. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter) —Combined Gender Coaches 50<70 Years

Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.14	0.27	0	0.57	1.2
Acetone	67-64-1	34	210	2700	3000	1400	8700	9300
Aniline	62-53-3	10	0	0.88	1.5	0	3.4	5.9
Anthracene	120-12-7	12	0	0.044	0.083	0	0.19	0.33
Anthracene, 2-methyl-	613-12-7	15	0	0.012	0.016	0	0.044	0.057
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.013	0.024	0	0.037	0.13
Anthracene, 9-phenyl	602-55-1	1	0	0.0012	0.0071	0	0	0.042
Benz[a]anthracene	56-55-3	1	0	0.00067	0.0039	0	0	0.023
Benzaldehyde	100-52-7	15	0	12	19	0	44	74
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	1.8	5.3	0	7.9	28
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.34	0.89	0	2.1	3.7
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	16	28	0	57	130
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.71	1.9	0	3.7	9.6
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	1	2	0	4.5	9.2
Benzene, butyl-	104-51-8	2	0	0.36	1.7	0	0.75	10
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.51	0.59	0.3	1.7	2
Benzo[b]fluoranthene	205-99-2	2	0	0.0033	0.014	0	0.02	0.057



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
7H-Benzo[c]fluorene	205-12-9	13	0	0.0073	0.011	0	0.019	0.048
Benzo[k]fluoranthene	207-08-9	4	0	0.0042	0.012	0	0.032	0.047
Benzothiazole	95-16-9	19	0	5	5.8	2.8	17	17
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.4	0.53	0.2	1.4	2.4
2-Benzothiazolone	934-34-9	5	0	0.63	1.6	0	4.1	6.1
Benzyl butyl phthalate	85-68-7	7	0	0.46	0.99	0	3.2	3.2
Butanal	123-72-8	13	0	41	100	0	240	510
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	22	37	6.7	78	180
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	7.9	13	0	31	44
p-Cymene	99-87-6	21	0	3.4	4.1	2.2	11	16
Decane	124-18-5	11	0	7.3	13	0	40	41
Dibenz[a,h]anthracene	53-70-3	14	0	0.019	0.039	0	0.097	0.16
Dibenzothiophene	132-65-0	14	0	0.17	0.26	0	0.71	0.93
Dibutyl phthalate	84-74-2	7	0	51	130	0	230	670
Diethyl phthalate	84-66-2	1	0	0.36	2.1	0	0	12
Diisobutyl phthalate	84-69-5	15	0	1.9	3.8	0	6.1	20
Diisooctylphthalate	27554-26-3	9	0	6.7	22	0	17	130
Di-n-octyl phthalate	117-84-0	5	0	0.013	0.034	0	0.074	0.15
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	4.4	6	2.3	17	20
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	2.3	4.5	0	11	15
Dodecane	112-40-3	4	0	0.92	3.1	0	5.2	16
Fluoranthene	206-44-0	21	0	0.51	0.65	0.18	1.8	2.3
Fluorene	86-73-7	15	0	0.81	1.6	0	2.9	8.3
Furan, 2-methyl	534-22-5	34	0	14	12	11	34	55
Heptanal	111-71-7	7	0	2	5	0	17	17
Hexadecane	544-76-3	11	0	4.3	7.8	0	23	31
2,5-Hexanedione	110-13-4	5	0	3.7	9.6	0	24	36
Indan	496-11-7	12	0	1.8	3.4	0	8	15



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Mesitylene	108-67-8	15	0	3.7	6.7	0	15	31
Methacrolein	78-85-3	20	0	10	13	5.9	30	58
Methyl Isobutyl Ketone	108-10-1	5	0	2.1	5.8	0	16	25
Naphthalene	91-20-3	9	0	3.6	7.5	0	16	35
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.05	0.14	0	0.49	0.49
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.36	0.56	0	1.7	2
Naphthalene, 1-methyl-	90-12-0	15	0	2.9	5	0	13	20
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.1	0.15	0	0.34	0.34
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.27	0.4	0	1.2	1.4
Naphthalene, 2-methyl-	91-57-6	10	0	4.5	9.5	0	22	43
1-Octadecene	112-88-9	16	0	0.6	0.77	0	2.1	2.5
Octanal	124-13-0	24	0	6.1	7	3.8	18	28
Octane	111-65-9	10	0	8.2	15	0	34	57
17-Pentatriacontene	6971-40-0	2	0	0.099	0.43	0	0.39	2.2
N-Phenylbenzamide	93-98-1	6	0	1.2	2.7	0	7.5	7.5
Phenanthrene	85-01-8	17	0	1.8	2.7	0.94	7.7	11
Phenanthrene, 1-methyl	832-69-9	17	0	0.11	0.14	0.05	0.4	0.46
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.2	0.26	0.14	0.74	0.85
Phenanthrene, 3-methyl	832-71-3	18	0	0.24	0.31	0.13	0.89	1
Propionaldehyde	123-38-6	12	0	25	50	0	99	240
Pyrene	129-00-0	20	0	0.43	0.55	0.17	1.6	1.9
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.0048	0.024	0	0.0084	0.14
Resorcinol	108-46-3	18	0	2.5	3.7	1	8.3	16
m-Tolualdehyde	620-23-5	19	0	37	39	30	100	120
TXIB "Kodaflex"	6846-50-0	1	0	0.27	1.6	0	0	9.4
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.51	0.72	0	1.8	2.6
Undecane	1120-21-4	5	0	1.8	4.6	0	14	18



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Valeraldehyde	110-62-3	11	0	130	220	0	570	620
Non-Field-Related Chemicals								
Benzene	71-43-2	35	12	82	66	58	210	330
Benzene, 1,4-dichloro	106-46-7	7	0	2.6	5.7	0	16	16
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	78	81	52	230	290
2-Butoxyethanol	111-76-2	1	0	0.67	4	0	0	23
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	9.9	59	0	0	350
Decanal	112-31-2	7	0	3.8	13	0	14	75
Ethylbenzene	100-41-4	13	0	23	39	0	95	160
Heptane	142-82-5	20	0	31	47	6	130	200
Hexanal	66-25-1	30	0	110	140	50	410	540
Hexane	110-54-3	26	0	91	200	40	240	1200
1-Hexanol, 2-ethyl-	104-76-7	6	0	1.1	2.7	0	5.3	12
Nonanal	124-19-6	2	0	1.1	4.3	0	5.5	18
Phenol	108-95-2	18	0	7.9	10	4.7	28	28
Tetrachloroethylene	127-18-4	7	0	6.5	15	0	43	57
Tetradecane	629-59-4	3	0	1.1	3.9	0	9.6	19
m/p-Xylene	106-42-3	25	0	78	110	32	270	470
o-Xylene	95-47-6	13	0	26	44	0	100	170

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-158. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter) —Combined Gender Referees 16<30 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	11	0	0.057	0.12	0	0.24	0.5
Acetone	67-64-1	34	89	1100	1200	600	3600	3900
Aniline	62-53-3	10	0	0.37	0.65	0	1.4	2.5
Anthracene	120-12-7	12	0	0.019	0.035	0	0.081	0.14
Anthracene, 2-methyl-	613-12-7	15	0	0.0049	0.0069	0	0.019	0.024
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.0055	0.0099	0	0.015	0.054
Anthracene, 9-phenyl	602-55-1	1	0	0.00051	0.003	0	0	0.017
Benz[a]anthracene	56-55-3	1	0	0.00028	0.0016	0	0	0.0095
Benzaldehyde	100-52-7	15	0	5.1	7.8	0	18	31
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	0.75	2.2	0	3.3	12
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.14	0.37	0	0.88	1.5
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	6.7	12	0	24	55
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.3	0.79	0	1.5	4
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	0.42	0.85	0	1.9	3.9
Benzene, butyl-	104-51-8	2	0	0.15	0.72	0	0.31	4.2
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.21	0.25	0.13	0.69	0.84
Benzo[b]fluoranthene	205-99-2	2	0	0.0014	0.0057	0	0.0083	0.024
7H-Benzo[c]fluorene	205-12-9	13	0	0.0031	0.0047	0	0.008	0.02
Benzo[k]fluoranthene	207-08-9	4	0	0.0017	0.0051	0	0.013	0.02
Benzothiazole	95-16-9	19	0	2.1	2.4	1.2	7	7
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.17	0.22	0.085	0.58	0.99
2-Benzothiazolone	934-34-9	5	0	0.26	0.68	0	1.7	2.5
Benzyl butyl phthalate	85-68-7	7	0	0.19	0.42	0	1.3	1.3
Butanal	123-72-8	13	0	17	43	0	100	210
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	9.1	15	2.8	33	74
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	3.3	5.3	0	13	19
p-Cymene	99-87-6	21	0	1.4	1.7	0.9	4.8	6.7



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Decane	124-18-5	11	0	3.1	5.5	0	17	17
Dibenz[a,h]anthracene	53-70-3	14	0	0.0081	0.016	0	0.041	0.069
Dibenzothiophene	132-65-0	14	0	0.07	0.11	0	0.3	0.39
Dibutyl phthalate	84-74-2	7	0	21	55	0	96	280
Diethyl phthalate	84-66-2	1	0	0.15	0.87	0	0	5.1
Diisobutyl phthalate	84-69-5	15	0	0.81	1.6	0	2.6	8.6
Diisooctylphthalate	27554-26-3	9	0	2.8	9.4	0	7.3	54
Di-n-octyl phthalate	117-84-0	5	0	0.0055	0.014	0	0.031	0.062
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	1.8	2.5	0.95	7.2	8.2
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	0.96	1.9	0	4.8	6.3
Dodecane	112-40-3	4	0	0.38	1.3	0	2.2	6.7
Fluoranthene	206-44-0	21	0	0.21	0.27	0.077	0.74	0.97
Fluorene	86-73-7	15	0	0.34	0.66	0	1.2	3.5
Furan, 2-methyl	534-22-5	34	0	6.1	5.1	4.5	14	23
Heptanal	111-71-7	7	0	0.84	2.1	0	7.3	7.3
Hexadecane	544-76-3	11	0	1.8	3.3	0	9.6	13
2,5-Hexanedione	110-13-4	5	0	1.5	4	0	10	15
Indan	496-11-7	12	0	0.77	1.4	0	3.3	6.2
Mesitylene	108-67-8	15	0	1.6	2.8	0	6.3	13
Methacrolein	78-85-3	20	0	4.3	5.5	2.5	13	24
Methyl Isobutyl Ketone	108-10-1	5	0	0.89	2.4	0	6.9	10
Naphthalene	91-20-3	9	0	1.5	3.2	0	6.5	15
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.021	0.061	0	0.2	0.2
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.15	0.23	0	0.72	0.85
Naphthalene, 1-methyl-	90-12-0	15	0	1.2	2.1	0	5.5	8.5
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.042	0.064	0	0.14	0.14
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.11	0.17	0	0.52	0.6
Naphthalene, 2-methyl-	91-57-6	10	0	1.9	4	0	9.1	18



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
1-Octadecene	112-88-9	16	0	0.25	0.32	0	0.87	1
Octanal	124-13-0	24	0	2.5	2.9	1.6	7.7	12
Octane	111-65-9	10	0	3.4	6.3	0	14	24
17-Pentatriacontene	6971-40-0	2	0	0.042	0.18	0	0.16	0.94
N-Phenylbenzamide	93-98-1	6	0	0.51	1.1	0	3.2	3.2
Phenanthrene	85-01-8	17	0	0.74	1.1	0.4	3.2	4.8
Phenanthrene, 1-methyl	832-69-9	17	0	0.047	0.061	0.021	0.17	0.19
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.085	0.11	0.057	0.31	0.36
Phenanthrene, 3-methyl	832-71-3	18	0	0.1	0.13	0.055	0.37	0.43
Propionaldehyde	123-38-6	12	0	10	21	0	41	100
Pyrene	129-00-0	20	0	0.18	0.23	0.072	0.65	0.78
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.002	0.01	0	0.0035	0.058
Resorcinol	108-46-3	18	0	1.1	1.5	0.43	3.5	6.6
m-Tolualdehyde	620-23-5	19	0	15	16	12	42	51
TXIB "Kodaflex"	6846-50-0	1	0	0.11	0.66	0	0	3.9
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.21	0.3	0	0.76	1.1
Undecane	1120-21-4	5	0	0.75	1.9	0	5.8	7.4
Valeraldehyde	110-62-3	11	0	53	92	0	240	260
Non-Field-Related Chemicals								
Benzene	71-43-2	35	5.1	34	28	24	86	140
Benzene, 1,4-dichloro	106-46-7	7	0	1.1	2.4	0	6.7	6.7
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	33	34	22	97	120
2-Butoxyethanol	111-76-2	1	0	0.28	1.7	0	0	9.9
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	4.2	25	0	0	150
Decanal	112-31-2	7	0	1.6	5.4	0	6	31
Ethylbenzene	100-41-4	13	0	9.8	16	0	40	66



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Heptane	142-82-5	20	0	13	20	2.5	54	85
Hexanal	66-25-1	30	0	45	57	21	170	230
Hexane	110-54-3	26	0	38	85	17	100	500
1-Hexanol, 2-ethyl-	104-76-7	6	0	0.44	1.1	0	2.2	5.2
Nonanal	124-19-6	2	0	0.44	1.8	0	2.3	7.7
Phenol	108-95-2	18	0	3.3	4.2	2	12	12
Tetrachloroethylene	127-18-4	7	0	2.7	6.4	0	18	24
Tetradecane	629-59-4	3	0	0.46	1.6	0	4	8.1
m/p-Xylene	106-42-3	25	0	33	47	13	110	200
o-Xylene	95-47-6	13	0	11	18	0	44	73

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-159. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter) —Combined Gender Referees 30<40 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.05	0.1	0	0.21	0.43
Acetone	67-64-1	34	78	970	1100	530	3200	3400
Aniline	62-53-3	10	0	0.32	0.57	0	1.3	2.2
Anthracene	120-12-7	12	0	0.016	0.03	0	0.071	0.12
Anthracene, 2-methyl-	613-12-7	15	0	0.0042	0.006	0	0.016	0.021
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.0048	0.0087	0	0.014	0.047
Anthracene, 9-phenyl	602-55-1	1	0	0.00045	0.0026	0	0	0.015
Benz[a]anthracene	56-55-3	1	0	0.00024	0.0014	0	0	0.0083
Benzaldehyde	100-52-7	15	0	4.4	6.9	0	16	27



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	0.65	1.9	0	2.9	10
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.12	0.33	0	0.77	1.3
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	5.8	10	0	21	48
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.26	0.69	0	1.3	3.5
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	0.37	0.74	0	1.6	3.4
Benzene, butyl-	104-51-8	2	0	0.13	0.63	0	0.27	3.6
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.19	0.22	0.11	0.6	0.73
Benzo[b]fluoranthene	205-99-2	2	0	0.0012	0.005	0	0.0073	0.021
7H-Benzo[c]fluorene	205-12-9	13	0	0.0027	0.0041	0	0.007	0.018
Benzo[k]fluoranthene	207-08-9	4	0	0.0015	0.0045	0	0.012	0.017
Benzothiazole	95-16-9	19	0	1.8	2.1	1	6.1	6.1
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.15	0.19	0.074	0.5	0.87
2-Benzothiazolone	934-34-9	5	0	0.23	0.6	0	1.5	2.2
Benzyl butyl phthalate	85-68-7	7	0	0.17	0.36	0	1.2	1.2
Butanal	123-72-8	13	0	15	37	0	89	190
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	8	14	2.5	28	65
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	2.9	4.6	0	11	16
p-Cymene	99-87-6	21	0	1.2	1.5	0.79	4.2	5.9
Decane	124-18-5	11	0	2.7	4.8	0	15	15
Dibenz[a,h]anthracene	53-70-3	14	0	0.0071	0.014	0	0.036	0.06
Dibenzothiophene	132-65-0	14	0	0.061	0.094	0	0.26	0.34
Dibutyl phthalate	84-74-2	7	0	19	48	0	84	240
Diethyl phthalate	84-66-2	1	0	0.13	0.76	0	0	4.4
Diisobutyl phthalate	84-69-5	15	0	0.71	1.4	0	2.2	7.5
Diisooctylphthalate	27554-26-3	9	0	2.5	8.2	0	6.4	47
Di-n-octyl phthalate	117-84-0	5	0	0.0048	0.012	0	0.027	0.054
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	1.6	2.2	0.83	6.3	7.2



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh} -field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	0.84	1.7	0	4.2	5.5
Dodecane	112-40-3	4	0	0.34	1.1	0	1.9	5.9
Fluoranthene	206-44-0	21	0	0.19	0.24	0.067	0.65	0.85
Fluorene	86-73-7	15	0	0.3	0.58	0	1	3
Furan, 2-methyl	534-22-5	34	0	5.3	4.4	3.9	12	20
Heptanal	111-71-7	7	0	0.73	1.8	0	6.4	6.4
Hexadecane	544-76-3	11	0	1.6	2.9	0	8.4	11
2,5-Hexanedione	110-13-4	5	0	1.3	3.5	0	8.8	13
Indan	496-11-7	12	0	0.68	1.3	0	2.9	5.4
Mesitylene	108-67-8	15	0	1.4	2.4	0	5.5	11
Methacrolein	78-85-3	20	0	3.8	4.8	2.1	11	21
Methyl Isobutyl Ketone	108-10-1	5	0	0.78	2.1	0	6	9.1
Naphthalene	91-20-3	9	0	1.3	2.8	0	5.7	13
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.018	0.053	0	0.18	0.18
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.13	0.2	0	0.62	0.74
Naphthalene, 1-methyl-	90-12-0	15	0	1.1	1.8	0	4.8	7.4
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.037	0.056	0	0.12	0.12
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.099	0.15	0	0.45	0.52
Naphthalene, 2-methyl-	91-57-6	10	0	1.6	3.5	0	8	16
1-Octadecene	112-88-9	16	0	0.22	0.28	0	0.76	0.9
Octanal	124-13-0	24	0	2.2	2.6	1.4	6.7	10
Octane	111-65-9	10	0	3	5.5	0	12	21
17-Pentatriacontene	6971-40-0	2	0	0.036	0.16	0	0.14	0.82
N-Phenylbenzamide	93-98-1	6	0	0.45	1	0	2.8	2.8
Phenanthrene	85-01-8	17	0	0.64	1	0.35	2.8	4.2
Phenanthrene, 1-methyl	832-69-9	17	0	0.041	0.053	0.018	0.15	0.17
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.074	0.097	0.05	0.27	0.31
Phenanthrene, 3-methyl	832-71-3	18	0	0.088	0.11	0.048	0.33	0.37



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh} -field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Propionaldehyde	123-38-6	12	0	9.1	18	0	36	89
Pyrene	129-00-0	20	0	0.16	0.2	0.063	0.57	0.68
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.0018	0.0088	0	0.0031	0.051
Resorcinol	108-46-3	18	0	0.93	1.3	0.38	3	5.8
m-Tolualdehyde	620-23-5	19	0	13	14	11	37	45
TXIB "Kodaflex"	6846-50-0	1	0	0.098	0.58	0	0	3.4
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.19	0.26	0	0.66	0.96
Undecane	1120-21-4	5	0	0.65	1.7	0	5	6.5
Valeraldehyde	110-62-3	11	0	46	80	0	210	230
Non-Field-Related Chemicals								
Benzene	71-43-2	35	4.5	30	24	21	75	120
Benzene, 1,4-dichloro	106-46-7	7	0	0.95	2.1	0	5.9	5.9
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	29	30	19	85	110
2-Butoxyethanol	111-76-2	1	0	0.25	1.5	0	0	8.6
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	3.6	22	0	0	130
Decanal	112-31-2	7	0	1.4	4.8	0	5.3	27
Ethylbenzene	100-41-4	13	0	8.5	14	0	35	57
Heptane	142-82-5	20	0	11	17	2.2	47	74
Hexanal	66-25-1	30	0	39	50	18	150	200
Hexane	110-54-3	26	0	33	74	15	88	430
1-Hexanol, 2-ethyl-	104-76-7	6	0	0.39	0.99	0	2	4.5
Nonanal	124-19-6	2	0	0.39	1.6	0	2	6.8
Phenol	108-95-2	18	0	2.9	3.7	1.7	10	10
Tetrachloroethylene	127-18-4	7	0	2.4	5.6	0	16	21
Tetradecane	629-59-4	3	0	0.4	1.4	0	3.5	7.1
m/p-Xylene	106-42-3	25	0	29	41	12	98	170



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
o-Xylene	95-47-6	13	0	9.6	16	0	38	63

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-160. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter) —Combined Gender Referees 40<50 Years

Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.05	0.1	0	0.21	0.44
Acetone	67-64-1	34	79	980	1100	530	3200	3500
Aniline	62-53-3	10	0	0.33	0.57	0	1.3	2.2
Anthracene	120-12-7	12	0	0.016	0.031	0	0.072	0.12
Anthracene, 2-methyl-	613-12-7	15	0	0.0043	0.0061	0	0.016	0.021
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.0049	0.0088	0	0.014	0.047
Anthracene, 9-phenyl	602-55-1	1	0	0.00045	0.0026	0	0	0.015
Benz[a]anthracene	56-55-3	1	0	0.00025	0.0014	0	0	0.0084
Benzaldehyde	100-52-7	15	0	4.5	6.9	0	16	27
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	0.66	2	0	2.9	10
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.13	0.33	0	0.78	1.4
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	5.9	10	0	21	49
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.26	0.69	0	1.4	3.6
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	0.37	0.75	0	1.7	3.4
Benzene, butyl-	104-51-8	2	0	0.13	0.64	0	0.28	3.7
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.19	0.22	0.11	0.61	0.74
Benzo[b]fluoranthene	205-99-2	2	0	0.0012	0.005	0	0.0074	0.021
7H-Benzo[c]fluorene	205-12-9	13	0	0.0027	0.0041	0	0.0071	0.018



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[k]fluoranthene	207-08-9	4	0	0.0015	0.0045	0	0.012	0.017
Benzothiazole	95-16-9	19	0	1.9	2.1	1	6.2	6.2
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.15	0.2	0.075	0.51	0.87
2-Benzothiazolone	934-34-9	5	0	0.23	0.6	0	1.5	2.2
Benzyl butyl phthalate	85-68-7	7	0	0.17	0.37	0	1.2	1.2
Butanal	123-72-8	13	0	15	38	0	90	190
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	8.1	14	2.5	29	66
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	2.9	4.6	0	11	16
p-Cymene	99-87-6	21	0	1.3	1.5	0.8	4.2	5.9
Decane	124-18-5	11	0	2.7	4.9	0	15	15
Dibenz[a,h]anthracene	53-70-3	14	0	0.0072	0.014	0	0.036	0.061
Dibenzothiophene	132-65-0	14	0	0.062	0.095	0	0.26	0.34
Dibutyl phthalate	84-74-2	7	0	19	48	0	85	250
Diethyl phthalate	84-66-2	1	0	0.13	0.77	0	0	4.5
Diisobutyl phthalate	84-69-5	15	0	0.72	1.4	0	2.3	7.6
Diisooctylphthalate	27554-26-3	9	0	2.5	8.3	0	6.5	48
Di-n-octyl phthalate	117-84-0	5	0	0.0048	0.013	0	0.027	0.055
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	1.6	2.2	0.84	6.4	7.3
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	0.85	1.7	0	4.3	5.6
Dodecane	112-40-3	4	0	0.34	1.1	0	1.9	5.9
Fluoranthene	206-44-0	21	0	0.19	0.24	0.068	0.66	0.86
Fluorene	86-73-7	15	0	0.3	0.58	0	1.1	3.1
Furan, 2-methyl	534-22-5	34	0	5.3	4.5	3.9	13	21
Heptanal	111-71-7	7	0	0.74	1.9	0	6.5	6.5
Hexadecane	544-76-3	11	0	1.6	2.9	0	8.5	11
2,5-Hexanedione	110-13-4	5	0	1.4	3.6	0	8.9	13
Indan	496-11-7	12	0	0.68	1.3	0	3	5.4
Mesitylene	108-67-8	15	0	1.4	2.5	0	5.5	11



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh} -field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Methacrolein	78-85-3	20	0	3.8	4.9	2.2	11	22
Methyl Isobutyl Ketone	108-10-1	5	0	0.79	2.1	0	6.1	9.2
Naphthalene	91-20-3	9	0	1.3	2.8	0	5.8	13
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.019	0.054	0	0.18	0.18
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.13	0.21	0	0.63	0.75
Naphthalene, 1-methyl-	90-12-0	15	0	1.1	1.9	0	4.9	7.5
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.037	0.056	0	0.13	0.13
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.1	0.15	0	0.46	0.53
Naphthalene, 2-methyl-	91-57-6	10	0	1.7	3.5	0	8.1	16
1-Octadecene	112-88-9	16	0	0.22	0.29	0	0.77	0.91
Octanal	124-13-0	24	0	2.2	2.6	1.4	6.8	10
Octane	111-65-9	10	0	3	5.5	0	13	21
17-Pentatriacontene	6971-40-0	2	0	0.037	0.16	0	0.15	0.83
N-Phenylbenzamide	93-98-1	6	0	0.45	1	0	2.8	2.8
Phenanthrene	85-01-8	17	0	0.65	1	0.35	2.8	4.2
Phenanthrene, 1-methyl	832-69-9	17	0	0.041	0.053	0.019	0.15	0.17
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.075	0.098	0.051	0.27	0.32
Phenanthrene, 3-methyl	832-71-3	18	0	0.089	0.12	0.048	0.33	0.38
Propionaldehyde	123-38-6	12	0	9.2	19	0	37	90
Pyrene	129-00-0	20	0	0.16	0.2	0.063	0.58	0.69
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.0018	0.0089	0	0.0031	0.051
Resorcinol	108-46-3	18	0	0.94	1.4	0.38	3.1	5.9
m-Tolualdehyde	620-23-5	19	0	14	14	11	37	45
TXIB "Kodaflex"	6846-50-0	1	0	0.099	0.59	0	0	3.5
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.19	0.27	0	0.67	0.97
Undecane	1120-21-4	5	0	0.66	1.7	0	5.1	6.6
Valeraldehyde	110-62-3	11	0	47	81	0	210	230
Non-Field-Related Chemicals								



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene	71-43-2	35	4.5	30	25	21	76	120
Benzene, 1,4-dichloro	106-46-7	7	0	0.96	2.1	0	5.9	5.9
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	29	30	19	86	110
2-Butoxyethanol	111-76-2	1	0	0.25	1.5	0	0	8.7
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	3.7	22	0	0	130
Decanal	112-31-2	7	0	1.4	4.8	0	5.3	28
Ethylbenzene	100-41-4	13	0	8.6	15	0	35	58
Heptane	142-82-5	20	0	12	17	2.2	48	75
Hexanal	66-25-1	30	0	40	51	18	150	200
Hexane	110-54-3	26	0	34	75	15	89	440
1-Hexanol, 2-ethyl-	104-76-7	6	0	0.39	1	0	2	4.6
Nonanal	124-19-6	2	0	0.39	1.6	0	2.1	6.8
Phenol	108-95-2	18	0	2.9	3.7	1.7	10	10
Tetrachloroethylene	127-18-4	7	0	2.4	5.7	0	16	21
Tetradecane	629-59-4	3	0	0.41	1.4	0	3.6	7.1
m/p-Xylene	106-42-3	25	0	29	42	12	99	170
o-Xylene	95-47-6	13	0	9.7	16	0	39	64

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Values are rounded to two significant figures.

Table F-161. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter) —Combined Gender Referees 50<70 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.05	0.1	0	0.21	0.44
Acetone	67-64-1	34	79	990	1100	530	3200	3500



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	10	0	0.33	0.57	0	1.3	2.2
Anthracene	120-12-7	12	0	0.016	0.031	0	0.072	0.12
Anthracene, 2-methyl-	613-12-7	15	0	0.0043	0.0061	0	0.017	0.021
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.0049	0.0088	0	0.014	0.048
Anthracene, 9-phenyl	602-55-1	1	0	0.00045	0.0027	0	0	0.015
Benz[a]anthracene	56-55-3	1	0	0.00025	0.0014	0	0	0.0084
Benzaldehyde	100-52-7	15	0	4.5	6.9	0	16	27
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	0.66	2	0	2.9	11
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.13	0.33	0	0.78	1.4
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	5.9	10	0	21	49
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.26	0.7	0	1.4	3.6
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	0.38	0.75	0	1.7	3.4
Benzene, butyl-	104-51-8	2	0	0.13	0.64	0	0.28	3.7
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.19	0.22	0.11	0.61	0.74
Benzo[b]fluoranthene	205-99-2	2	0	0.0012	0.005	0	0.0074	0.021
7H-Benzo[c]fluorene	205-12-9	13	0	0.0027	0.0041	0	0.0071	0.018
Benzo[k]fluoranthene	207-08-9	4	0	0.0015	0.0046	0	0.012	0.017
Benzothiazole	95-16-9	19	0	1.9	2.1	1	6.2	6.2
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.15	0.2	0.075	0.51	0.88
2-Benzothiazolone	934-34-9	5	0	0.23	0.61	0	1.5	2.3
Benzyl butyl phthalate	85-68-7	7	0	0.17	0.37	0	1.2	1.2
Butanal	123-72-8	13	0	15	38	0	91	190
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	8.1	14	2.5	29	66
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	2.9	4.7	0	11	16
p-Cymene	99-87-6	21	0	1.3	1.5	0.8	4.2	5.9
Decane	124-18-5	11	0	2.7	4.9	0	15	15
Dibenz[a,h]anthracene	53-70-3	14	0	0.0072	0.014	0	0.036	0.061
Dibenzothiophene	132-65-0	14	0	0.062	0.095	0	0.26	0.34
Dibutyl phthalate	84-74-2	7	0	19	48	0	85	250
Diethyl phthalate	84-66-2	1	0	0.13	0.77	0	0	4.5
Diisobutyl phthalate	84-69-5	15	0	0.72	1.4	0	2.3	7.6
Diisooctylphthalate	27554-26-3	9	0	2.5	8.3	0	6.5	48
Di-n-octyl phthalate	117-84-0	5	0	0.0049	0.013	0	0.027	0.055
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	1.6	2.2	0.85	6.4	7.3
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	0.85	1.7	0	4.3	5.6



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dodecane	112-40-3	4	0	0.34	1.1	0	1.9	5.9
Fluoranthene	206-44-0	21	0	0.19	0.24	0.068	0.66	0.86
Fluorene	86-73-7	15	0	0.3	0.59	0	1.1	3.1
Furan, 2-methyl	534-22-5	34	0	5.4	4.5	4	13	21
Heptanal	111-71-7	7	0	0.74	1.9	0	6.5	6.5
Hexadecane	544-76-3	11	0	1.6	2.9	0	8.5	11
2,5-Hexanedione	110-13-4	5	0	1.4	3.6	0	9	13
Indan	496-11-7	12	0	0.69	1.3	0	3	5.5
Mesitylene	108-67-8	15	0	1.4	2.5	0	5.5	11
Methacrolein	78-85-3	20	0	3.8	4.9	2.2	11	22
Methyl Isobutyl Ketone	108-10-1	5	0	0.79	2.1	0	6.1	9.2
Naphthalene	91-20-3	9	0	1.3	2.8	0	5.8	13
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.019	0.054	0	0.18	0.18
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.13	0.21	0	0.63	0.75
Naphthalene, 1-methyl-	90-12-0	15	0	1.1	1.9	0	4.9	7.5
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.037	0.056	0	0.13	0.13
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.1	0.15	0	0.46	0.53
Naphthalene, 2-methyl-	91-57-6	10	0	1.7	3.5	0	8.1	16
1-Octadecene	112-88-9	16	0	0.22	0.29	0	0.77	0.91
Octanal	124-13-0	24	0	2.3	2.6	1.4	6.8	10
Octane	111-65-9	10	0	3	5.6	0	13	21
17-Pentatriacontene	6971-40-0	2	0	0.037	0.16	0	0.15	0.83
N-Phenylbenzamide	93-98-1	6	0	0.45	1	0	2.8	2.8
Phenanthrene	85-01-8	17	0	0.65	1	0.35	2.8	4.2
Phenanthrene, 1-methyl	832-69-9	17	0	0.041	0.054	0.019	0.15	0.17
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.075	0.098	0.051	0.28	0.32
Phenanthrene, 3-methyl	832-71-3	18	0	0.09	0.12	0.048	0.33	0.38
Propionaldehyde	123-38-6	12	0	9.3	19	0	37	90
Pyrene	129-00-0	20	0	0.16	0.21	0.063	0.58	0.69
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.0018	0.0089	0	0.0031	0.052
Resorcinol	108-46-3	18	0	0.94	1.4	0.38	3.1	5.9
m-Tolualdehyde	620-23-5	19	0	14	14	11	37	45
TXIB "Kodaflex"	6846-50-0	1	0	0.099	0.59	0	0	3.5
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.19	0.27	0	0.67	0.98
Undecane	1120-21-4	5	0	0.66	1.7	0	5.1	6.6
Valeraldehyde	110-62-3	11	0	47	81	0	210	230



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field-Related Chemicals								
Benzene	71-43-2	35	4.5	30	25	21	76	120
Benzene, 1,4-dichloro	106-46-7	7	0	0.97	2.1	0	6	6
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	29	30	19	86	110
2-Butoxyethanol	111-76-2	1	0	0.25	1.5	0	0	8.7
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	3.7	22	0	0	130
Decanal	112-31-2	7	0	1.4	4.8	0	5.3	28
Ethylbenzene	100-41-4	13	0	8.6	15	0	35	58
Heptane	142-82-5	20	0	12	17	2.2	48	75
Hexanal	66-25-1	30	0	40	51	19	150	200
Hexane	110-54-3	26	0	34	75	15	89	440
1-Hexanol, 2-ethyl-	104-76-7	6	0	0.39	1	0	2	4.6
Nonanal	124-19-6	2	0	0.39	1.6	0	2.1	6.9
Phenol	108-95-2	18	0	2.9	3.7	1.7	10	10
Tetrachloroethylene	127-18-4	7	0	2.4	5.7	0	16	21
Tetradecane	629-59-4	3	0	0.41	1.4	0	3.6	7.2
m/p-Xylene	106-42-3	25	0	29	42	12	99	170
o-Xylene	95-47-6	13	0	9.7	16	0	39	64

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-162. **On-Field** Field-Specific^a Chronic Inhalation Exposure Concentration^b for **All General Chemicals** (Chronic $C_{inh-field}$, nanograms per cubic meter) —Combined Gender **Spectators Third Trimester<0 Years**

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.021	0.042	0	0.087	0.18
Acetone	67-64-1	34	32	400	450	220	1300	1400
Aniline	62-53-3	10	0	0.13	0.24	0	0.52	0.9
Anthracene	120-12-7	12	0	0.0067	0.013	0	0.029	0.05
Anthracene, 2-methyl-	613-12-7	15	0	0.0018	0.0025	0	0.0068	0.0086
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.002	0.0036	0	0.0056	0.019
Anthracene, 9-phenyl	602-55-1	1	0	0.00019	0.0011	0	0	0.0063
Benz[a]anthracene	56-55-3	1	0	0.0001	0.00059	0	0	0.0035
Benzaldehyde	100-52-7	15	0	1.8	2.8	0	6.6	11



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	0.27	0.8	0	1.2	4.3
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.052	0.14	0	0.32	0.56
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	2.4	4.3	0	8.6	20
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.11	0.29	0	0.56	1.5
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	0.15	0.31	0	0.68	1.4
Benzene, butyl-	104-51-8	2	0	0.054	0.26	0	0.11	1.5
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.077	0.09	0.046	0.25	0.3
Benzo[b]fluoranthene	205-99-2	2	0	0.00051	0.0021	0	0.003	0.0086
7H-Benzo[c]fluorene	205-12-9	13	0	0.0011	0.0017	0	0.0029	0.0073
Benzo[k]fluoranthene	207-08-9	4	0	0.00063	0.0019	0	0.0048	0.0072
Benzothiazole	95-16-9	19	0	0.76	0.88	0.42	2.5	2.5
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.061	0.08	0.031	0.21	0.36
2-Benzothiazolone	934-34-9	5	0	0.095	0.25	0	0.62	0.92
Benzyl butyl phthalate	85-68-7	7	0	0.071	0.15	0	0.48	0.48
Butanal	123-72-8	13	0	6.3	15	0	37	78
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	3.3	5.6	1	12	27
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	1.2	1.9	0	4.7	6.7
p-Cymene	99-87-6	21	0	0.52	0.63	0.33	1.7	2.4
Decane	124-18-5	11	0	1.1	2	0	6.1	6.3
Dibenz[a,h]anthracene	53-70-3	14	0	0.003	0.0059	0	0.015	0.025
Dibenzothiophene	132-65-0	14	0	0.025	0.039	0	0.11	0.14
Dibutyl phthalate	84-74-2	7	0	7.7	20	0	35	100
Diethyl phthalate	84-66-2	1	0	0.054	0.32	0	0	1.8
Diisobutyl phthalate	84-69-5	15	0	0.29	0.58	0	0.93	3.1
Diisooctylphthalate	27554-26-3	9	0	1	3.4	0	2.7	20
Di-n-octyl phthalate	117-84-0	5	0	0.002	0.0052	0	0.011	0.023
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	0.67	0.91	0.35	2.6	3
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	0.35	0.69	0	1.7	2.3
Dodecane	112-40-3	4	0	0.14	0.47	0	0.79	2.4
Fluoranthene	206-44-0	21	0	0.078	0.099	0.028	0.27	0.35
Fluorene	86-73-7	15	0	0.12	0.24	0	0.43	1.3
Furan, 2-methyl	534-22-5	34	0	2.2	1.8	1.6	5.2	8.4
Heptanal	111-71-7	7	0	0.3	0.76	0	2.7	2.7
Hexadecane	544-76-3	11	0	0.65	1.2	0	3.5	4.7



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2,5-Hexanedione	110-13-4	5	0	0.56	1.5	0	3.7	5.4
Indan	496-11-7	12	0	0.28	0.52	0	1.2	2.2
Mesitylene	108-67-8	15	0	0.57	1	0	2.3	4.7
Methacrolein	78-85-3	20	0	1.6	2	0.89	4.6	8.8
Methyl Isobutyl Ketone	108-10-1	5	0	0.32	0.88	0	2.5	3.8
Naphthalene	91-20-3	9	0	0.55	1.1	0	2.4	5.3
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.0077	0.022	0	0.074	0.074
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.055	0.084	0	0.26	0.31
Naphthalene, 1-methyl-	90-12-0	15	0	0.45	0.76	0	2	3.1
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.015	0.023	0	0.052	0.052
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.041	0.061	0	0.19	0.22
Naphthalene, 2-methyl-	91-57-6	10	0	0.68	1.4	0	3.3	6.5
1-Octadecene	112-88-9	16	0	0.091	0.12	0	0.31	0.37
Octanal	124-13-0	24	0	0.92	1.1	0.57	2.8	4.3
Octane	111-65-9	10	0	1.2	2.3	0	5.2	8.7
17-Pentatriacontene	6971-40-0	2	0	0.015	0.065	0	0.06	0.34
N-Phenylbenzamide	93-98-1	6	0	0.18	0.42	0	1.1	1.1
Phenanthrene	85-01-8	17	0	0.27	0.41	0.14	1.2	1.7
Phenanthrene, 1-methyl	832-69-9	17	0	0.017	0.022	0.0076	0.061	0.07
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.031	0.04	0.021	0.11	0.13
Phenanthrene, 3-methyl	832-71-3	18	0	0.037	0.047	0.02	0.14	0.15
Propionaldehyde	123-38-6	12	0	3.8	7.7	0	15	37
Pyrene	129-00-0	20	0	0.066	0.084	0.026	0.24	0.28
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.00073	0.0037	0	0.0013	0.021
Resorcinol	108-46-3	18	0	0.39	0.56	0.16	1.3	2.4
m-Tolualdehyde	620-23-5	19	0	5.6	5.9	4.5	15	19
TXIB "Kodaflex"	6846-50-0	1	0	0.041	0.24	0	0	1.4
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.078	0.11	0	0.28	0.4
Undecane	1120-21-4	5	0	0.27	0.71	0	2.1	2.7
Valeraldehyde	110-62-3	11	0	19	33	0	87	94
Non-Field-Related Chemicals								
Benzene	71-43-2	35	1.9	12	10	8.8	31	51
Benzene, 1,4-dichloro	106-46-7	7	0	0.4	0.87	0	2.4	2.4
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	12	12	7.9	35	44
2-Butoxyethanol	111-76-2	1	0	0.1	0.6	0	0	3.6



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	1.5	8.9	0	0	53
Decanal	112-31-2	7	0	0.58	2	0	2.2	11
Ethylbenzene	100-41-4	13	0	3.5	6	0	14	24
Heptane	142-82-5	20	0	4.8	7.1	0.91	20	31
Hexanal	66-25-1	30	0	16	21	7.6	63	82
Hexane	110-54-3	26	0	14	31	6.1	37	180
1-Hexanol, 2-ethyl-	104-76-7	6	0	0.16	0.41	0	0.81	1.9
Nonanal	124-19-6	2	0	0.16	0.66	0	0.84	2.8
Phenol	108-95-2	18	0	1.2	1.5	0.71	4.3	4.3
Tetrachloroethylene	127-18-4	7	0	0.98	2.3	0	6.6	8.6
Tetradecane	629-59-4	3	0	0.17	0.59	0	1.5	2.9
m/p-Xylene	106-42-3	25	0	12	17	4.8	41	72
o-Xylene	95-47-6	13	0	4	6.7	0	16	26

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-163. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter) —Combined Gender Spectators 0<2 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.17	0.34	0	0.7	1.5
Acetone	67-64-1	34	260	3200	3600	1800	11000	11000
Aniline	62-53-3	10	0	1.1	1.9	0	4.2	7.2
Anthracene	120-12-7	12	0	0.054	0.1	0	0.24	0.4
Anthracene, 2-methyl-	613-12-7	15	0	0.014	0.02	0	0.054	0.069
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.016	0.029	0	0.045	0.16
Anthracene, 9-phenyl	602-55-1	1	0	0.0015	0.0087	0	0	0.051
Benz[a]anthracene	56-55-3	1	0	0.00082	0.0048	0	0	0.028
Benzaldehyde	100-52-7	15	0	15	23	0	53	90
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	2.2	6.5	0	9.6	35
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.41	1.1	0	2.6	4.5
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	19	34	0	69	160
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.87	2.3	0	4.5	12



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	1.2	2.5	0	5.5	11
Benzene, butyl-	104-51-8	2	0	0.43	2.1	0	0.91	12
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.62	0.72	0.37	2	2.4
Benzo[b]fluoranthene	205-99-2	2	0	0.0041	0.017	0	0.024	0.069
7H-Benzo[c]fluorene	205-12-9	13	0	0.0089	0.014	0	0.023	0.059
Benzo[k]fluoranthene	207-08-9	4	0	0.0051	0.015	0	0.039	0.058
Benzothiazole	95-16-9	19	0	6.1	7	3.4	20	20
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.49	0.64	0.25	1.7	2.9
2-Benzothiazolone	934-34-9	5	0	0.76	2	0	5	7.4
Benzyl butyl phthalate	85-68-7	7	0	0.57	1.2	0	3.9	3.9
Butanal	123-72-8	13	0	51	120	0	300	630
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	27	45	8.2	95	220
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	9.6	15	0	38	54
p-Cymene	99-87-6	21	0	4.2	5.1	2.6	14	20
Decane	124-18-5	11	0	9	16	0	49	51
Dibenz[a,h]anthracene	53-70-3	14	0	0.024	0.047	0	0.12	0.2
Dibenzothiophene	132-65-0	14	0	0.2	0.31	0	0.87	1.1
Dibutyl phthalate	84-74-2	7	0	62	160	0	280	810
Diethyl phthalate	84-66-2	1	0	0.44	2.5	0	0	15
Diisobutyl phthalate	84-69-5	15	0	2.4	4.6	0	7.4	25
Diisooctylphthalate	27554-26-3	9	0	8.2	27	0	21	160
Di-n-octyl phthalate	117-84-0	5	0	0.016	0.042	0	0.091	0.18
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	5.4	7.3	2.8	21	24
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	2.8	5.5	0	14	18
Dodecane	112-40-3	4	0	1.1	3.8	0	6.4	20
Fluoranthene	206-44-0	21	0	0.63	0.8	0.22	2.2	2.8
Fluorene	86-73-7	15	0	0.99	1.9	0	3.5	10
Furan, 2-methyl	534-22-5	34	0	18	15	13	41	68
Heptanal	111-71-7	7	0	2.4	6.1	0	21	21
Hexadecane	544-76-3	11	0	5.3	9.6	0	28	38
2,5-Hexanedione	110-13-4	5	0	4.5	12	0	29	44
Indan	496-11-7	12	0	2.3	4.2	0	9.8	18
Mesitylene	108-67-8	15	0	4.6	8.1	0	18	38
Methacrolein	78-85-3	20	0	13	16	7.2	37	71
Methyl Isobutyl Ketone	108-10-1	5	0	2.6	7.1	0	20	30



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene	91-20-3	9	0	4.4	9.2	0	19	43
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.062	0.18	0	0.6	0.6
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.44	0.68	0	2.1	2.5
Naphthalene, 1-methyl-	90-12-0	15	0	3.6	6.1	0	16	25
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.12	0.19	0	0.42	0.42
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.33	0.49	0	1.5	1.7
Naphthalene, 2-methyl-	91-57-6	10	0	5.5	12	0	27	52
1-Octadecene	112-88-9	16	0	0.73	0.94	0	2.5	3
Octanal	124-13-0	24	0	7.4	8.6	4.6	22	34
Octane	111-65-9	10	0	10	18	0	42	70
17-Pentatriacontene	6971-40-0	2	0	0.12	0.52	0	0.48	2.7
N-Phenylbenzamide	93-98-1	6	0	1.5	3.3	0	9.2	9.2
Phenanthrene	85-01-8	17	0	2.2	3.3	1.2	9.4	14
Phenanthrene, 1-methyl	832-69-9	17	0	0.14	0.18	0.061	0.49	0.56
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.25	0.32	0.17	0.91	1
Phenanthrene, 3-methyl	832-71-3	18	0	0.3	0.38	0.16	1.1	1.2
Propionaldehyde	123-38-6	12	0	31	62	0	120	300
Pyrene	129-00-0	20	0	0.53	0.68	0.21	1.9	2.3
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.0059	0.029	0	0.01	0.17
Resorcinol	108-46-3	18	0	3.1	4.5	1.3	10	19
m-Tolualdehyde	620-23-5	19	0	45	47	36	120	150
TXIB "Kodaflex"	6846-50-0	1	0	0.33	1.9	0	0	11
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.63	0.88	0	2.2	3.2
Undecane	1120-21-4	5	0	2.2	5.7	0	17	22
Valeraldehyde	110-62-3	11	0	150	270	0	700	760
Non-Field-Related Chemicals								
Benzene	71-43-2	35	15	100	81	71	250	410
Benzene, 1,4-dichloro	106-46-7	7	0	3.2	7	0	20	20
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	95	99	63	280	350
2-Butoxyethanol	111-76-2	1	0	0.82	4.9	0	0	29
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	12	72	0	0	420
Decanal	112-31-2	7	0	4.6	16	0	18	91
Ethylbenzene	100-41-4	13	0	28	48	0	120	190
Heptane	142-82-5	20	0	38	57	7.4	160	250



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Hexanal	66-25-1	30	0	130	170	61	510	660
Hexane	110-54-3	26	0	110	250	49	290	1500
1-Hexanol, 2-ethyl-	104-76-7	6	0	1.3	3.3	0	6.5	15
Nonanal	124-19-6	2	0	1.3	5.3	0	6.8	23
Phenol	108-95-2	18	0	9.7	12	5.7	34	34
Tetrachloroethylene	127-18-4	7	0	7.9	19	0	53	69
Tetradecane	629-59-4	3	0	1.3	4.8	0	12	24
m/p-Xylene	106-42-3	25	0	96	140	39	330	580
o-Xylene	95-47-6	13	0	32	54	0	130	210

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-164. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter) —Combined Gender Spectators 2<6 Years

Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.087	0.18	0	0.37	0.76
Acetone	67-64-1	34	140	1700	1900	920	5600	6000
Aniline	62-53-3	10	0	0.56	0.99	0	2.2	3.8
Anthracene	120-12-7	12	0	0.028	0.053	0	0.12	0.21
Anthracene, 2-methyl-	613-12-7	15	0	0.0074	0.011	0	0.029	0.036
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.0085	0.015	0	0.024	0.082
Anthracene, 9-phenyl	602-55-1	1	0	0.00079	0.0046	0	0	0.027
Benz[a]anthracene	56-55-3	1	0	0.00043	0.0025	0	0	0.015
Benzaldehyde	100-52-7	15	0	7.8	12	0	28	47
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	1.1	3.4	0	5.1	18
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.22	0.57	0	1.3	2.3
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	10	18	0	36	84
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.46	1.2	0	2.4	6.2
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	0.65	1.3	0	2.9	5.9
Benzene, butyl-	104-51-8	2	0	0.23	1.1	0	0.48	6.4
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.33	0.38	0.2	1.1	1.3



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[b]fluoranthene	205-99-2	2	0	0.0021	0.0087	0	0.013	0.037
7H-Benzo[c]fluorene	205-12-9	13	0	0.0047	0.0071	0	0.012	0.031
Benzo[k]fluoranthene	207-08-9	4	0	0.0027	0.0079	0	0.02	0.03
Benzothiazole	95-16-9	19	0	3.2	3.7	1.8	11	11
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.26	0.34	0.13	0.88	1.5
2-Benzothiazolone	934-34-9	5	0	0.4	1.1	0	2.6	3.9
Benzyl butyl phthalate	85-68-7	7	0	0.3	0.64	0	2	2
Butanal	123-72-8	13	0	27	66	0	160	330
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	14	24	4.3	50	110
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	5	8.1	0	20	29
p-Cymene	99-87-6	21	0	2.2	2.7	1.4	7.3	10
Decane	124-18-5	11	0	4.7	8.4	0	26	27
Dibenz[a,h]anthracene	53-70-3	14	0	0.012	0.025	0	0.062	0.11
Dibenzothiophene	132-65-0	14	0	0.11	0.16	0	0.46	0.59
Dibutyl phthalate	84-74-2	7	0	33	84	0	150	430
Diethyl phthalate	84-66-2	1	0	0.23	1.3	0	0	7.8
Diisobutyl phthalate	84-69-5	15	0	1.2	2.4	0	3.9	13
Diisooctylphthalate	27554-26-3	9	0	4.3	14	0	11	83
Di-n-octyl phthalate	117-84-0	5	0	0.0084	0.022	0	0.048	0.095
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	2.8	3.8	1.5	11	13
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	1.5	2.9	0	7.4	9.6
Dodecane	112-40-3	4	0	0.59	2	0	3.3	10
Fluoranthene	206-44-0	21	0	0.33	0.42	0.12	1.1	1.5
Fluorene	86-73-7	15	0	0.52	1	0	1.8	5.3
Furan, 2-methyl	534-22-5	34	0	9.3	7.8	6.8	22	36
Heptanal	111-71-7	7	0	1.3	3.2	0	11	11
Hexadecane	544-76-3	11	0	2.8	5	0	15	20
2,5-Hexanedione	110-13-4	5	0	2.4	6.2	0	16	23
Indan	496-11-7	12	0	1.2	2.2	0	5.1	9.4
Mesitylene	108-67-8	15	0	2.4	4.3	0	9.6	20
Methacrolein	78-85-3	20	0	6.6	8.4	3.8	19	37
Methyl Isobutyl Ketone	108-10-1	5	0	1.4	3.7	0	11	16
Naphthalene	91-20-3	9	0	2.3	4.8	0	10	23
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.032	0.093	0	0.31	0.31
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.23	0.36	0	1.1	1.3
Naphthalene, 1-methyl-	90-12-0	15	0	1.9	3.2	0	8.4	13



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.064	0.097	0	0.22	0.22
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.17	0.26	0	0.8	0.91
Naphthalene, 2-methyl-	91-57-6	10	0	2.9	6.1	0	14	27
1-Octadecene	112-88-9	16	0	0.38	0.5	0	1.3	1.6
Octanal	124-13-0	24	0	3.9	4.5	2.4	12	18
Octane	111-65-9	10	0	5.3	9.6	0	22	37
17-Pentatriacontene	6971-40-0	2	0	0.064	0.27	0	0.25	1.4
N-Phenylbenzamide	93-98-1	6	0	0.78	1.8	0	4.8	4.8
Phenanthrene	85-01-8	17	0	1.1	1.8	0.61	4.9	7.3
Phenanthrene, 1-methyl	832-69-9	17	0	0.072	0.093	0.032	0.26	0.29
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.13	0.17	0.088	0.48	0.55
Phenanthrene, 3-methyl	832-71-3	18	0	0.16	0.2	0.084	0.57	0.65
Propionaldehyde	123-38-6	12	0	16	32	0	64	160
Pyrene	129-00-0	20	0	0.28	0.35	0.11	1	1.2
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.0031	0.015	0	0.0054	0.089
Resorcinol	108-46-3	18	0	1.6	2.4	0.66	5.3	10
m-Tolualdehyde	620-23-5	19	0	23	25	19	64	79
TXIB "Kodaflex"	6846-50-0	1	0	0.17	1	0	0	6
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.33	0.46	0	1.2	1.7
Undecane	1120-21-4	5	0	1.1	3	0	8.8	11
Valeraldehyde	110-62-3	11	0	81	140	0	370	400
Non-Field-Related Chemicals								
Benzene	71-43-2	35	7.9	53	43	37	130	210
Benzene, 1,4-dichloro	106-46-7	7	0	1.7	3.7	0	10	10
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	50	52	33	150	180
2-Butoxyethanol	111-76-2	1	0	0.43	2.6	0	0	15
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	6.4	38	0	0	220
Decanal	112-31-2	7	0	2.4	8.3	0	9.2	48
Ethylbenzene	100-41-4	13	0	15	25	0	61	100
Heptane	142-82-5	20	0	20	30	3.9	83	130
Hexanal	66-25-1	30	0	69	88	32	270	350
Hexane	110-54-3	26	0	58	130	26	150	760
1-Hexanol, 2-ethyl-	104-76-7	6	0	0.68	1.7	0	3.4	7.9
Nonanal	124-19-6	2	0	0.68	2.8	0	3.6	12



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Phenol	108-95-2	18	0	5.1	6.4	3	18	18
Tetrachloroethylene	127-18-4	7	0	4.2	9.9	0	28	36
Tetradecane	629-59-4	3	0	0.71	2.5	0	6.2	12
m/p-Xylene	106-42-3	25	0	50	72	20	170	300
o-Xylene	95-47-6	13	0	17	28	0	67	110

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-165. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter)—Combined Gender Spectators 6<11 Years

Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.083	0.17	0	0.35	0.73
Acetone	67-64-1	34	130	1600	1800	880	5300	5700
Aniline	62-53-3	10	0	0.54	0.95	0	2.1	3.6
Anthracene	120-12-7	12	0	0.027	0.051	0	0.12	0.2
Anthracene, 2-methyl-	613-12-7	15	0	0.0071	0.01	0	0.027	0.035
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.0081	0.015	0	0.023	0.079
Anthracene, 9-phenyl	602-55-1	1	0	0.00075	0.0044	0	0	0.026
Benz[a]anthracene	56-55-3	1	0	0.00041	0.0024	0	0	0.014
Benzaldehyde	100-52-7	15	0	7.4	11	0	27	45
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	1.1	3.2	0	4.8	17
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.21	0.54	0	1.3	2.2
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	9.8	17	0	35	80
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.44	1.2	0	2.3	5.9
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	0.62	1.2	0	2.7	5.7
Benzene, butyl-	104-51-8	2	0	0.22	1.1	0	0.46	6.1



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.31	0.36	0.19	1	1.2
Benzo[b]fluoranthene	205-99-2	2	0	0.002	0.0083	0	0.012	0.035
7H-Benzo[c]fluorene	205-12-9	13	0	0.0045	0.0068	0	0.012	0.03
Benzo[k]fluoranthene	207-08-9	4	0	0.0025	0.0075	0	0.019	0.029
Benzothiazole	95-16-9	19	0	3.1	3.5	1.7	10	10
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.24	0.32	0.12	0.84	1.4
2-Benzothiazolone	934-34-9	5	0	0.38	1	0	2.5	3.7
Benzyl butyl phthalate	85-68-7	7	0	0.29	0.61	0	1.9	1.9
Butanal	123-72-8	13	0	25	63	0	150	310
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	13	23	4.1	48	110
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	4.8	7.7	0	19	27
p-Cymene	99-87-6	21	0	2.1	2.5	1.3	7	9.8
Decane	124-18-5	11	0	4.5	8	0	25	25
Dibenz[a,h]anthracene	53-70-3	14	0	0.012	0.024	0	0.059	0.1
Dibenzothiophene	132-65-0	14	0	0.1	0.16	0	0.44	0.57
Dibutyl phthalate	84-74-2	7	0	31	80	0	140	410
Diethyl phthalate	84-66-2	1	0	0.22	1.3	0	0	7.4
Diisobutyl phthalate	84-69-5	15	0	1.2	2.3	0	3.7	13
Diisooctylphthalate	27554-26-3	9	0	4.1	14	0	11	79
Di-n-octyl phthalate	117-84-0	5	0	0.008	0.021	0	0.045	0.091
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	2.7	3.7	1.4	11	12
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	1.4	2.8	0	7	9.2
Dodecane	112-40-3	4	0	0.56	1.9	0	3.2	9.8
Fluoranthene	206-44-0	21	0	0.31	0.4	0.11	1.1	1.4
Fluorene	86-73-7	15	0	0.5	0.97	0	1.7	5.1
Furan, 2-methyl	534-22-5	34	0	8.9	7.4	6.5	21	34
Heptanal	111-71-7	7	0	1.2	3.1	0	11	11



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Hexadecane	544-76-3	11	0	2.6	4.8	0	14	19
2,5-Hexanedione	110-13-4	5	0	2.3	5.9	0	15	22
Indan	496-11-7	12	0	1.1	2.1	0	4.9	9
Mesitylene	108-67-8	15	0	2.3	4.1	0	9.1	19
Methacrolein	78-85-3	20	0	6.3	8	3.6	19	36
Methyl Isobutyl Ketone	108-10-1	5	0	1.3	3.5	0	10	15
Naphthalene	91-20-3	9	0	2.2	4.6	0	9.5	22
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.031	0.089	0	0.3	0.3
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.22	0.34	0	1	1.2
Naphthalene, 1-methyl-	90-12-0	15	0	1.8	3.1	0	8.1	12
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.061	0.093	0	0.21	0.21
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.17	0.25	0	0.76	0.87
Naphthalene, 2-methyl-	91-57-6	10	0	2.8	5.8	0	13	26
1-Octadecene	112-88-9	16	0	0.37	0.47	0	1.3	1.5
Octanal	124-13-0	24	0	3.7	4.3	2.3	11	17
Octane	111-65-9	10	0	5	9.2	0	21	35
17-Pentatriacontene	6971-40-0	2	0	0.061	0.26	0	0.24	1.4
N-Phenylbenzamide	93-98-1	6	0	0.75	1.7	0	4.6	4.6
Phenanthrene	85-01-8	17	0	1.1	1.7	0.58	4.7	7
Phenanthrene, 1-methyl	832-69-9	17	0	0.068	0.089	0.031	0.25	0.28
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.12	0.16	0.084	0.46	0.52
Phenanthrene, 3-methyl	832-71-3	18	0	0.15	0.19	0.08	0.55	0.62
Propionaldehyde	123-38-6	12	0	15	31	0	61	150
Pyrene	129-00-0	20	0	0.27	0.34	0.1	0.95	1.1
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.0029	0.015	0	0.0052	0.085
Resorcinol	108-46-3	18	0	1.6	2.3	0.63	5.1	9.7
m-Tolualdehyde	620-23-5	19	0	22	24	18	61	75



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
TXIB "Kodaflex"	6846-50-0	1	0	0.16	0.97	0	0	5.8
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.31	0.44	0	1.1	1.6
Undecane	1120-21-4	5	0	1.1	2.8	0	8.4	11
Valeraldehyde	110-62-3	11	0	78	130	0	350	380
Non-Field-Related Chemicals								
Benzene	71-43-2	35	7.5	50	41	35	130	200
Benzene, 1,4-dichloro	106-46-7	7	0	1.6	3.5	0	9.9	9.9
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	48	50	32	140	180
2-Butoxyethanol	111-76-2	1	0	0.41	2.4	0	0	14
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	6.1	36	0	0	210
Decanal	112-31-2	7	0	2.3	8	0	8.8	46
Ethylbenzene	100-41-4	13	0	14	24	0	58	96
Heptane	142-82-5	20	0	19	29	3.7	79	120
Hexanal	66-25-1	30	0	65	84	31	250	330
Hexane	110-54-3	26	0	56	120	25	150	730
1-Hexanol, 2-ethyl-	104-76-7	6	0	0.65	1.7	0	3.3	7.6
Nonanal	124-19-6	2	0	0.65	2.7	0	3.4	11
Phenol	108-95-2	18	0	4.9	6.1	2.9	17	17
Tetrachloroethylene	127-18-4	7	0	4	9.4	0	27	35
Tetradecane	629-59-4	3	0	0.67	2.4	0	5.9	12
m/p-Xylene	106-42-3	25	0	48	69	20	160	290
o-Xylene	95-47-6	13	0	16	27	0	64	110

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-166. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter)—Combined Gender Spectators 11<16 Years

Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.036	0.073	0	0.15	0.31
Acetone	67-64-1	34	56	700	790	380	2300	2500
Aniline	62-53-3	10	0	0.23	0.41	0	0.91	1.6
Anthracene	120-12-7	12	0	0.012	0.022	0	0.051	0.087
Anthracene, 2-methyl-	613-12-7	15	0	0.0031	0.0043	0	0.012	0.015
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.0035	0.0063	0	0.0098	0.034
Anthracene, 9-phenyl	602-55-1	1	0	0.00032	0.0019	0	0	0.011
Benz[a]anthracene	56-55-3	1	0	0.00018	0.001	0	0	0.006
Benzaldehyde	100-52-7	15	0	3.2	5	0	12	19
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	0.47	1.4	0	2.1	7.5
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.09	0.24	0	0.56	0.97
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	4.2	7.5	0	15	35
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.19	0.5	0	0.97	2.5
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	0.27	0.54	0	1.2	2.4
Benzene, butyl-	104-51-8	2	0	0.094	0.46	0	0.2	2.6
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.13	0.16	0.081	0.44	0.53
Benzo[b]fluoranthene	205-99-2	2	0	0.00088	0.0036	0	0.0053	0.015
7H-Benzo[c]fluorene	205-12-9	13	0	0.0019	0.0029	0	0.005	0.013
Benzo[k]fluoranthene	207-08-9	4	0	0.0011	0.0032	0	0.0084	0.012
Benzothiazole	95-16-9	19	0	1.3	1.5	0.74	4.4	4.4
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.11	0.14	0.054	0.36	0.63
2-Benzothiazolone	934-34-9	5	0	0.17	0.43	0	1.1	1.6
Benzyl butyl phthalate	85-68-7	7	0	0.12	0.26	0	0.84	0.84
Butanal	123-72-8	13	0	11	27	0	65	140
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	5.8	9.8	1.8	21	47
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	2.1	3.3	0	8.1	12
p-Cymene	99-87-6	21	0	0.9	1.1	0.57	3	4.2
Decane	124-18-5	11	0	1.9	3.5	0	11	11
Dibenz[a,h]anthracene	53-70-3	14	0	0.0051	0.01	0	0.026	0.043
Dibenzothiophene	132-65-0	14	0	0.044	0.068	0	0.19	0.24
Dibutyl phthalate	84-74-2	7	0	13	34	0	61	180



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Diethyl phthalate	84-66-2	1	0	0.094	0.55	0	0	3.2
Diisobutyl phthalate	84-69-5	15	0	0.51	1	0	1.6	5.4
Diisooctylphthalate	27554-26-3	9	0	1.8	5.9	0	4.6	34
Di-n-octyl phthalate	117-84-0	5	0	0.0035	0.009	0	0.02	0.039
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	1.2	1.6	0.6	4.6	5.2
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	0.61	1.2	0	3	4
Dodecane	112-40-3	4	0	0.24	0.82	0	1.4	4.2
Fluoranthene	206-44-0	21	0	0.14	0.17	0.049	0.47	0.61
Fluorene	86-73-7	15	0	0.21	0.42	0	0.76	2.2
Furan, 2-methyl	534-22-5	34	0	3.8	3.2	2.8	9	15
Heptanal	111-71-7	7	0	0.53	1.3	0	4.6	4.6
Hexadecane	544-76-3	11	0	1.1	2.1	0	6.1	8.1
2,5-Hexanedione	110-13-4	5	0	0.97	2.5	0	6.4	9.5
Indan	496-11-7	12	0	0.49	0.91	0	2.1	3.9
Mesitylene	108-67-8	15	0	0.99	1.8	0	4	8.2
Methacrolein	78-85-3	20	0	2.7	3.5	1.6	8	15
Methyl Isobutyl Ketone	108-10-1	5	0	0.56	1.5	0	4.4	6.6
Naphthalene	91-20-3	9	0	0.95	2	0	4.1	9.3
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.013	0.038	0	0.13	0.13
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.095	0.15	0	0.45	0.54
Naphthalene, 1-methyl-	90-12-0	15	0	0.78	1.3	0	3.5	5.4
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.026	0.04	0	0.09	0.09
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.072	0.11	0	0.33	0.38
Naphthalene, 2-methyl-	91-57-6	10	0	1.2	2.5	0	5.8	11
1-Octadecene	112-88-9	16	0	0.16	0.2	0	0.55	0.65
Octanal	124-13-0	24	0	1.6	1.9	0.99	4.8	7.4
Octane	111-65-9	10	0	2.2	4	0	9	15
17-Pentatriacontene	6971-40-0	2	0	0.026	0.11	0	0.1	0.6
N-Phenylbenzamide	93-98-1	6	0	0.32	0.72	0	2	2
Phenanthrene	85-01-8	17	0	0.47	0.72	0.25	2	3
Phenanthrene, 1-methyl	832-69-9	17	0	0.029	0.038	0.013	0.11	0.12
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.054	0.07	0.036	0.2	0.23
Phenanthrene, 3-methyl	832-71-3	18	0	0.064	0.082	0.035	0.24	0.27
Propionaldehyde	123-38-6	12	0	6.6	13	0	26	64
Pyrene	129-00-0	20	0	0.11	0.15	0.045	0.41	0.49



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.0013	0.0064	0	0.0022	0.037
Resorcinol	108-46-3	18	0	0.67	0.97	0.27	2.2	4.2
m-Tolualdehyde	620-23-5	19	0	9.7	10	7.8	27	32
TXIB "Kodaflex"	6846-50-0	1	0	0.071	0.42	0	0	2.5
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.14	0.19	0	0.48	0.7
Undecane	1120-21-4	5	0	0.47	1.2	0	3.6	4.7
Valeraldehyde	110-62-3	11	0	33	58	0	150	160
Non-Field-Related Chemicals								
Benzene	71-43-2	35	3.2	22	18	15	54	88
Benzene, 1,4-dichloro	106-46-7	7	0	0.69	1.5	0	4.3	4.3
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	21	21	14	61	76
2-Butoxyethanol	111-76-2	1	0	0.18	1.1	0	0	6.2
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	2.6	16	0	0	92
Decanal	112-31-2	7	0	1	3.4	0	3.8	20
Ethylbenzene	100-41-4	13	0	6.2	10	0	25	41
Heptane	142-82-5	20	0	8.3	12	1.6	34	54
Hexanal	66-25-1	30	0	28	36	13	110	140
Hexane	110-54-3	26	0	24	54	11	64	310
1-Hexanol, 2-ethyl-	104-76-7	6	0	0.28	0.72	0	1.4	3.3
Nonanal	124-19-6	2	0	0.28	1.2	0	1.5	4.9
Phenol	108-95-2	18	0	2.1	2.6	1.2	7.4	7.4
Tetrachloroethylene	127-18-4	7	0	1.7	4.1	0	12	15
Tetradecane	629-59-4	3	0	0.29	1	0	2.6	5.1
m/p-Xylene	106-42-3	25	0	21	30	8.4	71	120
o-Xylene	95-47-6	13	0	6.9	12	0	28	46

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-167. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter)—Combined Gender Spectators 16<30 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.024	0.048	0	0.1	0.21



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh} -field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acetone	67-64-1	34	37	460	520	250	1500	1600
Aniline	62-53-3	10	0	0.15	0.27	0	0.6	1
Anthracene	120-12-7	12	0	0.0077	0.015	0	0.034	0.057
Anthracene, 2-methyl-	613-12-7	15	0	0.002	0.0029	0	0.0078	0.0099
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.0023	0.0041	0	0.0064	0.022
Anthracene, 9-phenyl	602-55-1	1	0	0.00021	0.0012	0	0	0.0073
Benz[a]anthracene	56-55-3	1	0	0.00012	0.00068	0	0	0.004
Benzaldehyde	100-52-7	15	0	2.1	3.3	0	7.6	13
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	0.31	0.92	0	1.4	4.9
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.059	0.16	0	0.37	0.64
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	2.8	4.9	0	9.9	23
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.12	0.33	0	0.64	1.7
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	0.18	0.35	0	0.78	1.6
Benzene, butyl-	104-51-8	2	0	0.062	0.3	0	0.13	1.7
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.088	0.1	0.053	0.29	0.35
Benzo[b]fluoranthene	205-99-2	2	0	0.00058	0.0024	0	0.0035	0.0099
7H-Benzo[c]fluorene	205-12-9	13	0	0.0013	0.0019	0	0.0033	0.0084
Benzo[k]fluoranthene	207-08-9	4	0	0.00073	0.0021	0	0.0056	0.0082
Benzothiazole	95-16-9	19	0	0.88	1	0.49	2.9	2.9
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.07	0.092	0.035	0.24	0.41
2-Benzothiazolone	934-34-9	5	0	0.11	0.29	0	0.72	1.1
Benzyl butyl phthalate	85-68-7	7	0	0.081	0.17	0	0.55	0.55
Butanal	123-72-8	13	0	7.2	18	0	43	89
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	3.8	6.5	1.2	14	31
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	1.4	2.2	0	5.4	7.8
p-Cymene	99-87-6	21	0	0.59	0.72	0.38	2	2.8
Decane	124-18-5	11	0	1.3	2.3	0	7	7.2
Dibenz[a,h]anthracene	53-70-3	14	0	0.0034	0.0068	0	0.017	0.029
Dibenzothiophene	132-65-0	14	0	0.029	0.045	0	0.12	0.16
Dibutyl phthalate	84-74-2	7	0	8.9	23	0	40	120
Diethyl phthalate	84-66-2	1	0	0.062	0.36	0	0	2.1
Diisobutyl phthalate	84-69-5	15	0	0.34	0.66	0	1.1	3.6
Diisooctylphthalate	27554-26-3	9	0	1.2	3.9	0	3.1	23
Di-n-octyl phthalate	117-84-0	5	0	0.0023	0.0059	0	0.013	0.026
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	0.77	1	0.4	3	3.4



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh} -field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	0.4	0.79	0	2	2.6
Dodecane	112-40-3	4	0	0.16	0.54	0	0.91	2.8
Fluoranthene	206-44-0	21	0	0.089	0.11	0.032	0.31	0.4
Fluorene	86-73-7	15	0	0.14	0.28	0	0.5	1.4
Furan, 2-methyl	534-22-5	34	0	2.5	2.1	1.9	5.9	9.7
Heptanal	111-71-7	7	0	0.35	0.87	0	3	3
Hexadecane	544-76-3	11	0	0.75	1.4	0	4	5.4
2,5-Hexanedione	110-13-4	5	0	0.64	1.7	0	4.2	6.2
Indan	496-11-7	12	0	0.32	0.6	0	1.4	2.6
Mesitylene	108-67-8	15	0	0.65	1.2	0	2.6	5.4
Methacrolein	78-85-3	20	0	1.8	2.3	1	5.3	10
Methyl Isobutyl Ketone	108-10-1	5	0	0.37	1	0	2.9	4.3
Naphthalene	91-20-3	9	0	0.63	1.3	0	2.7	6.1
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.0088	0.025	0	0.085	0.085
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.063	0.097	0	0.3	0.35
Naphthalene, 1-methyl-	90-12-0	15	0	0.51	0.88	0	2.3	3.5
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.017	0.026	0	0.059	0.059
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.047	0.07	0	0.22	0.25
Naphthalene, 2-methyl-	91-57-6	10	0	0.78	1.7	0	3.8	7.5
1-Octadecene	112-88-9	16	0	0.1	0.13	0	0.36	0.43
Octanal	124-13-0	24	0	1.1	1.2	0.66	3.2	4.9
Octane	111-65-9	10	0	1.4	2.6	0	6	10
17-Pentatriacontene	6971-40-0	2	0	0.017	0.074	0	0.069	0.39
N-Phenylbenzamide	93-98-1	6	0	0.21	0.48	0	1.3	1.3
Phenanthrene	85-01-8	17	0	0.31	0.48	0.16	1.3	2
Phenanthrene, 1-methyl	832-69-9	17	0	0.019	0.025	0.0088	0.07	0.08
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.035	0.046	0.024	0.13	0.15
Phenanthrene, 3-methyl	832-71-3	18	0	0.042	0.054	0.023	0.16	0.18
Propionaldehyde	123-38-6	12	0	4.4	8.8	0	17	42
Pyrene	129-00-0	20	0	0.076	0.096	0.03	0.27	0.33
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.00084	0.0042	0	0.0015	0.024
Resorcinol	108-46-3	18	0	0.44	0.64	0.18	1.5	2.8
m-Tolualdehyde	620-23-5	19	0	6.4	6.7	5.2	17	21
TXIB "Kodaflex"	6846-50-0	1	0	0.047	0.28	0	0	1.6
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.09	0.13	0	0.32	0.46



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Undecane	1120-21-4	5	0	0.31	0.81	0	2.4	3.1
Valeraldehyde	110-62-3	11	0	22	38	0	100	110
Non-Field-Related Chemicals								
Benzene	71-43-2	35	2.1	14	12	10	36	58
Benzene, 1,4-dichloro	106-46-7	7	0	0.45	1	0	2.8	2.8
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	14	14	9	41	50
2-Butoxyethanol	111-76-2	1	0	0.12	0.69	0	0	4.1
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	1.7	10	0	0	61
Decanal	112-31-2	7	0	0.66	2.3	0	2.5	13
Ethylbenzene	100-41-4	13	0	4.1	6.9	0	17	27
Heptane	142-82-5	20	0	5.5	8.1	1.1	23	35
Hexanal	66-25-1	30	0	19	24	8.7	72	94
Hexane	110-54-3	26	0	16	35	7	42	210
1-Hexanol, 2-ethyl-	104-76-7	6	0	0.18	0.47	0	0.93	2.2
Nonanal	124-19-6	2	0	0.18	0.76	0	0.97	3.2
Phenol	108-95-2	18	0	1.4	1.7	0.82	4.9	4.9
Tetrachloroethylene	127-18-4	7	0	1.1	2.7	0	7.6	9.9
Tetradecane	629-59-4	3	0	0.19	0.68	0	1.7	3.4
m/p-Xylene	106-42-3	25	0	14	20	5.6	47	82
o-Xylene	95-47-6	13	0	4.6	7.7	0	18	30

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-168. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter)—Combined Gender Spectators 30<40 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.021	0.042	0	0.088	0.18
Acetone	67-64-1	34	32	400	450	220	1300	1400
Aniline	62-53-3	10	0	0.13	0.24	0	0.52	0.9
Anthracene	120-12-7	12	0	0.0068	0.013	0	0.03	0.05
Anthracene, 2-methyl-	613-12-7	15	0	0.0018	0.0025	0	0.0068	0.0087
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.002	0.0036	0	0.0056	0.02



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh} -field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Anthracene, 9-phenyl	602-55-1	1	0	0.00019	0.0011	0	0	0.0064
Benz[a]anthracene	56-55-3	1	0	0.0001	0.00059	0	0	0.0035
Benzaldehyde	100-52-7	15	0	1.8	2.9	0	6.7	11
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	0.27	0.81	0	1.2	4.3
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.052	0.14	0	0.32	0.56
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	2.4	4.3	0	8.6	20
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.11	0.29	0	0.56	1.5
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	0.15	0.31	0	0.68	1.4
Benzene, butyl-	104-51-8	2	0	0.054	0.26	0	0.11	1.5
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.077	0.09	0.046	0.25	0.3
Benzo[b]fluoranthene	205-99-2	2	0	0.00051	0.0021	0	0.003	0.0087
7H-Benzo[c]fluorene	205-12-9	13	0	0.0011	0.0017	0	0.0029	0.0074
Benzo[k]fluoranthene	207-08-9	4	0	0.00063	0.0019	0	0.0049	0.0072
Benzothiazole	95-16-9	19	0	0.77	0.88	0.43	2.6	2.6
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.061	0.08	0.031	0.21	0.36
2-Benzothiazolone	934-34-9	5	0	0.095	0.25	0	0.63	0.93
Benzyl butyl phthalate	85-68-7	7	0	0.071	0.15	0	0.48	0.48
Butanal	123-72-8	13	0	6.3	16	0	37	78
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	3.3	5.6	1	12	27
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	1.2	1.9	0	4.7	6.8
p-Cymene	99-87-6	21	0	0.52	0.63	0.33	1.7	2.4
Decane	124-18-5	11	0	1.1	2	0	6.1	6.3
Dibenz[a,h]anthracene	53-70-3	14	0	0.003	0.0059	0	0.015	0.025
Dibenzothiophene	132-65-0	14	0	0.026	0.039	0	0.11	0.14
Dibutyl phthalate	84-74-2	7	0	7.8	20	0	35	100
Diethyl phthalate	84-66-2	1	0	0.054	0.32	0	0	1.8
Diisobutyl phthalate	84-69-5	15	0	0.29	0.58	0	0.93	3.1
Diisooctylphthalate	27554-26-3	9	0	1	3.4	0	2.7	20
Di-n-octyl phthalate	117-84-0	5	0	0.002	0.0052	0	0.011	0.023
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	0.67	0.91	0.35	2.6	3
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	0.35	0.69	0	1.8	2.3
Dodecane	112-40-3	4	0	0.14	0.47	0	0.79	2.4
Fluoranthene	206-44-0	21	0	0.078	0.1	0.028	0.27	0.35
Fluorene	86-73-7	15	0	0.12	0.24	0	0.44	1.3



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh} -field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Furan, 2-methyl	534-22-5	34	0	2.2	1.8	1.6	5.2	8.5
Heptanal	111-71-7	7	0	0.3	0.76	0	2.7	2.7
Hexadecane	544-76-3	11	0	0.66	1.2	0	3.5	4.7
2,5-Hexanedione	110-13-4	5	0	0.56	1.5	0	3.7	5.5
Indan	496-11-7	12	0	0.28	0.53	0	1.2	2.2
Mesitylene	108-67-8	15	0	0.57	1	0	2.3	4.7
Methacrolein	78-85-3	20	0	1.6	2	0.89	4.6	8.9
Methyl Isobutyl Ketone	108-10-1	5	0	0.32	0.88	0	2.5	3.8
Naphthalene	91-20-3	9	0	0.55	1.1	0	2.4	5.4
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.0077	0.022	0	0.075	0.075
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.055	0.085	0	0.26	0.31
Naphthalene, 1-methyl-	90-12-0	15	0	0.45	0.77	0	2	3.1
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.015	0.023	0	0.052	0.052
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.041	0.061	0	0.19	0.22
Naphthalene, 2-methyl-	91-57-6	10	0	0.68	1.4	0	3.3	6.5
1-Octadecene	112-88-9	16	0	0.091	0.12	0	0.32	0.38
Octanal	124-13-0	24	0	0.93	1.1	0.57	2.8	4.3
Octane	111-65-9	10	0	1.3	2.3	0	5.2	8.7
17-Pentatriacontene	6971-40-0	2	0	0.015	0.065	0	0.06	0.34
N-Phenylbenzamide	93-98-1	6	0	0.19	0.42	0	1.1	1.1
Phenanthrene	85-01-8	17	0	0.27	0.42	0.14	1.2	1.7
Phenanthrene, 1-methyl	832-69-9	17	0	0.017	0.022	0.0077	0.061	0.07
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.031	0.04	0.021	0.11	0.13
Phenanthrene, 3-methyl	832-71-3	18	0	0.037	0.048	0.02	0.14	0.15
Propionaldehyde	123-38-6	12	0	3.8	7.7	0	15	37
Pyrene	129-00-0	20	0	0.066	0.084	0.026	0.24	0.28
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.00073	0.0037	0	0.0013	0.021
Resorcinol	108-46-3	18	0	0.39	0.56	0.16	1.3	2.4
m-Tolualdehyde	620-23-5	19	0	5.6	5.9	4.5	15	19
TXIB "Kodaflex"	6846-50-0	1	0	0.041	0.24	0	0	1.4
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.078	0.11	0	0.28	0.4
Undecane	1120-21-4	5	0	0.27	0.71	0	2.1	2.7
Valeraldehyde	110-62-3	11	0	19	33	0	87	95
Non-Field-Related Chemicals								



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene	71-43-2	35	1.9	13	10	8.8	31	51
Benzene, 1,4-dichloro	106-46-7	7	0	0.4	0.87	0	2.5	2.5
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	12	12	7.9	35	44
2-Butoxyethanol	111-76-2	1	0	0.1	0.61	0	0	3.6
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	1.5	9	0	0	53
Decanal	112-31-2	7	0	0.58	2	0	2.2	11
Ethylbenzene	100-41-4	13	0	3.6	6	0	15	24
Heptane	142-82-5	20	0	4.8	7.1	0.92	20	31
Hexanal	66-25-1	30	0	16	21	7.6	63	82
Hexane	110-54-3	26	0	14	31	6.1	37	180
1-Hexanol, 2-ethyl-	104-76-7	6	0	0.16	0.41	0	0.82	1.9
Nonanal	124-19-6	2	0	0.16	0.66	0	0.85	2.8
Phenol	108-95-2	18	0	1.2	1.5	0.71	4.3	4.3
Tetrachloroethylene	127-18-4	7	0	0.99	2.3	0	6.6	8.7
Tetradecane	629-59-4	3	0	0.17	0.59	0	1.5	2.9
m/p-Xylene	106-42-3	25	0	12	17	4.9	41	72
o-Xylene	95-47-6	13	0	4	6.7	0	16	26

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-169. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter)—Combined Gender Spectators 40<50 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.021	0.042	0	0.089	0.18
Acetone	67-64-1	34	33	410	460	220	1300	1400
Aniline	62-53-3	10	0	0.14	0.24	0	0.53	0.91
Anthracene	120-12-7	12	0	0.0068	0.013	0	0.03	0.05
Anthracene, 2-methyl-	613-12-7	15	0	0.0018	0.0025	0	0.0069	0.0087
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.002	0.0037	0	0.0057	0.02
Anthracene, 9-phenyl	602-55-1	1	0	0.00019	0.0011	0	0	0.0064
Benz[a]anthracene	56-55-3	1	0	0.0001	0.0006	0	0	0.0035
Benzaldehyde	100-52-7	15	0	1.9	2.9	0	6.7	11
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	0.28	0.81	0	1.2	4.4



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.052	0.14	0	0.32	0.56
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	2.5	4.3	0	8.7	20
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.11	0.29	0	0.57	1.5
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	0.16	0.31	0	0.69	1.4
Benzene, butyl-	104-51-8	2	0	0.055	0.27	0	0.12	1.5
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.078	0.091	0.047	0.25	0.31
Benzo[b]fluoranthene	205-99-2	2	0	0.00052	0.0021	0	0.0031	0.0088
7H-Benzo[c]fluorene	205-12-9	13	0	0.0011	0.0017	0	0.0029	0.0075
Benzo[k]fluoranthene	207-08-9	4	0	0.00064	0.0019	0	0.0049	0.0073
Benzothiazole	95-16-9	19	0	0.77	0.89	0.43	2.6	2.6
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.062	0.081	0.031	0.21	0.36
2-Benzothiazolone	934-34-9	5	0	0.096	0.25	0	0.63	0.94
Benzyl butyl phthalate	85-68-7	7	0	0.072	0.15	0	0.49	0.49
Butanal	123-72-8	13	0	6.4	16	0	38	79
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	3.4	5.7	1	12	27
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	1.2	1.9	0	4.7	6.8
p-Cymene	99-87-6	21	0	0.52	0.64	0.33	1.7	2.5
Decane	124-18-5	11	0	1.1	2	0	6.2	6.4
Dibenz[a,h]anthracene	53-70-3	14	0	0.003	0.006	0	0.015	0.025
Dibenzothiophene	132-65-0	14	0	0.026	0.04	0	0.11	0.14
Dibutyl phthalate	84-74-2	7	0	7.9	20	0	35	100
Diethyl phthalate	84-66-2	1	0	0.055	0.32	0	0	1.9
Diisobutyl phthalate	84-69-5	15	0	0.3	0.59	0	0.94	3.1
Diisooctylphthalate	27554-26-3	9	0	1	3.5	0	2.7	20
Di-n-octyl phthalate	117-84-0	5	0	0.002	0.0052	0	0.011	0.023
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	0.68	0.92	0.35	2.7	3
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	0.35	0.7	0	1.8	2.3
Dodecane	112-40-3	4	0	0.14	0.48	0	0.8	2.5
Fluoranthene	206-44-0	21	0	0.079	0.1	0.028	0.27	0.36
Fluorene	86-73-7	15	0	0.12	0.24	0	0.44	1.3
Furan, 2-methyl	534-22-5	34	0	2.2	1.9	1.6	5.2	8.5
Heptanal	111-71-7	7	0	0.31	0.77	0	2.7	2.7
Hexadecane	544-76-3	11	0	0.66	1.2	0	3.5	4.7
2,5-Hexanedione	110-13-4	5	0	0.57	1.5	0	3.7	5.5



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Indan	496-11-7	12	0	0.28	0.53	0	1.2	2.3
Mesitylene	108-67-8	15	0	0.58	1	0	2.3	4.8
Methacrolein	78-85-3	20	0	1.6	2	0.9	4.7	9
Methyl Isobutyl Ketone	108-10-1	5	0	0.33	0.89	0	2.5	3.8
Naphthalene	91-20-3	9	0	0.56	1.2	0	2.4	5.4
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.0078	0.022	0	0.075	0.075
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.056	0.086	0	0.26	0.31
Naphthalene, 1-methyl-	90-12-0	15	0	0.45	0.78	0	2	3.1
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.015	0.023	0	0.052	0.052
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.042	0.062	0	0.19	0.22
Naphthalene, 2-methyl-	91-57-6	10	0	0.69	1.5	0	3.4	6.6
1-Octadecene	112-88-9	16	0	0.092	0.12	0	0.32	0.38
Octanal	124-13-0	24	0	0.94	1.1	0.58	2.8	4.3
Octane	111-65-9	10	0	1.3	2.3	0	5.3	8.8
17-Pentatriacontene	6971-40-0	2	0	0.015	0.066	0	0.061	0.35
N-Phenylbenzamide	93-98-1	6	0	0.19	0.42	0	1.2	1.2
Phenanthrene	85-01-8	17	0	0.27	0.42	0.15	1.2	1.8
Phenanthrene, 1-methyl	832-69-9	17	0	0.017	0.022	0.0077	0.062	0.071
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.031	0.041	0.021	0.11	0.13
Phenanthrene, 3-methyl	832-71-3	18	0	0.037	0.048	0.02	0.14	0.16
Propionaldehyde	123-38-6	12	0	3.9	7.8	0	15	37
Pyrene	129-00-0	20	0	0.067	0.085	0.026	0.24	0.29
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.00074	0.0037	0	0.0013	0.021
Resorcinol	108-46-3	18	0	0.39	0.57	0.16	1.3	2.4
m-Tolualdehyde	620-23-5	19	0	5.6	5.9	4.6	15	19
TXIB "Kodaflex"	6846-50-0	1	0	0.041	0.24	0	0	1.4
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.079	0.11	0	0.28	0.41
Undecane	1120-21-4	5	0	0.27	0.72	0	2.1	2.7
Valeraldehyde	110-62-3	11	0	20	34	0	88	96
Non-Field-Related Chemicals								
Benzene	71-43-2	35	1.9	13	10	8.9	32	51
Benzene, 1,4-dichloro	106-46-7	7	0	0.4	0.88	0	2.5	2.5
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	12	12	8	36	44
2-Butoxyethanol	111-76-2	1	0	0.1	0.61	0	0	3.6



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	1.5	9.1	0	0	54
Decanal	112-31-2	7	0	0.59	2	0	2.2	12
Ethylbenzene	100-41-4	13	0	3.6	6.1	0	15	24
Heptane	142-82-5	20	0	4.8	7.2	0.93	20	31
Hexanal	66-25-1	30	0	16	21	7.7	64	83
Hexane	110-54-3	26	0	14	31	6.2	37	180
1-Hexanol, 2-ethyl-	104-76-7	6	0	0.16	0.42	0	0.82	1.9
Nonanal	124-19-6	2	0	0.16	0.67	0	0.85	2.8
Phenol	108-95-2	18	0	1.2	1.5	0.72	4.3	4.3
Tetrachloroethylene	127-18-4	7	0	1	2.4	0	6.7	8.7
Tetradecane	629-59-4	3	0	0.17	0.6	0	1.5	3
m/p-Xylene	106-42-3	25	0	12	17	4.9	41	73
o-Xylene	95-47-6	13	0	4	6.8	0	16	27

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-170. On-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter)—Combined Gender Spectators 50<70 Years

Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	11	0	0.021	0.043	0	0.089	0.18
Acetone	67-64-1	34	33	410	460	220	1300	1400
Aniline	62-53-3	10	0	0.14	0.24	0	0.53	0.91
Anthracene	120-12-7	12	0	0.0069	0.013	0	0.03	0.051
Anthracene, 2-methyl-	613-12-7	15	0	0.0018	0.0025	0	0.0069	0.0088
Anthracene, 9,10-dimethyl	781-43-1	16	0	0.002	0.0037	0	0.0057	0.02
Anthracene, 9-phenyl	602-55-1	1	0	0.00019	0.0011	0	0	0.0064
Benz[a]anthracene	56-55-3	1	0	0.0001	0.0006	0	0	0.0035
Benzaldehyde	100-52-7	15	0	1.9	2.9	0	6.8	11
Benzene, 1,2,3-trimethyl-	526-73-8	6	0	0.28	0.82	0	1.2	4.4
Benzene, 1,2,4,5-tetramethyl-	95-93-2	5	0	0.052	0.14	0	0.32	0.56
Benzene, 1,2,4-trimethyl-	95-63-6	13	0	2.5	4.4	0	8.8	20



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	7	0	0.11	0.29	0	0.57	1.5
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	12	0	0.16	0.31	0	0.69	1.4
Benzene, butyl-	104-51-8	2	0	0.055	0.27	0	0.12	1.5
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	26	0	0.078	0.092	0.047	0.26	0.31
Benzo[b]fluoranthene	205-99-2	2	0	0.00052	0.0021	0	0.0031	0.0088
7H-Benzo[c]fluorene	205-12-9	13	0	0.0011	0.0017	0	0.0029	0.0075
Benzo[k]fluoranthene	207-08-9	4	0	0.00064	0.0019	0	0.0049	0.0073
Benzothiazole	95-16-9	19	0	0.78	0.89	0.43	2.6	2.6
Benzothiazole, 2-phenyl-	883-93-2	27	0	0.062	0.081	0.031	0.21	0.37
2-Benzothiazolone	934-34-9	5	0	0.097	0.25	0	0.63	0.94
Benzyl butyl phthalate	85-68-7	7	0	0.072	0.15	0	0.49	0.49
Butanal	123-72-8	13	0	6.4	16	0	38	79
Cyclopentasiloxane, decamethyl-	541-02-6	18	0	3.4	5.7	1	12	27
Cyclotetrasiloxane, octamethyl-	556-67-2	13	0	1.2	1.9	0	4.7	6.9
p-Cymene	99-87-6	21	0	0.53	0.64	0.33	1.8	2.5
Decane	124-18-5	11	0	1.1	2	0	6.2	6.4
Dibenz[a,h]anthracene	53-70-3	14	0	0.003	0.006	0	0.015	0.025
Dibenzothiophene	132-65-0	14	0	0.026	0.04	0	0.11	0.14
Dibutyl phthalate	84-74-2	7	0	7.9	20	0	36	100
Diethyl phthalate	84-66-2	1	0	0.055	0.32	0	0	1.9
Diisobutyl phthalate	84-69-5	15	0	0.3	0.59	0	0.94	3.2
Diisooctylphthalate	27554-26-3	9	0	1	3.5	0	2.7	20
Di-n-octyl phthalate	117-84-0	5	0	0.002	0.0053	0	0.011	0.023
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	0.68	0.92	0.35	2.7	3
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8	0	0.36	0.7	0	1.8	2.3
Dodecane	112-40-3	4	0	0.14	0.48	0	0.81	2.5
Fluoranthene	206-44-0	21	0	0.079	0.1	0.028	0.27	0.36
Fluorene	86-73-7	15	0	0.13	0.24	0	0.44	1.3
Furan, 2-methyl	534-22-5	34	0	2.2	1.9	1.6	5.2	8.6
Heptanal	111-71-7	7	0	0.31	0.77	0	2.7	2.7
Hexadecane	544-76-3	11	0	0.67	1.2	0	3.6	4.8
2,5-Hexanedione	110-13-4	5	0	0.57	1.5	0	3.7	5.5
Indan	496-11-7	12	0	0.29	0.53	0	1.2	2.3
Mesitylene	108-67-8	15	0	0.58	1	0	2.3	4.8



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Methacrolein	78-85-3	20	0	1.6	2	0.91	4.7	9
Methyl Isobutyl Ketone	108-10-1	5	0	0.33	0.89	0	2.6	3.8
Naphthalene	91-20-3	9	0	0.56	1.2	0	2.4	5.4
Naphthalene, 1,2-dimethyl-	573-98-8	4	0	0.0078	0.022	0	0.076	0.076
Naphthalene, 1,6-dimethyl-	575-43-9	14	0	0.056	0.086	0	0.26	0.31
Naphthalene, 1-methyl-	90-12-0	15	0	0.45	0.78	0	2	3.1
Naphthalene, 2-(bromomethyl)-	939-26-4	11	0	0.015	0.023	0	0.053	0.053
Naphthalene, 2,3-dimethyl-	581-40-8	14	0	0.042	0.062	0	0.19	0.22
Naphthalene, 2-methyl-	91-57-6	10	0	0.69	1.5	0	3.4	6.6
1-Octadecene	112-88-9	16	0	0.093	0.12	0	0.32	0.38
Octanal	124-13-0	24	0	0.94	1.1	0.58	2.8	4.3
Octane	111-65-9	10	0	1.3	2.3	0	5.3	8.8
17-Pentatriacontene	6971-40-0	2	0	0.015	0.066	0	0.061	0.35
N-Phenylbenzamide	93-98-1	6	0	0.19	0.42	0	1.2	1.2
Phenanthrene	85-01-8	17	0	0.27	0.42	0.15	1.2	1.8
Phenanthrene, 1-methyl	832-69-9	17	0	0.017	0.022	0.0078	0.062	0.071
Phenanthrene, 2-methyl-	2531-84-2	18	0	0.031	0.041	0.021	0.11	0.13
Phenanthrene, 3-methyl	832-71-3	18	0	0.037	0.048	0.02	0.14	0.16
Propionaldehyde	123-38-6	12	0	3.9	7.8	0	15	38
Pyrene	129-00-0	20	0	0.067	0.085	0.026	0.24	0.29
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	0.00074	0.0037	0	0.0013	0.021
Resorcinol	108-46-3	18	0	0.39	0.57	0.16	1.3	2.5
m-Tolualdehyde	620-23-5	19	0	5.7	6	4.6	15	19
TXIB "Kodaflex"	6846-50-0	1	0	0.041	0.25	0	0	1.5
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	14	0	0.079	0.11	0	0.28	0.41
Undecane	1120-21-4	5	0	0.28	0.72	0	2.1	2.7
Valeraldehyde	110-62-3	11	0	20	34	0	88	96
Non-Field-Related Chemicals								
Benzene	71-43-2	35	1.9	13	10	8.9	32	52
Benzene, 1,4-dichloro	106-46-7	7	0	0.4	0.88	0	2.5	2.5
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	12	13	8	36	44
2-Butoxyethanol	111-76-2	1	0	0.1	0.61	0	0	3.6
Cyclotrisiloxane, hexamethyl-	541-05-9	1	0	1.5	9.1	0	0	54
Decanal	112-31-2	7	0	0.59	2	0	2.2	12



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Ethylbenzene	100-41-4	13	0	3.6	6.1	0	15	24
Heptane	142-82-5	20	0	4.8	7.2	0.93	20	31
Hexanal	66-25-1	30	0	17	21	7.7	64	83
Hexane	110-54-3	26	0	14	31	6.2	37	180
1-Hexanol, 2-ethyl-	104-76-7	6	0	0.16	0.42	0	0.83	1.9
Nonanal	124-19-6	2	0	0.16	0.67	0	0.86	2.9
Phenol	108-95-2	18	0	1.2	1.5	0.72	4.3	4.3
Tetrachloroethylene	127-18-4	7	0	1	2.4	0	6.7	8.8
Tetradecane	629-59-4	3	0	0.17	0.6	0	1.5	3
m/p-Xylene	106-42-3	25	0	12	17	4.9	41	73
o-Xylene	95-47-6	13	0	4.1	6.8	0	16	27

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b On-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-171. Off-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter)—Combined Gender Spectators Third Trimester<0 Years

Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	8	0	0.022	0.047	0	0.13	0.19
Acetone	67-64-1	8	0	0.15	0.29	0	0.52	1.2
Aniline	62-53-3	4	0	0.0041	0.013	0	0.028	0.057
Anthracene	120-12-7	11	0	0.0011	0.002	0	0.0054	0.0076
Anthracene, 2-methyl-	613-12-7	12	0	0.0017	0.0031	0	0.0072	0.014
Anthracene, 9,10-dimethyl	781-43-1	2	0	0.00077	0.0031	0	0.0051	0.013
Anthracene, 9-phenyl	602-55-1	0	0	0	0	0	0	0
Benz[a]anthracene	56-55-3	14	0	2	3.1	0	8	12
Benzaldehyde	100-52-7	6	0	0.34	0.83	0	2.1	3.4
Benzene, 1,2,3-trimethyl-	526-73-8	5	0	0.066	0.18	0	0.37	0.84
Benzene, 1,2,4,5-tetramethyl-	95-93-2	12	0	2.6	4.5	0	11	18
Benzene, 1,2,4-trimethyl-	95-63-6	7	0	0.12	0.28	0	0.7	1.1
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	11	0	0.17	0.34	0	0.71	1.6
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	3	0	0.089	0.31	0	0.76	1.5
Benzene, butyl-	104-51-8	3	0	0.00079	0.0025	0	0.0086	0.0086



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh} -field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	19	0	0.087	0.11	0.036	0.3	0.33
Benzo[b]fluoranthene	205-99-2	3	0	0.00065	0.0021	0	0.0072	0.0072
7H-Benzo[c]fluorene	205-12-9	6	0	0.00065	0.0015	0	0.0029	0.0068
Benzo[k]fluoranthene	207-08-9	2	0	0.1	0.4	0	0.59	1.7
Benzothiazole	95-16-9	15	0	0.0089	0.012	0	0.035	0.037
Benzothiazole, 2-phenyl-	883-93-2	5	0	0.09	0.24	0	0.48	1.1
2-Benzothiazolone	934-34-9	3	0	0.084	0.27	0	0.92	0.92
Benzyl butyl phthalate	85-68-7	12	0	5.9	15	0	41	71
Butanal	123-72-8	14	0	3.1	5.4	0	12	24
Cyclopentasiloxane, decamethyl-	541-02-6	13	0	1.1	1.7	0	4.5	6.5
Cyclotetrasiloxane, octamethyl-	556-67-2	12	0	1.2	2	0	5	7.3
p-Cymene	99-87-6	13	0	0.01	0.017	0	0.046	0.07
Decane	124-18-5	9	0	0.0019	0.0041	0	0.012	0.015
Dibenz[a,h]anthracene	53-70-3	12	0	0.024	0.042	0	0.12	0.18
Dibenzothiophene	132-65-0	5	0	6.2	22	0	24	120
Dibutyl phthalate	84-74-2	2	0	0.22	0.89	0	1.5	3.7
Diethyl phthalate	84-66-2	14	0	0.32	0.52	0	1.5	1.9
Diisobutyl phthalate	84-69-5	6	0	0.3	0.66	0	1.7	1.7
Diisooctylphthalate	27554-26-3	3	0	0.002	0.0066	0	0.023	0.023
Di-n-octyl phthalate	117-84-0	3	0	0.18	0.61	0	1.6	2.4
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	0.81	1.4	0.3	3.1	6.4
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	3	0	0.23	0.8	0	1.7	4
Dodecane	112-40-3	14	0	0.046	0.085	0	0.22	0.39
Fluoranthene	206-44-0	11	0	0.1	0.22	0	0.45	1.1
Fluorene	86-73-7	21	0	0.78	1.2	0.61	2.8	5.6
Furan, 2-methyl	534-22-5	5	0	0.31	0.8	0	2.7	2.7
Heptanal	111-71-7	9	0	0.94	1.9	0	4	8.6
Hexadecane	544-76-3	12	0	0.28	0.54	0	1.3	2.2
2,5-Hexanedione	110-13-4	3	0	0.73	2.6	0	5.4	13
Indan	496-11-7	13	0	0.59	1	0	2.6	4.1
Mesitylene	108-67-8	16	0	1.4	1.9	0	4.6	8.3
Methacrolein	78-85-3	1	0	0.074	0.43	0	0	2.5
Methyl Isobutyl Ketone	108-10-1	9	0	0.6	1.2	0	2.7	5.3
Naphthalene	91-20-3	12	0	0.057	0.1	0	0.28	0.42
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	0.44	0.73	0	2	2.4
Naphthalene, 1,6-dimethyl-	575-43-9	8	0	0.016	0.034	0	0.052	0.16



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh} -field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene, 1-methyl-	90-12-0	12	0	0.043	0.074	0	0.21	0.3
Naphthalene, 2-(bromomethyl)-	939-26-4	12	0	0.73	1.2	0	3.4	4
Naphthalene, 2,3-dimethyl-	581-40-8	4	0	0.18	0.55	0	1.1	2.6
Naphthalene, 2-methyl-	91-57-6	16	0	0.9	1.3	0	3.5	5
1-Octadecene	112-88-9	14	0	0.084	0.13	0	0.38	0.39
Octanal	124-13-0	18	0	0.54	0.68	0.33	1.9	2.4
Octane	111-65-9	12	0	0.22	0.46	0	1.4	1.9
17-Pentatriacontene	6971-40-0	1	0	0.01	0.059	0	0	0.34
N-Phenylbenzamide	93-98-1	9	0	1.5	2.8	0	7.3	10
Phenanthrene	85-01-8	13	0	0.02	0.034	0	0.1	0.13
Phenanthrene, 1-methyl	832-69-9	13	0	0.023	0.04	0	0.11	0.16
Phenanthrene, 2-methyl-	2531-84-2	14	0	0.033	0.056	0	0.16	0.21
Phenanthrene, 3-methyl	832-71-3	1	0	0.00011	0.00064	0	0	0.0037
Propionaldehyde	123-38-6	18	0	0.66	1.1	0.31	2.4	5.4
Pyrene	129-00-0	2	0	0.084	0.34	0	0.5	1.4
Pyridine, 2-(4-methylphenyl)-	4467-06-5	7	0	0.37	0.83	0	2.1	3.3
Resorcinol	108-46-3	34	0.58	5.4	5.9	2.7	16	22
m-Tolualdehyde	620-23-5	6	0	0.016	0.037	0	0.074	0.16
TXIB "Kodaflex"	6846-50-0	7	0	0.36	0.93	0	1.3	4.8
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	13	0	0.056	0.076	0	0.13	0.3
Undecane	1120-21-4	1	0	0.098	0.57	0	0	3.3
Valeraldehyde	110-62-3	34	3.7	13	11	8.2	31	56
Non-Field-Related Chemicals								
Benzene	71-43-2	1	0	0.35	2.1	0	0	12
Benzene, 1,4-dichloro	106-46-7	13	0	3.7	6.4	0	16	27
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	17	0	4.8	7.5	0.91	18	34
2-Butoxyethanol	111-76-2	3	0	1.6	6.1	0	11	33
Cyclotrisiloxane, hexamethyl-	541-05-9	27	0	18	25	8.8	59	120
Decanal	112-31-2	28	0	9.4	11	5.5	29	42
Ethylbenzene	100-41-4	24	0	13	18	4.8	45	71
Heptane	142-82-5	0	0	0	0	0	0	0
Hexanal	66-25-1	11	0	3.9	7.1	0	18	29
Hexane	110-54-3	12	0	1.1	1.7	0	4.3	4.3
1-Hexanol, 2-ethyl-	104-76-7	33	0	12	13	4.8	39	41
Nonanal	124-19-6	2	0	0.13	0.55	0	0.51	2.9
Phenol	108-95-2	34	4.1	29	32	15	83	150
Tetrachloroethylene	127-18-4	11	0	0.2	0.38	0	0.79	1.7



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Tetradecane	629-59-4	7	0	0.76	1.6	0	4.8	4.8
m/p-Xylene	106-42-3	7	0	1	2.3	0	6.7	8.6
o-Xylene	95-47-6	11	0	2.1	8.3	0	6.9	48

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b Off-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-172. Off-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter)—Combined Gender Spectators 0<2 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	8	0	0.18	0.38	0	1	1.6
Acetone	67-64-1	8	0	1.2	2.3	0	4.2	9.9
Aniline	62-53-3	4	0	0.033	0.1	0	0.23	0.46
Anthracene	120-12-7	11	0	0.0087	0.016	0	0.044	0.061
Anthracene, 2-methyl-	613-12-7	12	0	0.014	0.025	0	0.058	0.11
Anthracene, 9,10-dimethyl	781-43-1	2	0	0.0062	0.025	0	0.041	0.1
Anthracene, 9-phenyl	602-55-1	0	0	0	0	0	0	0
Benz[a]anthracene	56-55-3	14	0	16	25	0	64	97
Benzaldehyde	100-52-7	6	0	2.8	6.7	0	17	27
Benzene, 1,2,3-trimethyl-	526-73-8	5	0	0.53	1.5	0	3	6.8
Benzene, 1,2,4,5-tetramethyl-	95-93-2	12	0	21	36	0	90	150
Benzene, 1,2,4-trimethyl-	95-63-6	7	0	0.99	2.3	0	5.6	8.6
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	11	0	1.4	2.8	0	5.8	13
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	3	0	0.72	2.5	0	6.1	12
Benzene, butyl-	104-51-8	3	0	0.0063	0.02	0	0.069	0.069
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	19	0	0.7	0.9	0.29	2.4	2.7
Benzo[b]fluoranthene	205-99-2	3	0	0.0052	0.017	0	0.058	0.058
7H-Benzo[c]fluorene	205-12-9	6	0	0.0052	0.012	0	0.023	0.055
Benzo[k]fluoranthene	207-08-9	2	0	0.8	3.3	0	4.8	14
Benzothiazole	95-16-9	15	0	0.071	0.099	0	0.28	0.3
Benzothiazole, 2-phenyl-	883-93-2	5	0	0.73	1.9	0	3.9	8.5



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Benzothiazolone	934-34-9	3	0	0.67	2.2	0	7.4	7.4
Benzyl butyl phthalate	85-68-7	12	0	47	120	0	330	570
Butanal	123-72-8	14	0	25	44	0	99	200
Cyclopentasiloxane, decamethyl-	541-02-6	13	0	8.6	14	0	36	53
Cyclotetrasiloxane, octamethyl-	556-67-2	12	0	10	16	0	40	59
p-Cymene	99-87-6	13	0	0.082	0.14	0	0.37	0.57
Decane	124-18-5	9	0	0.015	0.033	0	0.096	0.12
Dibenz[a,h]anthracene	53-70-3	12	0	0.19	0.34	0	0.95	1.4
Dibenzothiophene	132-65-0	5	0	50	180	0	190	970
Dibutyl phthalate	84-74-2	2	0	1.8	7.2	0	12	30
Diethyl phthalate	84-66-2	14	0	2.5	4.2	0	12	15
Diisobutyl phthalate	84-69-5	6	0	2.4	5.3	0	13	13
Diisooctylphthalate	27554-26-3	3	0	0.016	0.053	0	0.18	0.18
Di-n-octyl phthalate	117-84-0	3	0	1.4	4.9	0	13	20
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	6.5	11	2.4	25	51
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	3	0	1.8	6.4	0	14	32
Dodecane	112-40-3	14	0	0.37	0.68	0	1.7	3.1
Fluoranthene	206-44-0	11	0	0.83	1.8	0	3.6	8.8
Fluorene	86-73-7	21	0	6.3	10	4.9	23	45
Furan, 2-methyl	534-22-5	5	0	2.5	6.5	0	21	21
Heptanal	111-71-7	9	0	7.6	16	0	32	69
Hexadecane	544-76-3	12	0	2.2	4.3	0	10	18
2,5-Hexanedione	110-13-4	3	0	5.9	21	0	44	110
Indan	496-11-7	13	0	4.7	8.2	0	21	33
Mesitylene	108-67-8	16	0	12	16	0	37	67
Methacrolein	78-85-3	1	0	0.59	3.5	0	0	20
Methyl Isobutyl Ketone	108-10-1	9	0	4.8	9.7	0	22	42
Naphthalene	91-20-3	12	0	0.46	0.82	0	2.3	3.4
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	3.6	5.8	0	16	19
Naphthalene, 1,6-dimethyl-	575-43-9	8	0	0.13	0.27	0	0.42	1.3
Naphthalene, 1-methyl-	90-12-0	12	0	0.34	0.6	0	1.7	2.4
Naphthalene, 2-(bromomethyl)-	939-26-4	12	0	5.8	9.8	0	28	32
Naphthalene, 2,3-dimethyl-	581-40-8	4	0	1.5	4.4	0	9.2	21
Naphthalene, 2-methyl-	91-57-6	16	0	7.2	10	0	28	40
1-Octadecene	112-88-9	14	0	0.67	1	0	3	3.2



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Octanal	124-13-0	18	0	4.3	5.5	2.6	15	20
Octane	111-65-9	12	0	1.8	3.7	0	11	15
17-Pentatriacontene	6971-40-0	1	0	0.083	0.48	0	0	2.7
N-Phenylbenzamide	93-98-1	9	0	12	22	0	58	80
Phenanthrene	85-01-8	13	0	0.16	0.28	0	0.8	1
Phenanthrene, 1-methyl	832-69-9	13	0	0.18	0.32	0	0.86	1.3
Phenanthrene, 2-methyl-	2531-84-2	14	0	0.26	0.45	0	1.3	1.7
Phenanthrene, 3-methyl	832-71-3	1	0	0.00089	0.0051	0	0	0.029
Propionaldehyde	123-38-6	18	0	5.3	9	2.5	20	43
Pyrene	129-00-0	2	0	0.67	2.7	0	4	11
Pyridine, 2-(4-methylphenyl)-	4467-06-5	7	0	3	6.6	0	17	27
Resorcinol	108-46-3	34	4.6	44	47	22	130	180
m-Tolualdehyde	620-23-5	6	0	0.13	0.3	0	0.6	1.3
TXIB "Kodaflex"	6846-50-0	7	0	2.9	7.5	0	10	39
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	13	0	0.45	0.61	0	1	2.4
Undecane	1120-21-4	1	0	0.79	4.6	0	0	27
Valeraldehyde	110-62-3	34	30	110	88	66	250	450
Non-Field-Related Chemicals								
Benzene	71-43-2	1	0	2.8	17	0	0	97
Benzene, 1,4-dichloro	106-46-7	13	0	30	52	0	130	220
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	17	0	38	60	7.4	150	280
2-Butoxyethanol	111-76-2	3	0	13	49	0	87	260
Cyclotrisiloxane, hexamethyl-	541-05-9	27	0	140	200	71	480	960
Decanal	112-31-2	28	0	76	85	44	240	340
Ethylbenzene	100-41-4	24	0	100	150	39	370	570
Heptane	142-82-5	0	0	0	0	0	0	0
Hexanal	66-25-1	11	0	32	57	0	140	230
Hexane	110-54-3	12	0	8.8	14	0	34	34
1-Hexanol, 2-ethyl-	104-76-7	33	0	96	110	38	320	330
Nonanal	124-19-6	2	0	1	4.5	0	4.1	24
Phenol	108-95-2	34	33	230	260	120	670	1200
Tetrachloroethylene	127-18-4	11	0	1.6	3.1	0	6.3	14
Tetradecane	629-59-4	7	0	6.1	13	0	39	39
m/p-Xylene	106-42-3	7	0	8.2	19	0	54	69
o-Xylene	95-47-6	11	0	17	66	0	55	380

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b Off-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures.

Table F-173. **Off-Field** Field-Specific^a Chronic Inhalation Exposure Concentration^b for **All General Chemicals** (Chronic C_{inh-field}, nanograms per cubic meter)—Combined Gender **Spectators 2<6 Years**

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	8	0	0.092	0.2	0	0.53	0.82
Acetone	67-64-1	8	0	0.62	1.2	0	2.2	5.2
Aniline	62-53-3	4	0	0.018	0.054	0	0.12	0.24
Anthracene	120-12-7	11	0	0.0046	0.0085	0	0.023	0.032
Anthracene, 2-methyl-	613-12-7	12	0	0.0073	0.013	0	0.031	0.058
Anthracene, 9,10-dimethyl	781-43-1	2	0	0.0032	0.013	0	0.021	0.054
Anthracene, 9-phenyl	602-55-1	0	0	0	0	0	0	0
Benz[a]anthracene	56-55-3	14	0	8.5	13	0	34	51
Benzaldehyde	100-52-7	6	0	1.5	3.5	0	8.8	14
Benzene, 1,2,3-trimethyl-	526-73-8	5	0	0.28	0.77	0	1.6	3.6
Benzene, 1,2,4,5-tetramethyl-	95-93-2	12	0	11	19	0	47	77
Benzene, 1,2,4-trimethyl-	95-63-6	7	0	0.52	1.2	0	3	4.5
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	11	0	0.73	1.5	0	3	6.8
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	3	0	0.38	1.3	0	3.2	6.4
Benzene, butyl-	104-51-8	3	0	0.0033	0.011	0	0.037	0.037
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	19	0	0.37	0.48	0.15	1.3	1.4
Benzo[b]fluoranthene	205-99-2	3	0	0.0028	0.0088	0	0.03	0.03
7H-Benzo[c]fluorene	205-12-9	6	0	0.0027	0.0065	0	0.012	0.029
Benzo[k]fluoranthene	207-08-9	2	0	0.42	1.7	0	2.5	7.2
Benzothiazole	95-16-9	15	0	0.037	0.052	0	0.15	0.16
Benzothiazole, 2-phenyl-	883-93-2	5	0	0.38	1	0	2	4.5
2-Benzothiazolone	934-34-9	3	0	0.35	1.1	0	3.9	3.9
Benzyl butyl phthalate	85-68-7	12	0	25	65	0	170	300
Butanal	123-72-8	14	0	13	23	0	52	100
Cyclopentasiloxane, decamethyl-	541-02-6	13	0	4.5	7.2	0	19	28
Cyclotetrasiloxane, octamethyl-	556-67-2	12	0	5.2	8.6	0	21	31
p-Cymene	99-87-6	13	0	0.043	0.073	0	0.2	0.3
Decane	124-18-5	9	0	0.0078	0.017	0	0.05	0.064
Dibenz[a,h]anthracene	53-70-3	12	0	0.1	0.18	0	0.5	0.75
Dibenzothiophene	132-65-0	5	0	26	92	0	100	510



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dibutyl phthalate	84-74-2	2	0	0.94	3.8	0	6.2	16
Diethyl phthalate	84-66-2	14	0	1.3	2.2	0	6.4	7.9
Diisobutyl phthalate	84-69-5	6	0	1.3	2.8	0	7.1	7.1
Diisooctylphthalate	27554-26-3	3	0	0.0087	0.028	0	0.095	0.095
Di-n-octyl phthalate	117-84-0	3	0	0.76	2.6	0	6.9	10
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	3.4	5.7	1.3	13	27
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	3	0	0.96	3.4	0	7.4	17
Dodecane	112-40-3	14	0	0.19	0.36	0	0.91	1.6
Fluoranthene	206-44-0	11	0	0.43	0.93	0	1.9	4.6
Fluorene	86-73-7	21	0	3.3	5.2	2.6	12	24
Furan, 2-methyl	534-22-5	5	0	1.3	3.4	0	11	11
Heptanal	111-71-7	9	0	4	8.2	0	17	36
Hexadecane	544-76-3	12	0	1.2	2.3	0	5.5	9.3
2,5-Hexanedione	110-13-4	3	0	3.1	11	0	23	56
Indan	496-11-7	13	0	2.5	4.3	0	11	17
Mesitylene	108-67-8	16	0	6.1	8.2	0	19	35
Methacrolein	78-85-3	1	0	0.31	1.8	0	0	11
Methyl Isobutyl Ketone	108-10-1	9	0	2.5	5.1	0	11	22
Naphthalene	91-20-3	12	0	0.24	0.43	0	1.2	1.8
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	1.9	3.1	0	8.4	10
Naphthalene, 1,6-dimethyl-	575-43-9	8	0	0.067	0.14	0	0.22	0.68
Naphthalene, 1-methyl-	90-12-0	12	0	0.18	0.31	0	0.87	1.3
Naphthalene, 2-(bromomethyl)-	939-26-4	12	0	3.1	5.1	0	15	17
Naphthalene, 2,3-dimethyl-	581-40-8	4	0	0.77	2.3	0	4.8	11
Naphthalene, 2-methyl-	91-57-6	16	0	3.8	5.5	0	15	21
1-Octadecene	112-88-9	14	0	0.35	0.53	0	1.6	1.7
Octanal	124-13-0	18	0	2.3	2.9	1.4	8	10
Octane	111-65-9	12	0	0.94	2	0	5.8	8
17-Pentatriacontene	6971-40-0	1	0	0.044	0.25	0	0	1.4
N-Phenylbenzamide	93-98-1	9	0	6.3	12	0	31	42
Phenanthrene	85-01-8	13	0	0.083	0.15	0	0.42	0.55
Phenanthrene, 1-methyl	832-69-9	13	0	0.095	0.17	0	0.45	0.67
Phenanthrene, 2-methyl-	2531-84-2	14	0	0.14	0.24	0	0.67	0.89
Phenanthrene, 3-methyl	832-71-3	1	0	0.00047	0.0027	0	0	0.015
Propionaldehyde	123-38-6	18	0	2.8	4.8	1.3	10	23
Pyrene	129-00-0	2	0	0.35	1.4	0	2.1	6
Pyridine, 2-(4-methylphenyl)-	4467-06-5	7	0	1.6	3.5	0	8.8	14
Resorcinol	108-46-3	34	2.4	23	25	11	69	95



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
m-Tolualdehyde	620-23-5	6	0	0.068	0.16	0	0.31	0.66
TXIB "Kodaflex"	6846-50-0	7	0	1.5	3.9	0	5.3	20
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	13	0	0.24	0.32	0	0.54	1.3
Undecane	1120-21-4	1	0	0.41	2.4	0	0	14
Valeraldehyde	110-62-3	34	16	55	46	34	130	240
Non-Field-Related Chemicals								
Benzene	71-43-2	1	0	1.5	8.7	0	0	51
Benzene, 1,4-dichloro	106-46-7	13	0	16	27	0	69	110
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	17	0	20	32	3.9	78	150
2-Butoxyethanol	111-76-2	3	0	6.8	26	0	46	140
Cyclotrisiloxane, hexamethyl-	541-05-9	27	0	75	110	37	250	500
Decanal	112-31-2	28	0	40	45	23	120	180
Ethylbenzene	100-41-4	24	0	54	77	20	190	300
Heptane	142-82-5	0	0	0	0	0	0	0
Hexanal	66-25-1	11	0	17	30	0	74	120
Hexane	110-54-3	12	0	4.6	7.1	0	18	18
1-Hexanol, 2-ethyl-	104-76-7	33	0	50	55	20	170	170
Nonanal	124-19-6	2	0	0.55	2.3	0	2.2	12
Phenol	108-95-2	34	17	120	140	62	350	620
Tetrachloroethylene	127-18-4	11	0	0.86	1.6	0	3.3	7.3
Tetradecane	629-59-4	7	0	3.2	7	0	20	20
m/p-Xylene	106-42-3	7	0	4.3	9.9	0	29	36
o-Xylene	95-47-6	11	0	8.8	35	0	29	200

^a 35 field-specific C_{inh-field} values are included in the table.

^b Off-Field field-specific Chronic C_{inh-field} equals the modified field-specific average concentration of a field (C_{air-field} × A_{F_{inh}}) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-174. Off-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic C_{inh-field}, nanograms per cubic meter)—Combined Gender Spectators 6<11 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	8	0	0.088	0.19	0	0.51	0.79
Acetone	67-64-1	8	0	0.6	1.2	0	2.1	5
Aniline	62-53-3	4	0	0.017	0.051	0	0.11	0.23
Anthracene	120-12-7	11	0	0.0044	0.0081	0	0.022	0.031



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Anthracene, 2-methyl-	613-12-7	12	0	0.007	0.012	0	0.029	0.056
Anthracene, 9,10-dimethyl	781-43-1	2	0	0.0031	0.012	0	0.02	0.051
Anthracene, 9-phenyl	602-55-1	0	0	0	0	0	0	0
Benz[a]anthracene	56-55-3	14	0	8.1	12	0	32	49
Benzaldehyde	100-52-7	6	0	1.4	3.3	0	8.4	14
Benzene, 1,2,3-trimethyl-	526-73-8	5	0	0.26	0.74	0	1.5	3.4
Benzene, 1,2,4,5-tetramethyl-	95-93-2	12	0	11	18	0	45	74
Benzene, 1,2,4-trimethyl-	95-63-6	7	0	0.5	1.1	0	2.8	4.3
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	11	0	0.7	1.4	0	2.9	6.4
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	3	0	0.36	1.2	0	3.1	6.1
Benzene, butyl-	104-51-8	3	0	0.0032	0.01	0	0.035	0.035
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	19	0	0.35	0.45	0.15	1.2	1.3
Benzo[b]fluoranthene	205-99-2	3	0	0.0026	0.0084	0	0.029	0.029
7H-Benzo[c]fluorene	205-12-9	6	0	0.0026	0.0062	0	0.012	0.028
Benzo[k]fluoranthene	207-08-9	2	0	0.4	1.6	0	2.4	6.8
Benzothiazole	95-16-9	15	0	0.036	0.049	0	0.14	0.15
Benzothiazole, 2-phenyl-	883-93-2	5	0	0.36	0.95	0	1.9	4.3
2-Benzothiazolone	934-34-9	3	0	0.34	1.1	0	3.7	3.7
Benzyl butyl phthalate	85-68-7	12	0	24	62	0	160	290
Butanal	123-72-8	14	0	12	22	0	50	98
Cyclopentasiloxane, decamethyl-	541-02-6	13	0	4.3	6.9	0	18	26
Cyclotetrasiloxane, octamethyl-	556-67-2	12	0	5	8.2	0	20	29
p-Cymene	99-87-6	13	0	0.041	0.069	0	0.19	0.28
Decane	124-18-5	9	0	0.0075	0.017	0	0.048	0.061
Dibenz[a,h]anthracene	53-70-3	12	0	0.098	0.17	0	0.47	0.71
Dibenzothiophene	132-65-0	5	0	25	88	0	97	480
Dibutyl phthalate	84-74-2	2	0	0.9	3.6	0	5.9	15
Diethyl phthalate	84-66-2	14	0	1.3	2.1	0	6.1	7.6
Diisobutyl phthalate	84-69-5	6	0	1.2	2.6	0	6.7	6.7
Diisooctylphthalate	27554-26-3	3	0	0.0083	0.027	0	0.091	0.091
Di-n-octyl phthalate	117-84-0	3	0	0.72	2.5	0	6.6	9.8
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	3.3	5.5	1.2	13	26
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	3	0	0.91	3.2	0	7	16
Dodecane	112-40-3	14	0	0.19	0.34	0	0.87	1.6



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Fluoranthene	206-44-0	11	0	0.41	0.89	0	1.8	4.4
Fluorene	86-73-7	21	0	3.2	5	2.5	11	22
Furan, 2-methyl	534-22-5	5	0	1.3	3.2	0	11	11
Heptanal	111-71-7	9	0	3.8	7.8	0	16	35
Hexadecane	544-76-3	12	0	1.1	2.2	0	5.2	8.8
2,5-Hexanedione	110-13-4	3	0	2.9	10	0	22	53
Indan	496-11-7	13	0	2.4	4.1	0	11	17
Mesitylene	108-67-8	16	0	5.8	7.8	0	19	33
Methacrolein	78-85-3	1	0	0.3	1.7	0	0	10
Methyl Isobutyl Ketone	108-10-1	9	0	2.4	4.9	0	11	21
Naphthalene	91-20-3	12	0	0.23	0.41	0	1.1	1.7
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	1.8	2.9	0	8	9.6
Naphthalene, 1,6-dimethyl-	575-43-9	8	0	0.064	0.14	0	0.21	0.65
Naphthalene, 1-methyl-	90-12-0	12	0	0.17	0.3	0	0.83	1.2
Naphthalene, 2-(bromomethyl)-	939-26-4	12	0	2.9	4.9	0	14	16
Naphthalene, 2,3-dimethyl-	581-40-8	4	0	0.74	2.2	0	4.6	11
Naphthalene, 2-methyl-	91-57-6	16	0	3.6	5.3	0	14	20
1-Octadecene	112-88-9	14	0	0.34	0.51	0	1.5	1.6
Octanal	124-13-0	18	0	2.2	2.7	1.3	7.6	9.9
Octane	111-65-9	12	0	0.9	1.9	0	5.6	7.7
17-Pentatriacontene	6971-40-0	1	0	0.042	0.24	0	0	1.4
N-Phenylbenzamide	93-98-1	9	0	6	11	0	29	40
Phenanthrene	85-01-8	13	0	0.08	0.14	0	0.4	0.52
Phenanthrene, 1-methyl	832-69-9	13	0	0.091	0.16	0	0.43	0.64
Phenanthrene, 2-methyl-	2531-84-2	14	0	0.13	0.23	0	0.64	0.85
Phenanthrene, 3-methyl	832-71-3	1	0	0.00045	0.0026	0	0	0.015
Propionaldehyde	123-38-6	18	0	2.7	4.5	1.3	9.9	22
Pyrene	129-00-0	2	0	0.34	1.4	0	2	5.8
Pyridine, 2-(4-methylphenyl)-	4467-06-5	7	0	1.5	3.3	0	8.4	13
Resorcinol	108-46-3	34	2.3	22	24	11	66	90
m-Tolualdehyde	620-23-5	6	0	0.065	0.15	0	0.3	0.63
TXIB "Kodaflex"	6846-50-0	7	0	1.5	3.7	0	5	20
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	13	0	0.22	0.31	0	0.51	1.2
Undecane	1120-21-4	1	0	0.4	2.3	0	0	13
Valeraldehyde	110-62-3	34	15	53	44	33	130	230
Non-Field-Related Chemicals								
Benzene	71-43-2	1	0	1.4	8.3	0	0	48
Benzene, 1,4-dichloro	106-46-7	13	0	15	26	0	66	110



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	17	0	19	30	3.7	75	140
2-Butoxyethanol	111-76-2	3	0	6.4	24	0	44	130
Cyclotrisiloxane, hexamethyl-	541-05-9	27	0	72	100	35	240	480
Decanal	112-31-2	28	0	38	43	22	120	170
Ethylbenzene	100-41-4	24	0	51	73	20	180	280
Heptane	142-82-5	0	0	0	0	0	0	0
Hexanal	66-25-1	11	0	16	29	0	71	120
Hexane	110-54-3	12	0	4.4	6.8	0	17	17
1-Hexanol, 2-ethyl-	104-76-7	33	0	48	53	19	160	160
Nonanal	124-19-6	2	0	0.52	2.2	0	2.1	12
Phenol	108-95-2	34	17	120	130	59	340	590
Tetrachloroethylene	127-18-4	11	0	0.82	1.6	0	3.2	7
Tetradecane	629-59-4	7	0	3.1	6.7	0	20	20
m/p-Xylene	106-42-3	7	0	4.1	9.4	0	27	35
o-Xylene	95-47-6	11	0	8.4	33	0	28	190

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b Off-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-175. Off-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter)—Combined Gender Spectators 11<16 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	8	0	0.038	0.082	0	0.22	0.34
Acetone	67-64-1	8	0	0.26	0.5	0	0.91	2.1
Aniline	62-53-3	4	0	0.0072	0.022	0	0.049	0.1
Anthracene	120-12-7	11	0	0.0019	0.0035	0	0.0095	0.013
Anthracene, 2-methyl-	613-12-7	12	0	0.003	0.0053	0	0.013	0.024
Anthracene, 9,10-dimethyl	781-43-1	2	0	0.0013	0.0053	0	0.0088	0.022
Anthracene, 9-phenyl	602-55-1	0	0	0	0	0	0	0
Benz[a]anthracene	56-55-3	14	0	3.5	5.4	0	14	21
Benzaldehyde	100-52-7	6	0	0.6	1.4	0	3.6	5.9
Benzene, 1,2,3-trimethyl-	526-73-8	5	0	0.11	0.32	0	0.65	1.5
Benzene, 1,2,4,5-tetramethyl-	95-93-2	12	0	4.6	7.8	0	20	32
Benzene, 1,2,4-trimethyl-	95-63-6	7	0	0.21	0.49	0	1.2	1.9



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	11	0	0.3	0.6	0	1.2	2.8
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	3	0	0.16	0.54	0	1.3	2.6
Benzene, butyl-	104-51-8	3	0	0.0014	0.0044	0	0.015	0.015
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	19	0	0.15	0.2	0.063	0.52	0.58
Benzo[b]fluoranthene	205-99-2	3	0	0.0011	0.0036	0	0.012	0.012
7H-Benzo[c]fluorene	205-12-9	6	0	0.0011	0.0027	0	0.005	0.012
Benzo[k]fluoranthene	207-08-9	2	0	0.17	0.7	0	1	3
Benzothiazole	95-16-9	15	0	0.015	0.021	0	0.061	0.064
Benzothiazole, 2-phenyl-	883-93-2	5	0	0.16	0.41	0	0.84	1.8
2-Benzothiazolone	934-34-9	3	0	0.15	0.47	0	1.6	1.6
Benzyl butyl phthalate	85-68-7	12	0	10	27	0	71	120
Butanal	123-72-8	14	0	5.3	9.5	0	21	42
Cyclopentasiloxane, decamethyl-	541-02-6	13	0	1.9	3	0	7.8	11
Cyclotetrasiloxane, octamethyl-	556-67-2	12	0	2.2	3.5	0	8.7	13
p-Cymene	99-87-6	13	0	0.018	0.03	0	0.081	0.12
Decane	124-18-5	9	0	0.0032	0.0072	0	0.021	0.026
Dibenz[a,h]anthracene	53-70-3	12	0	0.042	0.074	0	0.2	0.31
Dibenzothiophene	132-65-0	5	0	11	38	0	42	210
Dibutyl phthalate	84-74-2	2	0	0.39	1.6	0	2.6	6.4
Diethyl phthalate	84-66-2	14	0	0.55	0.91	0	2.6	3.3
Diisobutyl phthalate	84-69-5	6	0	0.53	1.1	0	2.9	2.9
Diisooctylphthalate	27554-26-3	3	0	0.0036	0.011	0	0.039	0.039
Di-n-octyl phthalate	117-84-0	3	0	0.31	1.1	0	2.9	4.2
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	1.4	2.4	0.52	5.4	11
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	3	0	0.4	1.4	0	3	7
Dodecane	112-40-3	14	0	0.08	0.15	0	0.38	0.67
Fluoranthene	206-44-0	11	0	0.18	0.38	0	0.78	1.9
Fluorene	86-73-7	21	0	1.4	2.2	1.1	4.9	9.7
Furan, 2-methyl	534-22-5	5	0	0.54	1.4	0	4.6	4.6
Heptanal	111-71-7	9	0	1.6	3.4	0	7	15
Hexadecane	544-76-3	12	0	0.48	0.94	0	2.3	3.8
2,5-Hexanedione	110-13-4	3	0	1.3	4.5	0	9.5	23
Indan	496-11-7	13	0	1	1.8	0	4.6	7.2
Mesitylene	108-67-8	16	0	2.5	3.4	0	8	14
Methacrolein	78-85-3	1	0	0.13	0.75	0	0	4.4
Methyl Isobutyl Ketone	108-10-1	9	0	1	2.1	0	4.7	9.2
Naphthalene	91-20-3	12	0	0.099	0.18	0	0.49	0.73



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	0.77	1.3	0	3.5	4.1
Naphthalene, 1,6-dimethyl-	575-43-9	8	0	0.028	0.059	0	0.09	0.28
Naphthalene, 1-methyl-	90-12-0	12	0	0.075	0.13	0	0.36	0.52
Naphthalene, 2-(bromomethyl)-	939-26-4	12	0	1.3	2.1	0	6	7
Naphthalene, 2,3-dimethyl-	581-40-8	4	0	0.32	0.96	0	2	4.6
Naphthalene, 2-methyl-	91-57-6	16	0	1.6	2.3	0	6.1	8.7
1-Octadecene	112-88-9	14	0	0.15	0.22	0	0.66	0.68
Octanal	124-13-0	18	0	0.94	1.2	0.57	3.3	4.3
Octane	111-65-9	12	0	0.39	0.81	0	2.4	3.3
17-Pentatriacontene	6971-40-0	1	0	0.018	0.1	0	0	0.6
N-Phenylbenzamide	93-98-1	9	0	2.6	4.8	0	13	17
Phenanthrene	85-01-8	13	0	0.034	0.06	0	0.17	0.23
Phenanthrene, 1-methyl	832-69-9	13	0	0.039	0.069	0	0.19	0.28
Phenanthrene, 2-methyl-	2531-84-2	14	0	0.057	0.097	0	0.28	0.37
Phenanthrene, 3-methyl	832-71-3	1	0	0.00019	0.0011	0	0	0.0064
Propionaldehyde	123-38-6	18	0	1.2	2	0.54	4.3	9.4
Pyrene	129-00-0	2	0	0.15	0.59	0	0.87	2.5
Pyridine, 2-(4-methylphenyl)-	4467-06-5	7	0	0.65	1.4	0	3.6	5.7
Resorcinol	108-46-3	34	1	9.5	10	4.7	29	39
m-Tolualdehyde	620-23-5	6	0	0.028	0.064	0	0.13	0.27
TXIB "Kodaflex"	6846-50-0	7	0	0.63	1.6	0	2.2	8.4
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	13	0	0.097	0.13	0	0.22	0.53
Undecane	1120-21-4	1	0	0.17	0.99	0	0	5.8
Valeraldehyde	110-62-3	34	6.5	23	19	14	54	98
Non-Field-Related Chemicals								
Benzene	71-43-2	1	0	0.62	3.6	0	0	21
Benzene, 1,4-dichloro	106-46-7	13	0	6.5	11	0	29	47
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	17	0	8.3	13	1.6	32	60
2-Butoxyethanol	111-76-2	3	0	2.8	11	0	19	57
Cyclotrisiloxane, hexamethyl-	541-05-9	27	0	31	43	15	100	210
Decanal	112-31-2	28	0	16	18	9.6	51	73
Ethylbenzene	100-41-4	24	0	22	32	8.4	79	120
Heptane	142-82-5	0	0	0	0	0	0	0
Hexanal	66-25-1	11	0	6.8	12	0	31	50
Hexane	110-54-3	12	0	1.9	2.9	0	7.4	7.4
1-Hexanol, 2-ethyl-	104-76-7	33	0	21	23	8.3	69	71
Nonanal	124-19-6	2	0	0.23	0.97	0	0.89	5.1
Phenol	108-95-2	34	7.2	50	56	25	140	260



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Tetrachloroethylene	127-18-4	11	0	0.35	0.67	0	1.4	3
Tetradecane	629-59-4	7	0	1.3	2.9	0	8.4	8.4
m/p-Xylene	106-42-3	7	0	1.8	4.1	0	12	15
o-Xylene	95-47-6	11	0	3.6	14	0	12	83

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b Off-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-176. Off-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter)—Combined Gender Spectators 16<30 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	8	0	0.025	0.054	0	0.14	0.22
Acetone	67-64-1	8	0	0.17	0.33	0	0.6	1.4
Aniline	62-53-3	4	0	0.0048	0.015	0	0.032	0.066
Anthracene	120-12-7	11	0	0.0012	0.0023	0	0.0063	0.0087
Anthracene, 2-methyl-	613-12-7	12	0	0.002	0.0035	0	0.0083	0.016
Anthracene, 9,10-dimethyl	781-43-1	2	0	0.00088	0.0035	0	0.0058	0.015
Anthracene, 9-phenyl	602-55-1	0	0	0	0	0	0	0
Benz[a]anthracene	56-55-3	14	0	2.3	3.5	0	9.2	14
Benzaldehyde	100-52-7	6	0	0.4	0.95	0	2.4	3.9
Benzene, 1,2,3-trimethyl-	526-73-8	5	0	0.075	0.21	0	0.43	0.97
Benzene, 1,2,4,5-tetramethyl-	95-93-2	12	0	3	5.2	0	13	21
Benzene, 1,2,4-trimethyl-	95-63-6	7	0	0.14	0.33	0	0.8	1.2
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	11	0	0.2	0.39	0	0.82	1.8
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	3	0	0.1	0.36	0	0.87	1.7
Benzene, butyl-	104-51-8	3	0	0.0009	0.0029	0	0.0099	0.0099
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	19	0	0.099	0.13	0.042	0.34	0.38
Benzo[b]fluoranthene	205-99-2	3	0	0.00075	0.0024	0	0.0082	0.0082
7H-Benzo[c]fluorene	205-12-9	6	0	0.00074	0.0018	0	0.0033	0.0079
Benzo[k]fluoranthene	207-08-9	2	0	0.11	0.46	0	0.68	1.9
Benzothiazole	95-16-9	15	0	0.01	0.014	0	0.041	0.042
Benzothiazole, 2-phenyl-	883-93-2	5	0	0.1	0.27	0	0.55	1.2



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Benzothiazolone	934-34-9	3	0	0.096	0.31	0	1.1	1.1
Benzyl butyl phthalate	85-68-7	12	0	6.7	18	0	47	81
Butanal	123-72-8	14	0	3.5	6.3	0	14	28
Cyclopentasiloxane, decamethyl-	541-02-6	13	0	1.2	2	0	5.1	7.5
Cyclotetrasiloxane, octamethyl-	556-67-2	12	0	1.4	2.3	0	5.8	8.4
p-Cymene	99-87-6	13	0	0.012	0.02	0	0.053	0.081
Decane	124-18-5	9	0	0.0021	0.0048	0	0.014	0.017
Dibenz[a,h]anthracene	53-70-3	12	0	0.028	0.049	0	0.14	0.2
Dibenzothiophene	132-65-0	5	0	7.2	25	0	28	140
Dibutyl phthalate	84-74-2	2	0	0.26	1	0	1.7	4.2
Diethyl phthalate	84-66-2	14	0	0.36	0.6	0	1.7	2.2
Diisobutyl phthalate	84-69-5	6	0	0.35	0.75	0	1.9	1.9
Diisooctylphthalate	27554-26-3	3	0	0.0024	0.0076	0	0.026	0.026
Di-n-octyl phthalate	117-84-0	3	0	0.21	0.7	0	1.9	2.8
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	0.93	1.6	0.34	3.6	7.3
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	3	0	0.26	0.92	0	2	4.6
Dodecane	112-40-3	14	0	0.053	0.097	0	0.25	0.44
Fluoranthene	206-44-0	11	0	0.12	0.25	0	0.52	1.3
Fluorene	86-73-7	21	0	0.9	1.4	0.7	3.2	6.4
Furan, 2-methyl	534-22-5	5	0	0.36	0.92	0	3	3
Heptanal	111-71-7	9	0	1.1	2.2	0	4.6	9.9
Hexadecane	544-76-3	12	0	0.32	0.62	0	1.5	2.5
2,5-Hexanedione	110-13-4	3	0	0.84	3	0	6.2	15
Indan	496-11-7	13	0	0.67	1.2	0	3	4.7
Mesitylene	108-67-8	16	0	1.6	2.2	0	5.3	9.5
Methacrolein	78-85-3	1	0	0.085	0.49	0	0	2.9
Methyl Isobutyl Ketone	108-10-1	9	0	0.69	1.4	0	3.1	6.1
Naphthalene	91-20-3	12	0	0.066	0.12	0	0.33	0.48
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	0.51	0.84	0	2.3	2.7
Naphthalene, 1,6-dimethyl-	575-43-9	8	0	0.018	0.039	0	0.059	0.18
Naphthalene, 1-methyl-	90-12-0	12	0	0.049	0.085	0	0.24	0.34
Naphthalene, 2-(bromomethyl)-	939-26-4	12	0	0.83	1.4	0	4	4.6
Naphthalene, 2,3-dimethyl-	581-40-8	4	0	0.21	0.63	0	1.3	3
Naphthalene, 2-methyl-	91-57-6	16	0	1	1.5	0	4	5.7
1-Octadecene	112-88-9	14	0	0.096	0.15	0	0.43	0.45
Octanal	124-13-0	18	0	0.62	0.78	0.38	2.2	2.8



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Octane	111-65-9	12	0	0.26	0.53	0	1.6	2.2
17-Pentatriacontene	6971-40-0	1	0	0.012	0.068	0	0	0.39
N-Phenylbenzamide	93-98-1	9	0	1.7	3.2	0	8.3	11
Phenanthrene	85-01-8	13	0	0.023	0.04	0	0.11	0.15
Phenanthrene, 1-methyl	832-69-9	13	0	0.026	0.046	0	0.12	0.18
Phenanthrene, 2-methyl-	2531-84-2	14	0	0.038	0.064	0	0.18	0.24
Phenanthrene, 3-methyl	832-71-3	1	0	0.00013	0.00073	0	0	0.0042
Propionaldehyde	123-38-6	18	0	0.76	1.3	0.36	2.8	6.2
Pyrene	129-00-0	2	0	0.096	0.39	0	0.57	1.6
Pyridine, 2-(4-methylphenyl)-	4467-06-5	7	0	0.43	0.95	0	2.4	3.8
Resorcinol	108-46-3	34	0.66	6.3	6.8	3.1	19	26
m-Tolualdehyde	620-23-5	6	0	0.018	0.042	0	0.085	0.18
TXIB "Kodaflex"	6846-50-0	7	0	0.42	1.1	0	1.4	5.6
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	13	0	0.064	0.088	0	0.15	0.35
Undecane	1120-21-4	1	0	0.11	0.66	0	0	3.8
Valeraldehyde	110-62-3	34	4.3	15	13	9.4	36	64
Non-Field-Related Chemicals								
Benzene	71-43-2	1	0	0.41	2.4	0	0	14
Benzene, 1,4-dichloro	106-46-7	13	0	4.3	7.4	0	19	31
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	17	0	5.5	8.6	1.1	21	40
2-Butoxyethanol	111-76-2	3	0	1.8	7	0	12	37
Cyclotrisiloxane, hexamethyl-	541-05-9	27	0	21	29	10	68	140
Decanal	112-31-2	28	0	11	12	6.3	34	48
Ethylbenzene	100-41-4	24	0	15	21	5.6	52	81
Heptane	142-82-5	0	0	0	0	0	0	0
Hexanal	66-25-1	11	0	4.5	8.2	0	20	33
Hexane	110-54-3	12	0	1.3	1.9	0	4.9	4.9
1-Hexanol, 2-ethyl-	104-76-7	33	0	14	15	5.5	45	47
Nonanal	124-19-6	2	0	0.15	0.64	0	0.59	3.4
Phenol	108-95-2	34	4.7	33	37	17	96	170
Tetrachloroethylene	127-18-4	11	0	0.23	0.44	0	0.9	2
Tetradecane	629-59-4	7	0	0.87	1.9	0	5.6	5.6
m/p-Xylene	106-42-3	7	0	1.2	2.7	0	7.8	9.9
o-Xylene	95-47-6	11	0	2.4	9.5	0	7.9	55

^a 35 field-specific C_{inh-field} values are included in the table.



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

^b Off-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-177. Off-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter)—Combined Gender Spectators 30<40 Years

Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	8	0	0.022	0.047	0	0.13	0.2
Acetone	67-64-1	8	0	0.15	0.29	0	0.52	1.2
Aniline	62-53-3	4	0	0.0042	0.013	0	0.028	0.058
Anthracene	120-12-7	11	0	0.0011	0.002	0	0.0055	0.0076
Anthracene, 2-methyl-	613-12-7	12	0	0.0017	0.0031	0	0.0073	0.014
Anthracene, 9,10-dimethyl	781-43-1	2	0	0.00077	0.0031	0	0.0051	0.013
Anthracene, 9-phenyl	602-55-1	0	0	0	0	0	0	0
Benz[a]anthracene	56-55-3	14	0	2	3.1	0	8	12
Benzaldehyde	100-52-7	6	0	0.35	0.83	0	2.1	3.4
Benzene, 1,2,3-trimethyl-	526-73-8	5	0	0.066	0.18	0	0.38	0.85
Benzene, 1,2,4,5-tetramethyl-	95-93-2	12	0	2.6	4.5	0	11	18
Benzene, 1,2,4-trimethyl-	95-63-6	7	0	0.12	0.28	0	0.7	1.1
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	11	0	0.17	0.34	0	0.72	1.6
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	3	0	0.089	0.31	0	0.76	1.5
Benzene, butyl-	104-51-8	3	0	0.00079	0.0025	0	0.0087	0.0087
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	19	0	0.087	0.11	0.036	0.3	0.34
Benzo[b]fluoranthene	205-99-2	3	0	0.00065	0.0021	0	0.0072	0.0072
7H-Benzo[c]fluorene	205-12-9	6	0	0.00065	0.0015	0	0.0029	0.0069
Benzo[k]fluoranthene	207-08-9	2	0	0.1	0.41	0	0.6	1.7
Benzothiazole	95-16-9	15	0	0.0089	0.012	0	0.035	0.037
Benzothiazole, 2-phenyl-	883-93-2	5	0	0.091	0.24	0	0.48	1.1
2-Benzothiazolone	934-34-9	3	0	0.084	0.27	0	0.93	0.93
Benzyl butyl phthalate	85-68-7	12	0	5.9	15	0	41	71
Butanal	123-72-8	14	0	3.1	5.5	0	12	24
Cyclopentasiloxane, decamethyl-	541-02-6	13	0	1.1	1.7	0	4.5	6.6
Cyclotetrasiloxane, octamethyl-	556-67-2	12	0	1.2	2	0	5	7.3
p-Cymene	99-87-6	13	0	0.01	0.017	0	0.046	0.071
Decane	124-18-5	9	0	0.0019	0.0042	0	0.012	0.015
Dibenz[a,h]anthracene	53-70-3	12	0	0.024	0.043	0	0.12	0.18



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dibenzothiophene	132-65-0	5	0	6.3	22	0	24	120
Dibutyl phthalate	84-74-2	2	0	0.22	0.9	0	1.5	3.7
Diethyl phthalate	84-66-2	14	0	0.32	0.52	0	1.5	1.9
Diisobutyl phthalate	84-69-5	6	0	0.31	0.66	0	1.7	1.7
Diisooctylphthalate	27554-26-3	3	0	0.0021	0.0066	0	0.023	0.023
Di-n-octyl phthalate	117-84-0	3	0	0.18	0.61	0	1.7	2.4
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	0.81	1.4	0.3	3.1	6.4
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	3	0	0.23	0.8	0	1.8	4
Dodecane	112-40-3	14	0	0.046	0.085	0	0.22	0.39
Fluoranthene	206-44-0	11	0	0.1	0.22	0	0.45	1.1
Fluorene	86-73-7	21	0	0.79	1.2	0.61	2.8	5.6
Furan, 2-methyl	534-22-5	5	0	0.31	0.81	0	2.7	2.7
Heptanal	111-71-7	9	0	0.94	1.9	0	4.1	8.7
Hexadecane	544-76-3	12	0	0.28	0.54	0	1.3	2.2
2,5-Hexanedione	110-13-4	3	0	0.73	2.6	0	5.5	13
Indan	496-11-7	13	0	0.59	1	0	2.6	4.1
Mesitylene	108-67-8	16	0	1.4	2	0	4.6	8.3
Methacrolein	78-85-3	1	0	0.074	0.43	0	0	2.5
Methyl Isobutyl Ketone	108-10-1	9	0	0.6	1.2	0	2.7	5.3
Naphthalene	91-20-3	12	0	0.057	0.1	0	0.28	0.42
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	0.45	0.73	0	2	2.4
Naphthalene, 1,6-dimethyl-	575-43-9	8	0	0.016	0.034	0	0.052	0.16
Naphthalene, 1-methyl-	90-12-0	12	0	0.043	0.074	0	0.21	0.3
Naphthalene, 2-(bromomethyl)-	939-26-4	12	0	0.73	1.2	0	3.5	4
Naphthalene, 2,3-dimethyl-	581-40-8	4	0	0.18	0.55	0	1.1	2.6
Naphthalene, 2-methyl-	91-57-6	16	0	0.9	1.3	0	3.5	5
1-Octadecene	112-88-9	14	0	0.084	0.13	0	0.38	0.39
Octanal	124-13-0	18	0	0.54	0.68	0.33	1.9	2.5
Octane	111-65-9	12	0	0.22	0.46	0	1.4	1.9
17-Pentatriacontene	6971-40-0	1	0	0.01	0.06	0	0	0.34
N-Phenylbenzamide	93-98-1	9	0	1.5	2.8	0	7.3	10
Phenanthrene	85-01-8	13	0	0.02	0.035	0	0.1	0.13
Phenanthrene, 1-methyl	832-69-9	13	0	0.023	0.04	0	0.11	0.16
Phenanthrene, 2-methyl-	2531-84-2	14	0	0.033	0.056	0	0.16	0.21
Phenanthrene, 3-methyl	832-71-3	1	0	0.00011	0.00064	0	0	0.0037
Propionaldehyde	123-38-6	18	0	0.66	1.1	0.31	2.5	5.4
Pyrene	129-00-0	2	0	0.084	0.34	0	0.5	1.4



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Pyridine, 2-(4-methylphenyl)-	4467-06-5	7	0	0.37	0.83	0	2.1	3.3
Resorcinol	108-46-3	34	0.58	5.5	5.9	2.7	16	22
m-Tolualdehyde	620-23-5	6	0	0.016	0.037	0	0.075	0.16
TXIB "Kodaflex"	6846-50-0	7	0	0.36	0.93	0	1.3	4.9
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	13	0	0.056	0.077	0	0.13	0.31
Undecane	1120-21-4	1	0	0.098	0.57	0	0	3.3
Valeraldehyde	110-62-3	34	3.7	13	11	8.2	31	56
Non-Field-Related Chemicals								
Benzene	71-43-2	1	0	0.35	2.1	0	0	12
Benzene, 1,4-dichloro	106-46-7	13	0	3.8	6.4	0	16	27
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	17	0	4.8	7.5	0.92	19	35
2-Butoxyethanol	111-76-2	3	0	1.6	6.1	0	11	33
Cyclotrisiloxane, hexamethyl-	541-05-9	27	0	18	25	8.8	59	120
Decanal	112-31-2	28	0	9.4	11	5.5	30	42
Ethylbenzene	100-41-4	24	0	13	18	4.9	46	71
Heptane	142-82-5	0	0	0	0	0	0	0
Hexanal	66-25-1	11	0	3.9	7.2	0	18	29
Hexane	110-54-3	12	0	1.1	1.7	0	4.3	4.3
1-Hexanol, 2-ethyl-	104-76-7	33	0	12	13	4.8	40	41
Nonanal	124-19-6	2	0	0.13	0.56	0	0.51	2.9
Phenol	108-95-2	34	4.1	29	32	15	84	150
Tetrachloroethylene	127-18-4	11	0	0.2	0.39	0	0.79	1.7
Tetradecane	629-59-4	7	0	0.76	1.7	0	4.9	4.9
m/p-Xylene	106-42-3	7	0	1	2.3	0	6.8	8.7
o-Xylene	95-47-6	11	0	2.1	8.3	0	6.9	48

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b Off-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-178. Off-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter)—Combined Gender Spectators 40<50 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	8	0	0.022	0.048	0	0.13	0.2
Acetone	67-64-1	8	0	0.15	0.29	0	0.53	1.2



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	0.0042	0.013	0	0.028	0.058
Anthracene	120-12-7	11	0	0.0011	0.002	0	0.0055	0.0077
Anthracene, 2-methyl-	613-12-7	12	0	0.0017	0.0031	0	0.0073	0.014
Anthracene, 9,10-dimethyl	781-43-1	2	0	0.00078	0.0031	0	0.0051	0.013
Anthracene, 9-phenyl	602-55-1	0	0	0	0	0	0	0
Benz[a]anthracene	56-55-3	14	0	2	3.1	0	8.1	12
Benzaldehyde	100-52-7	6	0	0.35	0.84	0	2.1	3.4
Benzene, 1,2,3-trimethyl-	526-73-8	5	0	0.067	0.19	0	0.38	0.86
Benzene, 1,2,4,5-tetramethyl-	95-93-2	12	0	2.7	4.6	0	11	19
Benzene, 1,2,4-trimethyl-	95-63-6	7	0	0.12	0.29	0	0.71	1.1
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	11	0	0.18	0.35	0	0.73	1.6
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	3	0	0.09	0.31	0	0.77	1.5
Benzene, butyl-	104-51-8	3	0	0.0008	0.0026	0	0.0088	0.0088
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	19	0	0.088	0.11	0.037	0.3	0.34
Benzo[b]fluoranthene	205-99-2	3	0	0.00066	0.0021	0	0.0073	0.0073
7H-Benzo[c]fluorene	205-12-9	6	0	0.00066	0.0016	0	0.0029	0.0069
Benzo[k]fluoranthene	207-08-9	2	0	0.1	0.41	0	0.6	1.7
Benzothiazole	95-16-9	15	0	0.009	0.012	0	0.036	0.037
Benzothiazole, 2-phenyl-	883-93-2	5	0	0.092	0.24	0	0.49	1.1
2-Benzothiazolone	934-34-9	3	0	0.085	0.27	0	0.94	0.94
Benzyl butyl phthalate	85-68-7	12	0	6	16	0	41	72
Butanal	123-72-8	14	0	3.1	5.5	0	12	25
Cyclopentasiloxane, decamethyl-	541-02-6	13	0	1.1	1.7	0	4.5	6.6
Cyclotetrasiloxane, octamethyl-	556-67-2	12	0	1.3	2.1	0	5.1	7.4
p-Cymene	99-87-6	13	0	0.01	0.017	0	0.047	0.071
Decane	124-18-5	9	0	0.0019	0.0042	0	0.012	0.015
Dibenz[a,h]anthracene	53-70-3	12	0	0.025	0.043	0	0.12	0.18
Dibenzothiophene	132-65-0	5	0	6.3	22	0	24	120
Dibutyl phthalate	84-74-2	2	0	0.23	0.9	0	1.5	3.7
Diethyl phthalate	84-66-2	14	0	0.32	0.53	0	1.5	1.9
Diisobutyl phthalate	84-69-5	6	0	0.31	0.66	0	1.7	1.7
Diisooctylphthalate	27554-26-3	3	0	0.0021	0.0067	0	0.023	0.023
Di-n-octyl phthalate	117-84-0	3	0	0.18	0.62	0	1.7	2.5
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	0.82	1.4	0.3	3.2	6.4



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	3	0	0.23	0.81	0	1.8	4.1
Dodecane	112-40-3	14	0	0.047	0.086	0	0.22	0.39
Fluoranthene	206-44-0	11	0	0.1	0.22	0	0.46	1.1
Fluorene	86-73-7	21	0	0.79	1.3	0.62	2.9	5.7
Furan, 2-methyl	534-22-5	5	0	0.32	0.82	0	2.7	2.7
Heptanal	111-71-7	9	0	0.95	2	0	4.1	8.8
Hexadecane	544-76-3	12	0	0.28	0.55	0	1.3	2.2
2,5-Hexanedione	110-13-4	3	0	0.74	2.6	0	5.5	13
Indan	496-11-7	13	0	0.59	1	0	2.7	4.2
Mesitylene	108-67-8	16	0	1.5	2	0	4.7	8.4
Methacrolein	78-85-3	1	0	0.075	0.44	0	0	2.5
Methyl Isobutyl Ketone	108-10-1	9	0	0.61	1.2	0	2.7	5.3
Naphthalene	91-20-3	12	0	0.058	0.1	0	0.29	0.42
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	0.45	0.74	0	2	2.4
Naphthalene, 1,6-dimethyl-	575-43-9	8	0	0.016	0.034	0	0.052	0.16
Naphthalene, 1-methyl-	90-12-0	12	0	0.043	0.075	0	0.21	0.3
Naphthalene, 2-(bromomethyl)-	939-26-4	12	0	0.74	1.2	0	3.5	4.1
Naphthalene, 2,3-dimethyl-	581-40-8	4	0	0.19	0.56	0	1.2	2.7
Naphthalene, 2-methyl-	91-57-6	16	0	0.91	1.3	0	3.6	5.1
1-Octadecene	112-88-9	14	0	0.085	0.13	0	0.38	0.4
Octanal	124-13-0	18	0	0.55	0.69	0.33	1.9	2.5
Octane	111-65-9	12	0	0.23	0.47	0	1.4	1.9
17-Pentatriacontene	6971-40-0	1	0	0.011	0.06	0	0	0.35
N-Phenylbenzamide	93-98-1	9	0	1.5	2.8	0	7.4	10
Phenanthrene	85-01-8	13	0	0.02	0.035	0	0.1	0.13
Phenanthrene, 1-methyl	832-69-9	13	0	0.023	0.04	0	0.11	0.16
Phenanthrene, 2-methyl-	2531-84-2	14	0	0.033	0.057	0	0.16	0.21
Phenanthrene, 3-methyl	832-71-3	1	0	0.00011	0.00065	0	0	0.0037
Propionaldehyde	123-38-6	18	0	0.67	1.1	0.32	2.5	5.5
Pyrene	129-00-0	2	0	0.085	0.35	0	0.51	1.4
Pyridine, 2-(4-methylphenyl)-	4467-06-5	7	0	0.38	0.84	0	2.1	3.3
Resorcinol	108-46-3	34	0.59	5.5	6	2.8	17	23
m-Tolualdehyde	620-23-5	6	0	0.016	0.038	0	0.075	0.16
TXIB "Kodaflex"	6846-50-0	7	0	0.37	0.94	0	1.3	4.9
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	13	0	0.056	0.077	0	0.13	0.31
Undecane	1120-21-4	1	0	0.099	0.58	0	0	3.4



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Valeraldehyde	110-62-3	34	3.8	13	11	8.3	32	57
Non-Field-Related Chemicals								
Benzene	71-43-2	1	0	0.36	2.1	0	0	12
Benzene, 1,4-dichloro	106-46-7	13	0	3.8	6.5	0	17	27
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	17	0	4.8	7.6	0.93	19	35
2-Butoxyethanol	111-76-2	3	0	1.6	6.1	0	11	33
Cyclotrisiloxane, hexamethyl-	541-05-9	27	0	18	25	8.9	60	120
Decanal	112-31-2	28	0	9.5	11	5.6	30	42
Ethylbenzene	100-41-4	24	0	13	18	4.9	46	72
Heptane	142-82-5	0	0	0	0	0	0	0
Hexanal	66-25-1	11	0	4	7.2	0	18	29
Hexane	110-54-3	12	0	1.1	1.7	0	4.3	4.3
1-Hexanol, 2-ethyl-	104-76-7	33	0	12	13	4.8	40	41
Nonanal	124-19-6	2	0	0.13	0.56	0	0.52	3
Phenol	108-95-2	34	4.2	29	33	15	84	150
Tetrachloroethylene	127-18-4	11	0	0.21	0.39	0	0.8	1.8
Tetradecane	629-59-4	7	0	0.77	1.7	0	4.9	4.9
m/p-Xylene	106-42-3	7	0	1	2.4	0	6.8	8.7
o-Xylene	95-47-6	11	0	2.1	8.4	0	7	48

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b Off-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-179. Off-Field Field-Specific^a Chronic Inhalation Exposure Concentration^b for All General Chemicals (Chronic $C_{inh-field}$, nanograms per cubic meter)—Combined Gender Spectators 50<70 Years

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Acenaphthylene	208-96-8	8	0	0.022	0.048	0	0.13	0.2
Acetone	67-64-1	8	0	0.15	0.29	0	0.53	1.3
Aniline	62-53-3	4	0	0.0042	0.013	0	0.029	0.058
Anthracene	120-12-7	11	0	0.0011	0.002	0	0.0055	0.0077
Anthracene, 2-methyl-	613-12-7	12	0	0.0018	0.0031	0	0.0074	0.014
Anthracene, 9,10-dimethyl	781-43-1	2	0	0.00078	0.0031	0	0.0052	0.013
Anthracene, 9-phenyl	602-55-1	0	0	0	0	0	0	0
Benz[a]anthracene	56-55-3	14	0	2	3.1	0	8.1	12



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh} -field					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzaldehyde	100-52-7	6	0	0.35	0.84	0	2.1	3.5
Benzene, 1,2,3-trimethyl-	526-73-8	5	0	0.067	0.19	0	0.38	0.86
Benzene, 1,2,4,5-tetramethyl-	95-93-2	12	0	2.7	4.6	0	11	19
Benzene, 1,2,4-trimethyl-	95-63-6	7	0	0.12	0.29	0	0.71	1.1
Benzene, 1-ethyl-2,4-dimethyl-	874-41-9	11	0	0.18	0.35	0	0.73	1.6
Benzene, 2-ethyl-1,4-dimethyl-	1758-88-9	3	0	0.091	0.32	0	0.77	1.5
Benzene, butyl-	104-51-8	3	0	0.0008	0.0026	0	0.0088	0.0088
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	19	0	0.088	0.11	0.037	0.3	0.34
Benzo[b]fluoranthene	205-99-2	3	0	0.00066	0.0021	0	0.0073	0.0073
7H-Benzo[c]fluorene	205-12-9	6	0	0.00066	0.0016	0	0.0029	0.007
Benzo[k]fluoranthene	207-08-9	2	0	0.1	0.41	0	0.6	1.7
Benzothiazole	95-16-9	15	0	0.009	0.012	0	0.036	0.037
Benzothiazole, 2-phenyl-	883-93-2	5	0	0.092	0.24	0	0.49	1.1
2-Benzothiazolone	934-34-9	3	0	0.085	0.27	0	0.94	0.94
Benzyl butyl phthalate	85-68-7	12	0	6	16	0	42	72
Butanal	123-72-8	14	0	3.1	5.5	0	13	25
Cyclopentasiloxane, decamethyl-	541-02-6	13	0	1.1	1.7	0	4.6	6.6
Cyclotetrasiloxane, octamethyl-	556-67-2	12	0	1.3	2.1	0	5.1	7.4
p-Cymene	99-87-6	13	0	0.01	0.017	0	0.047	0.072
Decane	124-18-5	9	0	0.0019	0.0042	0	0.012	0.015
Dibenz[a,h]anthracene	53-70-3	12	0	0.025	0.043	0	0.12	0.18
Dibenzothiophene	132-65-0	5	0	6.3	22	0	24	120
Dibutyl phthalate	84-74-2	2	0	0.23	0.91	0	1.5	3.7
Diethyl phthalate	84-66-2	14	0	0.32	0.53	0	1.5	1.9
Diisobutyl phthalate	84-69-5	6	0	0.31	0.67	0	1.7	1.7
Diisooctylphthalate	27554-26-3	3	0	0.0021	0.0067	0	0.023	0.023
Di-n-octyl phthalate	117-84-0	3	0	0.18	0.62	0	1.7	2.5
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	26	0	0.83	1.4	0.3	3.2	6.5
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	3	0	0.23	0.81	0	1.8	4.1
Dodecane	112-40-3	14	0	0.047	0.086	0	0.22	0.39
Fluoranthene	206-44-0	11	0	0.1	0.22	0	0.46	1.1
Fluorene	86-73-7	21	0	0.8	1.3	0.62	2.9	5.7
Furan, 2-methyl	534-22-5	5	0	0.32	0.82	0	2.7	2.7
Heptanal	111-71-7	9	0	0.96	2	0	4.1	8.8
Hexadecane	544-76-3	12	0	0.28	0.55	0	1.3	2.2



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	Chronic C _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2,5-Hexanedione	110-13-4	3	0	0.74	2.6	0	5.5	13
Indan	496-11-7	13	0	0.6	1	0	2.7	4.2
Mesitylene	108-67-8	16	0	1.5	2	0	4.7	8.4
Methacrolein	78-85-3	1	0	0.075	0.44	0	0	2.6
Methyl Isobutyl Ketone	108-10-1	9	0	0.61	1.2	0	2.7	5.4
Naphthalene	91-20-3	12	0	0.058	0.1	0	0.29	0.43
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	0.45	0.74	0	2	2.4
Naphthalene, 1,6-dimethyl-	575-43-9	8	0	0.016	0.034	0	0.053	0.16
Naphthalene, 1-methyl-	90-12-0	12	0	0.044	0.075	0	0.21	0.3
Naphthalene, 2-(bromomethyl)-	939-26-4	12	0	0.74	1.2	0	3.5	4.1
Naphthalene, 2,3-dimethyl-	581-40-8	4	0	0.19	0.56	0	1.2	2.7
Naphthalene, 2-methyl-	91-57-6	16	0	0.91	1.3	0	3.6	5.1
1-Octadecene	112-88-9	14	0	0.085	0.13	0	0.38	0.4
Octanal	124-13-0	18	0	0.55	0.69	0.33	1.9	2.5
Octane	111-65-9	12	0	0.23	0.47	0	1.4	1.9
17-Pentatriacontene	6971-40-0	1	0	0.011	0.061	0	0	0.35
N-Phenylbenzamide	93-98-1	9	0	1.5	2.8	0	7.4	10
Phenanthrene	85-01-8	13	0	0.02	0.035	0	0.1	0.13
Phenanthrene, 1-methyl	832-69-9	13	0	0.023	0.041	0	0.11	0.16
Phenanthrene, 2-methyl-	2531-84-2	14	0	0.033	0.057	0	0.16	0.21
Phenanthrene, 3-methyl	832-71-3	1	0	0.00011	0.00065	0	0	0.0037
Propionaldehyde	123-38-6	18	0	0.67	1.1	0.32	2.5	5.5
Pyrene	129-00-0	2	0	0.085	0.35	0	0.51	1.5
Pyridine, 2-(4-methylphenyl)-	4467-06-5	7	0	0.38	0.84	0	2.1	3.4
Resorcinol	108-46-3	34	0.59	5.5	6	2.8	17	23
m-Tolualdehyde	620-23-5	6	0	0.016	0.038	0	0.076	0.16
TXIB "Kodaflex"	6846-50-0	7	0	0.37	0.94	0	1.3	4.9
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	13	0	0.057	0.078	0	0.13	0.31
Undecane	1120-21-4	1	0	0.1	0.58	0	0	3.4
Valeraldehyde	110-62-3	34	3.8	13	11	8.3	32	57
Non-Field-Related Chemicals								
Benzene	71-43-2	1	0	0.36	2.1	0	0	12
Benzene, 1,4-dichloro	106-46-7	13	0	3.8	6.5	0	17	27
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	17	0	4.8	7.6	0.93	19	35
2-Butoxyethanol	111-76-2	3	0	1.6	6.2	0	11	33



Chemical	CASRN	Detection	Chronic $C_{inh-field}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Cyclotrisiloxane, hexamethyl-	541-05-9	27	0	18	25	8.9	60	120
Decanal	112-31-2	28	0	9.6	11	5.6	30	43
Ethylbenzene	100-41-4	24	0	13	18	4.9	46	72
Heptane	142-82-5	0	0	0	0	0	0	0
Hexanal	66-25-1	11	0	4	7.3	0	18	29
Hexane	110-54-3	12	0	1.1	1.7	0	4.3	4.3
1-Hexanol, 2-ethyl-	104-76-7	33	0	12	13	4.8	40	42
Nonanal	124-19-6	2	0	0.13	0.56	0	0.52	3
Phenol	108-95-2	34	4.2	29	33	15	85	150
Tetrachloroethylene	127-18-4	11	0	0.21	0.39	0	0.8	1.8
Tetradecane	629-59-4	7	0	0.77	1.7	0	4.9	4.9
m/p-Xylene	106-42-3	7	0	1	2.4	0	6.9	8.8
o-Xylene	95-47-6	11	0	2.1	8.4	0	7	49

^a 35 field-specific $C_{inh-field}$ values are included in the table.

^b Off-Field field-specific Chronic $C_{inh-field}$ equals the modified field-specific average concentration of a field ($C_{air-field} \times AF_{inh}$) of a general chemical (see Section F.4.3 of this Appendix F).

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

F.5. Average Daily Dose (ADD) and Average One-Day Dose (AD)

This section presents the average daily doses (ADD) or average one-day dose (AD) of all detected chemicals (with available toxicity criteria) on synthetic turf fields for the inhalation, dermal, and ingestion pathways. ADD and AD values are used to calculate the non-cancer hazard or cancer risk for exposure to chemicals through the inhalation, dermal, or ingestion pathway. Details of each exposure pathway and how to calculate the ADD or AD are presented in Main Report Sections 5.4.4 and 5.4.5 and Appendix Sections 3, 4, and 5. An example calculation for each pathway is presented in Appendix G.1.

F.5.1. Average Inhalation Daily Dose (ADD_{inh}) for Lifetime Cancer Risk Assessment of Chemicals

Table F-180. Average **On-Field** Inhalation Daily Dose^a (ADD_{inh} , milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender Athletes

Chemical	CASRN	ADD _{inh}						
		2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Field-Related Chemicals								
Aniline	62-53-3	2.9E-07	3.7E-07	3.6E-07	6.9E-07	4.5E-07	4.4E-07	4.3E-07
Benz[a]anthracene	56-55-3	2.2E-10	2.8E-10	2.8E-10	5.2E-10	3.4E-10	3.4E-10	3.3E-10



Chemical	CASRN	ADD _{inh}						
		2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Benzo[a]pyrene	50-32-8	5.1E-08	6.6E-08	6.4E-08	1.2E-07	7.9E-08	7.8E-08	7.6E-08
Benzo[b]fluoranthene	205-99-2	1.1E-09	1.4E-09	1.4E-09	2.6E-09	1.7E-09	1.7E-09	1.7E-09
Benzo[k]fluoranthene	207-08-9	1.4E-09	1.8E-09	1.7E-09	3.2E-09	2.1E-09	2.1E-09	2.1E-09
Chrysene	218-01-9	9.1E-09	1.2E-08	1.2E-08	2.2E-08	1.4E-08	1.4E-08	1.4E-08
Cyclopenta[cd]pyrene	27208-37-3	3.2E-09	4.2E-09	4.1E-09	7.7E-09	5.0E-09	5.0E-09	4.9E-09
Dibenz[a,h]anthracene	53-70-3	6.4E-09	8.2E-09	8.1E-09	1.5E-08	9.9E-09	9.8E-09	9.6E-09
Indeno[1,2,3-cd]pyrene	193-39-5	4.8E-09	6.2E-09	6.0E-09	1.1E-08	7.4E-09	7.3E-09	7.2E-09
Methyl Isobutyl Ketone	108-10-1	6.9E-07	9.0E-07	8.8E-07	1.7E-06	1.1E-06	1.1E-06	1.0E-06
Naphthalene	91-20-3	1.2E-06	1.5E-06	1.5E-06	2.8E-06	1.8E-06	1.8E-06	1.8E-06
Styrene	100-42-5	2.6E-06	3.4E-06	3.3E-06	6.2E-06	4.1E-06	4.0E-06	3.9E-06
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	1.1E-04	1.4E-04	1.4E-04	2.7E-04	1.7E-04	1.7E-04	1.7E-04
Benzene	71-43-2	2.7E-05	3.5E-05	3.4E-05	6.4E-05	4.2E-05	4.1E-05	4.1E-05
Benzene, 1,4-dichloro	106-46-7	8.5E-07	1.1E-06	1.1E-06	2.0E-06	1.3E-06	1.3E-06	1.3E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	2.5E-05	3.3E-05	3.2E-05	6.1E-05	4.0E-05	3.9E-05	3.9E-05
Ethylbenzene	100-41-4	7.6E-06	9.9E-06	9.6E-06	1.8E-05	1.2E-05	1.2E-05	1.1E-05
Formaldehyde	50-00-0	1.7E-04	2.2E-04	2.2E-04	4.1E-04	2.6E-04	2.6E-04	2.6E-04
Tetrachloroethylene	127-18-4	2.1E-06	2.7E-06	2.7E-06	5.1E-06	3.3E-06	3.3E-06	3.2E-06

^a On-Field ADD_{inh} was calculated from the 35 individual field average concentrations (C_{air}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-181. Average **On-Field** Inhalation Daily Dose^a (ADD_{inh}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender Coaches

Chemical	CASRN	ADD _{inh}			
		16<30 years	30<40 years	40<50 years	50<70 years
Field-Related Chemicals					
Aniline	62-53-3	2.8E-07	2.5E-07	2.5E-07	2.5E-07
Benz[a]anthracene	56-55-3	2.2E-10	1.9E-10	1.9E-10	1.9E-10
Benzo[a]pyrene	50-32-8	5.0E-08	4.4E-08	4.4E-08	4.4E-08
Benzo[b]fluoranthene	205-99-2	1.1E-09	9.4E-10	9.5E-10	9.5E-10
Benzo[k]fluoranthene	207-08-9	1.3E-09	1.2E-09	1.2E-09	1.2E-09
Chrysene	218-01-9	9.0E-09	7.8E-09	7.9E-09	7.9E-09
Cyclopenta[cd]pyrene	27208-37-3	3.2E-09	2.8E-09	2.8E-09	2.8E-09
Dibenz[a,h]anthracene	53-70-3	6.3E-09	5.5E-09	5.5E-09	5.6E-09
Indeno[1,2,3-cd]pyrene	193-39-5	4.7E-09	4.1E-09	4.2E-09	4.2E-09
Methyl Isobutyl Ketone	108-10-1	6.8E-07	6.0E-07	6.0E-07	6.1E-07
Naphthalene	91-20-3	1.2E-06	1.0E-06	1.0E-06	1.0E-06



Chemical	CASRN	ADD _{inh}			
		16<30 years	30<40 years	40<50 years	50<70 years
Styrene	100-42-5	2.6E-06	2.2E-06	2.3E-06	2.3E-06
Non-Field-Related Chemicals					
Acetaldehyde	75-07-0	1.1E-04	9.6E-05	9.7E-05	9.8E-05
Benzene	71-43-2	2.6E-05	2.3E-05	2.3E-05	2.3E-05
Benzene, 1,4-dichloro	106-46-7	8.4E-07	7.3E-07	7.4E-07	7.4E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	2.5E-05	2.2E-05	2.2E-05	2.2E-05
Ethylbenzene	100-41-4	7.5E-06	6.6E-06	6.6E-06	6.7E-06
Formaldehyde	50-00-0	1.7E-04	1.5E-04	1.5E-04	1.5E-04
Tetrachloroethylene	127-18-4	2.1E-06	1.8E-06	1.8E-06	1.9E-06

^a On-Field ADD_{inh} was calculated from the 35 individual field average concentrations (C_{air}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-182. Average **On-Field** Inhalation Daily Dose^a (ADD_{inh}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender
Referees

Chemical	CASRN	ADD _{inh}			
		16<30 years	30<40 years	40<50 years	50<70 years
Field-Related Chemicals					
Aniline	62-53-3	1.1E-07	9.2E-08	9.3E-08	9.3E-08
Benz[a]anthracene	56-55-3	8.0E-11	7.0E-11	7.1E-11	7.1E-11
Benzo[a]pyrene	50-32-8	1.9E-08	1.6E-08	1.6E-08	1.6E-08
Benzo[b]fluoranthene	205-99-2	4.0E-10	3.5E-10	3.5E-10	3.5E-10
Benzo[k]fluoranthene	207-08-9	5.0E-10	4.3E-10	4.4E-10	4.4E-10
Chrysene	218-01-9	3.3E-09	2.9E-09	2.9E-09	2.9E-09
Cyclopenta[cd]pyrene	27208-37-3	1.2E-09	1.0E-09	1.0E-09	1.0E-09
Dibenz[a,h]anthracene	53-70-3	2.3E-09	2.0E-09	2.1E-09	2.1E-09
Indeno[1,2,3-cd]pyrene	193-39-5	1.7E-09	1.5E-09	1.5E-09	1.5E-09
Methyl Isobutyl Ketone	108-10-1	2.5E-07	2.2E-07	2.2E-07	2.3E-07
Naphthalene	91-20-3	4.3E-07	3.8E-07	3.8E-07	3.8E-07
Styrene	100-42-5	9.6E-07	8.4E-07	8.4E-07	8.5E-07
Non-Field-Related Chemicals					
Acetaldehyde	75-07-0	4.1E-05	3.6E-05	3.6E-05	3.6E-05
Benzene	71-43-2	9.8E-06	8.6E-06	8.7E-06	8.7E-06
Benzene, 1,4-dichloro	106-46-7	3.1E-07	2.7E-07	2.8E-07	2.8E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	9.3E-06	8.2E-06	8.2E-06	8.3E-06
Ethylbenzene	100-41-4	2.8E-06	2.4E-06	2.5E-06	2.5E-06
Formaldehyde	50-00-0	6.2E-05	5.4E-05	5.5E-05	5.5E-05
Tetrachloroethylene	127-18-4	7.8E-07	6.8E-07	6.9E-07	6.9E-07

^a On-Field ADD_{inh} was calculated from the 35 individual field average concentrations (C_{air}) of a chemical



detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-183. Average **On-Field** Inhalation Daily Dose^a (ADD_{inh}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender Spectators

Chemical	CASRN	ADD _{inh}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Field-Related Chemicals										
Aniline	62-53-3	3.8E-08	3.1E-07	1.6E-07	1.5E-07	6.6E-08	4.4E-08	3.8E-08	3.9E-08	3.9E-08
Benz[a]anthracene	56-55-3	2.9E-11	2.3E-10	1.2E-10	1.2E-10	5.1E-11	3.3E-11	2.9E-11	2.9E-11	3.0E-11
Benzo[a]pyrene	50-32-8	6.7E-09	5.4E-08	2.8E-08	2.7E-08	1.2E-08	7.7E-09	6.7E-09	6.8E-09	6.8E-09
Benzo[b]fluoranthene	205-99-2	1.5E-10	1.2E-09	6.1E-10	5.9E-10	2.5E-10	1.7E-10	1.5E-10	1.5E-10	1.5E-10
Benzo[k]fluoranthene	207-08-9	1.8E-10	1.5E-09	7.6E-10	7.3E-10	3.1E-10	2.1E-10	1.8E-10	1.8E-10	1.8E-10
Chrysene	218-01-9	1.2E-09	9.7E-09	5.1E-09	4.9E-09	2.1E-09	1.4E-09	1.2E-09	1.2E-09	1.2E-09
Cyclopenta[cd]pyrene	27208-37-3	4.3E-10	3.4E-09	1.8E-09	1.7E-09	7.4E-10	4.9E-10	4.3E-10	4.3E-10	4.4E-10
Dibenz[a,h]anthracene	53-70-3	8.4E-10	6.8E-09	3.6E-09	3.4E-09	1.5E-09	9.7E-10	8.5E-10	8.6E-10	8.6E-10
Indeno[1,2,3-cd]pyrene	193-39-5	6.3E-10	5.1E-09	2.7E-09	2.6E-09	1.1E-09	7.3E-10	6.4E-10	6.4E-10	6.4E-10
Methyl Isobutyl Ketone	108-10-1	9.2E-08	7.4E-07	3.9E-07	3.7E-07	1.6E-07	1.1E-07	9.3E-08	9.3E-08	9.4E-08
Naphthalene	91-20-3	1.6E-07	1.3E-06	6.6E-07	6.3E-07	2.7E-07	1.8E-07	1.6E-07	1.6E-07	1.6E-07
Styrene	100-42-5	3.5E-07	2.8E-06	1.5E-06	1.4E-06	6.0E-07	4.0E-07	3.5E-07	3.5E-07	3.5E-07
Non-Field-Related Chemicals										
Acetaldehyde	75-07-0	1.5E-05	1.2E-04	6.3E-05	6.0E-05	2.6E-05	1.7E-05	1.5E-05	1.5E-05	1.5E-05
Benzene	71-43-2	3.6E-06	2.9E-05	1.5E-05	1.4E-05	6.2E-06	4.1E-06	3.6E-06	3.6E-06	3.6E-06
Benzene, 1,4-dichloro	106-46-7	1.1E-07	9.1E-07	4.8E-07	4.6E-07	2.0E-07	1.3E-07	1.1E-07	1.1E-07	1.1E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	3.4E-06	2.7E-05	1.4E-05	1.4E-05	5.9E-06	3.9E-06	3.4E-06	3.4E-06	3.4E-06
Ethylbenzene	100-41-4	1.0E-06	8.1E-06	4.3E-06	4.1E-06	1.8E-06	1.2E-06	1.0E-06	1.0E-06	1.0E-06
Formaldehyde	50-00-0	2.3E-05	1.8E-04	9.5E-05	9.1E-05	3.9E-05	2.6E-05	2.3E-05	2.3E-05	2.3E-05
Tetrachloroethylene	127-18-4	2.8E-07	2.3E-06	1.2E-06	1.1E-06	4.9E-07	3.2E-07	2.8E-07	2.9E-07	2.9E-07

^a On-Field ADD_{inh} was calculated from the 35 individual field average concentrations (C_{air}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-184. Average **Off-Field** Inhalation Daily Dose^a (ADD_{inh}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender Spectators

Chemical	CASRN	ADD _{inh}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Field-Related Chemicals										
Aniline	62-53-3	4.2E-08	3.4E-07	1.8E-07	1.7E-07	7.3E-08	4.8E-08	4.2E-08	4.3E-08	4.3E-08
Benz[a]anthracene	56-55-3	0	0	0	0	0	0	0	0	0



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	ADD _{inh}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Benzo[a]pyrene	50-32-8	8.3E-09	6.6E-08	3.5E-08	3.3E-08	1.4E-08	9.5E-09	8.3E-09	8.4E-09	8.4E-09
Benzo[b]fluoranthene	205-99-2	2.2E-10	1.8E-09	9.5E-10	9.0E-10	3.9E-10	2.6E-10	2.3E-10	2.3E-10	2.3E-10
Benzo[k]fluoranthene	207-08-9	1.9E-10	1.5E-09	7.9E-10	7.5E-10	3.2E-10	2.1E-10	1.9E-10	1.9E-10	1.9E-10
Chrysene	218-01-9	6.2E-10	5.0E-09	2.6E-09	2.5E-09	1.1E-09	7.1E-10	6.2E-10	6.3E-10	6.3E-10
Cyclopenta[cd]pyrene	27208-37-3	3.7E-10	3.0E-09	1.6E-09	1.5E-09	6.5E-10	4.3E-10	3.8E-10	3.8E-10	3.8E-10
Dibenz[a,h]anthracene	53-70-3	5.3E-10	4.3E-09	2.2E-09	2.1E-09	9.2E-10	6.1E-10	5.3E-10	5.4E-10	5.4E-10
Indeno[1,2,3-cd]pyrene	193-39-5	4.4E-10	3.5E-09	1.8E-09	1.8E-09	7.6E-10	5.0E-10	4.4E-10	4.4E-10	4.4E-10
Methyl Isobutyl Ketone	108-10-1	2.1E-08	1.7E-07	8.9E-08	8.5E-08	3.7E-08	2.4E-08	2.1E-08	2.1E-08	2.1E-08
Naphthalene	91-20-3	1.7E-07	1.4E-06	7.3E-07	6.9E-07	3.0E-07	2.0E-07	1.7E-07	1.7E-07	1.7E-07
Styrene	100-42-5	3.5E-07	2.8E-06	1.5E-06	1.4E-06	6.2E-07	4.1E-07	3.6E-07	3.6E-07	3.6E-07
Non-Field-Related Chemicals										
Benzene	71-43-2	3.6E-06	3.7E-06	3.0E-05	1.6E-05	1.5E-05	6.5E-06	4.3E-06	3.8E-06	3.8E-06
Benzene, 1,4-dichloro	106-46-7	1.1E-07	1.0E-07	8.3E-07	4.3E-07	4.1E-07	1.8E-07	1.2E-07	1.0E-07	1.0E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	3.4E-06	3.4E-06	2.7E-05	1.4E-05	1.4E-05	5.9E-06	3.9E-06	3.4E-06	3.5E-06
Ethylbenzene	100-41-4	1.0E-06	1.1E-06	8.6E-06	4.5E-06	4.3E-06	1.9E-06	1.2E-06	1.1E-06	1.1E-06
Tetrachloroethylene	127-18-4	2.8E-07	2.9E-07	2.3E-06	1.2E-06	1.2E-06	5.0E-07	3.3E-07	2.9E-07	2.9E-07

^a Off-Field ADD_{inh} was calculated from the 35 individual field average concentrations (C_{air}) of a chemical detected in air off the fields. Carbonyl samples were not collected at off-field locations.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

INDIVIDUAL FIELD ASSESSMENT (Table F-185 to Table F-217)

Table F-185. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Athletes 2<6 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	2.9E-07	5.1E-07	0	1.1E-06	1.9E-06
Benz[a]anthracene	56-55-3	1	0	2.2E-10	1.3E-09	0	0	7.4E-09
Benzo[a]pyrene	50-32-8	28	0	5.1E-08	5.3E-08	2.7E-08	1.4E-07	2.1E-07
Benzo[b]fluoranthene	205-99-2	2	0	1.1E-09	4.4E-09	0	6.5E-09	1.9E-08
Benzo[k]fluoranthene	207-08-9	4	0	1.4E-09	4.0E-09	0	1.0E-08	1.5E-08
Chrysene	218-01-9	13	0	9.1E-09	1.4E-08	0	3.4E-08	5.9E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	3.2E-09	3.9E-09	1.2E-09	9.3E-09	1.8E-08
Dibenz[a,h]anthracene	53-70-3	14	0	6.4E-09	1.3E-08	0	3.2E-08	5.4E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	4.8E-09	1.6E-08	0	5.4E-08	5.4E-08



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Methyl Isobutyl Ketone	108-10-1	5	0	6.9E-07	1.9E-06	0	5.4E-06	8.1E-06
Naphthalene	91-20-3	9	0	1.2E-06	2.5E-06	0	5.1E-06	1.1E-05
Styrene	100-42-5	17	0	2.6E-06	5.4E-06	0	8.8E-06	2.9E-05
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	1.2E-05	1.1E-04	8.2E-05	8.5E-05	2.1E-04	4.2E-04
Benzene	71-43-2	35	4.0E-06	2.7E-05	2.2E-05	1.9E-05	6.7E-05	1.1E-04
Benzene, 1,4-dichloro	106-46-7	7	0	8.5E-07	1.9E-06	0	5.3E-06	5.3E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	2.5E-05	2.6E-05	1.7E-05	7.6E-05	9.4E-05
Ethylbenzene	100-41-4	13	0	7.6E-06	1.3E-05	0	3.1E-05	5.1E-05
Formaldehyde	50-00-0	34	3.6E-05	1.7E-04	1.3E-04	1.4E-04	2.8E-04	7.3E-04
Tetrachloroethylene	127-18-4	7	0	2.1E-06	5.0E-06	0	1.4E-05	1.9E-05

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-186. Field-Specific Average **On-Field** Inhalation Daily Dose^a milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Athletes 6<11 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	3.7E-07	6.6E-07	0	1.5E-06	2.5E-06
Benz[a]anthracene	56-55-3	1	0	2.8E-10	1.7E-09	0	0	9.6E-09
Benzo[a]pyrene	50-32-8	28	0	6.6E-08	6.9E-08	3.5E-08	1.8E-07	2.7E-07
Benzo[b]fluoranthene	205-99-2	2	0	1.4E-09	5.8E-09	0	8.4E-09	2.4E-08
Benzo[k]fluoranthene	207-08-9	4	0	1.8E-09	5.2E-09	0	1.3E-08	2.0E-08
Chrysene	218-01-9	13	0	1.2E-08	1.8E-08	0	4.4E-08	7.6E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	4.2E-09	5.1E-09	1.6E-09	1.2E-08	2.4E-08
Dibenz[a,h]anthracene	53-70-3	14	0	8.2E-09	1.6E-08	0	4.1E-08	6.9E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	6.2E-09	2.0E-08	0	7.0E-08	7.0E-08
Methyl Isobutyl Ketone	108-10-1	5	0	9.0E-07	2.5E-06	0	7.0E-06	1.0E-05
Naphthalene	91-20-3	9	0	1.5E-06	3.2E-06	0	6.6E-06	1.5E-05
Styrene	100-42-5	17	0	3.4E-06	7.0E-06	0	1.1E-05	3.8E-05
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	1.5E-05	1.4E-04	1.1E-04	1.1E-04	2.8E-04	5.5E-04
Benzene	71-43-2	35	5.2E-06	3.5E-05	2.8E-05	2.4E-05	8.7E-05	1.4E-04



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene, 1,4-dichloro	106-46-7	7	0	1.1E-06	2.4E-06	0	6.8E-06	6.8E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	3.3E-05	3.4E-05	2.2E-05	9.8E-05	1.2E-04
Ethylbenzene	100-41-4	13	0	9.9E-06	1.7E-05	0	4.0E-05	6.6E-05
Formaldehyde	50-00-0	34	4.7E-05	2.2E-04	1.7E-04	1.8E-04	3.7E-04	9.4E-04
Tetrachloroethylene	127-18-4	7	0	2.7E-06	6.5E-06	0	1.8E-05	2.4E-05

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-187. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Athletes 11<16 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	3.6E-07	6.4E-07	0	1.4E-06	2.5E-06
Benz[a]anthracene	56-55-3	1	0	2.8E-10	1.6E-09	0	0	9.4E-09
Benzo[a]pyrene	50-32-8	28	0	6.4E-08	6.7E-08	3.4E-08	1.7E-07	2.7E-07
Benzo[b]fluoranthene	205-99-2	2	0	1.4E-09	5.6E-09	0	8.2E-09	2.4E-08
Benzo[k]fluoranthene	207-08-9	4	0	1.7E-09	5.1E-09	0	1.3E-08	2.0E-08
Chrysene	218-01-9	13	0	1.2E-08	1.8E-08	0	4.3E-08	7.4E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	4.1E-09	5.0E-09	1.5E-09	1.2E-08	2.3E-08
Dibenz[a,h]anthracene	53-70-3	14	0	8.1E-09	1.6E-08	0	4.0E-08	6.8E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	6.0E-09	2.0E-08	0	6.9E-08	6.9E-08
Methyl Isobutyl Ketone	108-10-1	5	0	8.8E-07	2.4E-06	0	6.8E-06	1.0E-05
Naphthalene	91-20-3	9	0	1.5E-06	3.1E-06	0	6.4E-06	1.5E-05
Styrene	100-42-5	17	0	3.3E-06	6.8E-06	0	1.1E-05	3.7E-05
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	1.5E-05	1.4E-04	1.0E-04	1.1E-04	2.7E-04	5.4E-04
Benzene	71-43-2	35	5.1E-06	3.4E-05	2.7E-05	2.4E-05	8.5E-05	1.4E-04
Benzene, 1,4-dichloro	106-46-7	7	0	1.1E-06	2.4E-06	0	6.7E-06	6.7E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	3.2E-05	3.4E-05	2.1E-05	9.6E-05	1.2E-04
Ethylbenzene	100-41-4	13	0	9.6E-06	1.6E-05	0	3.9E-05	6.5E-05
Formaldehyde	50-00-0	34	4.6E-05	2.2E-04	1.6E-04	1.8E-04	3.6E-04	9.2E-04
Tetrachloroethylene	127-18-4	7	0	2.7E-06	6.4E-06	0	1.8E-05	2.3E-05

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected



in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-188. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Athletes 16<30 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	6.9E-07	1.2E-06	0	2.7E-06	4.6E-06
Benz[a]anthracene	56-55-3	1	0	5.2E-10	3.0E-09	0	0	1.8E-08
Benzo[a]pyrene	50-32-8	28	0	1.2E-07	1.3E-07	6.5E-08	3.3E-07	5.0E-07
Benzo[b]fluoranthene	205-99-2	2	0	2.6E-09	1.1E-08	0	1.6E-08	4.4E-08
Benzo[k]fluoranthene	207-08-9	4	0	3.2E-09	9.6E-09	0	2.5E-08	3.7E-08
Chrysene	218-01-9	13	0	2.2E-08	3.4E-08	0	8.1E-08	1.4E-07
Cyclopenta[cd]pyrene	27208-37-3	28	0	7.7E-09	9.3E-09	2.9E-09	2.2E-08	4.4E-08
Dibenz[a,h]anthracene	53-70-3	14	0	1.5E-08	3.0E-08	0	7.6E-08	1.3E-07
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	1.1E-08	3.7E-08	0	1.3E-07	1.3E-07
Methyl Isobutyl Ketone	108-10-1	5	0	1.7E-06	4.5E-06	0	1.3E-05	1.9E-05
Naphthalene	91-20-3	9	0	2.8E-06	5.9E-06	0	1.2E-05	2.7E-05
Styrene	100-42-5	17	0	6.2E-06	1.3E-05	0	2.1E-05	7.0E-05
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	2.8E-05	2.7E-04	2.0E-04	2.0E-04	5.1E-04	1.0E-03
Benzene	71-43-2	35	9.6E-06	6.4E-05	5.2E-05	4.5E-05	1.6E-04	2.6E-04
Benzene, 1,4-dichloro	106-46-7	7	0	2.0E-06	4.5E-06	0	1.3E-05	1.3E-05
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	6.1E-05	6.3E-05	4.0E-05	1.8E-04	2.2E-04
Ethylbenzene	100-41-4	13	0	1.8E-05	3.1E-05	0	7.4E-05	1.2E-04
Formaldehyde	50-00-0	34	8.6E-05	4.1E-04	3.0E-04	3.4E-04	6.8E-04	1.7E-03
Tetrachloroethylene	127-18-4	7	0	5.1E-06	1.2E-05	0	3.4E-05	4.4E-05

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-189. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Athletes 30<40 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	10	0	4.5E-07	7.9E-07	0	1.7E-06	3.0E-06
Benz[a]anthracene	56-55-3	1	0	3.4E-10	2.0E-09	0	0	1.2E-08
Benzo[a]pyrene	50-32-8	28	0	7.9E-08	8.3E-08	4.2E-08	2.1E-07	3.3E-07
Benzo[b]fluoranthene	205-99-2	2	0	1.7E-09	6.9E-09	0	1.0E-08	2.9E-08
Benzo[k]fluoranthene	207-08-9	4	0	2.1E-09	6.3E-09	0	1.6E-08	2.4E-08
Chrysene	218-01-9	13	0	1.4E-08	2.2E-08	0	5.3E-08	9.1E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	5.0E-09	6.1E-09	1.9E-09	1.5E-08	2.9E-08
Dibenz[a,h]anthracene	53-70-3	14	0	9.9E-09	2.0E-08	0	4.9E-08	8.4E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	7.4E-09	2.4E-08	0	8.4E-08	8.4E-08
Methyl Isobutyl Ketone	108-10-1	5	0	1.1E-06	2.9E-06	0	8.4E-06	1.3E-05
Naphthalene	91-20-3	9	0	1.8E-06	3.8E-06	0	7.9E-06	1.8E-05
Styrene	100-42-5	17	0	4.1E-06	8.4E-06	0	1.4E-05	4.6E-05
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	1.8E-05	1.7E-04	1.3E-04	1.3E-04	3.3E-04	6.6E-04
Benzene	71-43-2	35	6.2E-06	4.2E-05	3.4E-05	2.9E-05	1.0E-04	1.7E-04
Benzene, 1,4-dichloro	106-46-7	7	0	1.3E-06	2.9E-06	0	8.2E-06	8.2E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	4.0E-05	4.1E-05	2.6E-05	1.2E-04	1.5E-04
Ethylbenzene	100-41-4	13	0	1.2E-05	2.0E-05	0	4.8E-05	8.0E-05
Formaldehyde	50-00-0	34	5.6E-05	2.6E-04	2.0E-04	2.2E-04	4.4E-04	1.1E-03
Tetrachloroethylene	127-18-4	7	0	3.3E-06	7.8E-06	0	2.2E-05	2.9E-05

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-190. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Athletes 40<50 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	4.4E-07	7.8E-07	0	1.7E-06	3.0E-06
Benz[a]anthracene	56-55-3	1	0	3.4E-10	2.0E-09	0	0	1.1E-08
Benzo[a]pyrene	50-32-8	28	0	7.8E-08	8.2E-08	4.2E-08	2.1E-07	3.2E-07
Benzo[b]fluoranthene	205-99-2	2	0	1.7E-09	6.8E-09	0	1.0E-08	2.9E-08
Benzo[k]fluoranthene	207-08-9	4	0	2.1E-09	6.2E-09	0	1.6E-08	2.4E-08
Chrysene	218-01-9	13	0	1.4E-08	2.2E-08	0	5.2E-08	9.0E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	5.0E-09	6.0E-09	1.8E-09	1.4E-08	2.8E-08



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dibenz[a,h]anthracene	53-70-3	14	0	9.8E-09	2.0E-08	0	4.9E-08	8.2E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	7.3E-09	2.4E-08	0	8.3E-08	8.3E-08
Methyl Isobutyl Ketone	108-10-1	5	0	1.1E-06	2.9E-06	0	8.3E-06	1.2E-05
Naphthalene	91-20-3	9	0	1.8E-06	3.8E-06	0	7.8E-06	1.8E-05
Styrene	100-42-5	17	0	4.0E-06	8.3E-06	0	1.4E-05	4.5E-05
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	1.8E-05	1.7E-04	1.3E-04	1.3E-04	3.3E-04	6.5E-04
Benzene	71-43-2	35	6.2E-06	4.1E-05	3.3E-05	2.9E-05	1.0E-04	1.7E-04
Benzene, 1,4-dichloro	106-46-7	7	0	1.3E-06	2.9E-06	0	8.1E-06	8.1E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	3.9E-05	4.1E-05	2.6E-05	1.2E-04	1.4E-04
Ethylbenzene	100-41-4	13	0	1.2E-05	2.0E-05	0	4.8E-05	7.9E-05
Formaldehyde	50-00-0	34	5.5E-05	2.6E-04	2.0E-04	2.2E-04	4.4E-04	1.1E-03
Tetrachloroethylene	127-18-4	7	0	3.3E-06	7.7E-06	0	2.2E-05	2.9E-05

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-191. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Athletes 50<70 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	4.3E-07	7.6E-07	0	1.7E-06	2.9E-06
Benz[a]anthracene	56-55-3	1	0	3.3E-10	1.9E-09	0	0	1.1E-08
Benzo[a]pyrene	50-32-8	28	0	7.6E-08	8.0E-08	4.1E-08	2.1E-07	3.2E-07
Benzo[b]fluoranthene	205-99-2	2	0	1.7E-09	6.7E-09	0	9.8E-09	2.8E-08
Benzo[k]fluoranthene	207-08-9	4	0	2.1E-09	6.1E-09	0	1.6E-08	2.3E-08
Chrysene	218-01-9	13	0	1.4E-08	2.1E-08	0	5.1E-08	8.9E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	4.9E-09	5.9E-09	1.8E-09	1.4E-08	2.8E-08
Dibenz[a,h]anthracene	53-70-3	14	0	9.6E-09	1.9E-08	0	4.8E-08	8.1E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	7.2E-09	2.4E-08	0	8.2E-08	8.2E-08
Methyl Isobutyl Ketone	108-10-1	5	0	1.0E-06	2.9E-06	0	8.1E-06	1.2E-05
Naphthalene	91-20-3	9	0	1.8E-06	3.7E-06	0	7.7E-06	1.7E-05
Styrene	100-42-5	17	0	3.9E-06	8.1E-06	0	1.3E-05	4.4E-05
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	1.7E-05	1.7E-04	1.2E-04	1.3E-04	3.2E-04	6.4E-04



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene	71-43-2	35	6.0E-06	4.1E-05	3.3E-05	2.9E-05	1.0E-04	1.6E-04
Benzene, 1,4-dichloro	106-46-7	7	0	1.3E-06	2.8E-06	0	7.9E-06	7.9E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	3.9E-05	4.0E-05	2.6E-05	1.1E-04	1.4E-04
Ethylbenzene	100-41-4	13	0	1.1E-05	1.9E-05	0	4.7E-05	7.7E-05
Formaldehyde	50-00-0	34	5.4E-05	2.6E-04	1.9E-04	2.1E-04	4.3E-04	1.1E-03
Tetrachloroethylene	127-18-4	7	0	3.2E-06	7.6E-06	0	2.1E-05	2.8E-05

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-192. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Coaches 16<30 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	2.8E-07	5.0E-07	0	1.1E-06	1.9E-06
Benz[a]anthracene	56-55-3	1	0	2.2E-10	1.3E-09	0	0	7.3E-09
Benzo[a]pyrene	50-32-8	28	0	5.0E-08	5.3E-08	2.7E-08	1.3E-07	2.1E-07
Benzo[b]fluoranthene	205-99-2	2	0	1.1E-09	4.4E-09	0	6.4E-09	1.8E-08
Benzo[k]fluoranthene	207-08-9	4	0	1.3E-09	4.0E-09	0	1.0E-08	1.5E-08
Chrysene	218-01-9	13	0	9.0E-09	1.4E-08	0	3.4E-08	5.8E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	3.2E-09	3.9E-09	1.2E-09	9.2E-09	1.8E-08
Dibenz[a,h]anthracene	53-70-3	14	0	6.3E-09	1.3E-08	0	3.1E-08	5.3E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	4.7E-09	1.5E-08	0	5.3E-08	5.3E-08
Methyl Isobutyl Ketone	108-10-1	5	0	6.8E-07	1.9E-06	0	5.3E-06	8.0E-06
Naphthalene	91-20-3	9	0	1.2E-06	2.4E-06	0	5.0E-06	1.1E-05
Styrene	100-42-5	17	0	2.6E-06	5.3E-06	0	8.7E-06	2.9E-05
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	1.1E-05	1.1E-04	8.1E-05	8.4E-05	2.1E-04	4.2E-04
Benzene	71-43-2	35	4.0E-06	2.6E-05	2.1E-05	1.9E-05	6.6E-05	1.1E-04
Benzene, 1,4-dichloro	106-46-7	7	0	8.4E-07	1.8E-06	0	5.2E-06	5.2E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	2.5E-05	2.6E-05	1.7E-05	7.5E-05	9.3E-05
Ethylbenzene	100-41-4	13	0	7.5E-06	1.3E-05	0	3.1E-05	5.1E-05
Formaldehyde	50-00-0	34	3.5E-05	1.7E-04	1.3E-04	1.4E-04	2.8E-04	7.2E-04
Tetrachloroethylene	127-18-4	7	0	2.1E-06	4.9E-06	0	1.4E-05	1.8E-05

^a 35 field-specific ADD_{inh-field} values are included in the table.



^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-193. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Coaches 30<40 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	2.5E-07	4.4E-07	0	9.6E-07	1.7E-06
Benz[a]anthracene	56-55-3	1	0	1.9E-10	1.1E-09	0	0	6.4E-09
Benzo[a]pyrene	50-32-8	28	0	4.4E-08	4.6E-08	2.3E-08	1.2E-07	1.8E-07
Benzo[b]fluoranthene	205-99-2	2	0	9.4E-10	3.8E-09	0	5.6E-09	1.6E-08
Benzo[k]fluoranthene	207-08-9	4	0	1.2E-09	3.5E-09	0	9.0E-09	1.3E-08
Chrysene	218-01-9	13	0	7.8E-09	1.2E-08	0	2.9E-08	5.1E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	2.8E-09	3.4E-09	1.0E-09	8.0E-09	1.6E-08
Dibenz[a,h]anthracene	53-70-3	14	0	5.5E-09	1.1E-08	0	2.7E-08	4.6E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	4.1E-09	1.3E-08	0	4.7E-08	4.7E-08
Methyl Isobutyl Ketone	108-10-1	5	0	6.0E-07	1.6E-06	0	4.7E-06	7.0E-06
Naphthalene	91-20-3	9	0	1.0E-06	2.1E-06	0	4.4E-06	9.9E-06
Styrene	100-42-5	17	0	2.2E-06	4.6E-06	0	7.6E-06	2.5E-05
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	1.0E-05	9.6E-05	7.1E-05	7.3E-05	1.8E-04	3.7E-04
Benzene	71-43-2	35	3.5E-06	2.3E-05	1.9E-05	1.6E-05	5.8E-05	9.4E-05
Benzene, 1,4-dichloro	106-46-7	7	0	7.3E-07	1.6E-06	0	4.5E-06	4.5E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	2.2E-05	2.3E-05	1.5E-05	6.5E-05	8.1E-05
Ethylbenzene	100-41-4	13	0	6.6E-06	1.1E-05	0	2.7E-05	4.4E-05
Formaldehyde	50-00-0	34	3.1E-05	1.5E-04	1.1E-04	1.2E-04	2.4E-04	6.3E-04
Tetrachloroethylene	127-18-4	7	0	1.8E-06	4.3E-06	0	1.2E-05	1.6E-05

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-194. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Coaches 40<50 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	2.5E-07	4.4E-07	0	9.7E-07	1.7E-06
Benz[a]anthracene	56-55-3	1	0	1.9E-10	1.1E-09	0	0	6.5E-09
Benzo[a]pyrene	50-32-8	28	0	4.4E-08	4.6E-08	2.4E-08	1.2E-07	1.8E-07
Benzo[b]fluoranthene	205-99-2	2	0	9.5E-10	3.9E-09	0	5.7E-09	1.6E-08
Benzo[k]fluoranthene	207-08-9	4	0	1.2E-09	3.5E-09	0	9.1E-09	1.3E-08
Chrysene	218-01-9	13	0	7.9E-09	1.2E-08	0	3.0E-08	5.1E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	2.8E-09	3.4E-09	1.0E-09	8.1E-09	1.6E-08
Dibenz[a,h]anthracene	53-70-3	14	0	5.5E-09	1.1E-08	0	2.8E-08	4.7E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	4.2E-09	1.4E-08	0	4.7E-08	4.7E-08
Methyl Isobutyl Ketone	108-10-1	5	0	6.0E-07	1.6E-06	0	4.7E-06	7.1E-06
Naphthalene	91-20-3	9	0	1.0E-06	2.1E-06	0	4.4E-06	1.0E-05
Styrene	100-42-5	17	0	2.3E-06	4.7E-06	0	7.7E-06	2.5E-05
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	1.0E-05	9.7E-05	7.2E-05	7.4E-05	1.9E-04	3.7E-04
Benzene	71-43-2	35	3.5E-06	2.3E-05	1.9E-05	1.6E-05	5.8E-05	9.5E-05
Benzene, 1,4-dichloro	106-46-7	7	0	7.4E-07	1.6E-06	0	4.6E-06	4.6E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	2.2E-05	2.3E-05	1.5E-05	6.6E-05	8.2E-05
Ethylbenzene	100-41-4	13	0	6.6E-06	1.1E-05	0	2.7E-05	4.5E-05
Formaldehyde	50-00-0	34	3.1E-05	1.5E-04	1.1E-04	1.2E-04	2.5E-04	6.3E-04
Tetrachloroethylene	127-18-4	7	0	1.8E-06	4.4E-06	0	1.2E-05	1.6E-05

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-195. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Coaches 50<70 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	2.5E-07	4.4E-07	0	9.8E-07	1.7E-06
Benz[a]anthracene	56-55-3	1	0	1.9E-10	1.1E-09	0	0	6.5E-09
Benzo[a]pyrene	50-32-8	28	0	4.4E-08	4.7E-08	2.4E-08	1.2E-07	1.8E-07



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[b]fluoranthene	205-99-2	2	0	9.5E-10	3.9E-09	0	5.7E-09	1.6E-08
Benzo[k]fluoranthene	207-08-9	4	0	1.2E-09	3.5E-09	0	9.1E-09	1.3E-08
Chrysene	218-01-9	13	0	7.9E-09	1.2E-08	0	3.0E-08	5.1E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	2.8E-09	3.4E-09	1.1E-09	8.1E-09	1.6E-08
Dibenz[a,h]anthracene	53-70-3	14	0	5.6E-09	1.1E-08	0	2.8E-08	4.7E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	4.2E-09	1.4E-08	0	4.7E-08	4.7E-08
Methyl Isobutyl Ketone	108-10-1	5	0	6.1E-07	1.7E-06	0	4.7E-06	7.1E-06
Naphthalene	91-20-3	9	0	1.0E-06	2.2E-06	0	4.4E-06	1.0E-05
Styrene	100-42-5	17	0	2.3E-06	4.7E-06	0	7.7E-06	2.5E-05
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	1.0E-05	9.8E-05	7.2E-05	7.4E-05	1.9E-04	3.7E-04
Benzene	71-43-2	35	3.5E-06	2.3E-05	1.9E-05	1.6E-05	5.9E-05	9.5E-05
Benzene, 1,4-dichloro	106-46-7	7	0	7.4E-07	1.6E-06	0	4.6E-06	4.6E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	2.2E-05	2.3E-05	1.5E-05	6.6E-05	8.2E-05
Ethylbenzene	100-41-4	13	0	6.7E-06	1.1E-05	0	2.7E-05	4.5E-05
Formaldehyde	50-00-0	34	3.1E-05	1.5E-04	1.1E-04	1.2E-04	2.5E-04	6.3E-04
Tetrachloroethylene	127-18-4	7	0	1.9E-06	4.4E-06	0	1.2E-05	1.6E-05

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-196. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Referees 16<30 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	1.1E-07	1.9E-07	0	4.1E-07	7.1E-07
Benz[a]anthracene	56-55-3	1	0	8.0E-11	4.7E-10	0	0	2.7E-09
Benzo[a]pyrene	50-32-8	28	0	1.9E-08	2.0E-08	9.9E-09	5.0E-08	7.7E-08
Benzo[b]fluoranthene	205-99-2	2	0	4.0E-10	1.6E-09	0	2.4E-09	6.8E-09
Benzo[k]fluoranthene	207-08-9	4	0	5.0E-10	1.5E-09	0	3.8E-09	5.6E-09
Chrysene	218-01-9	13	0	3.3E-09	5.2E-09	0	1.2E-08	2.1E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	1.2E-09	1.4E-09	4.4E-10	3.4E-09	6.7E-09
Dibenz[a,h]anthracene	53-70-3	14	0	2.3E-09	4.6E-09	0	1.2E-08	2.0E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	1.7E-09	5.7E-09	0	2.0E-08	2.0E-08
Methyl Isobutyl Ketone	108-10-1	5	0	2.5E-07	6.9E-07	0	2.0E-06	3.0E-06



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene	91-20-3	9	0	4.3E-07	9.0E-07	0	1.9E-06	4.2E-06
Styrene	100-42-5	17	0	9.6E-07	2.0E-06	0	3.2E-06	1.1E-05
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	4.2E-06	4.1E-05	3.0E-05	3.1E-05	7.8E-05	1.6E-04
Benzene	71-43-2	35	1.5E-06	9.8E-06	7.9E-06	6.9E-06	2.5E-05	4.0E-05
Benzene, 1,4-dichloro	106-46-7	7	0	3.1E-07	6.8E-07	0	1.9E-06	1.9E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	9.3E-06	9.7E-06	6.2E-06	2.8E-05	3.4E-05
Ethylbenzene	100-41-4	13	0	2.8E-06	4.7E-06	0	1.1E-05	1.9E-05
Formaldehyde	50-00-0	34	1.3E-05	6.2E-05	4.7E-05	5.2E-05	1.0E-04	2.7E-04
Tetrachloroethylene	127-18-4	7	0	7.8E-07	1.8E-06	0	5.2E-06	6.8E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-197. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Referees 30<40 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	9.2E-08	1.6E-07	0	3.6E-07	6.2E-07
Benz[a]anthracene	56-55-3	1	0	7.0E-11	4.1E-10	0	0	2.4E-09
Benzo[a]pyrene	50-32-8	28	0	1.6E-08	1.7E-08	8.7E-09	4.4E-08	6.7E-08
Benzo[b]fluoranthene	205-99-2	2	0	3.5E-10	1.4E-09	0	2.1E-09	5.9E-09
Benzo[k]fluoranthene	207-08-9	4	0	4.3E-10	1.3E-09	0	3.3E-09	4.9E-09
Chrysene	218-01-9	13	0	2.9E-09	4.5E-09	0	1.1E-08	1.9E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	1.0E-09	1.3E-09	3.8E-10	3.0E-09	5.9E-09
Dibenz[a,h]anthracene	53-70-3	14	0	2.0E-09	4.1E-09	0	1.0E-08	1.7E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	1.5E-09	5.0E-09	0	1.7E-08	1.7E-08
Methyl Isobutyl Ketone	108-10-1	5	0	2.2E-07	6.0E-07	0	1.7E-06	2.6E-06
Naphthalene	91-20-3	9	0	3.8E-07	7.9E-07	0	1.6E-06	3.7E-06
Styrene	100-42-5	17	0	8.4E-07	1.7E-06	0	2.8E-06	9.3E-06
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	3.7E-06	3.6E-05	2.6E-05	2.7E-05	6.8E-05	1.4E-04
Benzene	71-43-2	35	1.3E-06	8.6E-06	6.9E-06	6.0E-06	2.1E-05	3.5E-05
Benzene, 1,4-dichloro	106-46-7	7	0	2.7E-07	6.0E-07	0	1.7E-06	1.7E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	8.2E-06	8.5E-06	5.4E-06	2.4E-05	3.0E-05



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Ethylbenzene	100-41-4	13	0	2.4E-06	4.1E-06	0	9.9E-06	1.6E-05
Formaldehyde	50-00-0	34	1.2E-05	5.4E-05	4.1E-05	4.5E-05	9.1E-05	2.3E-04
Tetrachloroethylene	127-18-4	7	0	6.8E-07	1.6E-06	0	4.5E-06	5.9E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-198. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Referees 40<50 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	9.3E-08	1.6E-07	0	3.6E-07	6.3E-07
Benz[a]anthracene	56-55-3	1	0	7.1E-11	4.1E-10	0	0	2.4E-09
Benzo[a]pyrene	50-32-8	28	0	1.6E-08	1.7E-08	8.8E-09	4.4E-08	6.8E-08
Benzo[b]fluoranthene	205-99-2	2	0	3.5E-10	1.4E-09	0	2.1E-09	6.0E-09
Benzo[k]fluoranthene	207-08-9	4	0	4.4E-10	1.3E-09	0	3.4E-09	5.0E-09
Chrysene	218-01-9	13	0	2.9E-09	4.6E-09	0	1.1E-08	1.9E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	1.0E-09	1.3E-09	3.9E-10	3.0E-09	5.9E-09
Dibenz[a,h]anthracene	53-70-3	14	0	2.1E-09	4.1E-09	0	1.0E-08	1.7E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	1.5E-09	5.0E-09	0	1.7E-08	1.7E-08
Methyl Isobutyl Ketone	108-10-1	5	0	2.2E-07	6.1E-07	0	1.7E-06	2.6E-06
Naphthalene	91-20-3	9	0	3.8E-07	8.0E-07	0	1.6E-06	3.7E-06
Styrene	100-42-5	17	0	8.4E-07	1.7E-06	0	2.8E-06	9.4E-06
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	3.7E-06	3.6E-05	2.7E-05	2.7E-05	6.9E-05	1.4E-04
Benzene	71-43-2	35	1.3E-06	8.7E-06	7.0E-06	6.1E-06	2.2E-05	3.5E-05
Benzene, 1,4-dichloro	106-46-7	7	0	2.8E-07	6.0E-07	0	1.7E-06	1.7E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	8.2E-06	8.6E-06	5.5E-06	2.5E-05	3.0E-05
Ethylbenzene	100-41-4	13	0	2.5E-06	4.2E-06	0	1.0E-05	1.7E-05
Formaldehyde	50-00-0	34	1.2E-05	5.5E-05	4.1E-05	4.6E-05	9.2E-05	2.3E-04
Tetrachloroethylene	127-18-4	7	0	6.9E-07	1.6E-06	0	4.6E-06	6.0E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-199. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Referees 50<70 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	9.3E-08	1.6E-07	0	3.6E-07	6.3E-07
Benz[a]anthracene	56-55-3	1	0	7.1E-11	4.1E-10	0	0	2.4E-09
Benzo[a]pyrene	50-32-8	28	0	1.6E-08	1.7E-08	8.8E-09	4.4E-08	6.8E-08
Benzo[b]fluoranthene	205-99-2	2	0	3.5E-10	1.4E-09	0	2.1E-09	6.0E-09
Benzo[k]fluoranthene	207-08-9	4	0	4.4E-10	1.3E-09	0	3.4E-09	5.0E-09
Chrysene	218-01-9	13	0	2.9E-09	4.6E-09	0	1.1E-08	1.9E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	1.0E-09	1.3E-09	3.9E-10	3.0E-09	5.9E-09
Dibenz[a,h]anthracene	53-70-3	14	0	2.1E-09	4.1E-09	0	1.0E-08	1.7E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	1.5E-09	5.0E-09	0	1.8E-08	1.8E-08
Methyl Isobutyl Ketone	108-10-1	5	0	2.3E-07	6.1E-07	0	1.8E-06	2.6E-06
Naphthalene	91-20-3	9	0	3.8E-07	8.0E-07	0	1.7E-06	3.7E-06
Styrene	100-42-5	17	0	8.5E-07	1.7E-06	0	2.9E-06	9.5E-06
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	3.8E-06	3.6E-05	2.7E-05	2.8E-05	6.9E-05	1.4E-04
Benzene	71-43-2	35	1.3E-06	8.7E-06	7.0E-06	6.1E-06	2.2E-05	3.5E-05
Benzene, 1,4-dichloro	106-46-7	7	0	2.8E-07	6.1E-07	0	1.7E-06	1.7E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	8.3E-06	8.6E-06	5.5E-06	2.5E-05	3.0E-05
Ethylbenzene	100-41-4	13	0	2.5E-06	4.2E-06	0	1.0E-05	1.7E-05
Formaldehyde	50-00-0	34	1.2E-05	5.5E-05	4.1E-05	4.6E-05	9.2E-05	2.4E-04
Tetrachloroethylene	127-18-4	7	0	6.9E-07	1.6E-06	0	4.6E-06	6.0E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-200. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators Thrid Trimester Fetus<0 Year**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	3.8E-08	6.7E-08	0	1.5E-07	2.6E-07
Benz[a]anthracene	56-55-3	1	0	2.9E-11	1.7E-10	0	0	9.9E-10
Benzo[a]pyrene	50-32-8	28	0	6.7E-09	7.1E-09	3.6E-09	1.8E-08	2.8E-08



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[b]fluoranthene	205-99-2	2	0	1.5E-10	5.9E-10	0	8.6E-10	2.5E-09
Benzo[k]fluoranthene	207-08-9	4	0	1.8E-10	5.3E-10	0	1.4E-09	2.0E-09
Chrysene	218-01-9	13	0	1.2E-09	1.9E-09	0	4.5E-09	7.8E-09
Cyclopenta[cd]pyrene	27208-37-3	28	0	4.3E-10	5.2E-10	1.6E-10	1.2E-09	2.4E-09
Dibenz[a,h]anthracene	53-70-3	14	0	8.4E-10	1.7E-09	0	4.2E-09	7.1E-09
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	6.3E-10	2.1E-09	0	7.2E-09	7.2E-09
Methyl Isobutyl Ketone	108-10-1	5	0	9.2E-08	2.5E-07	0	7.2E-07	1.1E-06
Naphthalene	91-20-3	9	0	1.6E-07	3.3E-07	0	6.8E-07	1.5E-06
Styrene	100-42-5	17	0	3.5E-07	7.1E-07	0	1.2E-06	3.9E-06
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	1.5E-06	1.5E-05	1.1E-05	1.1E-05	2.8E-05	5.6E-05
Benzene	71-43-2	35	5.3E-07	3.6E-06	2.9E-06	2.5E-06	8.9E-06	1.4E-05
Benzene, 1,4-dichloro	106-46-7	7	0	1.1E-07	2.5E-07	0	7.0E-07	7.0E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	3.4E-06	3.5E-06	2.2E-06	1.0E-05	1.2E-05
Ethylbenzene	100-41-4	13	0	1.0E-06	1.7E-06	0	4.1E-06	6.8E-06
Formaldehyde	50-00-0	34	4.8E-06	2.3E-05	1.7E-05	1.9E-05	3.8E-05	9.6E-05
Tetrachloroethylene	127-18-4	7	0	2.8E-07	6.7E-07	0	1.9E-06	2.5E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-201. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators 0<2 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	3.1E-07	5.4E-07	0	1.2E-06	2.1E-06
Benz[a]anthracene	56-55-3	1	0	2.3E-10	1.4E-09	0	0	7.9E-09
Benzo[a]pyrene	50-32-8	28	0	5.4E-08	5.7E-08	2.9E-08	1.5E-07	2.2E-07
Benzo[b]fluoranthene	205-99-2	2	0	1.2E-09	4.7E-09	0	6.9E-09	2.0E-08
Benzo[k]fluoranthene	207-08-9	4	0	1.5E-09	4.3E-09	0	1.1E-08	1.6E-08
Chrysene	218-01-9	13	0	9.7E-09	1.5E-08	0	3.6E-08	6.3E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	3.4E-09	4.2E-09	1.3E-09	1.0E-08	2.0E-08
Dibenz[a,h]anthracene	53-70-3	14	0	6.8E-09	1.4E-08	0	3.4E-08	5.7E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	5.1E-09	1.7E-08	0	5.8E-08	5.8E-08
Methyl Isobutyl Ketone	108-10-1	5	0	7.4E-07	2.0E-06	0	5.8E-06	8.6E-06



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene	91-20-3	9	0	1.3E-06	2.6E-06	0	5.4E-06	1.2E-05
Styrene	100-42-5	17	0	2.8E-06	5.7E-06	0	9.4E-06	3.1E-05
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	1.2E-05	1.2E-04	8.8E-05	9.1E-05	2.3E-04	4.5E-04
Benzene	71-43-2	35	4.3E-06	2.9E-05	2.3E-05	2.0E-05	7.2E-05	1.2E-04
Benzene, 1,4-dichloro	106-46-7	7	0	9.1E-07	2.0E-06	0	5.6E-06	5.6E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	2.7E-05	2.8E-05	1.8E-05	8.1E-05	1.0E-04
Ethylbenzene	100-41-4	13	0	8.1E-06	1.4E-05	0	3.3E-05	5.5E-05
Formaldehyde	50-00-0	34	3.8E-05	1.8E-04	1.4E-04	1.5E-04	3.0E-04	7.8E-04
Tetrachloroethylene	127-18-4	7	0	2.3E-06	5.4E-06	0	1.5E-05	2.0E-05

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-202. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators 2<6 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	1.6E-07	2.8E-07	0	6.3E-07	1.1E-06
Benz[a]anthracene	56-55-3	1	0	1.2E-10	7.2E-10	0	0	4.2E-09
Benzo[a]pyrene	50-32-8	28	0	2.8E-08	3.0E-08	1.5E-08	7.6E-08	1.2E-07
Benzo[b]fluoranthene	205-99-2	2	0	6.1E-10	2.5E-09	0	3.7E-09	1.0E-08
Benzo[k]fluoranthene	207-08-9	4	0	7.6E-10	2.3E-09	0	5.8E-09	8.6E-09
Chrysene	218-01-9	13	0	5.1E-09	7.9E-09	0	1.9E-08	3.3E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	1.8E-09	2.2E-09	6.7E-10	5.2E-09	1.0E-08
Dibenz[a,h]anthracene	53-70-3	14	0	3.6E-09	7.1E-09	0	1.8E-08	3.0E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	2.7E-09	8.7E-09	0	3.0E-08	3.0E-08
Methyl Isobutyl Ketone	108-10-1	5	0	3.9E-07	1.1E-06	0	3.0E-06	4.5E-06
Naphthalene	91-20-3	9	0	6.6E-07	1.4E-06	0	2.9E-06	6.5E-06
Styrene	100-42-5	17	0	1.5E-06	3.0E-06	0	4.9E-06	1.6E-05
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	6.5E-06	6.3E-05	4.6E-05	4.8E-05	1.2E-04	2.4E-04
Benzene	71-43-2	35	2.2E-06	1.5E-05	1.2E-05	1.1E-05	3.8E-05	6.1E-05
Benzene, 1,4-dichloro	106-46-7	7	0	4.8E-07	1.0E-06	0	3.0E-06	3.0E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	1.4E-05	1.5E-05	9.5E-06	4.3E-05	5.3E-05



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Ethylbenzene	100-41-4	13	0	4.3E-06	7.2E-06	0	1.7E-05	2.9E-05
Formaldehyde	50-00-0	34	2.0E-05	9.5E-05	7.2E-05	8.0E-05	1.6E-04	4.1E-04
Tetrachloroethylene	127-18-4	7	0	1.2E-06	2.8E-06	0	8.0E-06	1.0E-05

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-203. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators 6<11 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	1.5E-07	2.7E-07	0	6.0E-07	1.0E-06
Benz[a]anthracene	56-55-3	1	0	1.2E-10	6.8E-10	0	0	4.0E-09
Benzo[a]pyrene	50-32-8	28	0	2.7E-08	2.9E-08	1.4E-08	7.3E-08	1.1E-07
Benzo[b]fluoranthene	205-99-2	2	0	5.9E-10	2.4E-09	0	3.5E-09	1.0E-08
Benzo[k]fluoranthene	207-08-9	4	0	7.3E-10	2.1E-09	0	5.6E-09	8.3E-09
Chrysene	218-01-9	13	0	4.9E-09	7.6E-09	0	1.8E-08	3.1E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	1.7E-09	2.1E-09	6.4E-10	5.0E-09	9.8E-09
Dibenz[a,h]anthracene	53-70-3	14	0	3.4E-09	6.8E-09	0	1.7E-08	2.9E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	2.6E-09	8.3E-09	0	2.9E-08	2.9E-08
Methyl Isobutyl Ketone	108-10-1	5	0	3.7E-07	1.0E-06	0	2.9E-06	4.3E-06
Naphthalene	91-20-3	9	0	6.3E-07	1.3E-06	0	2.7E-06	6.2E-06
Styrene	100-42-5	17	0	1.4E-06	2.9E-06	0	4.7E-06	1.6E-05
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	6.2E-06	6.0E-05	4.4E-05	4.6E-05	1.1E-04	2.3E-04
Benzene	71-43-2	35	2.1E-06	1.4E-05	1.2E-05	1.0E-05	3.6E-05	5.8E-05
Benzene, 1,4-dichloro	106-46-7	7	0	4.6E-07	1.0E-06	0	2.8E-06	2.8E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	1.4E-05	1.4E-05	9.1E-06	4.1E-05	5.0E-05
Ethylbenzene	100-41-4	13	0	4.1E-06	6.9E-06	0	1.7E-05	2.7E-05
Formaldehyde	50-00-0	34	1.9E-05	9.1E-05	6.8E-05	7.6E-05	1.5E-04	3.9E-04
Tetrachloroethylene	127-18-4	7	0	1.1E-06	2.7E-06	0	7.6E-06	9.9E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-204. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators 11<16 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	6.6E-08	1.2E-07	0	2.6E-07	4.5E-07
Benz[a]anthracene	56-55-3	1	0	5.1E-11	2.9E-10	0	0	1.7E-09
Benzo[a]pyrene	50-32-8	28	0	1.2E-08	1.2E-08	6.3E-09	3.2E-08	4.8E-08
Benzo[b]fluoranthene	205-99-2	2	0	2.5E-10	1.0E-09	0	1.5E-09	4.3E-09
Benzo[k]fluoranthene	207-08-9	4	0	3.1E-10	9.3E-10	0	2.4E-09	3.6E-09
Chrysene	218-01-9	13	0	2.1E-09	3.3E-09	0	7.9E-09	1.4E-08
Cyclopenta[cd]pyrene	27208-37-3	28	0	7.4E-10	9.1E-10	2.8E-10	2.2E-09	4.2E-09
Dibenz[a,h]anthracene	53-70-3	14	0	1.5E-09	2.9E-09	0	7.3E-09	1.2E-08
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	1.1E-09	3.6E-09	0	1.3E-08	1.3E-08
Methyl Isobutyl Ketone	108-10-1	5	0	1.6E-07	4.4E-07	0	1.2E-06	1.9E-06
Naphthalene	91-20-3	9	0	2.7E-07	5.7E-07	0	1.2E-06	2.7E-06
Styrene	100-42-5	17	0	6.0E-07	1.2E-06	0	2.0E-06	6.7E-06
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	2.7E-06	2.6E-05	1.9E-05	2.0E-05	5.0E-05	9.8E-05
Benzene	71-43-2	35	9.3E-07	6.2E-06	5.0E-06	4.4E-06	1.6E-05	2.5E-05
Benzene, 1,4-dichloro	106-46-7	7	0	2.0E-07	4.3E-07	0	1.2E-06	1.2E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	5.9E-06	6.1E-06	3.9E-06	1.8E-05	2.2E-05
Ethylbenzene	100-41-4	13	0	1.8E-06	3.0E-06	0	7.2E-06	1.2E-05
Formaldehyde	50-00-0	34	8.3E-06	3.9E-05	3.0E-05	3.3E-05	6.6E-05	1.7E-04
Tetrachloroethylene	127-18-4	7	0	4.9E-07	1.2E-06	0	3.3E-06	4.3E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-205. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators 16<30 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	4.4E-08	7.7E-08	0	1.7E-07	3.0E-07
Benz[a]anthracene	56-55-3	1	0	3.3E-11	1.9E-10	0	0	1.1E-09
Benzo[a]pyrene	50-32-8	28	0	7.7E-09	8.1E-09	4.1E-09	2.1E-08	3.2E-08



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[b]fluoranthene	205-99-2	2	0	1.7E-10	6.8E-10	0	9.9E-10	2.8E-09
Benzo[k]fluoranthene	207-08-9	4	0	2.1E-10	6.1E-10	0	1.6E-09	2.4E-09
Chrysene	218-01-9	13	0	1.4E-09	2.2E-09	0	5.2E-09	9.0E-09
Cyclopenta[cd]pyrene	27208-37-3	28	0	4.9E-10	6.0E-10	1.8E-10	1.4E-09	2.8E-09
Dibenz[a,h]anthracene	53-70-3	14	0	9.7E-10	1.9E-09	0	4.8E-09	8.2E-09
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	7.3E-10	2.4E-09	0	8.3E-09	8.3E-09
Methyl Isobutyl Ketone	108-10-1	5	0	1.1E-07	2.9E-07	0	8.2E-07	1.2E-06
Naphthalene	91-20-3	9	0	1.8E-07	3.8E-07	0	7.8E-07	1.8E-06
Styrene	100-42-5	17	0	4.0E-07	8.2E-07	0	1.3E-06	4.5E-06
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	1.8E-06	1.7E-05	1.3E-05	1.3E-05	3.3E-05	6.5E-05
Benzene	71-43-2	35	6.1E-07	4.1E-06	3.3E-06	2.9E-06	1.0E-05	1.7E-05
Benzene, 1,4-dichloro	106-46-7	7	0	1.3E-07	2.9E-07	0	8.0E-07	8.0E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	3.9E-06	4.0E-06	2.6E-06	1.2E-05	1.4E-05
Ethylbenzene	100-41-4	13	0	1.2E-06	2.0E-06	0	4.7E-06	7.8E-06
Formaldehyde	50-00-0	34	5.5E-06	2.6E-05	1.9E-05	2.2E-05	4.3E-05	1.1E-04
Tetrachloroethylene	127-18-4	7	0	3.2E-07	7.7E-07	0	2.2E-06	2.8E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-206. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators 30<40 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	3.8E-08	6.7E-08	0	1.5E-07	2.6E-07
Benz[a]anthracene	56-55-3	1	0	2.9E-11	1.7E-10	0	0	9.9E-10
Benzo[a]pyrene	50-32-8	28	0	6.7E-09	7.1E-09	3.6E-09	1.8E-08	2.8E-08
Benzo[b]fluoranthene	205-99-2	2	0	1.5E-10	5.9E-10	0	8.7E-10	2.5E-09
Benzo[k]fluoranthene	207-08-9	4	0	1.8E-10	5.3E-10	0	1.4E-09	2.1E-09
Chrysene	218-01-9	13	0	1.2E-09	1.9E-09	0	4.5E-09	7.8E-09
Cyclopenta[cd]pyrene	27208-37-3	28	0	4.3E-10	5.2E-10	1.6E-10	1.2E-09	2.4E-09
Dibenz[a,h]anthracene	53-70-3	14	0	8.5E-10	1.7E-09	0	4.2E-09	7.1E-09
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	6.4E-10	2.1E-09	0	7.2E-09	7.2E-09
Methyl Isobutyl Ketone	108-10-1	5	0	9.3E-08	2.5E-07	0	7.2E-07	1.1E-06



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene	91-20-3	9	0	1.6E-07	3.3E-07	0	6.8E-07	1.5E-06
Styrene	100-42-5	17	0	3.5E-07	7.2E-07	0	1.2E-06	3.9E-06
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	1.5E-06	1.5E-05	1.1E-05	1.1E-05	2.9E-05	5.7E-05
Benzene	71-43-2	35	5.3E-07	3.6E-06	2.9E-06	2.5E-06	8.9E-06	1.5E-05
Benzene, 1,4-dichloro	106-46-7	7	0	1.1E-07	2.5E-07	0	7.0E-07	7.0E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	3.4E-06	3.5E-06	2.3E-06	1.0E-05	1.3E-05
Ethylbenzene	100-41-4	13	0	1.0E-06	1.7E-06	0	4.1E-06	6.8E-06
Formaldehyde	50-00-0	34	4.8E-06	2.3E-05	1.7E-05	1.9E-05	3.8E-05	9.7E-05
Tetrachloroethylene	127-18-4	7	0	2.8E-07	6.7E-07	0	1.9E-06	2.5E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-207. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators 40<50 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	3.9E-08	6.8E-08	0	1.5E-07	2.6E-07
Benz[a]anthracene	56-55-3	1	0	2.9E-11	1.7E-10	0	0	1.0E-09
Benzo[a]pyrene	50-32-8	28	0	6.8E-09	7.2E-09	3.6E-09	1.8E-08	2.8E-08
Benzo[b]fluoranthene	205-99-2	2	0	1.5E-10	6.0E-10	0	8.8E-10	2.5E-09
Benzo[k]fluoranthene	207-08-9	4	0	1.8E-10	5.4E-10	0	1.4E-09	2.1E-09
Chrysene	218-01-9	13	0	1.2E-09	1.9E-09	0	4.6E-09	7.9E-09
Cyclopenta[cd]pyrene	27208-37-3	28	0	4.3E-10	5.3E-10	1.6E-10	1.3E-09	2.5E-09
Dibenz[a,h]anthracene	53-70-3	14	0	8.6E-10	1.7E-09	0	4.3E-09	7.2E-09
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	6.4E-10	2.1E-09	0	7.3E-09	7.3E-09
Methyl Isobutyl Ketone	108-10-1	5	0	9.3E-08	2.5E-07	0	7.3E-07	1.1E-06
Naphthalene	91-20-3	9	0	1.6E-07	3.3E-07	0	6.9E-07	1.5E-06
Styrene	100-42-5	17	0	3.5E-07	7.3E-07	0	1.2E-06	3.9E-06
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	1.6E-06	1.5E-05	1.1E-05	1.1E-05	2.9E-05	5.7E-05
Benzene	71-43-2	35	5.4E-07	3.6E-06	2.9E-06	2.5E-06	9.0E-06	1.5E-05
Benzene, 1,4-dichloro	106-46-7	7	0	1.1E-07	2.5E-07	0	7.1E-07	7.1E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	3.4E-06	3.6E-06	2.3E-06	1.0E-05	1.3E-05



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Ethylbenzene	100-41-4	13	0	1.0E-06	1.7E-06	0	4.2E-06	6.9E-06
Formaldehyde	50-00-0	34	4.8E-06	2.3E-05	1.7E-05	1.9E-05	3.8E-05	9.8E-05
Tetrachloroethylene	127-18-4	7	0	2.9E-07	6.8E-07	0	1.9E-06	2.5E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-208. Field-Specific Average **On-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators 50<70 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	10	0	3.9E-08	6.8E-08	0	1.5E-07	2.6E-07
Benz[a]anthracene	56-55-3	1	0	3.0E-11	1.7E-10	0	0	1.0E-09
Benzo[a]pyrene	50-32-8	28	0	6.8E-09	7.2E-09	3.7E-09	1.8E-08	2.8E-08
Benzo[b]fluoranthene	205-99-2	2	0	1.5E-10	6.0E-10	0	8.8E-10	2.5E-09
Benzo[k]fluoranthene	207-08-9	4	0	1.8E-10	5.4E-10	0	1.4E-09	2.1E-09
Chrysene	218-01-9	13	0	1.2E-09	1.9E-09	0	4.6E-09	7.9E-09
Cyclopenta[cd]pyrene	27208-37-3	28	0	4.4E-10	5.3E-10	1.6E-10	1.3E-09	2.5E-09
Dibenz[a,h]anthracene	53-70-3	14	0	8.6E-10	1.7E-09	0	4.3E-09	7.2E-09
Indeno[1,2,3-cd]pyrene	193-39-5	3	0	6.4E-10	2.1E-09	0	7.3E-09	7.3E-09
Methyl Isobutyl Ketone	108-10-1	5	0	9.4E-08	2.6E-07	0	7.3E-07	1.1E-06
Naphthalene	91-20-3	9	0	1.6E-07	3.3E-07	0	6.9E-07	1.6E-06
Styrene	100-42-5	17	0	3.5E-07	7.3E-07	0	1.2E-06	3.9E-06
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	34	1.6E-06	1.5E-05	1.1E-05	1.1E-05	2.9E-05	5.7E-05
Benzene	71-43-2	35	5.4E-07	3.6E-06	2.9E-06	2.6E-06	9.1E-06	1.5E-05
Benzene, 1,4-dichloro	106-46-7	7	0	1.1E-07	2.5E-07	0	7.1E-07	7.1E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	34	0	3.4E-06	3.6E-06	2.3E-06	1.0E-05	1.3E-05
Ethylbenzene	100-41-4	13	0	1.0E-06	1.7E-06	0	4.2E-06	6.9E-06
Formaldehyde	50-00-0	34	4.9E-06	2.3E-05	1.7E-05	1.9E-05	3.8E-05	9.8E-05
Tetrachloroethylene	127-18-4	7	0	2.9E-07	6.8E-07	0	1.9E-06	2.5E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-209. Field-Specific Average **Off-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators Third Trimester Fetus<0 Years**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	8	0	4.2E-08	8.3E-08	0	1.5E-07	3.5E-07
Benz[a]anthracene	56-55-3	0	0	0	0	0	0	0
Benzo[a]pyrene	50-32-8	24	0	8.3E-09	1.3E-08	3.4E-09	2.9E-08	5.8E-08
Benzo[b]fluoranthene	205-99-2	3	0	2.2E-10	7.2E-10	0	2.5E-09	2.5E-09
Benzo[k]fluoranthene	207-08-9	3	0	1.9E-10	6.0E-10	0	2.0E-09	2.0E-09
Chrysene	218-01-9	7	0	6.2E-10	1.3E-09	0	2.5E-09	5.5E-09
Cyclopenta[cd]pyrene	27208-37-3	22	0	3.7E-10	4.8E-10	1.6E-10	1.3E-09	1.5E-09
Dibenz[a,h]anthracene	53-70-3	9	0	5.3E-10	1.2E-09	0	3.4E-09	4.3E-09
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	4.4E-10	1.7E-09	0	2.9E-09	7.2E-09
Methyl Isobutyl Ketone	108-10-1	1	0	2.1E-08	1.2E-07	0	0	7.2E-07
Naphthalene	91-20-3	9	0	1.7E-07	3.5E-07	0	7.7E-07	1.5E-06
Styrene	100-42-5	13	0	3.5E-07	7.5E-07	0	1.3E-06	3.9E-06
Non-Field-Related Chemicals								
Benzene	71-43-2	34	1.1E-06	3.7E-06	3.1E-06	2.3E-06	8.9E-06	1.6E-05
Benzene, 1,4-dichloro	106-46-7	6	0	1.0E-07	2.4E-07	0	7.0E-07	7.0E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	33	0	3.4E-06	3.7E-06	1.4E-06	1.1E-05	1.2E-05
Ethylbenzene	100-41-4	13	0	1.1E-06	1.8E-06	0	4.7E-06	7.7E-06
Tetrachloroethylene	127-18-4	7	0	2.9E-07	6.7E-07	0	1.9E-06	2.5E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b Off-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air off the fields. Carbonyl samples were not collected at off-field locations.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-210. Field-Specific Average **Off-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators 0<2 Year**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	8	0	3.4E-07	6.7E-07	0	1.2E-06	2.8E-06
Benz[a]anthracene	56-55-3	0	0	0	0	0	0	0
Benzo[a]pyrene	50-32-8	24	0	6.6E-08	1.0E-07	2.8E-08	2.4E-07	4.7E-07
Benzo[b]fluoranthene	205-99-2	3	0	1.8E-09	5.8E-09	0	2.0E-08	2.0E-08
Benzo[k]fluoranthene	207-08-9	3	0	1.5E-09	4.8E-09	0	1.6E-08	1.6E-08



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Chrysene	218-01-9	7	0	5.0E-09	1.1E-08	0	2.0E-08	4.4E-08
Cyclopenta[cd]pyrene	27208-37-3	22	0	3.0E-09	3.9E-09	1.3E-09	1.1E-08	1.2E-08
Dibenz[a,h]anthracene	53-70-3	9	0	4.3E-09	9.5E-09	0	2.7E-08	3.5E-08
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	3.5E-09	1.4E-08	0	2.3E-08	5.8E-08
Methyl Isobutyl Ketone	108-10-1	1	0	1.7E-07	9.9E-07	0	0	5.8E-06
Naphthalene	91-20-3	9	0	1.4E-06	2.8E-06	0	6.2E-06	1.2E-05
Styrene	100-42-5	13	0	2.8E-06	6.1E-06	0	1.1E-05	3.2E-05
Non-Field-Related Chemicals								
Benzene	71-43-2	34	8.6E-06	3.0E-05	2.5E-05	1.9E-05	7.2E-05	1.3E-04
Benzene, 1,4-dichloro	106-46-7	6	0	8.3E-07	1.9E-06	0	5.6E-06	5.6E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	33	0	2.7E-05	3.0E-05	1.1E-05	9.1E-05	9.4E-05
Ethylbenzene	100-41-4	13	0	8.6E-06	1.5E-05	0	3.8E-05	6.2E-05
Tetrachloroethylene	127-18-4	7	0	2.3E-06	5.4E-06	0	1.6E-05	2.0E-05

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b Off-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air off the fields. Carbonyl samples were not collected at off-field locations.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-211. Field-Specific Average **Off-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators 2<6 Year**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	8	0	1.8E-07	3.5E-07	0	6.3E-07	1.5E-06
Benz[a]anthracene	56-55-3	0	0	0	0	0	0	0
Benzo[a]pyrene	50-32-8	24	0	3.5E-08	5.4E-08	1.4E-08	1.2E-07	2.5E-07
Benzo[b]fluoranthene	205-99-2	3	0	9.5E-10	3.0E-09	0	1.0E-08	1.0E-08
Benzo[k]fluoranthene	207-08-9	3	0	7.9E-10	2.5E-09	0	8.6E-09	8.6E-09
Chrysene	218-01-9	7	0	2.6E-09	5.5E-09	0	1.1E-08	2.3E-08
Cyclopenta[cd]pyrene	27208-37-3	22	0	1.6E-09	2.0E-09	6.7E-10	5.6E-09	6.5E-09
Dibenz[a,h]anthracene	53-70-3	9	0	2.2E-09	5.0E-09	0	1.4E-08	1.8E-08
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	1.8E-09	7.4E-09	0	1.2E-08	3.0E-08
Methyl Isobutyl Ketone	108-10-1	1	0	8.9E-08	5.2E-07	0	0	3.0E-06
Naphthalene	91-20-3	9	0	7.3E-07	1.5E-06	0	3.2E-06	6.4E-06
Styrene	100-42-5	13	0	1.5E-06	3.2E-06	0	5.6E-06	1.7E-05
Non-Field-Related Chemicals								
Benzene	71-43-2	34	4.5E-06	1.6E-05	1.3E-05	9.9E-06	3.8E-05	6.8E-05



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene, 1,4-dichloro	106-46-7	6	0	4.3E-07	1.0E-06	0	3.0E-06	3.0E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	33	0	1.4E-05	1.6E-05	5.8E-06	4.8E-05	4.9E-05
Ethylbenzene	100-41-4	13	0	4.5E-06	7.7E-06	0	2.0E-05	3.2E-05
Tetrachloroethylene	127-18-4	7	0	1.2E-06	2.8E-06	0	8.2E-06	1.0E-05

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b Off-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air off the fields. Carbonyl samples were not collected at off-field locations.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-212. Field-Specific Average **Off-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators 6<11 Year**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	8	0	1.7E-07	3.3E-07	0	6.0E-07	1.4E-06
Benz[a]anthracene	56-55-3	0	0	0	0	0	0	0
Benzo[a]pyrene	50-32-8	24	0	3.3E-08	5.2E-08	1.4E-08	1.2E-07	2.4E-07
Benzo[b]fluoranthene	205-99-2	3	0	9.0E-10	2.9E-09	0	1.0E-08	1.0E-08
Benzo[k]fluoranthene	207-08-9	3	0	7.5E-10	2.4E-09	0	8.3E-09	8.3E-09
Chrysene	218-01-9	7	0	2.5E-09	5.3E-09	0	1.0E-08	2.2E-08
Cyclopenta[cd]pyrene	27208-37-3	22	0	1.5E-09	1.9E-09	6.4E-10	5.4E-09	6.2E-09
Dibenz[a,h]anthracene	53-70-3	9	0	2.1E-09	4.8E-09	0	1.4E-08	1.7E-08
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	1.8E-09	7.0E-09	0	1.2E-08	2.9E-08
Methyl Isobutyl Ketone	108-10-1	1	0	8.5E-08	5.0E-07	0	0	2.9E-06
Naphthalene	91-20-3	9	0	6.9E-07	1.4E-06	0	3.1E-06	6.1E-06
Styrene	100-42-5	13	0	1.4E-06	3.0E-06	0	5.3E-06	1.6E-05
Non-Field-Related Chemicals								
Benzene	71-43-2	34	4.3E-06	1.5E-05	1.3E-05	9.4E-06	3.6E-05	6.5E-05
Benzene, 1,4-dichloro	106-46-7	6	0	4.1E-07	9.6E-07	0	2.8E-06	2.8E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	33	0	1.4E-05	1.5E-05	5.5E-06	4.5E-05	4.7E-05
Ethylbenzene	100-41-4	13	0	4.3E-06	7.4E-06	0	1.9E-05	3.1E-05
Tetrachloroethylene	127-18-4	7	0	1.2E-06	2.7E-06	0	7.8E-06	9.9E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b Off-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air off the fields. Carbonyl samples were not collected at off-field locations.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-213. Field-Specific Average **Off-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators 11<16 Year**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	8	0	7.3E-08	1.4E-07	0	2.6E-07	6.1E-07
Benz[a]anthracene	56-55-3	0	0	0	0	0	0	0
Benzo[a]pyrene	50-32-8	24	0	1.4E-08	2.2E-08	6.0E-09	5.1E-08	1.0E-07
Benzo[b]fluoranthene	205-99-2	3	0	3.9E-10	1.3E-09	0	4.3E-09	4.3E-09
Benzo[k]fluoranthene	207-08-9	3	0	3.2E-10	1.0E-09	0	3.6E-09	3.6E-09
Chrysene	218-01-9	7	0	1.1E-09	2.3E-09	0	4.3E-09	9.5E-09
Cyclopenta[cd]pyrene	27208-37-3	22	0	6.5E-10	8.4E-10	2.8E-10	2.3E-09	2.7E-09
Dibenz[a,h]anthracene	53-70-3	9	0	9.2E-10	2.1E-09	0	5.9E-09	7.5E-09
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	7.6E-10	3.0E-09	0	5.0E-09	1.3E-08
Methyl Isobutyl Ketone	108-10-1	1	0	3.7E-08	2.1E-07	0	0	1.2E-06
Naphthalene	91-20-3	9	0	3.0E-07	6.0E-07	0	1.3E-06	2.6E-06
Styrene	100-42-5	13	0	6.2E-07	1.3E-06	0	2.3E-06	6.9E-06
Non-Field-Related Chemicals								
Benzene	71-43-2	34	1.9E-06	6.5E-06	5.5E-06	4.1E-06	1.6E-05	2.8E-05
Benzene, 1,4-dichloro	106-46-7	6	0	1.8E-07	4.1E-07	0	1.2E-06	1.2E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	33	0	5.9E-06	6.5E-06	2.4E-06	2.0E-05	2.0E-05
Ethylbenzene	100-41-4	13	0	1.9E-06	3.2E-06	0	8.2E-06	1.3E-05
Tetrachloroethylene	127-18-4	7	0	5.0E-07	1.2E-06	0	3.4E-06	4.3E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b Off-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air off the fields. Carbonyl samples were not collected at off-field locations.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-214. Field-Specific Average **Off-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators 16<30 Year**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	8	0	4.8E-08	9.5E-08	0	1.7E-07	4.0E-07
Benz[a]anthracene	56-55-3	0	0	0	0	0	0	0
Benzo[a]pyrene	50-32-8	24	0	9.5E-09	1.5E-08	3.9E-09	3.4E-08	6.7E-08
Benzo[b]fluoranthene	205-99-2	3	0	2.6E-10	8.3E-10	0	2.8E-09	2.8E-09
Benzo[k]fluoranthene	207-08-9	3	0	2.1E-10	6.9E-10	0	2.4E-09	2.4E-09



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Chrysene	218-01-9	7	0	7.1E-10	1.5E-09	0	2.9E-09	6.3E-09
Cyclopenta[cd]pyrene	27208-37-3	22	0	4.3E-10	5.5E-10	1.8E-10	1.5E-09	1.8E-09
Dibenz[a,h]anthracene	53-70-3	9	0	6.1E-10	1.4E-09	0	3.9E-09	5.0E-09
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	5.0E-10	2.0E-09	0	3.3E-09	8.3E-09
Methyl Isobutyl Ketone	108-10-1	1	0	2.4E-08	1.4E-07	0	0	8.2E-07
Naphthalene	91-20-3	9	0	2.0E-07	4.0E-07	0	8.8E-07	1.7E-06
Styrene	100-42-5	13	0	4.1E-07	8.7E-07	0	1.5E-06	4.5E-06
Non-Field-Related Chemicals								
Benzene	71-43-2	34	1.2E-06	4.3E-06	3.6E-06	2.7E-06	1.0E-05	1.8E-05
Benzene, 1,4-dichloro	106-46-7	6	0	1.2E-07	2.7E-07	0	8.0E-07	8.0E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	33	0	3.9E-06	4.3E-06	1.6E-06	1.3E-05	1.3E-05
Ethylbenzene	100-41-4	13	0	1.2E-06	2.1E-06	0	5.4E-06	8.8E-06
Tetrachloroethylene	127-18-4	7	0	3.3E-07	7.7E-07	0	2.2E-06	2.8E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b Off-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air off the fields. Carbonyl samples were not collected at off-field locations.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-215. Field-Specific Average **Off-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators 30<40 Year**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	8	0	4.2E-08	8.3E-08	0	1.5E-07	3.5E-07
Benz[a]anthracene	56-55-3	0	0	0	0	0	0	0
Benzo[a]pyrene	50-32-8	24	0	8.3E-09	1.3E-08	3.4E-09	3.0E-08	5.9E-08
Benzo[b]fluoranthene	205-99-2	3	0	2.3E-10	7.2E-10	0	2.5E-09	2.5E-09
Benzo[k]fluoranthene	207-08-9	3	0	1.9E-10	6.0E-10	0	2.1E-09	2.1E-09
Chrysene	218-01-9	7	0	6.2E-10	1.3E-09	0	2.5E-09	5.5E-09
Cyclopenta[cd]pyrene	27208-37-3	22	0	3.8E-10	4.8E-10	1.6E-10	1.3E-09	1.5E-09
Dibenz[a,h]anthracene	53-70-3	9	0	5.3E-10	1.2E-09	0	3.4E-09	4.3E-09
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	4.4E-10	1.7E-09	0	2.9E-09	7.2E-09
Methyl Isobutyl Ketone	108-10-1	1	0	2.1E-08	1.2E-07	0	0	7.2E-07
Naphthalene	91-20-3	9	0	1.7E-07	3.5E-07	0	7.7E-07	1.5E-06
Styrene	100-42-5	13	0	3.6E-07	7.6E-07	0	1.3E-06	4.0E-06
Non-Field-Related Chemicals								
Benzene	71-43-2	34	1.1E-06	3.8E-06	3.1E-06	2.3E-06	9.0E-06	1.6E-05



Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene, 1,4-dichloro	106-46-7	6	0	1.0E-07	2.4E-07	0	7.0E-07	7.0E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	33	0	3.4E-06	3.7E-06	1.4E-06	1.1E-05	1.2E-05
Ethylbenzene	100-41-4	13	0	1.1E-06	1.8E-06	0	4.7E-06	7.7E-06
Tetrachloroethylene	127-18-4	7	0	2.9E-07	6.7E-07	0	1.9E-06	2.5E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b Off-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air off the fields. Carbonyl samples were not collected at off-field locations.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-216. Field-Specific Average **Off-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators 40<50 Year**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	8	0	4.3E-08	8.4E-08	0	1.5E-07	3.6E-07
Benz[a]anthracene	56-55-3	0	0	0	0	0	0	0
Benzo[a]pyrene	50-32-8	24	0	8.4E-09	1.3E-08	3.5E-09	3.0E-08	5.9E-08
Benzo[b]fluoranthene	205-99-2	3	0	2.3E-10	7.3E-10	0	2.5E-09	2.5E-09
Benzo[k]fluoranthene	207-08-9	3	0	1.9E-10	6.1E-10	0	2.1E-09	2.1E-09
Chrysene	218-01-9	7	0	6.3E-10	1.3E-09	0	2.5E-09	5.6E-09
Cyclopenta[cd]pyrene	27208-37-3	22	0	3.8E-10	4.9E-10	1.6E-10	1.3E-09	1.6E-09
Dibenz[a,h]anthracene	53-70-3	9	0	5.4E-10	1.2E-09	0	3.4E-09	4.4E-09
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	4.4E-10	1.8E-09	0	2.9E-09	7.3E-09
Methyl Isobutyl Ketone	108-10-1	1	0	2.1E-08	1.2E-07	0	0	7.3E-07
Naphthalene	91-20-3	9	0	1.7E-07	3.5E-07	0	7.8E-07	1.5E-06
Styrene	100-42-5	13	0	3.6E-07	7.6E-07	0	1.3E-06	4.0E-06
Non-Field-Related Chemicals								
Benzene	71-43-2	34	1.1E-06	3.8E-06	3.2E-06	2.4E-06	9.1E-06	1.6E-05
Benzene, 1,4-dichloro	106-46-7	6	0	1.0E-07	2.4E-07	0	7.1E-07	7.1E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	33	0	3.5E-06	3.8E-06	1.4E-06	1.1E-05	1.2E-05
Ethylbenzene	100-41-4	13	0	1.1E-06	1.9E-06	0	4.8E-06	7.8E-06
Tetrachloroethylene	127-18-4	7	0	2.9E-07	6.8E-07	0	2.0E-06	2.5E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b Off-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air off the fields. Carbonyl samples were not collected at off-field locations.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-217. Field-Specific Average **Off-Field** Inhalation Daily Dose^a (ADD_{inh-field}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of Chemicals—Combined Gender **Spectators 50<70 Year**

Chemical	CASRN	Detection	ADD _{inh-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals								
Aniline	62-53-3	8	0	4.3E-08	8.4E-08	0	1.5E-07	3.6E-07
Benz[a]anthracene	56-55-3	0	0	0	0	0	0	0
Benzo[a]pyrene	50-32-8	24	0	8.4E-09	1.3E-08	3.5E-09	3.0E-08	5.9E-08
Benzo[b]fluoranthene	205-99-2	3	0	2.3E-10	7.3E-10	0	2.5E-09	2.5E-09
Benzo[k]fluoranthene	207-08-9	3	0	1.9E-10	6.1E-10	0	2.1E-09	2.1E-09
Chrysene	218-01-9	7	0	6.3E-10	1.3E-09	0	2.5E-09	5.6E-09
Cyclopenta[cd]pyrene	27208-37-3	22	0	3.8E-10	4.9E-10	1.6E-10	1.3E-09	1.6E-09
Dibenz[a,h]anthracene	53-70-3	9	0	5.4E-10	1.2E-09	0	3.5E-09	4.4E-09
Indeno[1,2,3-cd]pyrene	193-39-5	2	0	4.4E-10	1.8E-09	0	2.9E-09	7.3E-09
Methyl Isobutyl Ketone	108-10-1	1	0	2.1E-08	1.3E-07	0	0	7.3E-07
Naphthalene	91-20-3	9	0	1.7E-07	3.5E-07	0	7.8E-07	1.5E-06
Styrene	100-42-5	13	0	3.6E-07	7.7E-07	0	1.3E-06	4.0E-06
Non-Field-Related Chemicals								
Benzene	71-43-2	34	1.1E-06	3.8E-06	3.2E-06	2.4E-06	9.1E-06	1.6E-05
Benzene, 1,4-dichloro	106-46-7	6	0	1.0E-07	2.4E-07	0	7.1E-07	7.1E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	33	0	3.5E-06	3.8E-06	1.4E-06	1.1E-05	1.2E-05
Ethylbenzene	100-41-4	13	0	1.1E-06	1.9E-06	0	4.8E-06	7.8E-06
Tetrachloroethylene	127-18-4	7	0	2.9E-07	6.8E-07	0	2.0E-06	2.5E-06

^a 35 field-specific ADD_{inh-field} values are included in the table.

^b Off-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air off the fields. Carbonyl samples were not collected at off-field locations.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

F.5.2. Average Dermal One-Day Dose (AD_{der-DART-field}) for One-Day Non-Cancer Hazard Assessment of DARTs

INDIVIDUAL FIELD ASSESSMENT (Table F-218 to Table F-241)

Table F-218. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Athletes 2<6 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	9.1E-10	4.1E-09	0	2.8E-09	2.2E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	1.3E-08	5.5E-08	0	5.0E-08	3.2E-07



Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	32	0	7.0E-10	6.9E-10	4.6E-10	1.8E-09	2.8E-09
Benzo[e]pyrene	192-97-2	35	2.0E-10	2.1E-09	1.5E-09	2.0E-09	5.0E-09	5.6E-09
Benzo[g,h,i]perylene	191-24-2	35	1.9E-10	1.6E-09	1.0E-09	1.4E-09	3.4E-09	4.1E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	2.0E-09	1.2E-08	0	0	6.9E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	6.9E-10	9.2E-09	1.5E-08	4.8E-09	2.1E-08	8.8E-08
Chrysene	218-01-9	35	6.5E-10	6.0E-09	3.8E-09	6.0E-09	1.2E-08	1.4E-08
Coronene	191-07-1	1	0	4.7E-10	2.8E-09	0	0	1.7E-08
Cyclopenta[cd]pyrene	27208-37-3	35	1.1E-10	1.1E-09	6.4E-10	1.0E-09	2.0E-09	2.4E-09
Dicyclohexylamine	101-83-7	30	0	1.8E-07	3.0E-07	1.4E-07	3.3E-07	1.8E-06
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	3.2E-10	6.4E-10	1.5E-10	1.3E-09	3.3E-09
Dimethyl phthalate	131-11-3	3	0	9.4E-10	3.1E-09	0	1.0E-08	1.2E-08
1,3-Diphenylguanidine	102-06-7	2	0	2.7E-09	1.2E-08	0	8.5E-09	6.7E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	6.8E-10	5.7E-10	6.1E-10	1.6E-09	2.5E-09
Methyl stearate	112-61-8	28	0	3.9E-09	5.6E-09	2.2E-09	1.0E-08	3.0E-08
4-tert-Octylphenol	140-66-9	34	0	1.7E-08	3.6E-08	4.7E-09	8.4E-08	1.6E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	4.3E-10	5.1E-09	4.1E-09	3.7E-09	1.3E-08	2.0E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	5.0E-10	5.5E-09	6.5E-09	3.1E-09	2.1E-08	2.7E-08

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-219. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Athletes 6<11 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	8.3E-10	3.7E-09	0	2.6E-09	2.0E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	1.2E-08	5.0E-08	0	4.6E-08	2.9E-07
Benzo[a]pyrene	50-32-8	32	0	6.3E-10	6.2E-10	4.1E-10	1.6E-09	2.6E-09
Benzo[e]pyrene	192-97-2	35	1.8E-10	1.9E-09	1.4E-09	1.8E-09	4.5E-09	5.1E-09
Benzo[g,h,i]perylene	191-24-2	35	1.8E-10	1.4E-09	9.2E-10	1.3E-09	3.1E-09	3.7E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	1.8E-09	1.1E-08	0	0	6.2E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	6.2E-10	8.4E-09	1.4E-08	4.3E-09	1.9E-08	8.0E-08
Chrysene	218-01-9	35	5.9E-10	5.4E-09	3.5E-09	5.4E-09	1.1E-08	1.2E-08
Coronene	191-07-1	1	0	4.3E-10	2.5E-09	0	0	1.5E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Cyclopenta[cd]pyrene	27208-37-3	35	9.9E-11	9.7E-10	5.8E-10	9.3E-10	1.9E-09	2.2E-09
Dicyclohexylamine	101-83-7	30	0	1.6E-07	2.7E-07	1.3E-07	3.0E-07	1.6E-06
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	2.9E-10	5.8E-10	1.3E-10	1.2E-09	3.0E-09
Dimethyl phthalate	131-11-3	3	0	8.5E-10	2.8E-09	0	9.3E-09	1.1E-08
1,3-Diphenylguanidine	102-06-7	2	0	2.5E-09	1.1E-08	0	7.7E-09	6.1E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	6.2E-10	5.2E-10	5.5E-10	1.4E-09	2.3E-09
Methyl stearate	112-61-8	28	0	3.6E-09	5.0E-09	2.0E-09	9.4E-09	2.8E-08
4-tert-Octylphenol	140-66-9	34	0	1.6E-08	3.3E-08	4.2E-09	7.6E-08	1.5E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	3.9E-10	4.6E-09	3.8E-09	3.3E-09	1.2E-08	1.8E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	4.6E-10	5.0E-09	5.9E-09	2.9E-09	1.9E-08	2.5E-08

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-220. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Athletes 11<16 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	7.1E-10	3.2E-09	0	2.2E-09	1.7E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	1.0E-08	4.3E-08	0	3.9E-08	2.5E-07
Benzo[a]pyrene	50-32-8	32	0	5.4E-10	5.3E-10	3.5E-10	1.4E-09	2.2E-09
Benzo[e]pyrene	192-97-2	35	1.5E-10	1.6E-09	1.2E-09	1.5E-09	3.9E-09	4.4E-09
Benzo[g,h,i]perylene	191-24-2	35	1.5E-10	1.2E-09	7.9E-10	1.1E-09	2.7E-09	3.2E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	1.5E-09	9.0E-09	0	0	5.3E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	5.3E-10	7.1E-09	1.2E-08	3.7E-09	1.7E-08	6.8E-08
Chrysene	218-01-9	35	5.1E-10	4.6E-09	3.0E-09	4.6E-09	9.3E-09	1.1E-08
Coronene	191-07-1	1	0	3.7E-10	2.2E-09	0	0	1.3E-08
Cyclopenta[cd]pyrene	27208-37-3	35	8.5E-11	8.3E-10	4.9E-10	8.0E-10	1.6E-09	1.9E-09
Dicyclohexylamine	101-83-7	30	0	1.4E-07	2.3E-07	1.1E-07	2.6E-07	1.4E-06
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	2.5E-10	4.9E-10	1.1E-10	1.0E-09	2.6E-09
Dimethyl phthalate	131-11-3	3	0	7.3E-10	2.4E-09	0	7.9E-09	9.3E-09
1,3-Diphenylguanidine	102-06-7	2	0	2.1E-09	9.4E-09	0	6.6E-09	5.2E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	5.3E-10	4.4E-10	4.7E-10	1.2E-09	1.9E-09
Methyl stearate	112-61-8	28	0	3.1E-09	4.3E-09	1.7E-09	8.1E-09	2.4E-08



Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
4-tert-Octylphenol	140-66-9	34	0	1.3E-08	2.8E-08	3.6E-09	6.5E-08	1.2E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	3.3E-10	3.9E-09	3.2E-09	2.8E-09	1.0E-08	1.5E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	3.9E-10	4.3E-09	5.1E-09	2.4E-09	1.6E-08	2.1E-08

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-221. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Athletes 16<30 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	6.2E-10	2.8E-09	0	1.9E-09	1.5E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	9.0E-09	3.8E-08	0	3.4E-08	2.2E-07
Benzo[a]pyrene	50-32-8	32	0	4.7E-10	4.7E-10	3.1E-10	1.2E-09	1.9E-09
Benzo[e]pyrene	192-97-2	35	1.3E-10	1.4E-09	1.0E-09	1.3E-09	3.4E-09	3.8E-09
Benzo[g,h,i]perylene	191-24-2	35	1.3E-10	1.1E-09	6.9E-10	9.8E-10	2.3E-09	2.8E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	1.3E-09	7.9E-09	0	0	4.7E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	4.7E-10	6.3E-09	1.0E-08	3.2E-09	1.4E-08	6.0E-08
Chrysene	218-01-9	35	4.4E-10	4.1E-09	2.6E-09	4.1E-09	8.2E-09	9.2E-09
Coronene	191-07-1	1	0	3.2E-10	1.9E-09	0	0	1.1E-08
Cyclopenta[cd]pyrene	27208-37-3	35	7.4E-11	7.3E-10	4.3E-10	7.0E-10	1.4E-09	1.7E-09
Dicyclohexylamine	101-83-7	30	0	1.2E-07	2.0E-07	9.4E-08	2.3E-07	1.2E-06
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	2.1E-10	4.3E-10	9.9E-11	8.9E-10	2.2E-09
Dimethyl phthalate	131-11-3	3	0	6.4E-10	2.1E-09	0	6.9E-09	8.2E-09
1,3-Diphenylguanidine	102-06-7	2	0	1.8E-09	8.3E-09	0	5.7E-09	4.6E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	4.7E-10	3.9E-10	4.1E-10	1.1E-09	1.7E-09
Methyl stearate	112-61-8	28	0	2.7E-09	3.8E-09	1.5E-09	7.1E-09	2.1E-08
4-tert-Octylphenol	140-66-9	34	0	1.2E-08	2.5E-08	3.2E-09	5.7E-08	1.1E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	2.9E-10	3.5E-09	2.8E-09	2.5E-09	8.8E-09	1.3E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	3.4E-10	3.8E-09	4.4E-09	2.1E-09	1.4E-08	1.9E-08

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number



Values are rounded to two significant figures.

Table F-222. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Athletes 30<40 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	5.8E-10	2.6E-09	0	1.8E-09	1.4E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	8.4E-09	3.5E-08	0	3.2E-08	2.1E-07
Benzo[a]pyrene	50-32-8	32	0	4.4E-10	4.4E-10	2.9E-10	1.2E-09	1.8E-09
Benzo[e]pyrene	192-97-2	35	1.3E-10	1.3E-09	9.7E-10	1.2E-09	3.2E-09	3.6E-09
Benzo[g,h,i]perylene	191-24-2	35	1.2E-10	9.9E-10	6.4E-10	9.2E-10	2.2E-09	2.6E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	1.2E-09	7.4E-09	0	0	4.4E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	4.4E-10	5.9E-09	9.5E-09	3.0E-09	1.4E-08	5.6E-08
Chrysene	218-01-9	35	4.1E-10	3.8E-09	2.4E-09	3.8E-09	7.6E-09	8.6E-09
Coronene	191-07-1	1	0	3.0E-10	1.8E-09	0	0	1.1E-08
Cyclopenta[cd]pyrene	27208-37-3	35	6.9E-11	6.8E-10	4.1E-10	6.6E-10	1.3E-09	1.6E-09
Dicyclohexylamine	101-83-7	30	0	1.1E-07	1.9E-07	8.8E-08	2.1E-07	1.1E-06
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	2.0E-10	4.1E-10	9.3E-11	8.4E-10	2.1E-09
Dimethyl phthalate	131-11-3	3	0	6.0E-10	2.0E-09	0	6.5E-09	7.7E-09
1,3-Diphenylguanidine	102-06-7	2	0	1.7E-09	7.7E-09	0	5.4E-09	4.3E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	4.4E-10	3.6E-10	3.9E-10	1.0E-09	1.6E-09
Methyl stearate	112-61-8	28	0	2.5E-09	3.5E-09	1.4E-09	6.6E-09	1.9E-08
4-tert-Octylphenol	140-66-9	34	0	1.1E-08	2.3E-08	3.0E-09	5.3E-08	1.0E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	2.7E-10	3.2E-09	2.6E-09	2.3E-09	8.2E-09	1.2E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	3.2E-10	3.5E-09	4.2E-09	2.0E-09	1.3E-08	1.8E-08

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-223. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Athletes 40<50 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	5.8E-10	2.6E-09	0	1.8E-09	1.4E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	8.5E-09	3.5E-08	0	3.2E-08	2.1E-07



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	32	0	4.5E-10	4.4E-10	2.9E-10	1.2E-09	1.8E-09
Benzo[e]pyrene	192-97-2	35	1.3E-10	1.3E-09	9.7E-10	1.2E-09	3.2E-09	3.6E-09
Benzo[g,h,i]perylene	191-24-2	35	1.2E-10	1.0E-09	6.5E-10	9.2E-10	2.2E-09	2.6E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	1.3E-09	7.4E-09	0	0	4.4E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	4.4E-10	5.9E-09	9.5E-09	3.0E-09	1.4E-08	5.6E-08
Chrysene	218-01-9	35	4.2E-10	3.8E-09	2.5E-09	3.8E-09	7.7E-09	8.7E-09
Coronene	191-07-1	1	0	3.0E-10	1.8E-09	0	0	1.1E-08
Cyclopenta[cd]pyrene	27208-37-3	35	7.0E-11	6.8E-10	4.1E-10	6.6E-10	1.3E-09	1.6E-09
Dicyclohexylamine	101-83-7	30	0	1.2E-07	1.9E-07	8.8E-08	2.1E-07	1.2E-06
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	2.0E-10	4.1E-10	9.3E-11	8.4E-10	2.1E-09
Dimethyl phthalate	131-11-3	3	0	6.0E-10	2.0E-09	0	6.5E-09	7.7E-09
1,3-Diphenylguanidine	102-06-7	2	0	1.7E-09	7.8E-09	0	5.4E-09	4.3E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	4.4E-10	3.7E-10	3.9E-10	1.0E-09	1.6E-09
Methyl stearate	112-61-8	28	0	2.5E-09	3.6E-09	1.4E-09	6.7E-09	1.9E-08
4-tert-Octylphenol	140-66-9	34	0	1.1E-08	2.3E-08	3.0E-09	5.4E-08	1.0E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	2.8E-10	3.3E-09	2.6E-09	2.3E-09	8.3E-09	1.2E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	3.2E-10	3.5E-09	4.2E-09	2.0E-09	1.3E-08	1.8E-08

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-224. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Athletes 50<70 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	5.8E-10	2.6E-09	0	1.8E-09	1.4E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	8.5E-09	3.6E-08	0	3.2E-08	2.1E-07
Benzo[a]pyrene	50-32-8	32	0	4.5E-10	4.4E-10	2.9E-10	1.2E-09	1.8E-09
Benzo[e]pyrene	192-97-2	35	1.3E-10	1.3E-09	9.7E-10	1.3E-09	3.2E-09	3.6E-09
Benzo[g,h,i]perylene	191-24-2	35	1.2E-10	1.0E-09	6.5E-10	9.3E-10	2.2E-09	2.6E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	1.3E-09	7.5E-09	0	0	4.4E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	4.4E-10	5.9E-09	9.6E-09	3.1E-09	1.4E-08	5.6E-08
Chrysene	218-01-9	35	4.2E-10	3.8E-09	2.5E-09	3.8E-09	7.7E-09	8.7E-09
Coronene	191-07-1	1	0	3.0E-10	1.8E-09	0	0	1.1E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Cyclopenta[cd]pyrene	27208-37-3	35	7.0E-11	6.8E-10	4.1E-10	6.6E-10	1.3E-09	1.6E-09
Dicyclohexylamine	101-83-7	30	0	1.2E-07	1.9E-07	8.8E-08	2.1E-07	1.2E-06
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	2.0E-10	4.1E-10	9.4E-11	8.4E-10	2.1E-09
Dimethyl phthalate	131-11-3	3	0	6.0E-10	2.0E-09	0	6.5E-09	7.7E-09
1,3-Diphenylguanidine	102-06-7	2	0	1.7E-09	7.8E-09	0	5.4E-09	4.3E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	4.4E-10	3.7E-10	3.9E-10	1.0E-09	1.6E-09
Methyl stearate	112-61-8	28	0	2.5E-09	3.6E-09	1.4E-09	6.7E-09	2.0E-08
4-tert-Octylphenol	140-66-9	34	0	1.1E-08	2.3E-08	3.0E-09	5.4E-08	1.0E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	2.8E-10	3.3E-09	2.7E-09	2.4E-09	8.3E-09	1.3E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	3.2E-10	3.6E-09	4.2E-09	2.0E-09	1.3E-08	1.8E-08

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-225. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Coaches 16<30 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	3.1E-10	1.4E-09	0	9.5E-10	7.5E-09
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	4.4E-09	1.9E-08	0	1.7E-08	1.1E-07
Benzo[a]pyrene	50-32-8	32	0	2.3E-10	2.3E-10	1.5E-10	6.1E-10	9.5E-10
Benzo[e]pyrene	192-97-2	35	6.7E-11	6.9E-10	5.1E-10	6.6E-10	1.7E-09	1.9E-09
Benzo[g,h,i]perylene	191-24-2	35	6.5E-11	5.2E-10	3.4E-10	4.8E-10	1.2E-09	1.4E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	6.6E-10	3.9E-09	0	0	2.3E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	2.3E-10	3.1E-09	5.0E-09	1.6E-09	7.2E-09	3.0E-08
Chrysene	218-01-9	35	2.2E-10	2.0E-09	1.3E-09	2.0E-09	4.0E-09	4.6E-09
Coronene	191-07-1	1	0	1.6E-10	9.4E-10	0	0	5.6E-09
Cyclopenta[cd]pyrene	27208-37-3	35	3.7E-11	3.6E-10	2.1E-10	3.5E-10	6.9E-10	8.2E-10
Dicyclohexylamine	101-83-7	30	0	6.1E-08	1.0E-07	4.6E-08	1.1E-07	6.1E-07
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	1.1E-10	2.1E-10	4.9E-11	4.4E-10	1.1E-09
Dimethyl phthalate	131-11-3	3	0	3.2E-10	1.0E-09	0	3.4E-09	4.0E-09
1,3-Diphenylguanidine	102-06-7	2	0	9.1E-10	4.1E-09	0	2.8E-09	2.3E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.3E-10	1.9E-10	2.0E-10	5.4E-10	8.4E-10
Methyl stearate	112-61-8	28	0	1.3E-09	1.9E-09	7.4E-10	3.5E-09	1.0E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
4-tert-Octylphenol	140-66-9	34	0	5.8E-09	1.2E-08	1.6E-09	2.8E-08	5.4E-08
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.4E-10	1.7E-09	1.4E-09	1.2E-09	4.3E-09	6.6E-09
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	1.7E-10	1.9E-09	2.2E-09	1.1E-09	6.9E-09	9.2E-09

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-226. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Coaches 30<40 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	2.9E-10	1.3E-09	0	8.9E-10	7.1E-09
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	4.2E-09	1.7E-08	0	1.6E-08	1.0E-07
Benzo[a]pyrene	50-32-8	32	0	2.2E-10	2.2E-10	1.4E-10	5.7E-10	8.9E-10
Benzo[e]pyrene	192-97-2	35	6.2E-11	6.5E-10	4.8E-10	6.1E-10	1.6E-09	1.8E-09
Benzo[g,h,i]perylene	191-24-2	35	6.1E-11	4.9E-10	3.2E-10	4.5E-10	1.1E-09	1.3E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	6.2E-10	3.7E-09	0	0	2.2E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	2.2E-10	2.9E-09	4.7E-09	1.5E-09	6.7E-09	2.8E-08
Chrysene	218-01-9	35	2.0E-10	1.9E-09	1.2E-09	1.9E-09	3.8E-09	4.3E-09
Coronene	191-07-1	1	0	1.5E-10	8.8E-10	0	0	5.2E-09
Cyclopenta[cd]pyrene	27208-37-3	35	3.4E-11	3.4E-10	2.0E-10	3.2E-10	6.4E-10	7.7E-10
Dicyclohexylamine	101-83-7	30	0	5.7E-08	9.4E-08	4.3E-08	1.0E-07	5.7E-07
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	9.9E-11	2.0E-10	4.6E-11	4.1E-10	1.0E-09
Dimethyl phthalate	131-11-3	3	0	3.0E-10	9.8E-10	0	3.2E-09	3.8E-09
1,3-Diphenylguanidine	102-06-7	2	0	8.6E-10	3.8E-09	0	2.7E-09	2.1E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.2E-10	1.8E-10	1.9E-10	5.0E-10	7.9E-10
Methyl stearate	112-61-8	28	0	1.2E-09	1.8E-09	6.9E-10	3.3E-09	9.6E-09
4-tert-Octylphenol	140-66-9	34	0	5.4E-09	1.1E-08	1.5E-09	2.6E-08	5.0E-08
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.4E-10	1.6E-09	1.3E-09	1.2E-09	4.1E-09	6.1E-09
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	1.6E-10	1.7E-09	2.1E-09	9.9E-10	6.5E-09	8.7E-09

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number



Values are rounded to two significant figures.

Table F-227. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Coaches 40<50 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	2.9E-10	1.3E-09	0	9.0E-10	7.1E-09
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	4.2E-09	1.8E-08	0	1.6E-08	1.0E-07
Benzo[a]pyrene	50-32-8	32	0	2.2E-10	2.2E-10	1.5E-10	5.8E-10	9.0E-10
Benzo[e]pyrene	192-97-2	35	6.3E-11	6.5E-10	4.8E-10	6.2E-10	1.6E-09	1.8E-09
Benzo[g,h,i]perylene	191-24-2	35	6.1E-11	5.0E-10	3.2E-10	4.6E-10	1.1E-09	1.3E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	6.2E-10	3.7E-09	0	0	2.2E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	2.2E-10	2.9E-09	4.7E-09	1.5E-09	6.8E-09	2.8E-08
Chrysene	218-01-9	35	2.1E-10	1.9E-09	1.2E-09	1.9E-09	3.8E-09	4.3E-09
Coronene	191-07-1	1	0	1.5E-10	8.9E-10	0	0	5.3E-09
Cyclopenta[cd]pyrene	27208-37-3	35	3.5E-11	3.4E-10	2.0E-10	3.3E-10	6.5E-10	7.8E-10
Dicyclohexylamine	101-83-7	30	0	5.7E-08	9.5E-08	4.4E-08	1.1E-07	5.7E-07
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	1.0E-10	2.0E-10	4.6E-11	4.2E-10	1.0E-09
Dimethyl phthalate	131-11-3	3	0	3.0E-10	9.9E-10	0	3.2E-09	3.8E-09
1,3-Diphenylguanidine	102-06-7	2	0	8.7E-10	3.9E-09	0	2.7E-09	2.1E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.2E-10	1.8E-10	1.9E-10	5.1E-10	7.9E-10
Methyl stearate	112-61-8	28	0	1.3E-09	1.8E-09	7.0E-10	3.3E-09	9.7E-09
4-tert-Octylphenol	140-66-9	34	0	5.5E-09	1.2E-08	1.5E-09	2.7E-08	5.1E-08
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.4E-10	1.6E-09	1.3E-09	1.2E-09	4.1E-09	6.2E-09
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	1.6E-10	1.8E-09	2.1E-09	1.0E-09	6.6E-09	8.7E-09

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-228. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Coaches 50<70 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	2.9E-10	1.3E-09	0	9.0E-10	7.2E-09
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	4.2E-09	1.8E-08	0	1.6E-08	1.0E-07



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	32	0	2.2E-10	2.2E-10	1.5E-10	5.8E-10	9.1E-10
Benzo[e]pyrene	192-97-2	35	6.3E-11	6.6E-10	4.8E-10	6.2E-10	1.6E-09	1.8E-09
Benzo[g,h,i]perylene	191-24-2	35	6.2E-11	5.0E-10	3.2E-10	4.6E-10	1.1E-09	1.3E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	6.3E-10	3.7E-09	0	0	2.2E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	2.2E-10	2.9E-09	4.8E-09	1.5E-09	6.8E-09	2.8E-08
Chrysene	218-01-9	35	2.1E-10	1.9E-09	1.2E-09	1.9E-09	3.8E-09	4.3E-09
Coronene	191-07-1	1	0	1.5E-10	8.9E-10	0	0	5.3E-09
Cyclopenta[cd]pyrene	27208-37-3	35	3.5E-11	3.4E-10	2.0E-10	3.3E-10	6.5E-10	7.8E-10
Dicyclohexylamine	101-83-7	30	0	5.7E-08	9.5E-08	4.4E-08	1.1E-07	5.7E-07
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	1.0E-10	2.0E-10	4.7E-11	4.2E-10	1.0E-09
Dimethyl phthalate	131-11-3	3	0	3.0E-10	1.0E-09	0	3.3E-09	3.8E-09
1,3-Diphenylguanidine	102-06-7	2	0	8.7E-10	3.9E-09	0	2.7E-09	2.1E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.2E-10	1.8E-10	1.9E-10	5.1E-10	8.0E-10
Methyl stearate	112-61-8	28	0	1.3E-09	1.8E-09	7.0E-10	3.3E-09	9.7E-09
4-tert-Octylphenol	140-66-9	34	0	5.5E-09	1.2E-08	1.5E-09	2.7E-08	5.1E-08
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.4E-10	1.6E-09	1.3E-09	1.2E-09	4.1E-09	6.2E-09
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	1.6E-10	1.8E-09	2.1E-09	1.0E-09	6.6E-09	8.8E-09

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-229. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Referees 16<30 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	3.1E-10	1.4E-09	0	9.5E-10	7.5E-09
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	4.4E-09	1.9E-08	0	1.7E-08	1.1E-07
Benzo[a]pyrene	50-32-8	32	0	2.3E-10	2.3E-10	1.5E-10	6.1E-10	9.5E-10
Benzo[e]pyrene	192-97-2	35	6.7E-11	6.9E-10	5.1E-10	6.6E-10	1.7E-09	1.9E-09
Benzo[g,h,i]perylene	191-24-2	35	6.5E-11	5.2E-10	3.4E-10	4.8E-10	1.2E-09	1.4E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	6.6E-10	3.9E-09	0	0	2.3E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	2.3E-10	3.1E-09	5.0E-09	1.6E-09	7.2E-09	3.0E-08
Chrysene	218-01-9	35	2.2E-10	2.0E-09	1.3E-09	2.0E-09	4.0E-09	4.6E-09
Coronene	191-07-1	1	0	1.6E-10	9.4E-10	0	0	5.6E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Cyclopenta[cd]pyrene	27208-37-3	35	3.7E-11	3.6E-10	2.1E-10	3.5E-10	6.9E-10	8.2E-10
Dicyclohexylamine	101-83-7	30	0	6.1E-08	1.0E-07	4.6E-08	1.1E-07	6.1E-07
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	1.1E-10	2.1E-10	4.9E-11	4.4E-10	1.1E-09
Dimethyl phthalate	131-11-3	3	0	3.2E-10	1.0E-09	0	3.4E-09	4.0E-09
1,3-Diphenylguanidine	102-06-7	2	0	9.1E-10	4.1E-09	0	2.8E-09	2.3E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.3E-10	1.9E-10	2.0E-10	5.4E-10	8.4E-10
Methyl stearate	112-61-8	28	0	1.3E-09	1.9E-09	7.4E-10	3.5E-09	1.0E-08
4-tert-Octylphenol	140-66-9	34	0	5.8E-09	1.2E-08	1.6E-09	2.8E-08	5.4E-08
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.4E-10	1.7E-09	1.4E-09	1.2E-09	4.3E-09	6.6E-09
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	1.7E-10	1.9E-09	2.2E-09	1.1E-09	6.9E-09	9.2E-09

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-230. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Referees 30<40 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	2.9E-10	1.3E-09	0	8.9E-10	7.1E-09
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	4.2E-09	1.7E-08	0	1.6E-08	1.0E-07
Benzo[a]pyrene	50-32-8	32	0	2.2E-10	2.2E-10	1.4E-10	5.7E-10	8.9E-10
Benzo[e]pyrene	192-97-2	35	6.2E-11	6.5E-10	4.8E-10	6.1E-10	1.6E-09	1.8E-09
Benzo[g,h,i]perylene	191-24-2	35	6.1E-11	4.9E-10	3.2E-10	4.5E-10	1.1E-09	1.3E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	6.2E-10	3.7E-09	0	0	2.2E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	2.2E-10	2.9E-09	4.7E-09	1.5E-09	6.7E-09	2.8E-08
Chrysene	218-01-9	35	2.0E-10	1.9E-09	1.2E-09	1.9E-09	3.8E-09	4.3E-09
Coronene	191-07-1	1	0	1.5E-10	8.8E-10	0	0	5.2E-09
Cyclopenta[cd]pyrene	27208-37-3	35	3.4E-11	3.4E-10	2.0E-10	3.2E-10	6.4E-10	7.7E-10
Dicyclohexylamine	101-83-7	30	0	5.7E-08	9.4E-08	4.3E-08	1.0E-07	5.7E-07
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	9.9E-11	2.0E-10	4.6E-11	4.1E-10	1.0E-09
Dimethyl phthalate	131-11-3	3	0	3.0E-10	9.8E-10	0	3.2E-09	3.8E-09
1,3-Diphenylguanidine	102-06-7	2	0	8.6E-10	3.8E-09	0	2.7E-09	2.1E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.2E-10	1.8E-10	1.9E-10	5.0E-10	7.9E-10
Methyl stearate	112-61-8	28	0	1.2E-09	1.8E-09	6.9E-10	3.3E-09	9.6E-09



Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
4-tert-Octylphenol	140-66-9	34	0	5.4E-09	1.1E-08	1.5E-09	2.6E-08	5.0E-08
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.4E-10	1.6E-09	1.3E-09	1.2E-09	4.1E-09	6.1E-09
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	1.6E-10	1.7E-09	2.1E-09	9.9E-10	6.5E-09	8.7E-09

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-231. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Referees 40<50 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	2.9E-10	1.3E-09	0	9.0E-10	7.1E-09
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	4.2E-09	1.8E-08	0	1.6E-08	1.0E-07
Benzo[a]pyrene	50-32-8	32	0	2.2E-10	2.2E-10	1.5E-10	5.8E-10	9.0E-10
Benzo[e]pyrene	192-97-2	35	6.3E-11	6.5E-10	4.8E-10	6.2E-10	1.6E-09	1.8E-09
Benzo[g,h,i]perylene	191-24-2	35	6.1E-11	5.0E-10	3.2E-10	4.6E-10	1.1E-09	1.3E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	6.2E-10	3.7E-09	0	0	2.2E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	2.2E-10	2.9E-09	4.7E-09	1.5E-09	6.8E-09	2.8E-08
Chrysene	218-01-9	35	2.1E-10	1.9E-09	1.2E-09	1.9E-09	3.8E-09	4.3E-09
Coronene	191-07-1	1	0	1.5E-10	8.9E-10	0	0	5.3E-09
Cyclopenta[cd]pyrene	27208-37-3	35	3.5E-11	3.4E-10	2.0E-10	3.3E-10	6.5E-10	7.8E-10
Dicyclohexylamine	101-83-7	30	0	5.7E-08	9.5E-08	4.4E-08	1.1E-07	5.7E-07
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	1.0E-10	2.0E-10	4.6E-11	4.2E-10	1.0E-09
Dimethyl phthalate	131-11-3	3	0	3.0E-10	9.9E-10	0	3.2E-09	3.8E-09
1,3-Diphenylguanidine	102-06-7	2	0	8.7E-10	3.9E-09	0	2.7E-09	2.1E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.2E-10	1.8E-10	1.9E-10	5.1E-10	7.9E-10
Methyl stearate	112-61-8	28	0	1.3E-09	1.8E-09	7.0E-10	3.3E-09	9.7E-09
4-tert-Octylphenol	140-66-9	34	0	5.5E-09	1.2E-08	1.5E-09	2.7E-08	5.1E-08
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.4E-10	1.6E-09	1.3E-09	1.2E-09	4.1E-09	6.2E-09
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	1.6E-10	1.8E-09	2.1E-09	1.0E-09	6.6E-09	8.7E-09

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number



Values are rounded to two significant figures.

Table F-232. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Referees 50<70 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	2.9E-10	1.3E-09	0	9.0E-10	7.2E-09
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	4.2E-09	1.8E-08	0	1.6E-08	1.0E-07
Benzo[a]pyrene	50-32-8	32	0	2.2E-10	2.2E-10	1.5E-10	5.8E-10	9.1E-10
Benzo[e]pyrene	192-97-2	35	6.3E-11	6.6E-10	4.8E-10	6.2E-10	1.6E-09	1.8E-09
Benzo[g,h,i]perylene	191-24-2	35	6.2E-11	5.0E-10	3.2E-10	4.6E-10	1.1E-09	1.3E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	6.3E-10	3.7E-09	0	0	2.2E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	2.2E-10	2.9E-09	4.8E-09	1.5E-09	6.8E-09	2.8E-08
Chrysene	218-01-9	35	2.1E-10	1.9E-09	1.2E-09	1.9E-09	3.8E-09	4.3E-09
Coronene	191-07-1	1	0	1.5E-10	8.9E-10	0	0	5.3E-09
Cyclopenta[cd]pyrene	27208-37-3	35	3.5E-11	3.4E-10	2.0E-10	3.3E-10	6.5E-10	7.8E-10
Dicyclohexylamine	101-83-7	30	0	5.7E-08	9.5E-08	4.4E-08	1.1E-07	5.7E-07
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	1.0E-10	2.0E-10	4.7E-11	4.2E-10	1.0E-09
Dimethyl phthalate	131-11-3	3	0	3.0E-10	1.0E-09	0	3.3E-09	3.8E-09
1,3-Diphenylguanidine	102-06-7	2	0	8.7E-10	3.9E-09	0	2.7E-09	2.1E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.2E-10	1.8E-10	1.9E-10	5.1E-10	8.0E-10
Methyl stearate	112-61-8	28	0	1.3E-09	1.8E-09	7.0E-10	3.3E-09	9.7E-09
4-tert-Octylphenol	140-66-9	34	0	5.5E-09	1.2E-08	1.5E-09	2.7E-08	5.1E-08
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.4E-10	1.6E-09	1.3E-09	1.2E-09	4.1E-09	6.2E-09
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	1.6E-10	1.8E-09	2.1E-09	1.0E-09	6.6E-09	8.8E-09

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-233. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Spectators Third Trimester Fetus<0 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	2.3E-10	1.0E-09	0	7.2E-10	5.7E-09
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	3.4E-09	1.4E-08	0	1.3E-08	8.2E-08



Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	32	0	1.8E-10	1.7E-10	1.2E-10	4.6E-10	7.2E-10
Benzo[e]pyrene	192-97-2	35	5.0E-11	5.2E-10	3.8E-10	4.9E-10	1.3E-09	1.4E-09
Benzo[g,h,i]perylene	191-24-2	35	4.9E-11	4.0E-10	2.6E-10	3.7E-10	8.7E-10	1.0E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	5.0E-10	2.9E-09	0	0	1.7E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	1.7E-10	2.3E-09	3.8E-09	1.2E-09	5.4E-09	2.2E-08
Chrysene	218-01-9	35	1.7E-10	1.5E-09	9.8E-10	1.5E-09	3.0E-09	3.4E-09
Coronene	191-07-1	1	0	1.2E-10	7.1E-10	0	0	4.2E-09
Cyclopenta[cd]pyrene	27208-37-3	35	2.8E-11	2.7E-10	1.6E-10	2.6E-10	5.2E-10	6.2E-10
Dicyclohexylamine	101-83-7	30	0	4.6E-08	7.6E-08	3.5E-08	8.5E-08	4.6E-07
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	8.0E-11	1.6E-10	3.7E-11	3.3E-10	8.3E-10
Dimethyl phthalate	131-11-3	3	0	2.4E-10	7.9E-10	0	2.6E-09	3.1E-09
1,3-Diphenylguanidine	102-06-7	2	0	6.9E-10	3.1E-09	0	2.1E-09	1.7E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	1.7E-10	1.4E-10	1.5E-10	4.0E-10	6.3E-10
Methyl stearate	112-61-8	28	0	1.0E-09	1.4E-09	5.6E-10	2.6E-09	7.7E-09
4-tert-Octylphenol	140-66-9	34	0	4.4E-09	9.2E-09	1.2E-09	2.1E-08	4.1E-08
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.1E-10	1.3E-09	1.0E-09	9.3E-10	3.3E-09	5.0E-09
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	1.3E-10	1.4E-09	1.7E-09	8.0E-10	5.2E-09	7.0E-09

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-234. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Spectators 0<2 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	1.2E-09	5.5E-09	0	3.8E-09	3.0E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	1.8E-08	7.5E-08	0	6.8E-08	4.4E-07
Benzo[a]pyrene	50-32-8	32	0	9.5E-10	9.3E-10	6.2E-10	2.5E-09	3.9E-09
Benzo[e]pyrene	192-97-2	35	2.7E-10	2.8E-09	2.1E-09	2.7E-09	6.7E-09	7.6E-09
Benzo[g,h,i]perylene	191-24-2	35	2.6E-10	2.1E-09	1.4E-09	2.0E-09	4.7E-09	5.6E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	2.7E-09	1.6E-08	0	0	9.3E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	9.3E-10	1.3E-08	2.0E-08	6.5E-09	2.9E-08	1.2E-07
Chrysene	218-01-9	35	8.9E-10	8.1E-09	5.2E-09	8.1E-09	1.6E-08	1.8E-08
Coronene	191-07-1	1	0	6.4E-10	3.8E-09	0	0	2.2E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Cyclopenta[cd]pyrene	27208-37-3	35	1.5E-10	1.5E-09	8.7E-10	1.4E-09	2.8E-09	3.3E-09
Dicyclohexylamine	101-83-7	30	0	2.4E-07	4.1E-07	1.9E-07	4.5E-07	2.5E-06
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	4.3E-10	8.7E-10	2.0E-10	1.8E-09	4.5E-09
Dimethyl phthalate	131-11-3	3	0	1.3E-09	4.2E-09	0	1.4E-08	1.6E-08
1,3-Diphenylguanidine	102-06-7	2	0	3.7E-09	1.7E-08	0	1.1E-08	9.1E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	9.3E-10	7.8E-10	8.2E-10	2.2E-09	3.4E-09
Methyl stearate	112-61-8	28	0	5.4E-09	7.6E-09	3.0E-09	1.4E-08	4.1E-08
4-tert-Octylphenol	140-66-9	34	0	2.3E-08	4.9E-08	6.4E-09	1.1E-07	2.2E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	5.9E-10	6.9E-09	5.6E-09	5.0E-09	1.8E-08	2.7E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	6.8E-10	7.5E-09	8.9E-09	4.3E-09	2.8E-08	3.7E-08

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-235. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Spectators 2<6 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	9.1E-10	4.1E-09	0	2.8E-09	2.2E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	1.3E-08	5.5E-08	0	5.0E-08	3.2E-07
Benzo[a]pyrene	50-32-8	32	0	7.0E-10	6.9E-10	4.6E-10	1.8E-09	2.8E-09
Benzo[e]pyrene	192-97-2	35	2.0E-10	2.1E-09	1.5E-09	2.0E-09	5.0E-09	5.6E-09
Benzo[g,h,i]perylene	191-24-2	35	1.9E-10	1.6E-09	1.0E-09	1.4E-09	3.4E-09	4.1E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	2.0E-09	1.2E-08	0	0	6.9E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	6.9E-10	9.2E-09	1.5E-08	4.8E-09	2.1E-08	8.8E-08
Chrysene	218-01-9	35	6.5E-10	6.0E-09	3.8E-09	6.0E-09	1.2E-08	1.4E-08
Coronene	191-07-1	1	0	4.7E-10	2.8E-09	0	0	1.7E-08
Cyclopenta[cd]pyrene	27208-37-3	35	1.1E-10	1.1E-09	6.4E-10	1.0E-09	2.0E-09	2.4E-09
Dicyclohexylamine	101-83-7	30	0	1.8E-07	3.0E-07	1.4E-07	3.3E-07	1.8E-06
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	3.2E-10	6.4E-10	1.5E-10	1.3E-09	3.3E-09
Dimethyl phthalate	131-11-3	3	0	9.4E-10	3.1E-09	0	1.0E-08	1.2E-08
1,3-Diphenylguanidine	102-06-7	2	0	2.7E-09	1.2E-08	0	8.5E-09	6.7E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	6.8E-10	5.7E-10	6.1E-10	1.6E-09	2.5E-09
Methyl stearate	112-61-8	28	0	3.9E-09	5.6E-09	2.2E-09	1.0E-08	3.0E-08



Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
4-tert-Octylphenol	140-66-9	34	0	1.7E-08	3.6E-08	4.7E-09	8.4E-08	1.6E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	4.3E-10	5.1E-09	4.1E-09	3.7E-09	1.3E-08	2.0E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	5.0E-10	5.5E-09	6.5E-09	3.1E-09	2.1E-08	2.7E-08

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-236. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Spectators 6<11 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	8.3E-10	3.7E-09	0	2.6E-09	2.0E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	1.2E-08	5.0E-08	0	4.6E-08	2.9E-07
Benzo[a]pyrene	50-32-8	32	0	6.3E-10	6.2E-10	4.1E-10	1.6E-09	2.6E-09
Benzo[e]pyrene	192-97-2	35	1.8E-10	1.9E-09	1.4E-09	1.8E-09	4.5E-09	5.1E-09
Benzo[g,h,i]perylene	191-24-2	35	1.8E-10	1.4E-09	9.2E-10	1.3E-09	3.1E-09	3.7E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	1.8E-09	1.1E-08	0	0	6.2E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	6.2E-10	8.4E-09	1.4E-08	4.3E-09	1.9E-08	8.0E-08
Chrysene	218-01-9	35	5.9E-10	5.4E-09	3.5E-09	5.4E-09	1.1E-08	1.2E-08
Coronene	191-07-1	1	0	4.3E-10	2.5E-09	0	0	1.5E-08
Cyclopenta[cd]pyrene	27208-37-3	35	9.9E-11	9.7E-10	5.8E-10	9.3E-10	1.9E-09	2.2E-09
Dicyclohexylamine	101-83-7	30	0	1.6E-07	2.7E-07	1.3E-07	3.0E-07	1.6E-06
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	2.9E-10	5.8E-10	1.3E-10	1.2E-09	3.0E-09
Dimethyl phthalate	131-11-3	3	0	8.5E-10	2.8E-09	0	9.3E-09	1.1E-08
1,3-Diphenylguanidine	102-06-7	2	0	2.5E-09	1.1E-08	0	7.7E-09	6.1E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	6.2E-10	5.2E-10	5.5E-10	1.4E-09	2.3E-09
Methyl stearate	112-61-8	28	0	3.6E-09	5.0E-09	2.0E-09	9.4E-09	2.8E-08
4-tert-Octylphenol	140-66-9	34	0	1.6E-08	3.3E-08	4.2E-09	7.6E-08	1.5E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	3.9E-10	4.6E-09	3.8E-09	3.3E-09	1.2E-08	1.8E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	4.6E-10	5.0E-09	5.9E-09	2.9E-09	1.9E-08	2.5E-08

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number



Values are rounded to two significant figures.

Table F-237. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Spectators 11<16 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	7.1E-10	3.2E-09	0	2.2E-09	1.7E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	1.0E-08	4.3E-08	0	3.9E-08	2.5E-07
Benzo[a]pyrene	50-32-8	32	0	5.4E-10	5.3E-10	3.5E-10	1.4E-09	2.2E-09
Benzo[e]pyrene	192-97-2	35	1.5E-10	1.6E-09	1.2E-09	1.5E-09	3.9E-09	4.4E-09
Benzo[g,h,i]perylene	191-24-2	35	1.5E-10	1.2E-09	7.9E-10	1.1E-09	2.7E-09	3.2E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	1.5E-09	9.0E-09	0	0	5.3E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	5.3E-10	7.1E-09	1.2E-08	3.7E-09	1.7E-08	6.8E-08
Chrysene	218-01-9	35	5.1E-10	4.6E-09	3.0E-09	4.6E-09	9.3E-09	1.1E-08
Coronene	191-07-1	1	0	3.7E-10	2.2E-09	0	0	1.3E-08
Cyclopenta[cd]pyrene	27208-37-3	35	8.5E-11	8.3E-10	4.9E-10	8.0E-10	1.6E-09	1.9E-09
Dicyclohexylamine	101-83-7	30	0	1.4E-07	2.3E-07	1.1E-07	2.6E-07	1.4E-06
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	2.5E-10	4.9E-10	1.1E-10	1.0E-09	2.6E-09
Dimethyl phthalate	131-11-3	3	0	7.3E-10	2.4E-09	0	7.9E-09	9.3E-09
1,3-Diphenylguanidine	102-06-7	2	0	2.1E-09	9.4E-09	0	6.6E-09	5.2E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	5.3E-10	4.4E-10	4.7E-10	1.2E-09	1.9E-09
Methyl stearate	112-61-8	28	0	3.1E-09	4.3E-09	1.7E-09	8.1E-09	2.4E-08
4-tert-Octylphenol	140-66-9	34	0	1.3E-08	2.8E-08	3.6E-09	6.5E-08	1.2E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	3.3E-10	3.9E-09	3.2E-09	2.8E-09	1.0E-08	1.5E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	3.9E-10	4.3E-09	5.1E-09	2.4E-09	1.6E-08	2.1E-08

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-238. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Spectators 16<30 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	3.1E-10	1.4E-09	0	9.6E-10	7.6E-09
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	4.5E-09	1.9E-08	0	1.7E-08	1.1E-07



Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	32	0	2.4E-10	2.3E-10	1.6E-10	6.2E-10	9.7E-10
Benzo[e]pyrene	192-97-2	35	6.7E-11	7.0E-10	5.2E-10	6.6E-10	1.7E-09	1.9E-09
Benzo[g,h,i]perylene	191-24-2	35	6.6E-11	5.3E-10	3.4E-10	4.9E-10	1.2E-09	1.4E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	6.7E-10	3.9E-09	0	0	2.3E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	2.3E-10	3.1E-09	5.1E-09	1.6E-09	7.2E-09	3.0E-08
Chrysene	218-01-9	35	2.2E-10	2.0E-09	1.3E-09	2.0E-09	4.1E-09	4.6E-09
Coronene	191-07-1	1	0	1.6E-10	9.5E-10	0	0	5.6E-09
Cyclopenta[cd]pyrene	27208-37-3	35	3.7E-11	3.6E-10	2.2E-10	3.5E-10	6.9E-10	8.3E-10
Dicyclohexylamine	101-83-7	30	0	6.1E-08	1.0E-07	4.7E-08	1.1E-07	6.1E-07
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	1.1E-10	2.2E-10	5.0E-11	4.5E-10	1.1E-09
Dimethyl phthalate	131-11-3	3	0	3.2E-10	1.1E-09	0	3.5E-09	4.1E-09
1,3-Diphenylguanidine	102-06-7	2	0	9.2E-10	4.1E-09	0	2.9E-09	2.3E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.3E-10	1.9E-10	2.1E-10	5.4E-10	8.5E-10
Methyl stearate	112-61-8	28	0	1.3E-09	1.9E-09	7.5E-10	3.5E-09	1.0E-08
4-tert-Octylphenol	140-66-9	34	0	5.9E-09	1.2E-08	1.6E-09	2.8E-08	5.4E-08
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.5E-10	1.7E-09	1.4E-09	1.2E-09	4.4E-09	6.6E-09
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	1.7E-10	1.9E-09	2.2E-09	1.1E-09	7.0E-09	9.3E-09

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-239. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Spectators 30<40 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	2.7E-10	1.2E-09	0	8.3E-10	6.6E-09
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	3.9E-09	1.6E-08	0	1.5E-08	9.5E-08
Benzo[a]pyrene	50-32-8	32	0	2.0E-10	2.0E-10	1.3E-10	5.3E-10	8.3E-10
Benzo[e]pyrene	192-97-2	35	5.8E-11	6.0E-10	4.5E-10	5.7E-10	1.5E-09	1.6E-09
Benzo[g,h,i]perylene	191-24-2	35	5.7E-11	4.6E-10	3.0E-10	4.2E-10	1.0E-09	1.2E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	5.8E-10	3.4E-09	0	0	2.0E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	2.0E-10	2.7E-09	4.4E-09	1.4E-09	6.3E-09	2.6E-08
Chrysene	218-01-9	35	1.9E-10	1.8E-09	1.1E-09	1.8E-09	3.5E-09	4.0E-09
Coronene	191-07-1	1	0	1.4E-10	8.2E-10	0	0	4.9E-09



Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Cyclopenta[cd]pyrene	27208-37-3	35	3.2E-11	3.1E-10	1.9E-10	3.0E-10	6.0E-10	7.2E-10
Dicyclohexylamine	101-83-7	30	0	5.3E-08	8.8E-08	4.0E-08	9.8E-08	5.3E-07
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	9.3E-11	1.9E-10	4.3E-11	3.9E-10	9.7E-10
Dimethyl phthalate	131-11-3	3	0	2.8E-10	9.2E-10	0	3.0E-09	3.5E-09
1,3-Diphenylguanidine	102-06-7	2	0	8.0E-10	3.6E-09	0	2.5E-09	2.0E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.0E-10	1.7E-10	1.8E-10	4.7E-10	7.3E-10
Methyl stearate	112-61-8	28	0	1.2E-09	1.6E-09	6.4E-10	3.1E-09	8.9E-09
4-tert-Octylphenol	140-66-9	34	0	5.1E-09	1.1E-08	1.4E-09	2.5E-08	4.7E-08
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.3E-10	1.5E-09	1.2E-09	1.1E-09	3.8E-09	5.7E-09
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	1.5E-10	1.6E-09	1.9E-09	9.2E-10	6.1E-09	8.1E-09

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-240. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Spectators 40<50 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	2.7E-10	1.2E-09	0	8.4E-10	6.7E-09
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	3.9E-09	1.6E-08	0	1.5E-08	9.6E-08
Benzo[a]pyrene	50-32-8	32	0	2.1E-10	2.0E-10	1.4E-10	5.4E-10	8.4E-10
Benzo[e]pyrene	192-97-2	35	5.9E-11	6.1E-10	4.5E-10	5.8E-10	1.5E-09	1.7E-09
Benzo[g,h,i]perylene	191-24-2	35	5.7E-11	4.6E-10	3.0E-10	4.3E-10	1.0E-09	1.2E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	5.8E-10	3.4E-09	0	0	2.0E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	2.0E-10	2.7E-09	4.4E-09	1.4E-09	6.3E-09	2.6E-08
Chrysene	218-01-9	35	1.9E-10	1.8E-09	1.1E-09	1.8E-09	3.6E-09	4.0E-09
Coronene	191-07-1	1	0	1.4E-10	8.3E-10	0	0	4.9E-09
Cyclopenta[cd]pyrene	27208-37-3	35	3.2E-11	3.2E-10	1.9E-10	3.1E-10	6.1E-10	7.2E-10
Dicyclohexylamine	101-83-7	30	0	5.3E-08	8.8E-08	4.1E-08	9.9E-08	5.3E-07
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	9.4E-11	1.9E-10	4.3E-11	3.9E-10	9.8E-10
Dimethyl phthalate	131-11-3	3	0	2.8E-10	9.3E-10	0	3.0E-09	3.6E-09
1,3-Diphenylguanidine	102-06-7	2	0	8.1E-10	3.6E-09	0	2.5E-09	2.0E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.0E-10	1.7E-10	1.8E-10	4.7E-10	7.4E-10
Methyl stearate	112-61-8	28	0	1.2E-09	1.7E-09	6.5E-10	3.1E-09	9.0E-09



Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
4-tert-Octylphenol	140-66-9	34	0	5.1E-09	1.1E-08	1.4E-09	2.5E-08	4.7E-08
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.3E-10	1.5E-09	1.2E-09	1.1E-09	3.8E-09	5.8E-09
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	1.5E-10	1.6E-09	1.9E-09	9.3E-10	6.1E-09	8.2E-09

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-241. Field-Specific Average Dermal One-Day Dose^a (AD_{der-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Spectators 50<70 Years**

Chemical	CASRN	Detection	AD _{der-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-Azacyclotridecanone	947-04-6	2	0	2.5E-10	1.1E-09	0	7.9E-10	6.3E-09
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	9	0	3.7E-09	1.5E-08	0	1.4E-08	9.1E-08
Benzo[a]pyrene	50-32-8	32	0	1.9E-10	1.9E-10	1.3E-10	5.1E-10	7.9E-10
Benzo[e]pyrene	192-97-2	35	5.5E-11	5.7E-10	4.2E-10	5.5E-10	1.4E-09	1.6E-09
Benzo[g,h,i]perylene	191-24-2	35	5.4E-11	4.4E-10	2.8E-10	4.0E-10	9.6E-10	1.1E-09
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	1	0	5.5E-10	3.2E-09	0	0	1.9E-08
Bis(2-Ethylhexyl)adipate	103-23-1	35	1.9E-10	2.6E-09	4.2E-09	1.3E-09	6.0E-09	2.5E-08
Chrysene	218-01-9	35	1.8E-10	1.7E-09	1.1E-09	1.7E-09	3.4E-09	3.8E-09
Coronene	191-07-1	1	0	1.3E-10	7.8E-10	0	0	4.6E-09
Cyclopenta[cd]pyrene	27208-37-3	35	3.0E-11	3.0E-10	1.8E-10	2.9E-10	5.7E-10	6.8E-10
Dicyclohexylamine	101-83-7	30	0	5.0E-08	8.3E-08	3.9E-08	9.3E-08	5.0E-07
N,N-Dicyclohexylmethylamine	7560-83-0	28	0	8.8E-11	1.8E-10	4.1E-11	3.7E-10	9.2E-10
Dimethyl phthalate	131-11-3	3	0	2.6E-10	8.7E-10	0	2.9E-09	3.4E-09
1,3-Diphenylguanidine	102-06-7	2	0	7.6E-10	3.4E-09	0	2.4E-09	1.9E-08
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	1.9E-10	1.6E-10	1.7E-10	4.5E-10	7.0E-10
Methyl stearate	112-61-8	28	0	1.1E-09	1.6E-09	6.1E-10	2.9E-09	8.5E-09
4-tert-Octylphenol	140-66-9	34	0	4.8E-09	1.0E-08	1.3E-09	2.3E-08	4.5E-08
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.2E-10	1.4E-09	1.2E-09	1.0E-09	3.6E-09	5.5E-09
Phenol, 4-(1-phenylethyl)-	1988-89-2	35	1.4E-10	1.6E-09	1.8E-09	8.8E-10	5.8E-09	7.7E-09

^a 35 field-specific AD_{der-DART-field} values are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number



Values are rounded to two significant figures.

F.5.3. Average Dermal Daily Dose (ADD_{der}) for Chronic Non-Cancer Hazard of General Chemicals

Table F-242. Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—
Combined Gender **Athletes**

Chemical Name	CASRN	ADD _{der}						
		2<6 Years	6<11 Years	11<16 Years	16<30 Years	30<40 Years	40<50 Years	50<70 Years
Acenaphthylene	208-96-8	8.0E-12	6.3E-12	6.7E-12	8.4E-12	6.0E-12	5.1E-12	5.1E-12
Aniline	62-53-3	2.6E-10	2.0E-10	2.1E-10	2.7E-10	1.9E-10	1.6E-10	1.6E-10
Anthracene	120-12-7	8.9E-11	7.0E-11	7.4E-11	9.3E-11	6.6E-11	5.6E-11	5.7E-11
Anthracene, 2-methyl-	613-12-7	1.3E-10	1.0E-10	1.1E-10	1.3E-10	9.5E-11	8.1E-11	8.1E-11
Anthracene, 9,10-diphenyl-	1499-10-1	1.7E-11	1.4E-11	1.5E-11	1.8E-11	1.3E-11	1.1E-11	1.1E-11
Anthracene, 9-phenyl	602-55-1	4.3E-11	3.4E-11	3.6E-11	4.6E-11	3.2E-11	2.7E-11	2.8E-11
Benz[a]anthracene	56-55-3	3.5E-10	2.8E-10	2.9E-10	3.7E-10	2.6E-10	2.2E-10	2.2E-10
Benzene, n-butyl-	104-51-8	5.3E-13	4.2E-13	4.4E-13	5.5E-13	3.9E-13	3.3E-13	3.4E-13
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	2.4E-08	1.9E-08	2.0E-08	2.5E-08	1.8E-08	1.5E-08	1.5E-08
Benzo[b]fluoranthene	205-99-2	5.0E-10	3.9E-10	4.2E-10	5.2E-10	3.7E-10	3.2E-10	3.2E-10
7H-Benzo[c]fluorene	205-12-9	4.4E-11	3.4E-11	3.6E-11	4.6E-11	3.2E-11	2.8E-11	2.8E-11
Benzo[k]fluoranthene	207-08-9	2.2E-10	1.7E-10	1.8E-10	2.3E-10	1.6E-10	1.4E-10	1.4E-10
Benzothiazole	95-16-9	2.8E-07	2.2E-07	2.4E-07	3.0E-07	2.1E-07	1.8E-07	1.8E-07
Benzothiazole, 2-phenyl-	883-93-2	6.9E-09	5.5E-09	5.8E-09	7.3E-09	5.2E-09	4.4E-09	4.4E-09
1,3-Benzothiazole-2-thiol	149-30-4	1.9E-09	1.5E-09	1.6E-09	2.0E-09	1.4E-09	1.2E-09	1.2E-09
Benzothiazolone	934-34-9	8.0E-07	6.3E-07	6.6E-07	8.4E-07	5.9E-07	5.0E-07	5.1E-07
Benzyl butyl phthalate	85-68-7	3.2E-09	2.6E-09	2.7E-09	3.4E-09	2.4E-09	2.1E-09	2.1E-09
Cyclohexyl isothiocyanate	1122-82-3	1.2E-11	9.5E-12	1.0E-11	1.3E-11	8.9E-12	7.6E-12	7.6E-12
Dibenz[a,h]anthracene	53-70-3	1.3E-10	1.0E-10	1.1E-10	1.4E-10	9.7E-11	8.2E-11	8.3E-11
Dibenzothiophene	132-65-0	7.8E-11	6.1E-11	6.5E-11	8.2E-11	5.8E-11	4.9E-11	5.0E-11
Diethyl Phthalate	84-66-2	5.9E-10	4.7E-10	4.9E-10	6.2E-10	4.4E-10	3.7E-10	3.8E-10
Diisobutyl Phthalate	84-69-5	4.9E-10	3.9E-10	4.1E-10	5.2E-10	3.7E-10	3.1E-10	3.2E-10
Diisooctylphthalate	27554-26-3	1.6E-08	1.3E-08	1.4E-08	1.7E-08	1.2E-08	1.0E-08	1.0E-08
Di-n-octyl phthalate	117-84-0	3.5E-09	2.7E-09	2.9E-09	3.6E-09	2.6E-09	2.2E-09	2.2E-09
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	3.0E-12	2.4E-12	2.5E-12	3.2E-12	2.3E-12	1.9E-12	1.9E-12
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	1.5E-08	1.2E-08	1.3E-08	1.6E-08	1.1E-08	9.6E-09	9.6E-09
Fluoranthene	206-44-0	2.7E-09	2.1E-09	2.3E-09	2.8E-09	2.0E-09	1.7E-09	1.7E-09
Fluorene	86-73-7	1.1E-11	9.0E-12	9.5E-12	1.2E-11	8.5E-12	7.3E-12	7.3E-12
Hexanoic Acid, 2-ethyl	149-57-5	1.3E-11	1.0E-11	1.1E-11	1.4E-11	9.8E-12	8.3E-12	8.4E-12
1-Hydroxypyrene	5315-79-7	7.6E-10	6.0E-10	6.3E-10	8.0E-10	5.7E-10	4.8E-10	4.8E-10



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical Name	CASRN	ADD _{der}						
		2<6 Years	6<11 Years	11<16 Years	16<30 Years	30<40 Years	40<50 Years	50<70 Years
2-(Methylthio)benzothiazole	615-22-5	1.0E-08	8.2E-09	8.7E-09	1.1E-08	7.8E-09	6.6E-09	6.7E-09
Naphthalene	91-20-3	1.7E-11	1.3E-11	1.4E-11	1.8E-11	1.2E-11	1.1E-11	1.1E-11
Naphthalene, 1,2-dimethyl-	573-98-8	1.2E-12	9.3E-13	9.8E-13	1.2E-12	8.8E-13	7.5E-13	7.5E-13
Naphthalene, 1,6-dimethyl-	575-43-9	1.2E-11	9.9E-12	1.0E-11	1.3E-11	9.3E-12	7.9E-12	8.0E-12
Naphthalene, 1-methyl-	90-12-0	2.6E-12	2.1E-12	2.2E-12	2.8E-12	2.0E-12	1.7E-12	1.7E-12
Naphthalene, 2-(bromomethyl)-	939-26-4	4.9E-10	3.9E-10	4.1E-10	5.2E-10	3.7E-10	3.1E-10	3.1E-10
Naphthalene, 2,3-dimethyl-	581-40-8	1.5E-10	1.1E-10	1.2E-10	1.5E-10	1.1E-10	9.2E-11	9.3E-11
Naphthalene, 2-methyl	91-57-6	2.6E-12	2.0E-12	2.1E-12	2.7E-12	1.9E-12	1.6E-12	1.6E-12
1-Octadecene	112-88-9	8.4E-10	6.6E-10	7.0E-10	8.8E-10	6.3E-10	5.3E-10	5.4E-10
Phenanthrene	85-01-8	5.8E-10	4.6E-10	4.8E-10	6.1E-10	4.3E-10	3.7E-10	3.7E-10
Phenanthrene, 1-methyl	832-69-9	3.0E-10	2.4E-10	2.5E-10	3.2E-10	2.3E-10	1.9E-10	1.9E-10
Phenanthrene, 2-methyl-	2531-84-2	3.7E-10	2.9E-10	3.1E-10	3.9E-10	2.7E-10	2.3E-10	2.3E-10
Phenanthrene, 3-methyl	832-71-3	1.3E-10	1.1E-10	1.1E-10	1.4E-10	1.0E-10	8.5E-11	8.5E-11
N-Phenylbenzamide	93-98-1	8.3E-10	6.6E-10	7.0E-10	8.8E-10	6.2E-10	5.3E-10	5.3E-10
Phthalimide	85-41-6	5.4E-09	4.3E-09	4.5E-09	5.7E-09	4.0E-09	3.4E-09	3.5E-09
Pyrene	129-00-0	7.3E-09	5.8E-09	6.1E-09	7.7E-09	5.5E-09	4.6E-09	4.7E-09
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2.7E-10	2.1E-10	2.2E-10	2.8E-10	2.0E-10	1.7E-10	1.7E-10
Triethylene glycol monobutyl ether	143-22-6	1.4E-09	1.1E-09	1.2E-09	1.5E-09	1.0E-09	8.9E-10	8.9E-10
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	1.9E-10	1.5E-10	1.6E-10	2.0E-10	1.4E-10	1.2E-10	1.2E-10

^a ADD_{der} was calculated from the mean of 35 individual field average concentrations (C_{der-crumb rubber}) of a chemical detected in artificial sweat extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-243. Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—
Combined Gender **Coaches**

Chemical Name	CASRN	ADD _{der}			
		16<30 Years	30<40 Years	40<50 Years	50<70 Years
Acenaphthylene	208-96-8	4.2E-12	3.9E-12	3.9E-12	3.9E-12
Aniline	62-53-3	1.3E-10	1.2E-10	1.3E-10	1.3E-10
Anthracene	120-12-7	4.6E-11	4.3E-11	4.4E-11	4.4E-11
Anthracene, 2-methyl-	613-12-7	6.6E-11	6.2E-11	6.3E-11	6.3E-11
Anthracene, 9,10-diphenyl-	1499-10-1	9.1E-12	8.5E-12	8.6E-12	8.6E-12
Anthracene, 9-phenyl	602-55-1	2.3E-11	2.1E-11	2.1E-11	2.1E-11
Benz[a]anthracene	56-55-3	1.8E-10	1.7E-10	1.7E-10	1.7E-10
Benzene, n-butyl-	104-51-8	2.7E-13	2.6E-13	2.6E-13	2.6E-13



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical Name	CASRN	ADD _{der}			
		16<30 Years	30<40 Years	40<50 Years	50<70 Years
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	1.2E-08	1.2E-08	1.2E-08	1.2E-08
Benzo[b]fluoranthene	205-99-2	2.6E-10	2.4E-10	2.5E-10	2.5E-10
7H-Benzo[c]fluorene	205-12-9	2.3E-11	2.1E-11	2.1E-11	2.1E-11
Benzo[k]fluoranthene	207-08-9	1.1E-10	1.1E-10	1.1E-10	1.1E-10
Benzothiazole	95-16-9	1.5E-07	1.4E-07	1.4E-07	1.4E-07
Benzothiazole, 2-phenyl-	883-93-2	3.6E-09	3.4E-09	3.4E-09	3.4E-09
1,3-Benzothiazole-2-thiol	149-30-4	9.8E-10	9.2E-10	9.3E-10	9.3E-10
Benzothiazolone	934-34-9	4.1E-07	3.9E-07	3.9E-07	3.9E-07
Benzyl butyl phthalate	85-68-7	1.7E-09	1.6E-09	1.6E-09	1.6E-09
Cyclohexyl isothiocyanate	1122-82-3	6.2E-12	5.8E-12	5.9E-12	5.9E-12
Dibenz[a,h]anthracene	53-70-3	6.7E-11	6.3E-11	6.4E-11	6.4E-11
Dibenzothiophene	132-65-0	4.0E-11	3.8E-11	3.8E-11	3.8E-11
Diethyl Phthalate	84-66-2	3.1E-10	2.9E-10	2.9E-10	2.9E-10
Diisobutyl Phthalate	84-69-5	2.6E-10	2.4E-10	2.4E-10	2.4E-10
Diisooctylphthalate	27554-26-3	8.5E-09	7.9E-09	8.0E-09	8.0E-09
Di-n-octyl phthalate	117-84-0	1.8E-09	1.7E-09	1.7E-09	1.7E-09
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	1.6E-12	1.5E-12	1.5E-12	1.5E-12
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	7.9E-09	7.4E-09	7.4E-09	7.5E-09
Fluoranthene	206-44-0	1.4E-09	1.3E-09	1.3E-09	1.3E-09
Fluorene	86-73-7	5.9E-12	5.6E-12	5.6E-12	5.6E-12
Hexanoic Acid, 2-ethyl	149-57-5	6.8E-12	6.4E-12	6.4E-12	6.5E-12
1-Hydroxypyrene	5315-79-7	3.9E-10	3.7E-10	3.7E-10	3.7E-10
2-(Methylthio)benzothiazole	615-22-5	5.4E-09	5.1E-09	5.1E-09	5.1E-09
Naphthalene	91-20-3	8.7E-12	8.1E-12	8.2E-12	8.3E-12
Naphthalene, 1,2-dimethyl-	573-98-8	6.1E-13	5.7E-13	5.8E-13	5.8E-13
Naphthalene, 1,6-dimethyl-	575-43-9	6.5E-12	6.1E-12	6.1E-12	6.2E-12
Naphthalene, 1-methyl-	90-12-0	1.4E-12	1.3E-12	1.3E-12	1.3E-12
Naphthalene, 2-(bromomethyl)-	939-26-4	2.5E-10	2.4E-10	2.4E-10	2.4E-10
Naphthalene, 2,3-dimethyl-	581-40-8	7.5E-11	7.1E-11	7.1E-11	7.2E-11
Naphthalene, 2-methyl	91-57-6	1.3E-12	1.3E-12	1.3E-12	1.3E-12
1-Octadecene	112-88-9	4.4E-10	4.1E-10	4.1E-10	4.1E-10
Phenanthrene	85-01-8	3.0E-10	2.8E-10	2.8E-10	2.9E-10
Phenanthrene, 1-methyl	832-69-9	1.6E-10	1.5E-10	1.5E-10	1.5E-10
Phenanthrene, 2-methyl-	2531-84-2	1.9E-10	1.8E-10	1.8E-10	1.8E-10
Phenanthrene, 3-methyl	832-71-3	6.9E-11	6.5E-11	6.6E-11	6.6E-11
N-Phenylbenzamide	93-98-1	4.3E-10	4.1E-10	4.1E-10	4.1E-10
Phthalimide	85-41-6	2.8E-09	2.6E-09	2.7E-09	2.7E-09
Pyrene	129-00-0	3.8E-09	3.6E-09	3.6E-09	3.6E-09



Chemical Name	CASRN	ADD _{der}			
		16<30 Years	30<40 Years	40<50 Years	50<70 Years
Pyridine, 2-(4-methylphenyl)-	4467-06-5	1.4E-10	1.3E-10	1.3E-10	1.3E-10
Triethylene glycol monobutyl ether	143-22-6	7.3E-10	6.8E-10	6.9E-10	6.9E-10
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	9.8E-11	9.2E-11	9.3E-11	9.3E-11

^a ADD_{der} was calculated from the mean of 35 individual field average concentrations (C_{der-crumb rubber}) of a chemical detected in artificial sweat extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-244. Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—
Combined Gender Referees

Chemical Name	CASRN	ADD _{der}			
		16<30 Years	30<40 Years	40<50 Years	50<70 Years
Acenaphthylene	208-96-8	1.7E-12	1.6E-12	1.6E-12	1.6E-12
Aniline	62-53-3	5.4E-11	5.0E-11	5.1E-11	5.1E-11
Anthracene	120-12-7	1.9E-11	1.7E-11	1.8E-11	1.8E-11
Anthracene, 2-methyl-	613-12-7	2.7E-11	2.5E-11	2.5E-11	2.5E-11
Anthracene, 9,10-diphenyl-	1499-10-1	3.7E-12	3.4E-12	3.5E-12	3.5E-12
Anthracene, 9-phenyl	602-55-1	9.1E-12	8.5E-12	8.6E-12	8.6E-12
Benz[a]anthracene	56-55-3	7.3E-11	6.9E-11	7.0E-11	7.0E-11
Benzene, n-butyl-	104-51-8	1.1E-13	1.0E-13	1.0E-13	1.0E-13
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	5.0E-09	4.7E-09	4.7E-09	4.7E-09
Benzo[b]fluoranthene	205-99-2	1.0E-10	9.8E-11	9.9E-11	9.9E-11
7H-Benzo[c]fluorene	205-12-9	9.1E-12	8.5E-12	8.6E-12	8.6E-12
Benzo[k]fluoranthene	207-08-9	4.6E-11	4.3E-11	4.3E-11	4.3E-11
Benzothiazole	95-16-9	5.9E-08	5.6E-08	5.6E-08	5.6E-08
Benzothiazole, 2-phenyl-	883-93-2	1.4E-09	1.4E-09	1.4E-09	1.4E-09
1,3-Benzothiazole-2-thiol	149-30-4	3.9E-10	3.7E-10	3.7E-10	3.7E-10
Benzothiazolone	934-34-9	1.7E-07	1.6E-07	1.6E-07	1.6E-07
Benzyl butyl phthalate	85-68-7	6.8E-10	6.3E-10	6.4E-10	6.4E-10
Cyclohexyl isothiocyanate	1122-82-3	2.5E-12	2.3E-12	2.4E-12	2.4E-12
Dibenz[a,h]anthracene	53-70-3	2.7E-11	2.5E-11	2.6E-11	2.6E-11
Dibenzothiophene	132-65-0	1.6E-11	1.5E-11	1.5E-11	1.5E-11
Diethyl Phthalate	84-66-2	1.2E-10	1.2E-10	1.2E-10	1.2E-10
Diisobutyl Phthalate	84-69-5	1.0E-10	9.7E-11	9.8E-11	9.8E-11
Diisooctylphthalate	27554-26-3	3.4E-09	3.2E-09	3.2E-09	3.2E-09
Di-n-octyl phthalate	117-84-0	7.2E-10	6.8E-10	6.8E-10	6.9E-10



Chemical Name	CASRN	ADD _{der}			
		16<30 Years	30<40 Years	40<50 Years	50<70 Years
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	6.4E-13	6.0E-13	6.0E-13	6.0E-13
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	3.2E-09	3.0E-09	3.0E-09	3.0E-09
Fluoranthene	206-44-0	5.6E-10	5.3E-10	5.3E-10	5.4E-10
Fluorene	86-73-7	2.4E-12	2.2E-12	2.3E-12	2.3E-12
Hexanoic Acid, 2-ethyl	149-57-5	2.7E-12	2.6E-12	2.6E-12	2.6E-12
1-Hydroxypyrene	5315-79-7	1.6E-10	1.5E-10	1.5E-10	1.5E-10
2-(Methylthio)benzothiazole	615-22-5	2.2E-09	2.0E-09	2.1E-09	2.1E-09
Naphthalene	91-20-3	3.5E-12	3.3E-12	3.3E-12	3.3E-12
Naphthalene, 1,2-dimethyl-	573-98-8	2.5E-13	2.3E-13	2.3E-13	2.3E-13
Naphthalene, 1,6-dimethyl-	575-43-9	2.6E-12	2.4E-12	2.5E-12	2.5E-12
Naphthalene, 1-methyl-	90-12-0	5.5E-13	5.2E-13	5.2E-13	5.3E-13
Naphthalene, 2-(bromomethyl)-	939-26-4	1.0E-10	9.6E-11	9.7E-11	9.7E-11
Naphthalene, 2,3-dimethyl-	581-40-8	3.0E-11	2.8E-11	2.9E-11	2.9E-11
Naphthalene, 2-methyl	91-57-6	5.4E-13	5.0E-13	5.1E-13	5.1E-13
1-Octadecene	112-88-9	1.8E-10	1.6E-10	1.7E-10	1.7E-10
Phenanthrene	85-01-8	1.2E-10	1.1E-10	1.1E-10	1.1E-10
Phenanthrene, 1-methyl	832-69-9	6.3E-11	5.9E-11	6.0E-11	6.0E-11
Phenanthrene, 2-methyl-	2531-84-2	7.7E-11	7.2E-11	7.3E-11	7.3E-11
Phenanthrene, 3-methyl	832-71-3	2.8E-11	2.6E-11	2.6E-11	2.6E-11
N-Phenylbenzamide	93-98-1	1.7E-10	1.6E-10	1.7E-10	1.7E-10
Phthalimide	85-41-6	1.1E-09	1.1E-09	1.1E-09	1.1E-09
Pyrene	129-00-0	1.5E-09	1.4E-09	1.4E-09	1.4E-09
Pyridine, 2-(4-methylphenyl)-	4467-06-5	5.6E-11	5.3E-11	5.3E-11	5.3E-11
Triethylene glycol monobutyl ether	143-22-6	2.9E-10	2.7E-10	2.8E-10	2.8E-10
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	3.9E-11	3.7E-11	3.7E-11	3.7E-11

^a ADD_{der} was calculated from the mean of 35 individual field average concentrations (C_{der-crumb rubber}) of a chemical detected in artificial sweat extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-245. Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—
Combined Gender **Spectators**

Chemical Name	CASRN	ADD _{der}								
		Third Trimester Fetus	0<2 Years	2<6 Years	6<11 Years	11<16 Years	16<30 Years	30<40 Years	40<50 Years	50<70 Years
Acenaphthylene	208-96-8	2.4E-12	1.3E-11	9.3E-12	8.4E-12	7.2E-12	3.2E-12	2.7E-12	2.8E-12	2.6E-12
Aniline	62-53-3	7.6E-11	4.1E-10	3.0E-10	2.7E-10	2.3E-10	1.0E-10	8.7E-11	8.8E-11	8.3E-11
Anthracene	120-12-7	2.6E-11	1.4E-10	1.0E-10	9.3E-11	8.0E-11	3.5E-11	3.0E-11	3.1E-11	2.9E-11



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical Name	CASRN	ADD _{der}								
		Third Trimester Fetus	0<2 Years	2<6 Years	6<11 Years	11<16 Years	16<30 Years	30<40 Years	40<50 Years	50<70 Years
Anthracene, 2-methyl-	613-12-7	3.8E-11	2.0E-10	1.5E-10	1.3E-10	1.2E-10	5.0E-11	4.4E-11	4.4E-11	4.1E-11
Anthracene, 9,10-diphenyl-	1499-10-1	5.1E-12	2.8E-11	2.0E-11	1.8E-11	1.6E-11	6.9E-12	6.0E-12	6.0E-12	5.7E-12
Anthracene, 9-phenyl	602-55-1	1.3E-11	6.9E-11	5.0E-11	4.6E-11	3.9E-11	1.7E-11	1.5E-11	1.5E-11	1.4E-11
Benz[a]anthracene	56-55-3	1.0E-10	5.6E-10	4.1E-10	3.7E-10	3.2E-10	1.4E-10	1.2E-10	1.2E-10	1.1E-10
Benzene, n-butyl-	104-51-8	1.6E-13	8.3E-13	6.1E-13	5.6E-13	4.8E-13	2.1E-13	1.8E-13	1.8E-13	1.7E-13
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	7.0E-09	3.8E-08	2.8E-08	2.5E-08	2.1E-08	9.4E-09	8.1E-09	8.2E-09	7.7E-09
Benzo[b]fluoranthene	205-99-2	1.5E-10	7.9E-10	5.8E-10	5.3E-10	4.5E-10	2.0E-10	1.7E-10	1.7E-10	1.6E-10
7H-Benzo[c]fluorene	205-12-9	1.3E-11	6.9E-11	5.1E-11	4.6E-11	3.9E-11	1.7E-11	1.5E-11	1.5E-11	1.4E-11
Benzo[k]fluoranthene	207-08-9	6.4E-11	3.4E-10	2.5E-10	2.3E-10	2.0E-10	8.6E-11	7.4E-11	7.5E-11	7.1E-11
Benzothiazole	95-16-9	8.4E-08	4.5E-07	3.3E-07	3.0E-07	2.6E-07	1.1E-07	9.7E-08	9.8E-08	9.2E-08
Benzothiazole, 2-phenyl-	883-93-2	2.0E-09	1.1E-08	8.0E-09	7.3E-09	6.2E-09	2.7E-09	2.4E-09	2.4E-09	2.2E-09
1,3-Benzothiazole-2-thiol	149-30-4	5.6E-10	3.0E-09	2.2E-09	2.0E-09	1.7E-09	7.4E-10	6.4E-10	6.5E-10	6.1E-10
Benzothiazolone	934-34-9	2.3E-07	1.3E-06	9.3E-07	8.4E-07	7.2E-07	3.1E-07	2.7E-07	2.7E-07	2.6E-07
Benzyl butyl phthalate	85-68-7	9.5E-10	5.1E-09	3.8E-09	3.4E-09	2.9E-09	1.3E-09	1.1E-09	1.1E-09	1.1E-09
Cyclohexyl isothiocyanate	1122-82-3	3.5E-12	1.9E-11	1.4E-11	1.3E-11	1.1E-11	4.7E-12	4.1E-12	4.1E-12	3.9E-12
Dibenz[a,h]anthracene	53-70-3	3.8E-11	2.0E-10	1.5E-10	1.4E-10	1.2E-10	5.1E-11	4.4E-11	4.5E-11	4.2E-11
Dibenzothiophene	132-65-0	2.3E-11	1.2E-10	9.0E-11	8.2E-11	7.0E-11	3.1E-11	2.7E-11	2.7E-11	2.5E-11
Diethyl Phthalate	84-66-2	1.7E-10	9.3E-10	6.9E-10	6.2E-10	5.3E-10	2.3E-10	2.0E-10	2.0E-10	1.9E-10
Diisobutyl Phthalate	84-69-5	1.5E-10	7.8E-10	5.7E-10	5.2E-10	4.5E-10	2.0E-10	1.7E-10	1.7E-10	1.6E-10
Diisooctylphthalate	27554-26-3	4.8E-09	2.6E-08	1.9E-08	1.7E-08	1.5E-08	6.4E-09	5.6E-09	5.6E-09	5.3E-09
Di-n-octyl phthalate	117-84-0	1.0E-09	5.5E-09	4.0E-09	3.6E-09	3.1E-09	1.4E-09	1.2E-09	1.2E-09	1.1E-09
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	9.0E-13	4.8E-12	3.5E-12	3.2E-12	2.7E-12	1.2E-12	1.0E-12	1.0E-12	9.9E-13
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	4.5E-09	2.4E-08	1.8E-08	1.6E-08	1.4E-08	6.0E-09	5.2E-09	5.2E-09	4.9E-09
Fluoranthene	206-44-0	8.0E-10	4.3E-09	3.1E-09	2.8E-09	2.4E-09	1.1E-09	9.2E-10	9.3E-10	8.8E-10
Fluorene	86-73-7	3.4E-12	1.8E-11	1.3E-11	1.2E-11	1.0E-11	4.5E-12	3.9E-12	3.9E-12	3.7E-12
Hexanoic Acid, 2-ethyl	149-57-5	3.9E-12	2.1E-11	1.5E-11	1.4E-11	1.2E-11	5.2E-12	4.5E-12	4.5E-12	4.2E-12
1-Hydroxypyrene	5315-79-7	2.2E-10	1.2E-09	8.8E-10	8.0E-10	6.9E-10	3.0E-10	2.6E-10	2.6E-10	2.5E-10
2-(Methylthio)benzothiazole	615-22-5	3.1E-09	1.6E-08	1.2E-08	1.1E-08	9.4E-09	4.1E-09	3.6E-09	3.6E-09	3.4E-09
Naphthalene	91-20-3	4.9E-12	2.6E-11	1.9E-11	1.8E-11	1.5E-11	6.6E-12	5.7E-12	5.8E-12	5.4E-12
Naphthalene, 1,2-dimethyl-	573-98-8	3.5E-13	1.9E-12	1.4E-12	1.2E-12	1.1E-12	4.7E-13	4.0E-13	4.1E-13	3.8E-13
Naphthalene, 1,6-dimethyl-	575-43-9	3.7E-12	2.0E-11	1.4E-11	1.3E-11	1.1E-11	4.9E-12	4.3E-12	4.3E-12	4.1E-12
Naphthalene, 1-methyl-	90-12-0	7.8E-13	4.2E-12	3.1E-12	2.8E-12	2.4E-12	1.0E-12	9.0E-13	9.1E-13	8.6E-13
Naphthalene, 2-(bromomethyl)-	939-26-4	1.4E-10	7.7E-10	5.7E-10	5.2E-10	4.4E-10	1.9E-10	1.7E-10	1.7E-10	1.6E-10
Naphthalene, 2,3-dimethyl-	581-40-8	4.3E-11	2.3E-10	1.7E-10	1.5E-10	1.3E-10	5.7E-11	5.0E-11	5.0E-11	4.7E-11
Naphthalene, 2-methyl	91-57-6	7.6E-13	4.1E-12	3.0E-12	2.7E-12	2.3E-12	1.0E-12	8.8E-13	8.9E-13	8.3E-13
1-Octadecene	112-88-9	2.5E-10	1.3E-09	9.7E-10	8.8E-10	7.6E-10	3.3E-10	2.9E-10	2.9E-10	2.7E-10
Phenanthrene	85-01-8	1.7E-10	9.1E-10	6.7E-10	6.1E-10	5.2E-10	2.3E-10	2.0E-10	2.0E-10	1.9E-10
Phenanthrene, 1-methyl	832-69-9	8.9E-11	4.8E-10	3.5E-10	3.2E-10	2.7E-10	1.2E-10	1.0E-10	1.0E-10	9.8E-11



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical Name	CASRN	ADD _{der}								
		Third Trimester Fetus	0<2 Years	2<6 Years	6<11 Years	11<16 Years	16<30 Years	30<40 Years	40<50 Years	50<70 Years
Phenanthrene, 2-methyl-	2531-84-2	1.1E-10	5.8E-10	4.3E-10	3.9E-10	3.3E-10	1.4E-10	1.2E-10	1.3E-10	1.2E-10
Phenanthrene, 3-methyl	832-71-3	3.9E-11	2.1E-10	1.6E-10	1.4E-10	1.2E-10	5.3E-11	4.5E-11	4.6E-11	4.3E-11
N-Phenylbenzamide	93-98-1	2.5E-10	1.3E-09	9.7E-10	8.8E-10	7.5E-10	3.3E-10	2.8E-10	2.9E-10	2.7E-10
Phthalimide	85-41-6	1.6E-09	8.6E-09	6.3E-09	5.7E-09	4.9E-09	2.1E-09	1.9E-09	1.9E-09	1.8E-09
Pyrene	129-00-0	2.2E-09	1.2E-08	8.5E-09	7.7E-09	6.6E-09	2.9E-09	2.5E-09	2.5E-09	2.4E-09
Pyridine, 2-(4-methylphenyl)-	4467-06-5	7.9E-11	4.3E-10	3.1E-10	2.8E-10	2.4E-10	1.1E-10	9.2E-11	9.3E-11	8.7E-11
Triethylene glycol monobutyl ether	143-22-6	4.1E-10	2.2E-09	1.6E-09	1.5E-09	1.3E-09	5.5E-10	4.8E-10	4.8E-10	4.5E-10
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	5.5E-11	3.0E-10	2.2E-10	2.0E-10	1.7E-10	7.4E-11	6.4E-11	6.5E-11	6.1E-11

^a ADD_{der} was calculated from the mean of 35 individual field average concentrations (C_{der-crumb rubber}) of a chemical detected in artificial sweat extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

INDIVIDUAL FIELD ASSESSMENT (Table F-246 to Table F-269)

Table F-246. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Athletes 2<6 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	8.0E-12	1.4E-11	0	3.3E-11	5.0E-11
Aniline	62-53-3	4	0	2.6E-10	8.0E-10	0	2.0E-09	3.8E-09
Anthracene	120-12-7	20	0	8.9E-11	1.2E-10	6.1E-11	2.2E-10	5.9E-10
Anthracene, 2-methyl-	613-12-7	35	3.1E-11	1.3E-10	1.0E-10	8.4E-11	2.9E-10	5.0E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	1.7E-11	4.9E-11	0	1.1E-10	2.4E-10
Anthracene, 9-phenyl	602-55-1	9	0	4.3E-11	9.8E-11	0	2.4E-10	4.3E-10
Benz[a]anthracene	56-55-3	31	0	3.5E-10	4.1E-10	1.9E-10	9.9E-10	1.9E-09
Benzene, n-butyl-	104-51-8	1	0	5.3E-13	3.1E-12	0	0	1.8E-11
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	2.4E-08	3.8E-08	9.9E-09	7.3E-08	2.0E-07
Benzo[b]fluoranthene	205-99-2	34	0	5.0E-10	4.4E-10	3.9E-10	1.2E-09	1.9E-09
7H-Benzo[c]fluorene	205-12-9	26	0	4.4E-11	5.5E-11	2.2E-11	1.4E-10	2.3E-10
Benzo[k]fluoranthene	207-08-9	29	0	2.2E-10	2.3E-10	1.3E-10	7.1E-10	8.8E-10
Benzothiazole	95-16-9	35	6.5E-08	2.8E-07	1.5E-07	2.4E-07	5.6E-07	6.4E-07
Benzothiazole, 2-phenyl-	883-93-2	35	2.0E-09	6.9E-09	5.4E-09	4.9E-09	1.9E-08	2.7E-08
1,3-Benzothiazole-2-thiol	149-30-4	1	0	1.9E-09	1.1E-08	0	0	6.6E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzothiazolone	934-34-9	35	1.7E-07	8.0E-07	2.5E-07	9.0E-07	1.1E-06	1.1E-06
Benzyl butyl phthalate	85-68-7	35	2.2E-10	3.2E-09	2.3E-09	2.7E-09	8.0E-09	9.2E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	1.2E-11	7.1E-11	0	0	4.2E-10
Dibenz[a,h]anthracene	53-70-3	14	0	1.3E-10	2.4E-10	0	6.3E-10	9.9E-10
Dibenzothiophene	132-65-0	20	0	7.8E-11	9.6E-11	6.5E-11	2.1E-10	4.6E-10
Diethyl Phthalate	84-66-2	5	0	5.9E-10	1.8E-09	0	3.3E-09	8.9E-09
Diisobutyl Phthalate	84-69-5	28	0	4.9E-10	8.3E-10	3.3E-10	1.3E-09	4.8E-09
Diisooctylphthalate	27554-26-3	35	1.4E-10	1.6E-08	1.1E-08	1.4E-08	4.0E-08	4.7E-08
Di-n-octyl phthalate	117-84-0	33	0	3.5E-09	4.9E-09	1.9E-09	1.2E-08	2.3E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	3.0E-12	9.0E-12	0	2.4E-11	4.0E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	1.5E-08	3.9E-08	3.5E-09	4.1E-08	2.3E-07
Fluoranthene	206-44-0	35	4.4E-10	2.7E-09	2.1E-09	1.9E-09	7.0E-09	7.4E-09
Fluorene	86-73-7	6	0	1.1E-11	3.5E-11	0	4.9E-11	2.0E-10
Hexanoic Acid, 2-ethyl	149-57-5	1	0	1.3E-11	7.7E-11	0	0	4.6E-10
1-Hydroxypyrene	5315-79-7	1	0	7.6E-10	4.5E-09	0	0	2.7E-08
2-(Methylthio)benzothiazole	615-22-5	9	0	1.0E-08	2.3E-08	0	4.8E-08	1.0E-07
Naphthalene	91-20-3	2	0	1.7E-11	6.9E-11	0	8.4E-11	3.0E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	1.2E-12	7.0E-12	0	0	4.1E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	1.2E-11	3.0E-11	0	9.0E-11	9.2E-11
Naphthalene, 1-methyl-	90-12-0	1	0	2.6E-12	1.6E-11	0	0	9.3E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	4.9E-10	3.9E-10	3.0E-10	1.2E-09	1.5E-09
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	1.5E-10	1.3E-10	8.7E-11	4.2E-10	5.8E-10
Naphthalene, 2-methyl	91-57-6	1	0	2.6E-12	1.5E-11	0	0	9.0E-11
1-Octadecene	112-88-9	27	0	8.4E-10	2.3E-09	2.3E-10	1.5E-09	1.4E-08
Phenanthrene	85-01-8	29	0	5.8E-10	7.8E-10	2.8E-10	1.9E-09	3.6E-09
Phenanthrene, 1-methyl	832-69-9	35	5.6E-11	3.0E-10	2.3E-10	2.4E-10	7.3E-10	9.2E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	3.7E-10	3.4E-10	2.8E-10	1.0E-09	1.4E-09
Phenanthrene, 3-methyl	832-71-3	3	0	1.3E-10	5.0E-10	0	9.5E-10	2.7E-09
N-Phenylbenzamide	93-98-1	8	0	8.3E-10	1.7E-09	0	4.0E-09	6.0E-09
Phthalimide	85-41-6	26	0	5.4E-09	1.3E-08	7.4E-11	3.1E-08	5.4E-08
Pyrene	129-00-0	35	1.3E-09	7.3E-09	4.7E-09	6.4E-09	1.7E-08	1.9E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	2.7E-10	5.1E-10	0	1.1E-09	2.3E-09



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Triethylene glycol monobutyl ether	143-22-6	7	0	1.4E-09	3.4E-09	0	8.8E-09	1.5E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	1.9E-10	2.3E-10	6.7E-11	5.6E-10	9.0E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-247. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Athletes 6<11 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	6.3E-12	1.1E-11	0	2.6E-11	3.9E-11
Aniline	62-53-3	4	0	2.0E-10	6.3E-10	0	1.6E-09	3.0E-09
Anthracene	120-12-7	20	0	7.0E-11	9.2E-11	4.8E-11	1.7E-10	4.7E-10
Anthracene, 2-methyl-	613-12-7	35	2.4E-11	1.0E-10	8.2E-11	6.7E-11	2.3E-10	4.0E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	1.4E-11	3.9E-11	0	8.4E-11	1.9E-10
Anthracene, 9-phenyl	602-55-1	9	0	3.4E-11	7.8E-11	0	1.9E-10	3.4E-10
Benz[a]anthracene	56-55-3	31	0	2.8E-10	3.2E-10	1.5E-10	7.8E-10	1.5E-09
Benzene, n-butyl-	104-51-8	1	0	4.2E-13	2.5E-12	0	0	1.5E-11
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	1.9E-08	3.0E-08	7.8E-09	5.8E-08	1.6E-07
Benzo[b]fluoranthene	205-99-2	34	0	3.9E-10	3.5E-10	3.1E-10	9.9E-10	1.5E-09
7H-Benzo[c]fluorene	205-12-9	26	0	3.4E-11	4.3E-11	1.8E-11	1.1E-10	1.8E-10
Benzo[k]fluoranthene	207-08-9	29	0	1.7E-10	1.8E-10	1.0E-10	5.6E-10	6.9E-10
Benzothiazole	95-16-9	35	5.2E-08	2.2E-07	1.2E-07	1.9E-07	4.4E-07	5.0E-07
Benzothiazole, 2-phenyl-	883-93-2	35	1.6E-09	5.5E-09	4.3E-09	3.9E-09	1.5E-08	2.1E-08
1,3-Benzothiazole-2-thiol	149-30-4	1	0	1.5E-09	8.8E-09	0	0	5.2E-08
Benzothiazolone	934-34-9	35	1.4E-07	6.3E-07	2.0E-07	7.1E-07	8.8E-07	9.0E-07
Benzyl butyl phthalate	85-68-7	35	1.7E-10	2.6E-09	1.8E-09	2.2E-09	6.4E-09	7.3E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	9.5E-12	5.6E-11	0	0	3.3E-10
Dibenz[a,h]anthracene	53-70-3	14	0	1.0E-10	1.9E-10	0	5.0E-10	7.8E-10
Dibenzothiophene	132-65-0	20	0	6.1E-11	7.6E-11	5.1E-11	1.7E-10	3.6E-10
Diethyl Phthalate	84-66-2	5	0	4.7E-10	1.4E-09	0	2.6E-09	7.0E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Diisobutyl Phthalate	84-69-5	28	0	3.9E-10	6.6E-10	2.6E-10	1.0E-09	3.8E-09
Diisooctylphthalate	27554-26-3	35	1.1E-10	1.3E-08	8.9E-09	1.1E-08	3.2E-08	3.7E-08
Di-n-octyl phthalate	117-84-0	33	0	2.7E-09	3.9E-09	1.5E-09	9.4E-09	1.9E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	2.4E-12	7.1E-12	0	1.9E-11	3.1E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	1.2E-08	3.1E-08	2.8E-09	3.2E-08	1.8E-07
Fluoranthene	206-44-0	35	3.5E-10	2.1E-09	1.7E-09	1.5E-09	5.5E-09	5.9E-09
Fluorene	86-73-7	6	0	9.0E-12	2.8E-11	0	3.8E-11	1.5E-10
Hexanoic Acid, 2-ethyl	149-57-5	1	0	1.0E-11	6.1E-11	0	0	3.6E-10
1-Hydroxypyrene	5315-79-7	1	0	6.0E-10	3.6E-09	0	0	2.1E-08
2-(Methylthio)benzothiazole	615-22-5	9	0	8.2E-09	1.8E-08	0	3.8E-08	8.3E-08
Naphthalene	91-20-3	2	0	1.3E-11	5.5E-11	0	6.7E-11	2.4E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	9.3E-13	5.5E-12	0	0	3.3E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	9.9E-12	2.3E-11	0	7.1E-11	7.3E-11
Naphthalene, 1-methyl-	90-12-0	1	0	2.1E-12	1.2E-11	0	0	7.3E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	3.9E-10	3.1E-10	2.4E-10	9.2E-10	1.2E-09
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	1.1E-10	1.0E-10	6.8E-11	3.3E-10	4.6E-10
Naphthalene, 2-methyl	91-57-6	1	0	2.0E-12	1.2E-11	0	0	7.1E-11
1-Octadecene	112-88-9	27	0	6.6E-10	1.8E-09	1.8E-10	1.2E-09	1.1E-08
Phenanthrene	85-01-8	29	0	4.6E-10	6.2E-10	2.2E-10	1.5E-09	2.9E-09
Phenanthrene, 1-methyl	832-69-9	35	4.5E-11	2.4E-10	1.8E-10	1.9E-10	5.8E-10	7.3E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	2.9E-10	2.7E-10	2.2E-10	8.1E-10	1.1E-09
Phenanthrene, 3-methyl	832-71-3	3	0	1.1E-10	4.0E-10	0	7.5E-10	2.1E-09
N-Phenylbenzamide	93-98-1	8	0	6.6E-10	1.3E-09	0	3.2E-09	4.8E-09
Phthalimide	85-41-6	26	0	4.3E-09	9.9E-09	5.8E-11	2.4E-08	4.3E-08
Pyrene	129-00-0	35	1.1E-09	5.8E-09	3.7E-09	5.0E-09	1.4E-08	1.5E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	2.1E-10	4.1E-10	0	9.0E-10	1.8E-09
Triethylene glycol monobutyl ether	143-22-6	7	0	1.1E-09	2.7E-09	0	7.0E-09	1.2E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	1.5E-10	1.8E-10	5.3E-11	4.4E-10	7.2E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.



CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures.

Table F-248. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Athletes 11<16 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	6.7E-12	1.1E-11	0	2.7E-11	4.1E-11
Aniline	62-53-3	4	0	2.1E-10	6.6E-10	0	1.6E-09	3.2E-09
Anthracene	120-12-7	20	0	7.4E-11	9.7E-11	5.0E-11	1.8E-10	4.9E-10
Anthracene, 2-methyl-	613-12-7	35	2.6E-11	1.1E-10	8.7E-11	7.0E-11	2.5E-10	4.2E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	1.5E-11	4.1E-11	0	8.9E-11	2.0E-10
Anthracene, 9-phenyl	602-55-1	9	0	3.6E-11	8.2E-11	0	2.0E-10	3.6E-10
Benz[a]anthracene	56-55-3	31	0	2.9E-10	3.4E-10	1.5E-10	8.2E-10	1.6E-09
Benzene, n-butyl-	104-51-8	1	0	4.4E-13	2.6E-12	0	0	1.5E-11
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	2.0E-08	3.2E-08	8.2E-09	6.1E-08	1.7E-07
Benzo[b]fluoranthene	205-99-2	34	0	4.2E-10	3.7E-10	3.3E-10	1.0E-09	1.6E-09
7H-Benzo[c]fluorene	205-12-9	26	0	3.6E-11	4.6E-11	1.9E-11	1.2E-10	1.9E-10
Benzo[k]fluoranthene	207-08-9	29	0	1.8E-10	1.9E-10	1.1E-10	5.9E-10	7.3E-10
Benzothiazole	95-16-9	35	5.4E-08	2.4E-07	1.2E-07	2.0E-07	4.6E-07	5.3E-07
Benzothiazole, 2-phenyl-	883-93-2	35	1.7E-09	5.8E-09	4.5E-09	4.1E-09	1.5E-08	2.3E-08
1,3-Benzothiazole-2-thiol	149-30-4	1	0	1.6E-09	9.3E-09	0	0	5.5E-08
Benzothiazolone	934-34-9	35	1.4E-07	6.6E-07	2.1E-07	7.5E-07	9.2E-07	9.5E-07
Benzyl butyl phthalate	85-68-7	35	1.8E-10	2.7E-09	1.9E-09	2.3E-09	6.7E-09	7.7E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	1.0E-11	5.9E-11	0	0	3.5E-10
Dibenz[a,h]anthracene	53-70-3	14	0	1.1E-10	2.0E-10	0	5.3E-10	8.2E-10
Dibenzothiophene	132-65-0	20	0	6.5E-11	8.0E-11	5.4E-11	1.8E-10	3.8E-10
Diethyl Phthalate	84-66-2	5	0	4.9E-10	1.5E-09	0	2.7E-09	7.4E-09
Diisobutyl Phthalate	84-69-5	28	0	4.1E-10	6.9E-10	2.8E-10	1.1E-09	4.0E-09
Diisooctylphthalate	27554-26-3	35	1.2E-10	1.4E-08	9.4E-09	1.2E-08	3.4E-08	3.9E-08
Di-n-octyl phthalate	117-84-0	33	0	2.9E-09	4.1E-09	1.6E-09	9.9E-09	2.0E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	2.5E-12	7.5E-12	0	2.0E-11	3.3E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	1.3E-08	3.3E-08	2.9E-09	3.4E-08	1.9E-07
Fluoranthene	206-44-0	35	3.7E-10	2.3E-09	1.8E-09	1.6E-09	5.8E-09	6.2E-09
Fluorene	86-73-7	6	0	9.5E-12	3.0E-11	0	4.1E-11	1.6E-10
Hexanoic Acid, 2-ethyl	149-57-5	1	0	1.1E-11	6.5E-11	0	0	3.8E-10



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
1-Hydroxypyrene	5315-79-7	1	0	6.3E-10	3.7E-09	0	0	2.2E-08
2-(Methylthio)benzothiazole	615-22-5	9	0	8.7E-09	1.9E-08	0	4.0E-08	8.7E-08
Naphthalene	91-20-3	2	0	1.4E-11	5.8E-11	0	7.0E-11	2.5E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	9.8E-13	5.8E-12	0	0	3.4E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	1.0E-11	2.5E-11	0	7.5E-11	7.7E-11
Naphthalene, 1-methyl-	90-12-0	1	0	2.2E-12	1.3E-11	0	0	7.7E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	4.1E-10	3.3E-10	2.5E-10	9.8E-10	1.3E-09
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	1.2E-10	1.1E-10	7.2E-11	3.5E-10	4.9E-10
Naphthalene, 2-methyl	91-57-6	1	0	2.1E-12	1.3E-11	0	0	7.5E-11
1-Octadecene	112-88-9	27	0	7.0E-10	1.9E-09	1.9E-10	1.2E-09	1.2E-08
Phenanthrene	85-01-8	29	0	4.8E-10	6.5E-10	2.3E-10	1.6E-09	3.0E-09
Phenanthrene, 1-methyl	832-69-9	35	4.7E-11	2.5E-10	1.9E-10	2.0E-10	6.1E-10	7.7E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	3.1E-10	2.8E-10	2.3E-10	8.5E-10	1.1E-09
Phenanthrene, 3-methyl	832-71-3	3	0	1.1E-10	4.2E-10	0	7.9E-10	2.2E-09
N-Phenylbenzamide	93-98-1	8	0	7.0E-10	1.4E-09	0	3.3E-09	5.0E-09
Phthalimide	85-41-6	26	0	4.5E-09	1.0E-08	6.2E-11	2.6E-08	4.5E-08
Pyrene	129-00-0	35	1.1E-09	6.1E-09	4.0E-09	5.3E-09	1.4E-08	1.6E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	2.2E-10	4.3E-10	0	9.5E-10	1.9E-09
Triethylene glycol monobutyl ether	143-22-6	7	0	1.2E-09	2.8E-09	0	7.3E-09	1.3E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	1.6E-10	1.9E-10	5.6E-11	4.7E-10	7.5E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-249. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Athletes 16<30 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	8.4E-12	1.4E-11	0	3.5E-11	5.2E-11



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	2.7E-10	8.4E-10	0	2.1E-09	4.0E-09
Anthracene	120-12-7	20	0	9.3E-11	1.2E-10	6.4E-11	2.3E-10	6.2E-10
Anthracene, 2-methyl-	613-12-7	35	3.2E-11	1.3E-10	1.1E-10	8.9E-11	3.1E-10	5.3E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	1.8E-11	5.2E-11	0	1.1E-10	2.5E-10
Anthracene, 9-phenyl	602-55-1	9	0	4.6E-11	1.0E-10	0	2.6E-10	4.6E-10
Benz[a]anthracene	56-55-3	31	0	3.7E-10	4.3E-10	2.0E-10	1.0E-09	2.0E-09
Benzene, n-butyl-	104-51-8	1	0	5.5E-13	3.3E-12	0	0	1.9E-11
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	2.5E-08	4.0E-08	1.0E-08	7.7E-08	2.2E-07
Benzo[b]fluoranthene	205-99-2	34	0	5.2E-10	4.6E-10	4.1E-10	1.3E-09	2.0E-09
7H-Benzo[c]fluorene	205-12-9	26	0	4.6E-11	5.8E-11	2.4E-11	1.5E-10	2.4E-10
Benzo[k]fluoranthene	207-08-9	29	0	2.3E-10	2.4E-10	1.4E-10	7.4E-10	9.2E-10
Benzothiazole	95-16-9	35	6.9E-08	3.0E-07	1.5E-07	2.5E-07	5.8E-07	6.7E-07
Benzothiazole, 2-phenyl-	883-93-2	35	2.1E-09	7.3E-09	5.7E-09	5.2E-09	2.0E-08	2.8E-08
1,3-Benzothiazole-2-thiol	149-30-4	1	0	2.0E-09	1.2E-08	0	0	6.9E-08
Benzothiazolone	934-34-9	35	1.8E-07	8.4E-07	2.7E-07	9.4E-07	1.2E-06	1.2E-06
Benzyl butyl phthalate	85-68-7	35	2.3E-10	3.4E-09	2.4E-09	2.9E-09	8.5E-09	9.7E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	1.3E-11	7.4E-11	0	0	4.4E-10
Dibenz[a,h]anthracene	53-70-3	14	0	1.4E-10	2.5E-10	0	6.6E-10	1.0E-09
Dibenzothiophene	132-65-0	20	0	8.2E-11	1.0E-10	6.9E-11	2.3E-10	4.9E-10
Diethyl Phthalate	84-66-2	5	0	6.2E-10	1.8E-09	0	3.4E-09	9.3E-09
Diisobutyl Phthalate	84-69-5	28	0	5.2E-10	8.8E-10	3.5E-10	1.4E-09	5.1E-09
Diisooctylphthalate	27554-26-3	35	1.5E-10	1.7E-08	1.2E-08	1.5E-08	4.2E-08	4.9E-08
Di-n-octyl phthalate	117-84-0	33	0	3.6E-09	5.1E-09	2.0E-09	1.3E-08	2.5E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	3.2E-12	9.5E-12	0	2.5E-11	4.2E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	1.6E-08	4.1E-08	3.7E-09	4.3E-08	2.4E-07
Fluoranthene	206-44-0	35	4.6E-10	2.8E-09	2.2E-09	2.0E-09	7.3E-09	7.8E-09
Fluorene	86-73-7	6	0	1.2E-11	3.7E-11	0	5.1E-11	2.1E-10
Hexanoic Acid, 2-ethyl	149-57-5	1	0	1.4E-11	8.1E-11	0	0	4.8E-10
1-Hydroxypyrene	5315-79-7	1	0	8.0E-10	4.7E-09	0	0	2.8E-08
2-(Methylthio)benzothiazole	615-22-5	9	0	1.1E-08	2.4E-08	0	5.1E-08	1.1E-07
Naphthalene	91-20-3	2	0	1.8E-11	7.3E-11	0	8.9E-11	3.2E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	1.2E-12	7.3E-12	0	0	4.3E-11



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	1.3E-11	3.1E-11	0	9.4E-11	9.7E-11
Naphthalene, 1-methyl-	90-12-0	1	0	2.8E-12	1.6E-11	0	0	9.7E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	5.2E-10	4.2E-10	3.2E-10	1.2E-09	1.6E-09
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	1.5E-10	1.3E-10	9.1E-11	4.4E-10	6.1E-10
Naphthalene, 2-methyl	91-57-6	1	0	2.7E-12	1.6E-11	0	0	9.5E-11
1-Octadecene	112-88-9	27	0	8.8E-10	2.4E-09	2.4E-10	1.5E-09	1.5E-08
Phenanthrene	85-01-8	29	0	6.1E-10	8.2E-10	3.0E-10	2.0E-09	3.8E-09
Phenanthrene, 1-methyl	832-69-9	35	5.9E-11	3.2E-10	2.4E-10	2.5E-10	7.7E-10	9.7E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	3.9E-10	3.6E-10	2.9E-10	1.1E-09	1.4E-09
Phenanthrene, 3-methyl	832-71-3	3	0	1.4E-10	5.3E-10	0	1.0E-09	2.8E-09
N-Phenylbenzamide	93-98-1	8	0	8.8E-10	1.8E-09	0	4.2E-09	6.4E-09
Phthalimide	85-41-6	26	0	5.7E-09	1.3E-08	7.8E-11	3.2E-08	5.7E-08
Pyrene	129-00-0	35	1.4E-09	7.7E-09	5.0E-09	6.7E-09	1.8E-08	2.0E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	2.8E-10	5.4E-10	0	1.2E-09	2.5E-09
Triethylene glycol monobutyl ether	143-22-6	7	0	1.5E-09	3.6E-09	0	9.3E-09	1.6E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	2.0E-10	2.4E-10	7.0E-11	5.9E-10	9.5E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-250. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Athletes 30<40 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	6.0E-12	1.0E-11	0	2.5E-11	3.7E-11
Aniline	62-53-3	4	0	1.9E-10	5.9E-10	0	1.5E-09	2.8E-09
Anthracene	120-12-7	20	0	6.6E-11	8.7E-11	4.5E-11	1.6E-10	4.4E-10
Anthracene, 2-methyl-	613-12-7	35	2.3E-11	9.5E-11	7.8E-11	6.3E-11	2.2E-10	3.7E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	1.3E-11	3.7E-11	0	7.9E-11	1.8E-10
Anthracene, 9-phenyl	602-55-1	9	0	3.2E-11	7.3E-11	0	1.8E-10	3.2E-10
Benz[a]anthracene	56-55-3	31	0	2.6E-10	3.0E-10	1.4E-10	7.4E-10	1.4E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene, n-butyl-	104-51-8	1	0	3.9E-13	2.3E-12	0	0	1.4E-11
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	1.8E-08	2.8E-08	7.4E-09	5.5E-08	1.5E-07
Benzo[b]fluoranthene	205-99-2	34	0	3.7E-10	3.3E-10	2.9E-10	9.3E-10	1.4E-09
7H-Benzo[c]fluorene	205-12-9	26	0	3.2E-11	4.1E-11	1.7E-11	1.0E-10	1.7E-10
Benzo[k]fluoranthene	207-08-9	29	0	1.6E-10	1.7E-10	9.9E-11	5.3E-10	6.6E-10
Benzothiazole	95-16-9	35	4.9E-08	2.1E-07	1.1E-07	1.8E-07	4.2E-07	4.8E-07
Benzothiazole, 2-phenyl-	883-93-2	35	1.5E-09	5.2E-09	4.0E-09	3.7E-09	1.4E-08	2.0E-08
1,3-Benzothiazole-2-thiol	149-30-4	1	0	1.4E-09	8.3E-09	0	0	4.9E-08
Benzothiazolone	934-34-9	35	1.3E-07	5.9E-07	1.9E-07	6.7E-07	8.3E-07	8.5E-07
Benzyl butyl phthalate	85-68-7	35	1.6E-10	2.4E-09	1.7E-09	2.0E-09	6.0E-09	6.9E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	8.9E-12	5.3E-11	0	0	3.1E-10
Dibenz[a,h]anthracene	53-70-3	14	0	9.7E-11	1.8E-10	0	4.7E-10	7.4E-10
Dibenzothiophene	132-65-0	20	0	5.8E-11	7.2E-11	4.9E-11	1.6E-10	3.4E-10
Diethyl Phthalate	84-66-2	5	0	4.4E-10	1.3E-09	0	2.4E-09	6.6E-09
Diisobutyl Phthalate	84-69-5	28	0	3.7E-10	6.2E-10	2.5E-10	9.7E-10	3.6E-09
Diisooctylphthalate	27554-26-3	35	1.0E-10	1.2E-08	8.4E-09	1.0E-08	3.0E-08	3.5E-08
Di-n-octyl phthalate	117-84-0	33	0	2.6E-09	3.6E-09	1.4E-09	8.9E-09	1.8E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	2.3E-12	6.7E-12	0	1.8E-11	3.0E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	1.1E-08	2.9E-08	2.6E-09	3.1E-08	1.7E-07
Fluoranthene	206-44-0	35	3.3E-10	2.0E-09	1.6E-09	1.4E-09	5.2E-09	5.5E-09
Fluorene	86-73-7	6	0	8.5E-12	2.6E-11	0	3.6E-11	1.5E-10
Hexanoic Acid, 2-ethyl	149-57-5	1	0	9.8E-12	5.8E-11	0	0	3.4E-10
1-Hydroxypyrene	5315-79-7	1	0	5.7E-10	3.4E-09	0	0	2.0E-08
2-(Methylthio)benzothiazole	615-22-5	9	0	7.8E-09	1.7E-08	0	3.6E-08	7.8E-08
Naphthalene	91-20-3	2	0	1.2E-11	5.1E-11	0	6.3E-11	2.3E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	8.8E-13	5.2E-12	0	0	3.1E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	9.3E-12	2.2E-11	0	6.7E-11	6.9E-11
Naphthalene, 1-methyl-	90-12-0	1	0	2.0E-12	1.2E-11	0	0	6.9E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	3.7E-10	2.9E-10	2.3E-10	8.7E-10	1.1E-09
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	1.1E-10	9.6E-11	6.5E-11	3.1E-10	4.4E-10
Naphthalene, 2-methyl	91-57-6	1	0	1.9E-12	1.1E-11	0	0	6.7E-11



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
1-Octadecene	112-88-9	27	0	6.3E-10	1.7E-09	1.7E-10	1.1E-09	1.0E-08
Phenanthrene	85-01-8	29	0	4.3E-10	5.8E-10	2.1E-10	1.4E-09	2.7E-09
Phenanthrene, 1-methyl	832-69-9	35	4.2E-11	2.3E-10	1.7E-10	1.8E-10	5.5E-10	6.9E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	2.7E-10	2.5E-10	2.1E-10	7.6E-10	1.0E-09
Phenanthrene, 3-methyl	832-71-3	3	0	1.0E-10	3.7E-10	0	7.1E-10	2.0E-09
N-Phenylbenzamide	93-98-1	8	0	6.2E-10	1.3E-09	0	3.0E-09	4.5E-09
Phthalimide	85-41-6	26	0	4.0E-09	9.3E-09	5.5E-11	2.3E-08	4.0E-08
Pyrene	129-00-0	35	1.0E-09	5.5E-09	3.5E-09	4.7E-09	1.3E-08	1.4E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	2.0E-10	3.8E-10	0	8.5E-10	1.7E-09
Triethylene glycol monobutyl ether	143-22-6	7	0	1.0E-09	2.5E-09	0	6.6E-09	1.1E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	1.4E-10	1.7E-10	5.0E-11	4.2E-10	6.8E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-251. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Athletes 40<50 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	5.1E-12	8.7E-12	0	2.1E-11	3.2E-11
Aniline	62-53-3	4	0	1.6E-10	5.0E-10	0	1.2E-09	2.4E-09
Anthracene	120-12-7	20	0	5.6E-11	7.4E-11	3.8E-11	1.4E-10	3.8E-10
Anthracene, 2-methyl-	613-12-7	35	1.9E-11	8.1E-11	6.6E-11	5.3E-11	1.9E-10	3.2E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	1.1E-11	3.1E-11	0	6.7E-11	1.5E-10
Anthracene, 9-phenyl	602-55-1	9	0	2.7E-11	6.2E-11	0	1.5E-10	2.7E-10
Benz[a]anthracene	56-55-3	31	0	2.2E-10	2.6E-10	1.2E-10	6.3E-10	1.2E-09
Benzene, n-butyl-	104-51-8	1	0	3.3E-13	2.0E-12	0	0	1.2E-11
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	1.5E-08	2.4E-08	6.3E-09	4.6E-08	1.3E-07
Benzo[b]fluoranthene	205-99-2	34	0	3.2E-10	2.8E-10	2.5E-10	7.9E-10	1.2E-09
7H-Benzo[c]fluorene	205-12-9	26	0	2.8E-11	3.5E-11	1.4E-11	8.8E-11	1.5E-10
Benzo[k]fluoranthene	207-08-9	29	0	1.4E-10	1.5E-10	8.4E-11	4.5E-10	5.6E-10



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzothiazole	95-16-9	35	4.1E-08	1.8E-07	9.3E-08	1.5E-07	3.5E-07	4.0E-07
Benzothiazole, 2-phenyl-	883-93-2	35	1.3E-09	4.4E-09	3.4E-09	3.1E-09	1.2E-08	1.7E-08
1,3-Benzothiazole-2-thiol	149-30-4	1	0	1.2E-09	7.1E-09	0	0	4.2E-08
Benzothiazolone	934-34-9	35	1.1E-07	5.0E-07	1.6E-07	5.7E-07	7.0E-07	7.2E-07
Benzyl butyl phthalate	85-68-7	35	1.4E-10	2.1E-09	1.5E-09	1.7E-09	5.1E-09	5.8E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	7.6E-12	4.5E-11	0	0	2.7E-10
Dibenz[a,h]anthracene	53-70-3	14	0	8.2E-11	1.5E-10	0	4.0E-10	6.3E-10
Dibenzothiophene	132-65-0	20	0	4.9E-11	6.1E-11	4.1E-11	1.4E-10	2.9E-10
Diethyl Phthalate	84-66-2	5	0	3.7E-10	1.1E-09	0	2.1E-09	5.6E-09
Diisobutyl Phthalate	84-69-5	28	0	3.1E-10	5.3E-10	2.1E-10	8.2E-10	3.1E-09
Diisooctylphthalate	27554-26-3	35	8.8E-11	1.0E-08	7.2E-09	8.9E-09	2.6E-08	3.0E-08
Di-n-octyl phthalate	117-84-0	33	0	2.2E-09	3.1E-09	1.2E-09	7.5E-09	1.5E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.9E-12	5.7E-12	0	1.5E-11	2.5E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	9.6E-09	2.5E-08	2.2E-09	2.6E-08	1.4E-07
Fluoranthene	206-44-0	35	2.8E-10	1.7E-09	1.3E-09	1.2E-09	4.4E-09	4.7E-09
Fluorene	86-73-7	6	0	7.3E-12	2.2E-11	0	3.1E-11	1.2E-10
Hexanoic Acid, 2-ethyl	149-57-5	1	0	8.3E-12	4.9E-11	0	0	2.9E-10
1-Hydroxypyrene	5315-79-7	1	0	4.8E-10	2.8E-09	0	0	1.7E-08
2-(Methylthio)benzothiazole	615-22-5	9	0	6.6E-09	1.5E-08	0	3.1E-08	6.6E-08
Naphthalene	91-20-3	2	0	1.1E-11	4.4E-11	0	5.3E-11	1.9E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	7.5E-13	4.4E-12	0	0	2.6E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	7.9E-12	1.9E-11	0	5.7E-11	5.9E-11
Naphthalene, 1-methyl-	90-12-0	1	0	1.7E-12	9.9E-12	0	0	5.9E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	3.1E-10	2.5E-10	1.9E-10	7.4E-10	9.8E-10
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	9.2E-11	8.1E-11	5.5E-11	2.7E-10	3.7E-10
Naphthalene, 2-methyl	91-57-6	1	0	1.6E-12	9.6E-12	0	0	5.7E-11
1-Octadecene	112-88-9	27	0	5.3E-10	1.5E-09	1.5E-10	9.3E-10	8.8E-09
Phenanthrene	85-01-8	29	0	3.7E-10	5.0E-10	1.8E-10	1.2E-09	2.3E-09
Phenanthrene, 1-methyl	832-69-9	35	3.6E-11	1.9E-10	1.5E-10	1.5E-10	4.7E-10	5.9E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	2.3E-10	2.1E-10	1.8E-10	6.5E-10	8.7E-10
Phenanthrene, 3-methyl	832-71-3	3	0	8.5E-11	3.2E-10	0	6.0E-10	1.7E-09
N-Phenylbenzamide	93-98-1	8	0	5.3E-10	1.1E-09	0	2.5E-09	3.8E-09



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Phthalimide	85-41-6	26	0	3.4E-09	7.9E-09	4.7E-11	2.0E-08	3.4E-08
Pyrene	129-00-0	35	8.5E-10	4.6E-09	3.0E-09	4.0E-09	1.1E-08	1.2E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	1.7E-10	3.3E-10	0	7.2E-10	1.5E-09
Triethylene glycol monobutyl ether	143-22-6	7	0	8.9E-10	2.2E-09	0	5.6E-09	9.6E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	1.2E-10	1.4E-10	4.2E-11	3.5E-10	5.7E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-252. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Athletes 50<70 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	5.1E-12	8.8E-12	0	2.1E-11	3.2E-11
Aniline	62-53-3	4	0	1.6E-10	5.1E-10	0	1.3E-09	2.4E-09
Anthracene	120-12-7	20	0	5.7E-11	7.4E-11	3.9E-11	1.4E-10	3.8E-10
Anthracene, 2-methyl-	613-12-7	35	2.0E-11	8.1E-11	6.6E-11	5.4E-11	1.9E-10	3.2E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	1.1E-11	3.1E-11	0	6.8E-11	1.5E-10
Anthracene, 9-phenyl	602-55-1	9	0	2.8E-11	6.3E-11	0	1.6E-10	2.8E-10
Benz[a]anthracene	56-55-3	31	0	2.2E-10	2.6E-10	1.2E-10	6.3E-10	1.2E-09
Benzene, n-butyl-	104-51-8	1	0	3.4E-13	2.0E-12	0	0	1.2E-11
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	1.5E-08	2.4E-08	6.3E-09	4.7E-08	1.3E-07
Benzo[b]fluoranthene	205-99-2	34	0	3.2E-10	2.8E-10	2.5E-10	8.0E-10	1.2E-09
7H-Benzo[c]fluorene	205-12-9	26	0	2.8E-11	3.5E-11	1.4E-11	8.9E-11	1.5E-10
Benzo[k]fluoranthene	207-08-9	29	0	1.4E-10	1.5E-10	8.4E-11	4.5E-10	5.6E-10
Benzothiazole	95-16-9	35	4.2E-08	1.8E-07	9.3E-08	1.5E-07	3.6E-07	4.1E-07
Benzothiazole, 2-phenyl-	883-93-2	35	1.3E-09	4.4E-09	3.5E-09	3.1E-09	1.2E-08	1.7E-08
1,3-Benzothiazole-2-thiol	149-30-4	1	0	1.2E-09	7.1E-09	0	0	4.2E-08
Benzothiazolone	934-34-9	35	1.1E-07	5.1E-07	1.6E-07	5.7E-07	7.1E-07	7.2E-07
Benzyl butyl phthalate	85-68-7	35	1.4E-10	2.1E-09	1.5E-09	1.7E-09	5.1E-09	5.9E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Cyclohexyl isothiocyanate	1122-82-3	1	0	7.6E-12	4.5E-11	0	0	2.7E-10
Dibenz[a,h]anthracene	53-70-3	14	0	8.3E-11	1.5E-10	0	4.0E-10	6.3E-10
Dibenzothiophene	132-65-0	20	0	5.0E-11	6.1E-11	4.2E-11	1.4E-10	2.9E-10
Diethyl Phthalate	84-66-2	5	0	3.8E-10	1.1E-09	0	2.1E-09	5.7E-09
Diisobutyl Phthalate	84-69-5	28	0	3.2E-10	5.3E-10	2.1E-10	8.3E-10	3.1E-09
Diisooctylphthalate	27554-26-3	35	8.9E-11	1.0E-08	7.2E-09	8.9E-09	2.6E-08	3.0E-08
Di-n-octyl phthalate	117-84-0	33	0	2.2E-09	3.1E-09	1.2E-09	7.6E-09	1.5E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.9E-12	5.7E-12	0	1.5E-11	2.5E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	9.6E-09	2.5E-08	2.2E-09	2.6E-08	1.4E-07
Fluoranthene	206-44-0	35	2.8E-10	1.7E-09	1.4E-09	1.2E-09	4.5E-09	4.7E-09
Fluorene	86-73-7	6	0	7.3E-12	2.3E-11	0	3.1E-11	1.2E-10
Hexanoic Acid, 2-ethyl	149-57-5	1	0	8.4E-12	4.9E-11	0	0	2.9E-10
1-Hydroxypyrene	5315-79-7	1	0	4.8E-10	2.9E-09	0	0	1.7E-08
2-(Methylthio)benzothiazole	615-22-5	9	0	6.7E-09	1.5E-08	0	3.1E-08	6.7E-08
Naphthalene	91-20-3	2	0	1.1E-11	4.4E-11	0	5.4E-11	1.9E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	7.5E-13	4.5E-12	0	0	2.6E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	8.0E-12	1.9E-11	0	5.7E-11	5.9E-11
Naphthalene, 1-methyl-	90-12-0	1	0	1.7E-12	1.0E-11	0	0	5.9E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	3.1E-10	2.5E-10	1.9E-10	7.5E-10	9.8E-10
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	9.3E-11	8.2E-11	5.5E-11	2.7E-10	3.7E-10
Naphthalene, 2-methyl	91-57-6	1	0	1.6E-12	9.7E-12	0	0	5.7E-11
1-Octadecene	112-88-9	27	0	5.4E-10	1.5E-09	1.5E-10	9.3E-10	8.9E-09
Phenanthrene	85-01-8	29	0	3.7E-10	5.0E-10	1.8E-10	1.2E-09	2.3E-09
Phenanthrene, 1-methyl	832-69-9	35	3.6E-11	1.9E-10	1.5E-10	1.5E-10	4.7E-10	5.9E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	2.3E-10	2.2E-10	1.8E-10	6.5E-10	8.8E-10
Phenanthrene, 3-methyl	832-71-3	3	0	8.5E-11	3.2E-10	0	6.1E-10	1.7E-09
N-Phenylbenzamide	93-98-1	8	0	5.3E-10	1.1E-09	0	2.6E-09	3.9E-09
Phthalimide	85-41-6	26	0	3.5E-09	8.0E-09	4.7E-11	2.0E-08	3.5E-08
Pyrene	129-00-0	35	8.6E-10	4.7E-09	3.0E-09	4.1E-09	1.1E-08	1.2E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	1.7E-10	3.3E-10	0	7.3E-10	1.5E-09
Triethylene glycol monobutyl ether	143-22-6	7	0	8.9E-10	2.2E-09	0	5.6E-09	9.7E-09



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	1.2E-10	1.4E-10	4.3E-11	3.6E-10	5.8E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-253. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Coaches 16<30 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	4.2E-12	7.2E-12	0	1.7E-11	2.6E-11
Aniline	62-53-3	4	0	1.3E-10	4.1E-10	0	1.0E-09	2.0E-09
Anthracene	120-12-7	20	0	4.6E-11	6.0E-11	3.1E-11	1.1E-10	3.1E-10
Anthracene, 2-methyl-	613-12-7	35	1.6E-11	6.6E-11	5.4E-11	4.4E-11	1.5E-10	2.6E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	9.1E-12	2.6E-11	0	5.5E-11	1.2E-10
Anthracene, 9-phenyl	602-55-1	9	0	2.3E-11	5.1E-11	0	1.3E-10	2.3E-10
Benz[a]anthracene	56-55-3	31	0	1.8E-10	2.1E-10	9.6E-11	5.1E-10	9.7E-10
Benzene, n-butyl-	104-51-8	1	0	2.7E-13	1.6E-12	0	0	9.6E-12
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	1.2E-08	2.0E-08	5.1E-09	3.8E-08	1.1E-07
Benzo[b]fluoranthene	205-99-2	34	0	2.6E-10	2.3E-10	2.0E-10	6.5E-10	1.0E-09
7H-Benzo[c]fluorene	205-12-9	26	0	2.3E-11	2.9E-11	1.2E-11	7.2E-11	1.2E-10
Benzo[k]fluoranthene	207-08-9	29	0	1.1E-10	1.2E-10	6.9E-11	3.7E-10	4.6E-10
Benzothiazole	95-16-9	35	3.4E-08	1.5E-07	7.6E-08	1.2E-07	2.9E-07	3.3E-07
Benzothiazole, 2-phenyl-	883-93-2	35	1.1E-09	3.6E-09	2.8E-09	2.5E-09	9.7E-09	1.4E-08
1,3-Benzothiazole-2-thiol	149-30-4	1	0	9.8E-10	5.8E-09	0	0	3.4E-08
Benzothiazolone	934-34-9	35	8.9E-08	4.1E-07	1.3E-07	4.7E-07	5.8E-07	5.9E-07
Benzyl butyl phthalate	85-68-7	35	1.1E-10	1.7E-09	1.2E-09	1.4E-09	4.2E-09	4.8E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	6.2E-12	3.7E-11	0	0	2.2E-10
Dibenz[a,h]anthracene	53-70-3	14	0	6.7E-11	1.2E-10	0	3.3E-10	5.1E-10
Dibenzothiophene	132-65-0	20	0	4.0E-11	5.0E-11	3.4E-11	1.1E-10	2.4E-10
Diethyl Phthalate	84-66-2	5	0	3.1E-10	9.1E-10	0	1.7E-09	4.6E-09
Diisobutyl Phthalate	84-69-5	28	0	2.6E-10	4.3E-10	1.7E-10	6.7E-10	2.5E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Diisooctylphthalate	27554-26-3	35	7.3E-11	8.5E-09	5.9E-09	7.3E-09	2.1E-08	2.4E-08
Di-n-octyl phthalate	117-84-0	33	0	1.8E-09	2.5E-09	1.0E-09	6.2E-09	1.2E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.6E-12	4.7E-12	0	1.2E-11	2.1E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	7.9E-09	2.0E-08	1.8E-09	2.1E-08	1.2E-07
Fluoranthene	206-44-0	35	2.3E-10	1.4E-09	1.1E-09	9.8E-10	3.6E-09	3.8E-09
Fluorene	86-73-7	6	0	5.9E-12	1.8E-11	0	2.5E-11	1.0E-10
Hexanoic Acid, 2-ethyl	149-57-5	1	0	6.8E-12	4.0E-11	0	0	2.4E-10
1-Hydroxypyrene	5315-79-7	1	0	3.9E-10	2.3E-09	0	0	1.4E-08
2-(Methylthio)benzothiazole	615-22-5	9	0	5.4E-09	1.2E-08	0	2.5E-08	5.4E-08
Naphthalene	91-20-3	2	0	8.7E-12	3.6E-11	0	4.4E-11	1.6E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	6.1E-13	3.6E-12	0	0	2.1E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	6.5E-12	1.5E-11	0	4.7E-11	4.8E-11
Naphthalene, 1-methyl-	90-12-0	1	0	1.4E-12	8.1E-12	0	0	4.8E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	2.5E-10	2.1E-10	1.6E-10	6.1E-10	8.0E-10
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	7.5E-11	6.7E-11	4.5E-11	2.2E-10	3.0E-10
Naphthalene, 2-methyl	91-57-6	1	0	1.3E-12	7.9E-12	0	0	4.7E-11
1-Octadecene	112-88-9	27	0	4.4E-10	1.2E-09	1.2E-10	7.6E-10	7.2E-09
Phenanthrene	85-01-8	29	0	3.0E-10	4.1E-10	1.5E-10	1.0E-09	1.9E-09
Phenanthrene, 1-methyl	832-69-9	35	2.9E-11	1.6E-10	1.2E-10	1.2E-10	3.8E-10	4.8E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	1.9E-10	1.8E-10	1.4E-10	5.3E-10	7.1E-10
Phenanthrene, 3-methyl	832-71-3	3	0	6.9E-11	2.6E-10	0	4.9E-10	1.4E-09
N-Phenylbenzamide	93-98-1	8	0	4.3E-10	8.7E-10	0	2.1E-09	3.1E-09
Phthalimide	85-41-6	26	0	2.8E-09	6.5E-09	3.8E-11	1.6E-08	2.8E-08
Pyrene	129-00-0	35	7.0E-10	3.8E-09	2.5E-09	3.3E-09	8.9E-09	1.0E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	1.4E-10	2.7E-10	0	5.9E-10	1.2E-09
Triethylene glycol monobutyl ether	143-22-6	7	0	7.3E-10	1.8E-09	0	4.6E-09	7.9E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	9.8E-11	1.2E-10	3.5E-11	2.9E-10	4.7E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number



Values are rounded to two significant figures.

Table F-254. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Coaches 30<40 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	3.9E-12	6.7E-12	0	1.6E-11	2.4E-11
Aniline	62-53-3	4	0	1.2E-10	3.9E-10	0	9.6E-10	1.9E-09
Anthracene	120-12-7	20	0	4.3E-11	5.7E-11	2.9E-11	1.1E-10	2.9E-10
Anthracene, 2-methyl-	613-12-7	35	1.5E-11	6.2E-11	5.1E-11	4.1E-11	1.4E-10	2.4E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	8.5E-12	2.4E-11	0	5.2E-11	1.2E-10
Anthracene, 9-phenyl	602-55-1	9	0	2.1E-11	4.8E-11	0	1.2E-10	2.1E-10
Benz[a]anthracene	56-55-3	31	0	1.7E-10	2.0E-10	9.0E-11	4.8E-10	9.1E-10
Benzene, n-butyl-	104-51-8	1	0	2.6E-13	1.5E-12	0	0	9.0E-12
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	1.2E-08	1.9E-08	4.8E-09	3.6E-08	1.0E-07
Benzo[b]fluoranthene	205-99-2	34	0	2.4E-10	2.1E-10	1.9E-10	6.1E-10	9.4E-10
7H-Benzo[c]fluorene	205-12-9	26	0	2.1E-11	2.7E-11	1.1E-11	6.8E-11	1.1E-10
Benzo[k]fluoranthene	207-08-9	29	0	1.1E-10	1.1E-10	6.4E-11	3.4E-10	4.3E-10
Benzothiazole	95-16-9	35	3.2E-08	1.4E-07	7.1E-08	1.2E-07	2.7E-07	3.1E-07
Benzothiazole, 2-phenyl-	883-93-2	35	9.9E-10	3.4E-09	2.6E-09	2.4E-09	9.0E-09	1.3E-08
1,3-Benzothiazole-2-thiol	149-30-4	1	0	9.2E-10	5.4E-09	0	0	3.2E-08
Benzothiazolone	934-34-9	35	8.3E-08	3.9E-07	1.2E-07	4.4E-07	5.4E-07	5.5E-07
Benzyl butyl phthalate	85-68-7	35	1.1E-10	1.6E-09	1.1E-09	1.3E-09	3.9E-09	4.5E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	5.8E-12	3.4E-11	0	0	2.0E-10
Dibenz[a,h]anthracene	53-70-3	14	0	6.3E-11	1.2E-10	0	3.1E-10	4.8E-10
Dibenzothiophene	132-65-0	20	0	3.8E-11	4.7E-11	3.2E-11	1.0E-10	2.2E-10
Diethyl Phthalate	84-66-2	5	0	2.9E-10	8.5E-10	0	1.6E-09	4.3E-09
Diisobutyl Phthalate	84-69-5	28	0	2.4E-10	4.1E-10	1.6E-10	6.3E-10	2.4E-09
Diisooctylphthalate	27554-26-3	35	6.8E-11	7.9E-09	5.5E-09	6.8E-09	2.0E-08	2.3E-08
Di-n-octyl phthalate	117-84-0	33	0	1.7E-09	2.4E-09	9.4E-10	5.8E-09	1.1E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.5E-12	4.4E-12	0	1.2E-11	1.9E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	7.4E-09	1.9E-08	1.7E-09	2.0E-08	1.1E-07
Fluoranthene	206-44-0	35	2.1E-10	1.3E-09	1.0E-09	9.2E-10	3.4E-09	3.6E-09
Fluorene	86-73-7	6	0	5.6E-12	1.7E-11	0	2.4E-11	9.5E-11
Hexanoic Acid, 2-ethyl	149-57-5	1	0	6.4E-12	3.8E-11	0	0	2.2E-10
1-Hydroxypyrene	5315-79-7	1	0	3.7E-10	2.2E-09	0	0	1.3E-08



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-(Methylthio)benzothiazole	615-22-5	9	0	5.1E-09	1.1E-08	0	2.4E-08	5.1E-08
Naphthalene	91-20-3	2	0	8.1E-12	3.4E-11	0	4.1E-11	1.5E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	5.7E-13	3.4E-12	0	0	2.0E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	6.1E-12	1.4E-11	0	4.4E-11	4.5E-11
Naphthalene, 1-methyl-	90-12-0	1	0	1.3E-12	7.6E-12	0	0	4.5E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	2.4E-10	1.9E-10	1.5E-10	5.7E-10	7.5E-10
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	7.1E-11	6.2E-11	4.2E-11	2.0E-10	2.8E-10
Naphthalene, 2-methyl	91-57-6	1	0	1.3E-12	7.4E-12	0	0	4.4E-11
1-Octadecene	112-88-9	27	0	4.1E-10	1.1E-09	1.1E-10	7.1E-10	6.8E-09
Phenanthrene	85-01-8	29	0	2.8E-10	3.8E-10	1.4E-10	9.3E-10	1.8E-09
Phenanthrene, 1-methyl	832-69-9	35	2.7E-11	1.5E-10	1.1E-10	1.2E-10	3.6E-10	4.5E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	1.8E-10	1.6E-10	1.4E-10	5.0E-10	6.7E-10
Phenanthrene, 3-methyl	832-71-3	3	0	6.5E-11	2.4E-10	0	4.6E-10	1.3E-09
N-Phenylbenzamide	93-98-1	8	0	4.1E-10	8.2E-10	0	1.9E-09	2.9E-09
Phthalimide	85-41-6	26	0	2.6E-09	6.1E-09	3.6E-11	1.5E-08	2.6E-08
Pyrene	129-00-0	35	6.5E-10	3.6E-09	2.3E-09	3.1E-09	8.3E-09	9.4E-09
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	1.3E-10	2.5E-10	0	5.5E-10	1.1E-09
Triethylene glycol monobutyl ether	143-22-6	7	0	6.8E-10	1.7E-09	0	4.3E-09	7.4E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	9.2E-11	1.1E-10	3.3E-11	2.7E-10	4.4E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-255. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Coaches 40<50 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	3.9E-12	6.8E-12	0	1.6E-11	2.4E-11
Aniline	62-53-3	4	0	1.3E-10	3.9E-10	0	9.7E-10	1.9E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Anthracene	120-12-7	20	0	4.4E-11	5.7E-11	3.0E-11	1.1E-10	2.9E-10
Anthracene, 2-methyl-	613-12-7	35	1.5E-11	6.3E-11	5.1E-11	4.1E-11	1.4E-10	2.5E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	8.6E-12	2.4E-11	0	5.2E-11	1.2E-10
Anthracene, 9-phenyl	602-55-1	9	0	2.1E-11	4.8E-11	0	1.2E-10	2.1E-10
Benz[a]anthracene	56-55-3	31	0	1.7E-10	2.0E-10	9.1E-11	4.9E-10	9.2E-10
Benzene, n-butyl-	104-51-8	1	0	2.6E-13	1.5E-12	0	0	9.1E-12
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	1.2E-08	1.9E-08	4.9E-09	3.6E-08	1.0E-07
Benzo[b]fluoranthene	205-99-2	34	0	2.5E-10	2.2E-10	1.9E-10	6.1E-10	9.5E-10
7H-Benzo[c]fluorene	205-12-9	26	0	2.1E-11	2.7E-11	1.1E-11	6.9E-11	1.1E-10
Benzo[k]fluoranthene	207-08-9	29	0	1.1E-10	1.1E-10	6.5E-11	3.5E-10	4.3E-10
Benzothiazole	95-16-9	35	3.2E-08	1.4E-07	7.2E-08	1.2E-07	2.7E-07	3.1E-07
Benzothiazole, 2-phenyl-	883-93-2	35	1.0E-09	3.4E-09	2.7E-09	2.4E-09	9.1E-09	1.3E-08
1,3-Benzothiazole-2-thiol	149-30-4	1	0	9.3E-10	5.5E-09	0	0	3.2E-08
Benzothiazolone	934-34-9	35	8.4E-08	3.9E-07	1.3E-07	4.4E-07	5.5E-07	5.6E-07
Benzyl butyl phthalate	85-68-7	35	1.1E-10	1.6E-09	1.1E-09	1.3E-09	4.0E-09	4.5E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	5.9E-12	3.5E-11	0	0	2.1E-10
Dibenz[a,h]anthracene	53-70-3	14	0	6.4E-11	1.2E-10	0	3.1E-10	4.9E-10
Dibenzothiophene	132-65-0	20	0	3.8E-11	4.7E-11	3.2E-11	1.1E-10	2.3E-10
Diethyl Phthalate	84-66-2	5	0	2.9E-10	8.6E-10	0	1.6E-09	4.4E-09
Diisobutyl Phthalate	84-69-5	28	0	2.4E-10	4.1E-10	1.6E-10	6.4E-10	2.4E-09
Diisooctylphthalate	27554-26-3	35	6.9E-11	8.0E-09	5.6E-09	6.9E-09	2.0E-08	2.3E-08
Di-n-octyl phthalate	117-84-0	33	0	1.7E-09	2.4E-09	9.5E-10	5.9E-09	1.2E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.5E-12	4.4E-12	0	1.2E-11	1.9E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	7.4E-09	1.9E-08	1.7E-09	2.0E-08	1.1E-07
Fluoranthene	206-44-0	35	2.2E-10	1.3E-09	1.0E-09	9.3E-10	3.4E-09	3.6E-09
Fluorene	86-73-7	6	0	5.6E-12	1.7E-11	0	2.4E-11	9.6E-11
Hexanoic Acid, 2-ethyl	149-57-5	1	0	6.4E-12	3.8E-11	0	0	2.3E-10
1-Hydroxypyrene	5315-79-7	1	0	3.7E-10	2.2E-09	0	0	1.3E-08
2-(Methylthio)benzothiazole	615-22-5	9	0	5.1E-09	1.1E-08	0	2.4E-08	5.1E-08
Naphthalene	91-20-3	2	0	8.2E-12	3.4E-11	0	4.2E-11	1.5E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	5.8E-13	3.4E-12	0	0	2.0E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	6.1E-12	1.5E-11	0	4.4E-11	4.5E-11



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene, 1-methyl-	90-12-0	1	0	1.3E-12	7.7E-12	0	0	4.6E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	2.4E-10	1.9E-10	1.5E-10	5.8E-10	7.6E-10
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	7.1E-11	6.3E-11	4.3E-11	2.1E-10	2.9E-10
Naphthalene, 2-methyl	91-57-6	1	0	1.3E-12	7.5E-12	0	0	4.4E-11
1-Octadecene	112-88-9	27	0	4.1E-10	1.1E-09	1.1E-10	7.2E-10	6.8E-09
Phenanthrene	85-01-8	29	0	2.8E-10	3.8E-10	1.4E-10	9.4E-10	1.8E-09
Phenanthrene, 1-methyl	832-69-9	35	2.8E-11	1.5E-10	1.1E-10	1.2E-10	3.6E-10	4.5E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	1.8E-10	1.7E-10	1.4E-10	5.0E-10	6.8E-10
Phenanthrene, 3-methyl	832-71-3	3	0	6.6E-11	2.5E-10	0	4.7E-10	1.3E-09
N-Phenylbenzamide	93-98-1	8	0	4.1E-10	8.3E-10	0	2.0E-09	3.0E-09
Phthalimide	85-41-6	26	0	2.7E-09	6.2E-09	3.6E-11	1.5E-08	2.7E-08
Pyrene	129-00-0	35	6.6E-10	3.6E-09	2.3E-09	3.1E-09	8.4E-09	9.5E-09
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	1.3E-10	2.5E-10	0	5.6E-10	1.1E-09
Triethylene glycol monobutyl ether	143-22-6	7	0	6.9E-10	1.7E-09	0	4.3E-09	7.5E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	9.3E-11	1.1E-10	3.3E-11	2.8E-10	4.5E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-256. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender **Coaches 50<70 Years****

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	3.9E-12	6.8E-12	0	1.6E-11	2.5E-11
Aniline	62-53-3	4	0	1.3E-10	3.9E-10	0	9.7E-10	1.9E-09
Anthracene	120-12-7	20	0	4.4E-11	5.7E-11	3.0E-11	1.1E-10	2.9E-10
Anthracene, 2-methyl-	613-12-7	35	1.5E-11	6.3E-11	5.1E-11	4.2E-11	1.5E-10	2.5E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	8.6E-12	2.4E-11	0	5.2E-11	1.2E-10
Anthracene, 9-phenyl	602-55-1	9	0	2.1E-11	4.8E-11	0	1.2E-10	2.1E-10
Benz[a]anthracene	56-55-3	31	0	1.7E-10	2.0E-10	9.2E-11	4.9E-10	9.2E-10
Benzene, n-butyl-	104-51-8	1	0	2.6E-13	1.5E-12	0	0	9.1E-12



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	1.2E-08	1.9E-08	4.9E-09	3.6E-08	1.0E-07
Benzo[b]fluoranthene	205-99-2	34	0	2.5E-10	2.2E-10	1.9E-10	6.2E-10	9.5E-10
7H-Benzo[c]fluorene	205-12-9	26	0	2.1E-11	2.7E-11	1.1E-11	6.9E-11	1.1E-10
Benzo[k]fluoranthene	207-08-9	29	0	1.1E-10	1.1E-10	6.5E-11	3.5E-10	4.3E-10
Benzothiazole	95-16-9	35	3.2E-08	1.4E-07	7.2E-08	1.2E-07	2.7E-07	3.1E-07
Benzothiazole, 2-phenyl-	883-93-2	35	1.0E-09	3.4E-09	2.7E-09	2.4E-09	9.2E-09	1.3E-08
1,3-Benzothiazole-2-thiol	149-30-4	1	0	9.3E-10	5.5E-09	0	0	3.3E-08
Benzothiazolone	934-34-9	35	8.5E-08	3.9E-07	1.3E-07	4.4E-07	5.5E-07	5.6E-07
Benzyl butyl phthalate	85-68-7	35	1.1E-10	1.6E-09	1.1E-09	1.3E-09	4.0E-09	4.5E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	5.9E-12	3.5E-11	0	0	2.1E-10
Dibenz[a,h]anthracene	53-70-3	14	0	6.4E-11	1.2E-10	0	3.1E-10	4.9E-10
Dibenzothiophene	132-65-0	20	0	3.8E-11	4.7E-11	3.2E-11	1.1E-10	2.3E-10
Diethyl Phthalate	84-66-2	5	0	2.9E-10	8.6E-10	0	1.6E-09	4.4E-09
Diisobutyl Phthalate	84-69-5	28	0	2.4E-10	4.1E-10	1.6E-10	6.4E-10	2.4E-09
Diisooctylphthalate	27554-26-3	35	6.9E-11	8.0E-09	5.6E-09	6.9E-09	2.0E-08	2.3E-08
Di-n-octyl phthalate	117-84-0	33	0	1.7E-09	2.4E-09	9.5E-10	5.9E-09	1.2E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.5E-12	4.4E-12	0	1.2E-11	2.0E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	7.5E-09	1.9E-08	1.7E-09	2.0E-08	1.1E-07
Fluoranthene	206-44-0	35	2.2E-10	1.3E-09	1.0E-09	9.3E-10	3.4E-09	3.7E-09
Fluorene	86-73-7	6	0	5.6E-12	1.8E-11	0	2.4E-11	9.6E-11
Hexanoic Acid, 2-ethyl	149-57-5	1	0	6.5E-12	3.8E-11	0	0	2.3E-10
1-Hydroxypyrene	5315-79-7	1	0	3.7E-10	2.2E-09	0	0	1.3E-08
2-(Methylthio)benzothiazole	615-22-5	9	0	5.1E-09	1.1E-08	0	2.4E-08	5.2E-08
Naphthalene	91-20-3	2	0	8.3E-12	3.4E-11	0	4.2E-11	1.5E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	5.8E-13	3.4E-12	0	0	2.0E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	6.2E-12	1.5E-11	0	4.4E-11	4.6E-11
Naphthalene, 1-methyl-	90-12-0	1	0	1.3E-12	7.7E-12	0	0	4.6E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	2.4E-10	1.9E-10	1.5E-10	5.8E-10	7.6E-10
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	7.2E-11	6.3E-11	4.3E-11	2.1E-10	2.9E-10
Naphthalene, 2-methyl	91-57-6	1	0	1.3E-12	7.5E-12	0	0	4.4E-11
1-Octadecene	112-88-9	27	0	4.1E-10	1.1E-09	1.1E-10	7.2E-10	6.9E-09



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Phenanthrene	85-01-8	29	0	2.9E-10	3.9E-10	1.4E-10	9.5E-10	1.8E-09
Phenanthrene, 1-methyl	832-69-9	35	2.8E-11	1.5E-10	1.1E-10	1.2E-10	3.6E-10	4.6E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	1.8E-10	1.7E-10	1.4E-10	5.1E-10	6.8E-10
Phenanthrene, 3-methyl	832-71-3	3	0	6.6E-11	2.5E-10	0	4.7E-10	1.3E-09
N-Phenylbenzamide	93-98-1	8	0	4.1E-10	8.3E-10	0	2.0E-09	3.0E-09
Phthalimide	85-41-6	26	0	2.7E-09	6.2E-09	3.6E-11	1.5E-08	2.7E-08
Pyrene	129-00-0	35	6.6E-10	3.6E-09	2.3E-09	3.1E-09	8.5E-09	9.5E-09
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	1.3E-10	2.5E-10	0	5.6E-10	1.2E-09
Triethylene glycol monobutyl ether	143-22-6	7	0	6.9E-10	1.7E-09	0	4.3E-09	7.5E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	9.3E-11	1.1E-10	3.3E-11	2.8E-10	4.5E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-257. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Referees 16<30 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	1.7E-12	2.9E-12	0	6.9E-12	1.0E-11
Aniline	62-53-3	4	0	5.4E-11	1.7E-10	0	4.1E-10	8.0E-10
Anthracene	120-12-7	20	0	1.9E-11	2.4E-11	1.3E-11	4.6E-11	1.2E-10
Anthracene, 2-methyl-	613-12-7	35	6.4E-12	2.7E-11	2.2E-11	1.8E-11	6.2E-11	1.0E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	3.7E-12	1.0E-11	0	2.2E-11	5.0E-11
Anthracene, 9-phenyl	602-55-1	9	0	9.1E-12	2.0E-11	0	5.1E-11	9.1E-11
Benz[a]anthracene	56-55-3	31	0	7.3E-11	8.5E-11	3.9E-11	2.1E-10	3.9E-10
Benzene, n-butyl-	104-51-8	1	0	1.1E-13	6.5E-13	0	0	3.9E-12
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	5.0E-09	8.0E-09	2.1E-09	1.5E-08	4.3E-08
Benzo[b]fluoranthene	205-99-2	34	0	1.0E-10	9.2E-11	8.2E-11	2.6E-10	4.0E-10
7H-Benzo[c]fluorene	205-12-9	26	0	9.1E-12	1.1E-11	4.7E-12	2.9E-11	4.8E-11
Benzo[k]fluoranthene	207-08-9	29	0	4.6E-11	4.8E-11	2.8E-11	1.5E-10	1.8E-10
Benzothiazole	95-16-9	35	1.4E-08	5.9E-08	3.1E-08	5.0E-08	1.2E-07	1.3E-07



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzothiazole, 2-phenyl-	883-93-2	35	4.2E-10	1.4E-09	1.1E-09	1.0E-09	3.9E-09	5.7E-09
1,3-Benzothiazole-2-thiol	149-30-4	1	0	3.9E-10	2.3E-09	0	0	1.4E-08
Benzothiazolone	934-34-9	35	3.6E-08	1.7E-07	5.3E-08	1.9E-07	2.3E-07	2.4E-07
Benzyl butyl phthalate	85-68-7	35	4.6E-11	6.8E-10	4.8E-10	5.7E-10	1.7E-09	1.9E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	2.5E-12	1.5E-11	0	0	8.8E-11
Dibenz[a,h]anthracene	53-70-3	14	0	2.7E-11	4.9E-11	0	1.3E-10	2.1E-10
Dibenzothiophene	132-65-0	20	0	1.6E-11	2.0E-11	1.4E-11	4.5E-11	9.6E-11
Diethyl Phthalate	84-66-2	5	0	1.2E-10	3.7E-10	0	6.8E-10	1.9E-09
Diisobutyl Phthalate	84-69-5	28	0	1.0E-10	1.7E-10	6.9E-11	2.7E-10	1.0E-09
Diisooctylphthalate	27554-26-3	35	2.9E-11	3.4E-09	2.4E-09	2.9E-09	8.4E-09	9.7E-09
Di-n-octyl phthalate	117-84-0	33	0	7.2E-10	1.0E-09	4.0E-10	2.5E-09	4.9E-09
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	6.4E-13	1.9E-12	0	5.0E-12	8.3E-12
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	3.2E-09	8.2E-09	7.3E-10	8.6E-09	4.7E-08
Fluoranthene	206-44-0	35	9.2E-11	5.6E-10	4.4E-10	4.0E-10	1.5E-09	1.5E-09
Fluorene	86-73-7	6	0	2.4E-12	7.4E-12	0	1.0E-11	4.1E-11
Hexanoic Acid, 2-ethyl	149-57-5	1	0	2.7E-12	1.6E-11	0	0	9.6E-11
1-Hydroxypyrene	5315-79-7	1	0	1.6E-10	9.4E-10	0	0	5.6E-09
2-(Methylthio)benzothiazole	615-22-5	9	0	2.2E-09	4.8E-09	0	1.0E-08	2.2E-08
Naphthalene	91-20-3	2	0	3.5E-12	1.4E-11	0	1.8E-11	6.4E-11
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	2.5E-13	1.5E-12	0	0	8.6E-12
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	2.6E-12	6.2E-12	0	1.9E-11	1.9E-11
Naphthalene, 1-methyl-	90-12-0	1	0	5.5E-13	3.3E-12	0	0	1.9E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	1.0E-10	8.2E-11	6.3E-11	2.4E-10	3.2E-10
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	3.0E-11	2.7E-11	1.8E-11	8.7E-11	1.2E-10
Naphthalene, 2-methyl	91-57-6	1	0	5.4E-13	3.2E-12	0	0	1.9E-11
1-Octadecene	112-88-9	27	0	1.8E-10	4.9E-10	4.9E-11	3.1E-10	2.9E-09
Phenanthrene	85-01-8	29	0	1.2E-10	1.6E-10	5.9E-11	4.0E-10	7.6E-10
Phenanthrene, 1-methyl	832-69-9	35	1.2E-11	6.3E-11	4.8E-11	5.0E-11	1.5E-10	1.9E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	7.7E-11	7.1E-11	5.8E-11	2.1E-10	2.9E-10
Phenanthrene, 3-methyl	832-71-3	3	0	2.8E-11	1.0E-10	0	2.0E-10	5.6E-10
N-Phenylbenzamide	93-98-1	8	0	1.7E-10	3.5E-10	0	8.4E-10	1.3E-09
Phthalimide	85-41-6	26	0	1.1E-09	2.6E-09	1.5E-11	6.5E-09	1.1E-08



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Pyrene	129-00-0	35	2.8E-10	1.5E-09	9.9E-10	1.3E-09	3.6E-09	4.0E-09
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	5.6E-11	1.1E-10	0	2.4E-10	4.9E-10
Triethylene glycol monobutyl ether	143-22-6	7	0	2.9E-10	7.1E-10	0	1.8E-09	3.2E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	3.9E-11	4.7E-11	1.4E-11	1.2E-10	1.9E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-258. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Referees 30<40 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	1.7E-12	2.9E-12	0	6.9E-12	1.0E-11
Aniline	62-53-3	4	0	5.4E-11	1.7E-10	0	4.1E-10	8.0E-10
Anthracene	120-12-7	20	0	1.9E-11	2.4E-11	1.3E-11	4.6E-11	1.2E-10
Anthracene, 2-methyl-	613-12-7	35	6.4E-12	2.7E-11	2.2E-11	1.8E-11	6.2E-11	1.0E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	3.7E-12	1.0E-11	0	2.2E-11	5.0E-11
Anthracene, 9-phenyl	602-55-1	9	0	9.1E-12	2.0E-11	0	5.1E-11	9.1E-11
Benz[a]anthracene	56-55-3	31	0	7.3E-11	8.5E-11	3.9E-11	2.1E-10	3.9E-10
Benzene, n-butyl-	104-51-8	1	0	1.1E-13	6.5E-13	0	0	3.9E-12
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	5.0E-09	8.0E-09	2.1E-09	1.5E-08	4.3E-08
Benzo[b]fluoranthene	205-99-2	34	0	1.0E-10	9.2E-11	8.2E-11	2.6E-10	4.0E-10
7H-Benzo[c]fluorene	205-12-9	26	0	9.1E-12	1.1E-11	4.7E-12	2.9E-11	4.8E-11
Benzo[k]fluoranthene	207-08-9	29	0	4.6E-11	4.8E-11	2.8E-11	1.5E-10	1.8E-10
Benzothiazole	95-16-9	35	1.4E-08	5.9E-08	3.1E-08	5.0E-08	1.2E-07	1.3E-07
Benzothiazole, 2-phenyl-	883-93-2	35	4.2E-10	1.4E-09	1.1E-09	1.0E-09	3.9E-09	5.7E-09
1,3-Benzothiazole-2-thiol	149-30-4	1	0	3.9E-10	2.3E-09	0	0	1.4E-08
Benzothiazolone	934-34-9	35	3.6E-08	1.7E-07	5.3E-08	1.9E-07	2.3E-07	2.4E-07
Benzyl butyl phthalate	85-68-7	35	4.6E-11	6.8E-10	4.8E-10	5.7E-10	1.7E-09	1.9E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	2.5E-12	1.5E-11	0	0	8.8E-11



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dibenz[a,h]anthracene	53-70-3	14	0	2.7E-11	4.9E-11	0	1.3E-10	2.1E-10
Dibenzothiophene	132-65-0	20	0	1.6E-11	2.0E-11	1.4E-11	4.5E-11	9.6E-11
Diethyl Phthalate	84-66-2	5	0	1.2E-10	3.7E-10	0	6.8E-10	1.9E-09
Diisobutyl Phthalate	84-69-5	28	0	1.0E-10	1.7E-10	6.9E-11	2.7E-10	1.0E-09
Diisooctylphthalate	27554-26-3	35	2.9E-11	3.4E-09	2.4E-09	2.9E-09	8.4E-09	9.7E-09
Di-n-octyl phthalate	117-84-0	33	0	7.2E-10	1.0E-09	4.0E-10	2.5E-09	4.9E-09
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	6.4E-13	1.9E-12	0	5.0E-12	8.3E-12
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	3.2E-09	8.2E-09	7.3E-10	8.6E-09	4.7E-08
Fluoranthene	206-44-0	35	9.2E-11	5.6E-10	4.4E-10	4.0E-10	1.5E-09	1.5E-09
Fluorene	86-73-7	6	0	2.4E-12	7.4E-12	0	1.0E-11	4.1E-11
Hexanoic Acid, 2-ethyl	149-57-5	1	0	2.7E-12	1.6E-11	0	0	9.6E-11
1-Hydroxypyrene	5315-79-7	1	0	1.6E-10	9.4E-10	0	0	5.6E-09
2-(Methylthio)benzothiazole	615-22-5	9	0	2.2E-09	4.8E-09	0	1.0E-08	2.2E-08
Naphthalene	91-20-3	2	0	3.5E-12	1.4E-11	0	1.8E-11	6.4E-11
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	2.5E-13	1.5E-12	0	0	8.6E-12
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	2.6E-12	6.2E-12	0	1.9E-11	1.9E-11
Naphthalene, 1-methyl-	90-12-0	1	0	5.5E-13	3.3E-12	0	0	1.9E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	1.0E-10	8.2E-11	6.3E-11	2.4E-10	3.2E-10
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	3.0E-11	2.7E-11	1.8E-11	8.7E-11	1.2E-10
Naphthalene, 2-methyl	91-57-6	1	0	5.4E-13	3.2E-12	0	0	1.9E-11
1-Octadecene	112-88-9	27	0	1.8E-10	4.9E-10	4.9E-11	3.1E-10	2.9E-09
Phenanthrene	85-01-8	29	0	1.2E-10	1.6E-10	5.9E-11	4.0E-10	7.6E-10
Phenanthrene, 1-methyl	832-69-9	35	1.2E-11	6.3E-11	4.8E-11	5.0E-11	1.5E-10	1.9E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	7.7E-11	7.1E-11	5.8E-11	2.1E-10	2.9E-10
Phenanthrene, 3-methyl	832-71-3	3	0	2.8E-11	1.0E-10	0	2.0E-10	5.6E-10
N-Phenylbenzamide	93-98-1	8	0	1.7E-10	3.5E-10	0	8.4E-10	1.3E-09
Phthalimide	85-41-6	26	0	1.1E-09	2.6E-09	1.5E-11	6.5E-09	1.1E-08
Pyrene	129-00-0	35	2.8E-10	1.5E-09	9.9E-10	1.3E-09	3.6E-09	4.0E-09
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	5.6E-11	1.1E-10	0	2.4E-10	4.9E-10
Triethylene glycol monobutyl ether	143-22-6	7	0	2.9E-10	7.1E-10	0	1.8E-09	3.2E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	3.9E-11	4.7E-11	1.4E-11	1.2E-10	1.9E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average



concentrations ($C_{\text{der-crumb rubber-field}}$) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-259. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard ($ADD_{\text{der-field}}$, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Referees 40<50 Years**

Chemical	CASRN	Detection	$ADD_{\text{der-field}}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	1.6E-12	2.7E-12	0	6.5E-12	9.8E-12
Aniline	62-53-3	4	0	5.1E-11	1.6E-10	0	3.9E-10	7.6E-10
Anthracene	120-12-7	20	0	1.8E-11	2.3E-11	1.2E-11	4.4E-11	1.2E-10
Anthracene, 2-methyl-	613-12-7	35	6.1E-12	2.5E-11	2.1E-11	1.7E-11	5.8E-11	9.9E-11
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	3.5E-12	9.7E-12	0	2.1E-11	4.7E-11
Anthracene, 9-phenyl	602-55-1	9	0	8.6E-12	1.9E-11	0	4.8E-11	8.6E-11
Benz[a]anthracene	56-55-3	31	0	7.0E-11	8.0E-11	3.7E-11	2.0E-10	3.7E-10
Benzene, n-butyl-	104-51-8	1	0	1.0E-13	6.2E-13	0	0	3.7E-12
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	4.7E-09	7.6E-09	2.0E-09	1.5E-08	4.1E-08
Benzo[b]fluoranthene	205-99-2	34	0	9.9E-11	8.7E-11	7.8E-11	2.5E-10	3.8E-10
7H-Benzo[c]fluorene	205-12-9	26	0	8.6E-12	1.1E-11	4.4E-12	2.8E-11	4.6E-11
Benzo[k]fluoranthene	207-08-9	29	0	4.3E-11	4.6E-11	2.6E-11	1.4E-10	1.7E-10
Benzothiazole	95-16-9	35	1.3E-08	5.6E-08	2.9E-08	4.7E-08	1.1E-07	1.3E-07
Benzothiazole, 2-phenyl-	883-93-2	35	4.0E-10	1.4E-09	1.1E-09	9.7E-10	3.7E-09	5.4E-09
1,3-Benzothiazole-2-thiol	149-30-4	1	0	3.7E-10	2.2E-09	0	0	1.3E-08
Benzothiazolone	934-34-9	35	3.4E-08	1.6E-07	5.0E-08	1.8E-07	2.2E-07	2.2E-07
Benzyl butyl phthalate	85-68-7	35	4.4E-11	6.4E-10	4.6E-10	5.4E-10	1.6E-09	1.8E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	2.4E-12	1.4E-11	0	0	8.3E-11
Dibenz[a,h]anthracene	53-70-3	14	0	2.6E-11	4.7E-11	0	1.2E-10	2.0E-10
Dibenzothiophene	132-65-0	20	0	1.5E-11	1.9E-11	1.3E-11	4.2E-11	9.1E-11
Diethyl Phthalate	84-66-2	5	0	1.2E-10	3.5E-10	0	6.4E-10	1.8E-09
Diisobutyl Phthalate	84-69-5	28	0	9.8E-11	1.6E-10	6.5E-11	2.6E-10	9.6E-10
Diisooctylphthalate	27554-26-3	35	2.8E-11	3.2E-09	2.2E-09	2.8E-09	8.0E-09	9.2E-09
Di-n-octyl phthalate	117-84-0	33	0	6.8E-10	9.7E-10	3.8E-10	2.4E-09	4.7E-09
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	6.0E-13	1.8E-12	0	4.7E-12	7.8E-12



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	3.0E-09	7.7E-09	6.9E-10	8.1E-09	4.5E-08
Fluoranthene	206-44-0	35	8.7E-11	5.3E-10	4.2E-10	3.7E-10	1.4E-09	1.5E-09
Fluorene	86-73-7	6	0	2.3E-12	7.0E-12	0	9.6E-12	3.9E-11
Hexanoic Acid, 2-ethyl	149-57-5	1	0	2.6E-12	1.5E-11	0	0	9.1E-11
1-Hydroxypyrene	5315-79-7	1	0	1.5E-10	8.9E-10	0	0	5.3E-09
2-(Methylthio)benzothiazole	615-22-5	9	0	2.1E-09	4.6E-09	0	9.6E-09	2.1E-08
Naphthalene	91-20-3	2	0	3.3E-12	1.4E-11	0	1.7E-11	6.0E-11
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	2.3E-13	1.4E-12	0	0	8.2E-12
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	2.5E-12	5.9E-12	0	1.8E-11	1.8E-11
Naphthalene, 1-methyl-	90-12-0	1	0	5.2E-13	3.1E-12	0	0	1.8E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	9.7E-11	7.8E-11	6.0E-11	2.3E-10	3.0E-10
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	2.9E-11	2.5E-11	1.7E-11	8.3E-11	1.2E-10
Naphthalene, 2-methyl	91-57-6	1	0	5.1E-13	3.0E-12	0	0	1.8E-11
1-Octadecene	112-88-9	27	0	1.7E-10	4.6E-10	4.6E-11	2.9E-10	2.8E-09
Phenanthrene	85-01-8	29	0	1.1E-10	1.5E-10	5.6E-11	3.8E-10	7.2E-10
Phenanthrene, 1-methyl	832-69-9	35	1.1E-11	6.0E-11	4.6E-11	4.7E-11	1.5E-10	1.8E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	7.3E-11	6.7E-11	5.5E-11	2.0E-10	2.7E-10
Phenanthrene, 3-methyl	832-71-3	3	0	2.6E-11	9.9E-11	0	1.9E-10	5.3E-10
N-Phenylbenzamide	93-98-1	8	0	1.7E-10	3.3E-10	0	7.9E-10	1.2E-09
Phthalimide	85-41-6	26	0	1.1E-09	2.5E-09	1.5E-11	6.1E-09	1.1E-08
Pyrene	129-00-0	35	2.7E-10	1.4E-09	9.4E-10	1.3E-09	3.4E-09	3.8E-09
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	5.3E-11	1.0E-10	0	2.3E-10	4.6E-10
Triethylene glycol monobutyl ether	143-22-6	7	0	2.8E-10	6.8E-10	0	1.7E-09	3.0E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	3.7E-11	4.5E-11	1.3E-11	1.1E-10	1.8E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-260. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Referees 50<70 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	1.6E-12	2.7E-12	0	6.5E-12	9.9E-12
Aniline	62-53-3	4	0	5.1E-11	1.6E-10	0	3.9E-10	7.6E-10
Anthracene	120-12-7	20	0	1.8E-11	2.3E-11	1.2E-11	4.4E-11	1.2E-10
Anthracene, 2-methyl-	613-12-7	35	6.1E-12	2.5E-11	2.1E-11	1.7E-11	5.8E-11	9.9E-11
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	3.5E-12	9.8E-12	0	2.1E-11	4.7E-11
Anthracene, 9-phenyl	602-55-1	9	0	8.6E-12	1.9E-11	0	4.8E-11	8.6E-11
Benz[a]anthracene	56-55-3	31	0	7.0E-11	8.1E-11	3.7E-11	2.0E-10	3.7E-10
Benzene, n-butyl-	104-51-8	1	0	1.0E-13	6.2E-13	0	0	3.7E-12
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	4.7E-09	7.6E-09	2.0E-09	1.5E-08	4.1E-08
Benzo[b]fluoranthene	205-99-2	34	0	9.9E-11	8.7E-11	7.8E-11	2.5E-10	3.8E-10
7H-Benzo[c]fluorene	205-12-9	26	0	8.6E-12	1.1E-11	4.5E-12	2.8E-11	4.6E-11
Benzo[k]fluoranthene	207-08-9	29	0	4.3E-11	4.6E-11	2.6E-11	1.4E-10	1.7E-10
Benzothiazole	95-16-9	35	1.3E-08	5.6E-08	2.9E-08	4.7E-08	1.1E-07	1.3E-07
Benzothiazole, 2-phenyl-	883-93-2	35	4.0E-10	1.4E-09	1.1E-09	9.7E-10	3.7E-09	5.4E-09
1,3-Benzothiazole-2-thiol	149-30-4	1	0	3.7E-10	2.2E-09	0	0	1.3E-08
Benzothiazolone	934-34-9	35	3.4E-08	1.6E-07	5.1E-08	1.8E-07	2.2E-07	2.3E-07
Benzyl butyl phthalate	85-68-7	35	4.4E-11	6.4E-10	4.6E-10	5.4E-10	1.6E-09	1.8E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	2.4E-12	1.4E-11	0	0	8.3E-11
Dibenz[a,h]anthracene	53-70-3	14	0	2.6E-11	4.7E-11	0	1.3E-10	2.0E-10
Dibenzothiophene	132-65-0	20	0	1.5E-11	1.9E-11	1.3E-11	4.3E-11	9.2E-11
Diethyl Phthalate	84-66-2	5	0	1.2E-10	3.5E-10	0	6.5E-10	1.8E-09
Diisobutyl Phthalate	84-69-5	28	0	9.8E-11	1.7E-10	6.6E-11	2.6E-10	9.6E-10
Diisooctylphthalate	27554-26-3	35	2.8E-11	3.2E-09	2.2E-09	2.8E-09	8.0E-09	9.3E-09
Di-n-octyl phthalate	117-84-0	33	0	6.9E-10	9.7E-10	3.8E-10	2.4E-09	4.7E-09
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	6.0E-13	1.8E-12	0	4.7E-12	7.9E-12
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	3.0E-09	7.8E-09	6.9E-10	8.1E-09	4.5E-08
Fluoranthene	206-44-0	35	8.7E-11	5.4E-10	4.2E-10	3.8E-10	1.4E-09	1.5E-09
Fluorene	86-73-7	6	0	2.3E-12	7.0E-12	0	9.6E-12	3.9E-11
Hexanoic Acid, 2-ethyl	149-57-5	1	0	2.6E-12	1.5E-11	0	0	9.1E-11
1-Hydroxypyrene	5315-79-7	1	0	1.5E-10	8.9E-10	0	0	5.3E-09



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-(Methylthio)benzothiazole	615-22-5	9	0	2.1E-09	4.6E-09	0	9.6E-09	2.1E-08
Naphthalene	91-20-3	2	0	3.3E-12	1.4E-11	0	1.7E-11	6.0E-11
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	2.3E-13	1.4E-12	0	0	8.2E-12
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	2.5E-12	5.9E-12	0	1.8E-11	1.8E-11
Naphthalene, 1-methyl-	90-12-0	1	0	5.3E-13	3.1E-12	0	0	1.8E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	9.7E-11	7.8E-11	6.0E-11	2.3E-10	3.1E-10
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	2.9E-11	2.5E-11	1.7E-11	8.3E-11	1.2E-10
Naphthalene, 2-methyl	91-57-6	1	0	5.1E-13	3.0E-12	0	0	1.8E-11
1-Octadecene	112-88-9	27	0	1.7E-10	4.6E-10	4.6E-11	2.9E-10	2.8E-09
Phenanthrene	85-01-8	29	0	1.1E-10	1.6E-10	5.6E-11	3.8E-10	7.2E-10
Phenanthrene, 1-methyl	832-69-9	35	1.1E-11	6.0E-11	4.6E-11	4.7E-11	1.5E-10	1.8E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	7.3E-11	6.7E-11	5.5E-11	2.0E-10	2.7E-10
Phenanthrene, 3-methyl	832-71-3	3	0	2.6E-11	1.0E-10	0	1.9E-10	5.3E-10
N-Phenylbenzamide	93-98-1	8	0	1.7E-10	3.3E-10	0	7.9E-10	1.2E-09
Phthalimide	85-41-6	26	0	1.1E-09	2.5E-09	1.5E-11	6.1E-09	1.1E-08
Pyrene	129-00-0	35	2.7E-10	1.4E-09	9.4E-10	1.3E-09	3.4E-09	3.8E-09
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	5.3E-11	1.0E-10	0	2.3E-10	4.6E-10
Triethylene glycol monobutyl ether	143-22-6	7	0	2.8E-10	6.8E-10	0	1.7E-09	3.0E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	3.7E-11	4.5E-11	1.3E-11	1.1E-10	1.8E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-261. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Spectators Third Trimester Fetus<0 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	2.4E-12	4.1E-12	0	9.7E-12	1.5E-11
Aniline	62-53-3	4	0	7.6E-11	2.3E-10	0	5.8E-10	1.1E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Anthracene	120-12-7	20	0	2.6E-11	3.4E-11	1.8E-11	6.5E-11	1.7E-10
Anthracene, 2-methyl-	613-12-7	35	9.1E-12	3.8E-11	3.1E-11	2.5E-11	8.7E-11	1.5E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	5.1E-12	1.4E-11	0	3.1E-11	7.0E-11
Anthracene, 9-phenyl	602-55-1	9	0	1.3E-11	2.9E-11	0	7.2E-11	1.3E-10
Benz[a]anthracene	56-55-3	31	0	1.0E-10	1.2E-10	5.5E-11	2.9E-10	5.5E-10
Benzene, n-butyl-	104-51-8	1	0	1.6E-13	9.2E-13	0	0	5.4E-12
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	7.0E-09	1.1E-08	2.9E-09	2.2E-08	6.0E-08
Benzo[b]fluoranthene	205-99-2	34	0	1.5E-10	1.3E-10	1.2E-10	3.7E-10	5.7E-10
7H-Benzo[c]fluorene	205-12-9	26	0	1.3E-11	1.6E-11	6.6E-12	4.1E-11	6.8E-11
Benzo[k]fluoranthene	207-08-9	29	0	6.4E-11	6.8E-11	3.9E-11	2.1E-10	2.6E-10
Benzothiazole	95-16-9	35	1.9E-08	8.4E-08	4.3E-08	7.0E-08	1.6E-07	1.9E-07
Benzothiazole, 2-phenyl-	883-93-2	35	6.0E-10	2.0E-09	1.6E-09	1.4E-09	5.5E-09	8.0E-09
1,3-Benzothiazole-2-thiol	149-30-4	1	0	5.6E-10	3.3E-09	0	0	1.9E-08
Benzothiazolone	934-34-9	35	5.0E-08	2.3E-07	7.5E-08	2.6E-07	3.3E-07	3.3E-07
Benzyl butyl phthalate	85-68-7	35	6.5E-11	9.5E-10	6.8E-10	8.1E-10	2.4E-09	2.7E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	3.5E-12	2.1E-11	0	0	1.2E-10
Dibenz[a,h]anthracene	53-70-3	14	0	3.8E-11	7.0E-11	0	1.9E-10	2.9E-10
Dibenzothiophene	132-65-0	20	0	2.3E-11	2.8E-11	1.9E-11	6.3E-11	1.4E-10
Diethyl Phthalate	84-66-2	5	0	1.7E-10	5.2E-10	0	9.6E-10	2.6E-09
Diisobutyl Phthalate	84-69-5	28	0	1.5E-10	2.5E-10	9.7E-11	3.8E-10	1.4E-09
Diisooctylphthalate	27554-26-3	35	4.1E-11	4.8E-09	3.3E-09	4.1E-09	1.2E-08	1.4E-08
Di-n-octyl phthalate	117-84-0	33	0	1.0E-09	1.4E-09	5.7E-10	3.5E-09	6.9E-09
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	9.0E-13	2.7E-12	0	7.0E-12	1.2E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	4.5E-09	1.2E-08	1.0E-09	1.2E-08	6.7E-08
Fluoranthene	206-44-0	35	1.3E-10	8.0E-10	6.2E-10	5.6E-10	2.1E-09	2.2E-09
Fluorene	86-73-7	6	0	3.4E-12	1.0E-11	0	1.4E-11	5.8E-11
Hexanoic Acid, 2-ethyl	149-57-5	1	0	3.9E-12	2.3E-11	0	0	1.3E-10
1-Hydroxypyrene	5315-79-7	1	0	2.2E-10	1.3E-09	0	0	7.8E-09
2-(Methylthio)benzothiazole	615-22-5	9	0	3.1E-09	6.8E-09	0	1.4E-08	3.1E-08
Naphthalene	91-20-3	2	0	4.9E-12	2.0E-11	0	2.5E-11	9.0E-11
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	3.5E-13	2.1E-12	0	0	1.2E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	3.7E-12	8.7E-12	0	2.6E-11	2.7E-11



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene, 1-methyl-	90-12-0	1	0	7.8E-13	4.6E-12	0	0	2.7E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	1.4E-10	1.2E-10	8.9E-11	3.4E-10	4.5E-10
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	4.3E-11	3.8E-11	2.6E-11	1.2E-10	1.7E-10
Naphthalene, 2-methyl	91-57-6	1	0	7.6E-13	4.5E-12	0	0	2.6E-11
1-Octadecene	112-88-9	27	0	2.5E-10	6.9E-10	6.8E-11	4.3E-10	4.1E-09
Phenanthrene	85-01-8	29	0	1.7E-10	2.3E-10	8.3E-11	5.6E-10	1.1E-09
Phenanthrene, 1-methyl	832-69-9	35	1.7E-11	8.9E-11	6.8E-11	7.0E-11	2.2E-10	2.7E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	1.1E-10	1.0E-10	8.2E-11	3.0E-10	4.1E-10
Phenanthrene, 3-methyl	832-71-3	3	0	3.9E-11	1.5E-10	0	2.8E-10	7.9E-10
N-Phenylbenzamide	93-98-1	8	0	2.5E-10	5.0E-10	0	1.2E-09	1.8E-09
Phthalimide	85-41-6	26	0	1.6E-09	3.7E-09	2.2E-11	9.1E-09	1.6E-08
Pyrene	129-00-0	35	4.0E-10	2.2E-09	1.4E-09	1.9E-09	5.0E-09	5.7E-09
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	7.9E-11	1.5E-10	0	3.4E-10	6.9E-10
Triethylene glycol monobutyl ether	143-22-6	7	0	4.1E-10	1.0E-09	0	2.6E-09	4.5E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	5.5E-11	6.7E-11	2.0E-11	1.6E-10	2.7E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-262. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender **Spectators 0<2 Years****

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	1.3E-11	2.2E-11	0	5.2E-11	7.9E-11
Aniline	62-53-3	4	0	4.1E-10	1.3E-09	0	3.1E-09	6.0E-09
Anthracene	120-12-7	20	0	1.4E-10	1.8E-10	9.6E-11	3.5E-10	9.4E-10
Anthracene, 2-methyl-	613-12-7	35	4.9E-11	2.0E-10	1.6E-10	1.3E-10	4.7E-10	7.9E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	2.8E-11	7.8E-11	0	1.7E-10	3.8E-10
Anthracene, 9-phenyl	602-55-1	9	0	6.9E-11	1.5E-10	0	3.8E-10	6.8E-10
Benz[a]anthracene	56-55-3	31	0	5.6E-10	6.4E-10	2.9E-10	1.6E-09	3.0E-09
Benzene, n-butyl-	104-51-8	1	0	8.3E-13	4.9E-12	0	0	2.9E-11



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	3.8E-08	6.0E-08	1.6E-08	1.2E-07	3.2E-07
Benzo[b]fluoranthene	205-99-2	34	0	7.9E-10	7.0E-10	6.2E-10	2.0E-09	3.1E-09
7H-Benzo[c]fluorene	205-12-9	26	0	6.9E-11	8.7E-11	3.5E-11	2.2E-10	3.6E-10
Benzo[k]fluoranthene	207-08-9	29	0	3.4E-10	3.6E-10	2.1E-10	1.1E-09	1.4E-09
Benzothiazole	95-16-9	35	1.0E-07	4.5E-07	2.3E-07	3.8E-07	8.8E-07	1.0E-06
Benzothiazole, 2-phenyl-	883-93-2	35	3.2E-09	1.1E-08	8.6E-09	7.8E-09	2.9E-08	4.3E-08
1,3-Benzothiazole-2-thiol	149-30-4	1	0	3.0E-09	1.8E-08	0	0	1.0E-07
Benzothiazolone	934-34-9	35	2.7E-07	1.3E-06	4.0E-07	1.4E-06	1.8E-06	1.8E-06
Benzyl butyl phthalate	85-68-7	35	3.5E-10	5.1E-09	3.6E-09	4.3E-09	1.3E-08	1.5E-08
Cyclohexyl isothiocyanate	1122-82-3	1	0	1.9E-11	1.1E-10	0	0	6.6E-10
Dibenz[a,h]anthracene	53-70-3	14	0	2.0E-10	3.7E-10	0	1.0E-09	1.6E-09
Dibenzothiophene	132-65-0	20	0	1.2E-10	1.5E-10	1.0E-10	3.4E-10	7.3E-10
Diethyl Phthalate	84-66-2	5	0	9.3E-10	2.8E-09	0	5.1E-09	1.4E-08
Diisobutyl Phthalate	84-69-5	28	0	7.8E-10	1.3E-09	5.2E-10	2.0E-09	7.7E-09
Diisooctylphthalate	27554-26-3	35	2.2E-10	2.6E-08	1.8E-08	2.2E-08	6.4E-08	7.4E-08
Di-n-octyl phthalate	117-84-0	33	0	5.5E-09	7.7E-09	3.0E-09	1.9E-08	3.7E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	4.8E-12	1.4E-11	0	3.7E-11	6.3E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	2.4E-08	6.2E-08	5.5E-09	6.5E-08	3.6E-07
Fluoranthene	206-44-0	35	6.9E-10	4.3E-09	3.3E-09	3.0E-09	1.1E-08	1.2E-08
Fluorene	86-73-7	6	0	1.8E-11	5.6E-11	0	7.7E-11	3.1E-10
Hexanoic Acid, 2-ethyl	149-57-5	1	0	2.1E-11	1.2E-10	0	0	7.2E-10
1-Hydroxypyrene	5315-79-7	1	0	1.2E-09	7.1E-09	0	0	4.2E-08
2-(Methylthio)benzothiazole	615-22-5	9	0	1.6E-08	3.6E-08	0	7.7E-08	1.7E-07
Naphthalene	91-20-3	2	0	2.6E-11	1.1E-10	0	1.3E-10	4.8E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	1.9E-12	1.1E-11	0	0	6.5E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	2.0E-11	4.7E-11	0	1.4E-10	1.5E-10
Naphthalene, 1-methyl-	90-12-0	1	0	4.2E-12	2.5E-11	0	0	1.5E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	7.7E-10	6.2E-10	4.8E-10	1.8E-09	2.4E-09
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	2.3E-10	2.0E-10	1.4E-10	6.6E-10	9.2E-10
Naphthalene, 2-methyl	91-57-6	1	0	4.1E-12	2.4E-11	0	0	1.4E-10
1-Octadecene	112-88-9	27	0	1.3E-09	3.7E-09	3.7E-10	2.3E-09	2.2E-08



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Phenanthrene	85-01-8	29	0	9.1E-10	1.2E-09	4.4E-10	3.0E-09	5.7E-09
Phenanthrene, 1-methyl	832-69-9	35	8.9E-11	4.8E-10	3.6E-10	3.8E-10	1.2E-09	1.5E-09
Phenanthrene, 2-methyl-	2531-84-2	34	0	5.8E-10	5.4E-10	4.4E-10	1.6E-09	2.2E-09
Phenanthrene, 3-methyl	832-71-3	3	0	2.1E-10	7.9E-10	0	1.5E-09	4.2E-09
N-Phenylbenzamide	93-98-1	8	0	1.3E-09	2.7E-09	0	6.3E-09	9.5E-09
Phthalimide	85-41-6	26	0	8.6E-09	2.0E-08	1.2E-10	4.9E-08	8.6E-08
Pyrene	129-00-0	35	2.1E-09	1.2E-08	7.5E-09	1.0E-08	2.7E-08	3.1E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	4.3E-10	8.1E-10	0	1.8E-09	3.7E-09
Triethylene glycol monobutyl ether	143-22-6	7	0	2.2E-09	5.4E-09	0	1.4E-08	2.4E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	3.0E-10	3.6E-10	1.1E-10	8.8E-10	1.4E-09

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-263. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Spectators 2<6 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	9.3E-12	1.6E-11	0	3.8E-11	5.8E-11
Aniline	62-53-3	4	0	3.0E-10	9.3E-10	0	2.3E-09	4.4E-09
Anthracene	120-12-7	20	0	1.0E-10	1.3E-10	7.0E-11	2.6E-10	6.9E-10
Anthracene, 2-methyl-	613-12-7	35	3.6E-11	1.5E-10	1.2E-10	9.8E-11	3.4E-10	5.8E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	2.0E-11	5.7E-11	0	1.2E-10	2.8E-10
Anthracene, 9-phenyl	602-55-1	9	0	5.0E-11	1.1E-10	0	2.8E-10	5.0E-10
Benz[a]anthracene	56-55-3	31	0	4.1E-10	4.7E-10	2.2E-10	1.1E-09	2.2E-09
Benzene, n-butyl-	104-51-8	1	0	6.1E-13	3.6E-12	0	0	2.1E-11
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	2.8E-08	4.4E-08	1.1E-08	8.5E-08	2.4E-07
Benzo[b]fluoranthene	205-99-2	34	0	5.8E-10	5.1E-10	4.6E-10	1.4E-09	2.2E-09
7H-Benzo[c]fluorene	205-12-9	26	0	5.1E-11	6.4E-11	2.6E-11	1.6E-10	2.7E-10
Benzo[k]fluoranthene	207-08-9	29	0	2.5E-10	2.7E-10	1.5E-10	8.2E-10	1.0E-09
Benzothiazole	95-16-9	35	7.6E-08	3.3E-07	1.7E-07	2.8E-07	6.5E-07	7.4E-07



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzothiazole, 2-phenyl-	883-93-2	35	2.4E-09	8.0E-09	6.3E-09	5.7E-09	2.2E-08	3.1E-08
1,3-Benzothiazole-2-thiol	149-30-4	1	0	2.2E-09	1.3E-08	0	0	7.7E-08
Benzothiazolone	934-34-9	35	2.0E-07	9.3E-07	3.0E-07	1.0E-06	1.3E-06	1.3E-06
Benzyl butyl phthalate	85-68-7	35	2.6E-10	3.8E-09	2.7E-09	3.2E-09	9.3E-09	1.1E-08
Cyclohexyl isothiocyanate	1122-82-3	1	0	1.4E-11	8.2E-11	0	0	4.9E-10
Dibenz[a,h]anthracene	53-70-3	14	0	1.5E-10	2.7E-10	0	7.3E-10	1.1E-09
Dibenzothiophene	132-65-0	20	0	9.0E-11	1.1E-10	7.6E-11	2.5E-10	5.4E-10
Diethyl Phthalate	84-66-2	5	0	6.9E-10	2.0E-09	0	3.8E-09	1.0E-08
Diisobutyl Phthalate	84-69-5	28	0	5.7E-10	9.7E-10	3.8E-10	1.5E-09	5.6E-09
Diisooctylphthalate	27554-26-3	35	1.6E-10	1.9E-08	1.3E-08	1.6E-08	4.7E-08	5.4E-08
Di-n-octyl phthalate	117-84-0	33	0	4.0E-09	5.7E-09	2.2E-09	1.4E-08	2.7E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	3.5E-12	1.0E-11	0	2.8E-11	4.6E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	1.8E-08	4.5E-08	4.1E-09	4.8E-08	2.6E-07
Fluoranthene	206-44-0	35	5.1E-10	3.1E-09	2.5E-09	2.2E-09	8.1E-09	8.6E-09
Fluorene	86-73-7	6	0	1.3E-11	4.1E-11	0	5.6E-11	2.3E-10
Hexanoic Acid, 2-ethyl	149-57-5	1	0	1.5E-11	9.0E-11	0	0	5.3E-10
1-Hydroxypyrene	5315-79-7	1	0	8.8E-10	5.2E-09	0	0	3.1E-08
2-(Methylthio)benzothiazole	615-22-5	9	0	1.2E-08	2.7E-08	0	5.6E-08	1.2E-07
Naphthalene	91-20-3	2	0	1.9E-11	8.0E-11	0	9.8E-11	3.5E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	1.4E-12	8.1E-12	0	0	4.8E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	1.4E-11	3.4E-11	0	1.0E-10	1.1E-10
Naphthalene, 1-methyl-	90-12-0	1	0	3.1E-12	1.8E-11	0	0	1.1E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	5.7E-10	4.6E-10	3.5E-10	1.4E-09	1.8E-09
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	1.7E-10	1.5E-10	1.0E-10	4.9E-10	6.8E-10
Naphthalene, 2-methyl	91-57-6	1	0	3.0E-12	1.8E-11	0	0	1.0E-10
1-Octadecene	112-88-9	27	0	9.7E-10	2.7E-09	2.7E-10	1.7E-09	1.6E-08
Phenanthrene	85-01-8	29	0	6.7E-10	9.1E-10	3.3E-10	2.2E-09	4.2E-09
Phenanthrene, 1-methyl	832-69-9	35	6.6E-11	3.5E-10	2.7E-10	2.8E-10	8.5E-10	1.1E-09
Phenanthrene, 2-methyl-	2531-84-2	34	0	4.3E-10	3.9E-10	3.2E-10	1.2E-09	1.6E-09
Phenanthrene, 3-methyl	832-71-3	3	0	1.6E-10	5.8E-10	0	1.1E-09	3.1E-09
N-Phenylbenzamide	93-98-1	8	0	9.7E-10	2.0E-09	0	4.7E-09	7.0E-09
Phthalimide	85-41-6	26	0	6.3E-09	1.5E-08	8.6E-11	3.6E-08	6.3E-08



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Pyrene	129-00-0	35	1.6E-09	8.5E-09	5.5E-09	7.4E-09	2.0E-08	2.2E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	3.1E-10	6.0E-10	0	1.3E-09	2.7E-09
Triethylene glycol monobutyl ether	143-22-6	7	0	1.6E-09	4.0E-09	0	1.0E-08	1.8E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	2.2E-10	2.6E-10	7.8E-11	6.5E-10	1.1E-09

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-264. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Spectators 6<11 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	8.4E-12	1.5E-11	0	3.5E-11	5.2E-11
Aniline	62-53-3	4	0	2.7E-10	8.4E-10	0	2.1E-09	4.0E-09
Anthracene	120-12-7	20	0	9.3E-11	1.2E-10	6.4E-11	2.3E-10	6.2E-10
Anthracene, 2-methyl-	613-12-7	35	3.2E-11	1.3E-10	1.1E-10	8.9E-11	3.1E-10	5.3E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	1.8E-11	5.2E-11	0	1.1E-10	2.5E-10
Anthracene, 9-phenyl	602-55-1	9	0	4.6E-11	1.0E-10	0	2.6E-10	4.6E-10
Benz[a]anthracene	56-55-3	31	0	3.7E-10	4.3E-10	2.0E-10	1.0E-09	2.0E-09
Benzene, n-butyl-	104-51-8	1	0	5.6E-13	3.3E-12	0	0	1.9E-11
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	2.5E-08	4.0E-08	1.0E-08	7.7E-08	2.2E-07
Benzo[b]fluoranthene	205-99-2	34	0	5.3E-10	4.6E-10	4.1E-10	1.3E-09	2.0E-09
7H-Benzo[c]fluorene	205-12-9	26	0	4.6E-11	5.8E-11	2.4E-11	1.5E-10	2.4E-10
Benzo[k]fluoranthene	207-08-9	29	0	2.3E-10	2.4E-10	1.4E-10	7.5E-10	9.3E-10
Benzothiazole	95-16-9	35	6.9E-08	3.0E-07	1.5E-07	2.5E-07	5.9E-07	6.7E-07
Benzothiazole, 2-phenyl-	883-93-2	35	2.1E-09	7.3E-09	5.7E-09	5.2E-09	2.0E-08	2.9E-08
1,3-Benzothiazole-2-thiol	149-30-4	1	0	2.0E-09	1.2E-08	0	0	7.0E-08
Benzothiazolone	934-34-9	35	1.8E-07	8.4E-07	2.7E-07	9.4E-07	1.2E-06	1.2E-06
Benzyl butyl phthalate	85-68-7	35	2.3E-10	3.4E-09	2.4E-09	2.9E-09	8.5E-09	9.7E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	1.3E-11	7.5E-11	0	0	4.4E-10



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dibenz[a,h]anthracene	53-70-3	14	0	1.4E-10	2.5E-10	0	6.7E-10	1.0E-09
Dibenzothiophene	132-65-0	20	0	8.2E-11	1.0E-10	6.9E-11	2.3E-10	4.9E-10
Diethyl Phthalate	84-66-2	5	0	6.2E-10	1.8E-09	0	3.4E-09	9.4E-09
Diisobutyl Phthalate	84-69-5	28	0	5.2E-10	8.8E-10	3.5E-10	1.4E-09	5.1E-09
Diisooctylphthalate	27554-26-3	35	1.5E-10	1.7E-08	1.2E-08	1.5E-08	4.2E-08	4.9E-08
Di-n-octyl phthalate	117-84-0	33	0	3.6E-09	5.1E-09	2.0E-09	1.3E-08	2.5E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	3.2E-12	9.5E-12	0	2.5E-11	4.2E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	1.6E-08	4.1E-08	3.7E-09	4.3E-08	2.4E-07
Fluoranthene	206-44-0	35	4.6E-10	2.8E-09	2.2E-09	2.0E-09	7.4E-09	7.8E-09
Fluorene	86-73-7	6	0	1.2E-11	3.7E-11	0	5.1E-11	2.1E-10
Hexanoic Acid, 2-ethyl	149-57-5	1	0	1.4E-11	8.2E-11	0	0	4.8E-10
1-Hydroxypyrene	5315-79-7	1	0	8.0E-10	4.7E-09	0	0	2.8E-08
2-(Methylthio)benzothiazole	615-22-5	9	0	1.1E-08	2.4E-08	0	5.1E-08	1.1E-07
Naphthalene	91-20-3	2	0	1.8E-11	7.3E-11	0	8.9E-11	3.2E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	1.2E-12	7.4E-12	0	0	4.4E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	1.3E-11	3.1E-11	0	9.5E-11	9.7E-11
Naphthalene, 1-methyl-	90-12-0	1	0	2.8E-12	1.7E-11	0	0	9.8E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	5.2E-10	4.2E-10	3.2E-10	1.2E-09	1.6E-09
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	1.5E-10	1.4E-10	9.1E-11	4.4E-10	6.2E-10
Naphthalene, 2-methyl	91-57-6	1	0	2.7E-12	1.6E-11	0	0	9.5E-11
1-Octadecene	112-88-9	27	0	8.8E-10	2.5E-09	2.4E-10	1.5E-09	1.5E-08
Phenanthrene	85-01-8	29	0	6.1E-10	8.2E-10	3.0E-10	2.0E-09	3.8E-09
Phenanthrene, 1-methyl	832-69-9	35	5.9E-11	3.2E-10	2.4E-10	2.5E-10	7.7E-10	9.7E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	3.9E-10	3.6E-10	2.9E-10	1.1E-09	1.4E-09
Phenanthrene, 3-methyl	832-71-3	3	0	1.4E-10	5.3E-10	0	1.0E-09	2.8E-09
N-Phenylbenzamide	93-98-1	8	0	8.8E-10	1.8E-09	0	4.2E-09	6.4E-09
Phthalimide	85-41-6	26	0	5.7E-09	1.3E-08	7.8E-11	3.3E-08	5.7E-08
Pyrene	129-00-0	35	1.4E-09	7.7E-09	5.0E-09	6.7E-09	1.8E-08	2.0E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	2.8E-10	5.4E-10	0	1.2E-09	2.5E-09
Triethylene glycol monobutyl ether	143-22-6	7	0	1.5E-09	3.6E-09	0	9.3E-09	1.6E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	2.0E-10	2.4E-10	7.0E-11	5.9E-10	9.5E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average



concentrations ($C_{\text{der-crumb rubber-field}}$) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-265. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard ($ADD_{\text{der-field}}$, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Spectators 11<16 Years**

Chemical	CASRN	Detection	$ADD_{\text{der-field}}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	7.2E-12	1.2E-11	0	3.0E-11	4.5E-11
Aniline	62-53-3	4	0	2.3E-10	7.2E-10	0	1.8E-09	3.4E-09
Anthracene	120-12-7	20	0	8.0E-11	1.0E-10	5.5E-11	2.0E-10	5.3E-10
Anthracene, 2-methyl-	613-12-7	35	2.8E-11	1.2E-10	9.4E-11	7.6E-11	2.7E-10	4.5E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	1.6E-11	4.4E-11	0	9.6E-11	2.2E-10
Anthracene, 9-phenyl	602-55-1	9	0	3.9E-11	8.8E-11	0	2.2E-10	3.9E-10
Benz[a]anthracene	56-55-3	31	0	3.2E-10	3.7E-10	1.7E-10	8.9E-10	1.7E-09
Benzene, n-butyl-	104-51-8	1	0	4.8E-13	2.8E-12	0	0	1.7E-11
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	2.1E-08	3.4E-08	8.9E-09	6.6E-08	1.8E-07
Benzo[b]fluoranthene	205-99-2	34	0	4.5E-10	4.0E-10	3.5E-10	1.1E-09	1.7E-09
7H-Benzo[c]fluorene	205-12-9	26	0	3.9E-11	5.0E-11	2.0E-11	1.3E-10	2.1E-10
Benzo[k]fluoranthene	207-08-9	29	0	2.0E-10	2.1E-10	1.2E-10	6.4E-10	7.9E-10
Benzothiazole	95-16-9	35	5.9E-08	2.6E-07	1.3E-07	2.1E-07	5.0E-07	5.7E-07
Benzothiazole, 2-phenyl-	883-93-2	35	1.8E-09	6.2E-09	4.9E-09	4.4E-09	1.7E-08	2.4E-08
1,3-Benzothiazole-2-thiol	149-30-4	1	0	1.7E-09	1.0E-08	0	0	6.0E-08
Benzothiazolone	934-34-9	35	1.5E-07	7.2E-07	2.3E-07	8.1E-07	1.0E-06	1.0E-06
Benzyl butyl phthalate	85-68-7	35	2.0E-10	2.9E-09	2.1E-09	2.5E-09	7.3E-09	8.3E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	1.1E-11	6.4E-11	0	0	3.8E-10
Dibenz[a,h]anthracene	53-70-3	14	0	1.2E-10	2.1E-10	0	5.7E-10	8.9E-10
Dibenzothiophene	132-65-0	20	0	7.0E-11	8.7E-11	5.9E-11	1.9E-10	4.2E-10
Diethyl Phthalate	84-66-2	5	0	5.3E-10	1.6E-09	0	2.9E-09	8.0E-09
Diisobutyl Phthalate	84-69-5	28	0	4.5E-10	7.5E-10	3.0E-10	1.2E-09	4.4E-09
Diisooctylphthalate	27554-26-3	35	1.3E-10	1.5E-08	1.0E-08	1.3E-08	3.6E-08	4.2E-08
Di-n-octyl phthalate	117-84-0	33	0	3.1E-09	4.4E-09	1.7E-09	1.1E-08	2.1E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	2.7E-12	8.1E-12	0	2.1E-11	3.6E-11



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	1.4E-08	3.5E-08	3.2E-09	3.7E-08	2.0E-07
Fluoranthene	206-44-0	35	4.0E-10	2.4E-09	1.9E-09	1.7E-09	6.3E-09	6.7E-09
Fluorene	86-73-7	6	0	1.0E-11	3.2E-11	0	4.4E-11	1.8E-10
Hexanoic Acid, 2-ethyl	149-57-5	1	0	1.2E-11	7.0E-11	0	0	4.1E-10
1-Hydroxypyrene	5315-79-7	1	0	6.9E-10	4.1E-09	0	0	2.4E-08
2-(Methylthio)benzothiazole	615-22-5	9	0	9.4E-09	2.1E-08	0	4.4E-08	9.4E-08
Naphthalene	91-20-3	2	0	1.5E-11	6.2E-11	0	7.6E-11	2.7E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	1.1E-12	6.3E-12	0	0	3.7E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	1.1E-11	2.7E-11	0	8.1E-11	8.3E-11
Naphthalene, 1-methyl-	90-12-0	1	0	2.4E-12	1.4E-11	0	0	8.4E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	4.4E-10	3.6E-10	2.7E-10	1.1E-09	1.4E-09
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	1.3E-10	1.2E-10	7.8E-11	3.8E-10	5.3E-10
Naphthalene, 2-methyl	91-57-6	1	0	2.3E-12	1.4E-11	0	0	8.1E-11
1-Octadecene	112-88-9	27	0	7.6E-10	2.1E-09	2.1E-10	1.3E-09	1.3E-08
Phenanthrene	85-01-8	29	0	5.2E-10	7.1E-10	2.5E-10	1.7E-09	3.3E-09
Phenanthrene, 1-methyl	832-69-9	35	5.1E-11	2.7E-10	2.1E-10	2.1E-10	6.6E-10	8.3E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	3.3E-10	3.1E-10	2.5E-10	9.2E-10	1.2E-09
Phenanthrene, 3-methyl	832-71-3	3	0	1.2E-10	4.5E-10	0	8.6E-10	2.4E-09
N-Phenylbenzamide	93-98-1	8	0	7.5E-10	1.5E-09	0	3.6E-09	5.5E-09
Phthalimide	85-41-6	26	0	4.9E-09	1.1E-08	6.7E-11	2.8E-08	4.9E-08
Pyrene	129-00-0	35	1.2E-09	6.6E-09	4.3E-09	5.7E-09	1.5E-08	1.7E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	2.4E-10	4.6E-10	0	1.0E-09	2.1E-09
Triethylene glycol monobutyl ether	143-22-6	7	0	1.3E-09	3.1E-09	0	7.9E-09	1.4E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	1.7E-10	2.0E-10	6.0E-11	5.0E-10	8.2E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-266. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Spectators 16<30 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	3.2E-12	5.4E-12	0	1.3E-11	2.0E-11
Aniline	62-53-3	4	0	1.0E-10	3.1E-10	0	7.8E-10	1.5E-09
Anthracene	120-12-7	20	0	3.5E-11	4.6E-11	2.4E-11	8.7E-11	2.3E-10
Anthracene, 2-methyl-	613-12-7	35	1.2E-11	5.0E-11	4.1E-11	3.3E-11	1.2E-10	2.0E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	6.9E-12	1.9E-11	0	4.2E-11	9.4E-11
Anthracene, 9-phenyl	602-55-1	9	0	1.7E-11	3.9E-11	0	9.6E-11	1.7E-10
Benz[a]anthracene	56-55-3	31	0	1.4E-10	1.6E-10	7.3E-11	3.9E-10	7.4E-10
Benzene, n-butyl-	104-51-8	1	0	2.1E-13	1.2E-12	0	0	7.3E-12
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	9.4E-09	1.5E-08	3.9E-09	2.9E-08	8.1E-08
Benzo[b]fluoranthene	205-99-2	34	0	2.0E-10	1.7E-10	1.6E-10	4.9E-10	7.6E-10
7H-Benzo[c]fluorene	205-12-9	26	0	1.7E-11	2.2E-11	8.9E-12	5.5E-11	9.1E-11
Benzo[k]fluoranthene	207-08-9	29	0	8.6E-11	9.1E-11	5.2E-11	2.8E-10	3.5E-10
Benzothiazole	95-16-9	35	2.6E-08	1.1E-07	5.8E-08	9.4E-08	2.2E-07	2.5E-07
Benzothiazole, 2-phenyl-	883-93-2	35	8.0E-10	2.7E-09	2.1E-09	1.9E-09	7.3E-09	1.1E-08
1,3-Benzothiazole-2-thiol	149-30-4	1	0	7.4E-10	4.4E-09	0	0	2.6E-08
Benzothiazolone	934-34-9	35	6.8E-08	3.1E-07	1.0E-07	3.5E-07	4.4E-07	4.5E-07
Benzyl butyl phthalate	85-68-7	35	8.7E-11	1.3E-09	9.1E-10	1.1E-09	3.2E-09	3.6E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	4.7E-12	2.8E-11	0	0	1.7E-10
Dibenz[a,h]anthracene	53-70-3	14	0	5.1E-11	9.3E-11	0	2.5E-10	3.9E-10
Dibenzothiophene	132-65-0	20	0	3.1E-11	3.8E-11	2.6E-11	8.5E-11	1.8E-10
Diethyl Phthalate	84-66-2	5	0	2.3E-10	6.9E-10	0	1.3E-09	3.5E-09
Diisobutyl Phthalate	84-69-5	28	0	2.0E-10	3.3E-10	1.3E-10	5.1E-10	1.9E-09
Diisooctylphthalate	27554-26-3	35	5.5E-11	6.4E-09	4.5E-09	5.5E-09	1.6E-08	1.8E-08
Di-n-octyl phthalate	117-84-0	33	0	1.4E-09	1.9E-09	7.6E-10	4.7E-09	9.3E-09
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.2E-12	3.6E-12	0	9.4E-12	1.6E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	6.0E-09	1.5E-08	1.4E-09	1.6E-08	9.0E-08
Fluoranthene	206-44-0	35	1.7E-10	1.1E-09	8.4E-10	7.5E-10	2.8E-09	2.9E-09
Fluorene	86-73-7	6	0	4.5E-12	1.4E-11	0	1.9E-11	7.7E-11
Hexanoic Acid, 2-ethyl	149-57-5	1	0	5.2E-12	3.1E-11	0	0	1.8E-10
1-Hydroxypyrene	5315-79-7	1	0	3.0E-10	1.8E-09	0	0	1.1E-08



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-(Methylthio)benzothiazole	615-22-5	9	0	4.1E-09	9.1E-09	0	1.9E-08	4.1E-08
Naphthalene	91-20-3	2	0	6.6E-12	2.7E-11	0	3.3E-11	1.2E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	4.7E-13	2.8E-12	0	0	1.6E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	4.9E-12	1.2E-11	0	3.5E-11	3.6E-11
Naphthalene, 1-methyl-	90-12-0	1	0	1.0E-12	6.2E-12	0	0	3.7E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	1.9E-10	1.6E-10	1.2E-10	4.6E-10	6.1E-10
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	5.7E-11	5.1E-11	3.4E-11	1.7E-10	2.3E-10
Naphthalene, 2-methyl	91-57-6	1	0	1.0E-12	6.0E-12	0	0	3.6E-11
1-Octadecene	112-88-9	27	0	3.3E-10	9.2E-10	9.2E-11	5.8E-10	5.5E-09
Phenanthrene	85-01-8	29	0	2.3E-10	3.1E-10	1.1E-10	7.6E-10	1.4E-09
Phenanthrene, 1-methyl	832-69-9	35	2.2E-11	1.2E-10	9.1E-11	9.4E-11	2.9E-10	3.7E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	1.4E-10	1.3E-10	1.1E-10	4.0E-10	5.4E-10
Phenanthrene, 3-methyl	832-71-3	3	0	5.3E-11	2.0E-10	0	3.8E-10	1.1E-09
N-Phenylbenzamide	93-98-1	8	0	3.3E-10	6.6E-10	0	1.6E-09	2.4E-09
Phthalimide	85-41-6	26	0	2.1E-09	4.9E-09	2.9E-11	1.2E-08	2.1E-08
Pyrene	129-00-0	35	5.3E-10	2.9E-09	1.9E-09	2.5E-09	6.8E-09	7.6E-09
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	1.1E-10	2.0E-10	0	4.5E-10	9.2E-10
Triethylene glycol monobutyl ether	143-22-6	7	0	5.5E-10	1.3E-09	0	3.5E-09	6.0E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	7.4E-11	8.9E-11	2.6E-11	2.2E-10	3.6E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-267. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Spectators 30<40 Years**

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	2.7E-12	4.7E-12	0	1.1E-11	1.7E-11
Aniline	62-53-3	4	0	8.7E-11	2.7E-10	0	6.7E-10	1.3E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Anthracene	120-12-7	20	0	3.0E-11	4.0E-11	2.1E-11	7.5E-11	2.0E-10
Anthracene, 2-methyl-	613-12-7	35	1.0E-11	4.4E-11	3.6E-11	2.9E-11	1.0E-10	1.7E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	6.0E-12	1.7E-11	0	3.6E-11	8.2E-11
Anthracene, 9-phenyl	602-55-1	9	0	1.5E-11	3.3E-11	0	8.3E-11	1.5E-10
Benz[a]anthracene	56-55-3	31	0	1.2E-10	1.4E-10	6.3E-11	3.4E-10	6.4E-10
Benzene, n-butyl-	104-51-8	1	0	1.8E-13	1.1E-12	0	0	6.3E-12
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	8.1E-09	1.3E-08	3.4E-09	2.5E-08	7.0E-08
Benzo[b]fluoranthene	205-99-2	34	0	1.7E-10	1.5E-10	1.3E-10	4.3E-10	6.6E-10
7H-Benzo[c]fluorene	205-12-9	26	0	1.5E-11	1.9E-11	7.7E-12	4.8E-11	7.9E-11
Benzo[k]fluoranthene	207-08-9	29	0	7.4E-11	7.9E-11	4.5E-11	2.4E-10	3.0E-10
Benzothiazole	95-16-9	35	2.2E-08	9.7E-08	5.0E-08	8.1E-08	1.9E-07	2.2E-07
Benzothiazole, 2-phenyl-	883-93-2	35	6.9E-10	2.4E-09	1.8E-09	1.7E-09	6.3E-09	9.2E-09
1,3-Benzothiazole-2-thiol	149-30-4	1	0	6.4E-10	3.8E-09	0	0	2.3E-08
Benzothiazolone	934-34-9	35	5.8E-08	2.7E-07	8.7E-08	3.1E-07	3.8E-07	3.9E-07
Benzyl butyl phthalate	85-68-7	35	7.5E-11	1.1E-09	7.8E-10	9.3E-10	2.7E-09	3.1E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	4.1E-12	2.4E-11	0	0	1.4E-10
Dibenz[a,h]anthracene	53-70-3	14	0	4.4E-11	8.1E-11	0	2.2E-10	3.4E-10
Dibenzothiophene	132-65-0	20	0	2.7E-11	3.3E-11	2.2E-11	7.3E-11	1.6E-10
Diethyl Phthalate	84-66-2	5	0	2.0E-10	6.0E-10	0	1.1E-09	3.0E-09
Diisobutyl Phthalate	84-69-5	28	0	1.7E-10	2.8E-10	1.1E-10	4.4E-10	1.7E-09
Diisooctylphthalate	27554-26-3	35	4.8E-11	5.6E-09	3.9E-09	4.8E-09	1.4E-08	1.6E-08
Di-n-octyl phthalate	117-84-0	33	0	1.2E-09	1.7E-09	6.6E-10	4.1E-09	8.0E-09
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.0E-12	3.1E-12	0	8.1E-12	1.4E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	5.2E-09	1.3E-08	1.2E-09	1.4E-08	7.7E-08
Fluoranthene	206-44-0	35	1.5E-10	9.2E-10	7.2E-10	6.5E-10	2.4E-09	2.5E-09
Fluorene	86-73-7	6	0	3.9E-12	1.2E-11	0	1.7E-11	6.7E-11
Hexanoic Acid, 2-ethyl	149-57-5	1	0	4.5E-12	2.6E-11	0	0	1.6E-10
1-Hydroxypyrene	5315-79-7	1	0	2.6E-10	1.5E-09	0	0	9.1E-09
2-(Methylthio)benzothiazole	615-22-5	9	0	3.6E-09	7.9E-09	0	1.7E-08	3.6E-08
Naphthalene	91-20-3	2	0	5.7E-12	2.4E-11	0	2.9E-11	1.0E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	4.0E-13	2.4E-12	0	0	1.4E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	4.3E-12	1.0E-11	0	3.1E-11	3.1E-11



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene, 1-methyl-	90-12-0	1	0	9.0E-13	5.3E-12	0	0	3.2E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	1.7E-10	1.3E-10	1.0E-10	4.0E-10	5.3E-10
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	5.0E-11	4.4E-11	3.0E-11	1.4E-10	2.0E-10
Naphthalene, 2-methyl	91-57-6	1	0	8.8E-13	5.2E-12	0	0	3.1E-11
1-Octadecene	112-88-9	27	0	2.9E-10	7.9E-10	7.9E-11	5.0E-10	4.7E-09
Phenanthrene	85-01-8	29	0	2.0E-10	2.7E-10	9.6E-11	6.5E-10	1.2E-09
Phenanthrene, 1-methyl	832-69-9	35	1.9E-11	1.0E-10	7.9E-11	8.1E-11	2.5E-10	3.2E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	1.2E-10	1.2E-10	9.5E-11	3.5E-10	4.7E-10
Phenanthrene, 3-methyl	832-71-3	3	0	4.5E-11	1.7E-10	0	3.2E-10	9.2E-10
N-Phenylbenzamide	93-98-1	8	0	2.8E-10	5.7E-10	0	1.4E-09	2.1E-09
Phthalimide	85-41-6	26	0	1.9E-09	4.3E-09	2.5E-11	1.1E-08	1.8E-08
Pyrene	129-00-0	35	4.6E-10	2.5E-09	1.6E-09	2.2E-09	5.8E-09	6.6E-09
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	9.2E-11	1.8E-10	0	3.9E-10	8.0E-10
Triethylene glycol monobutyl ether	143-22-6	7	0	4.8E-10	1.2E-09	0	3.0E-09	5.2E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	6.4E-11	7.7E-11	2.3E-11	1.9E-10	3.1E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-268. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender **Spectators 40<50 Years****

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	2.8E-12	4.7E-12	0	1.1E-11	1.7E-11
Aniline	62-53-3	4	0	8.8E-11	2.7E-10	0	6.8E-10	1.3E-09
Anthracene	120-12-7	20	0	3.1E-11	4.0E-11	2.1E-11	7.6E-11	2.0E-10
Anthracene, 2-methyl-	613-12-7	35	1.1E-11	4.4E-11	3.6E-11	2.9E-11	1.0E-10	1.7E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	6.0E-12	1.7E-11	0	3.7E-11	8.2E-11
Anthracene, 9-phenyl	602-55-1	9	0	1.5E-11	3.4E-11	0	8.4E-11	1.5E-10
Benz[a]anthracene	56-55-3	31	0	1.2E-10	1.4E-10	6.4E-11	3.4E-10	6.5E-10
Benzene, n-butyl-	104-51-8	1	0	1.8E-13	1.1E-12	0	0	6.4E-12



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	8.2E-09	1.3E-08	3.4E-09	2.5E-08	7.1E-08
Benzo[b]fluoranthene	205-99-2	34	0	1.7E-10	1.5E-10	1.4E-10	4.3E-10	6.7E-10
7H-Benzo[c]fluorene	205-12-9	26	0	1.5E-11	1.9E-11	7.7E-12	4.8E-11	7.9E-11
Benzo[k]fluoranthene	207-08-9	29	0	7.5E-11	7.9E-11	4.6E-11	2.4E-10	3.0E-10
Benzothiazole	95-16-9	35	2.3E-08	9.8E-08	5.0E-08	8.2E-08	1.9E-07	2.2E-07
Benzothiazole, 2-phenyl-	883-93-2	35	7.0E-10	2.4E-09	1.9E-09	1.7E-09	6.4E-09	9.3E-09
1,3-Benzothiazole-2-thiol	149-30-4	1	0	6.5E-10	3.8E-09	0	0	2.3E-08
Benzothiazolone	934-34-9	35	5.9E-08	2.7E-07	8.8E-08	3.1E-07	3.8E-07	3.9E-07
Benzyl butyl phthalate	85-68-7	35	7.6E-11	1.1E-09	7.9E-10	9.4E-10	2.8E-09	3.2E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	4.1E-12	2.4E-11	0	0	1.4E-10
Dibenz[a,h]anthracene	53-70-3	14	0	4.5E-11	8.1E-11	0	2.2E-10	3.4E-10
Dibenzothiophene	132-65-0	20	0	2.7E-11	3.3E-11	2.2E-11	7.4E-11	1.6E-10
Diethyl Phthalate	84-66-2	5	0	2.0E-10	6.0E-10	0	1.1E-09	3.1E-09
Diisobutyl Phthalate	84-69-5	28	0	1.7E-10	2.9E-10	1.1E-10	4.5E-10	1.7E-09
Diisooctylphthalate	27554-26-3	35	4.8E-11	5.6E-09	3.9E-09	4.8E-09	1.4E-08	1.6E-08
Di-n-octyl phthalate	117-84-0	33	0	1.2E-09	1.7E-09	6.6E-10	4.1E-09	8.1E-09
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.0E-12	3.1E-12	0	8.2E-12	1.4E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	5.2E-09	1.3E-08	1.2E-09	1.4E-08	7.8E-08
Fluoranthene	206-44-0	35	1.5E-10	9.3E-10	7.3E-10	6.5E-10	2.4E-09	2.6E-09
Fluorene	86-73-7	6	0	3.9E-12	1.2E-11	0	1.7E-11	6.7E-11
Hexanoic Acid, 2-ethyl	149-57-5	1	0	4.5E-12	2.7E-11	0	0	1.6E-10
1-Hydroxypyrene	5315-79-7	1	0	2.6E-10	1.5E-09	0	0	9.2E-09
2-(Methylthio)benzothiazole	615-22-5	9	0	3.6E-09	7.9E-09	0	1.7E-08	3.6E-08
Naphthalene	91-20-3	2	0	5.8E-12	2.4E-11	0	2.9E-11	1.0E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	4.1E-13	2.4E-12	0	0	1.4E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	4.3E-12	1.0E-11	0	3.1E-11	3.2E-11
Naphthalene, 1-methyl-	90-12-0	1	0	9.1E-13	5.4E-12	0	0	3.2E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	1.7E-10	1.4E-10	1.0E-10	4.0E-10	5.3E-10
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	5.0E-11	4.4E-11	3.0E-11	1.4E-10	2.0E-10
Naphthalene, 2-methyl	91-57-6	1	0	8.9E-13	5.2E-12	0	0	3.1E-11
1-Octadecene	112-88-9	27	0	2.9E-10	8.0E-10	8.0E-11	5.0E-10	4.8E-09



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Phenanthrene	85-01-8	29	0	2.0E-10	2.7E-10	9.7E-11	6.6E-10	1.2E-09
Phenanthrene, 1-methyl	832-69-9	35	1.9E-11	1.0E-10	7.9E-11	8.2E-11	2.5E-10	3.2E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	1.3E-10	1.2E-10	9.6E-11	3.5E-10	4.7E-10
Phenanthrene, 3-methyl	832-71-3	3	0	4.6E-11	1.7E-10	0	3.3E-10	9.3E-10
N-Phenylbenzamide	93-98-1	8	0	2.9E-10	5.8E-10	0	1.4E-09	2.1E-09
Phthalimide	85-41-6	26	0	1.9E-09	4.3E-09	2.5E-11	1.1E-08	1.9E-08
Pyrene	129-00-0	35	4.6E-10	2.5E-09	1.6E-09	2.2E-09	5.9E-09	6.7E-09
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	9.3E-11	1.8E-10	0	3.9E-10	8.0E-10
Triethylene glycol monobutyl ether	143-22-6	7	0	4.8E-10	1.2E-09	0	3.0E-09	5.2E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	6.5E-11	7.8E-11	2.3E-11	1.9E-10	3.1E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-269. Field-Specific Average Dermal Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender Spectators 50<70 Years

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	10	0	2.6E-12	4.5E-12	0	1.1E-11	1.6E-11
Aniline	62-53-3	4	0	8.3E-11	2.6E-10	0	6.4E-10	1.2E-09
Anthracene	120-12-7	20	0	2.9E-11	3.8E-11	2.0E-11	7.2E-11	1.9E-10
Anthracene, 2-methyl-	613-12-7	35	1.0E-11	4.1E-11	3.4E-11	2.7E-11	9.6E-11	1.6E-10
Anthracene, 9,10-diphenyl-	1499-10-1	5	0	5.7E-12	1.6E-11	0	3.4E-11	7.8E-11
Anthracene, 9-phenyl	602-55-1	9	0	1.4E-11	3.2E-11	0	7.9E-11	1.4E-10
Benz[a]anthracene	56-55-3	31	0	1.1E-10	1.3E-10	6.0E-11	3.2E-10	6.1E-10
Benzene, n-butyl-	104-51-8	1	0	1.7E-13	1.0E-12	0	0	6.0E-12
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	32	0	7.7E-09	1.2E-08	3.2E-09	2.4E-08	6.7E-08
Benzo[b]fluoranthene	205-99-2	34	0	1.6E-10	1.4E-10	1.3E-10	4.0E-10	6.3E-10
7H-Benzo[c]fluorene	205-12-9	26	0	1.4E-11	1.8E-11	7.3E-12	4.5E-11	7.5E-11
Benzo[k]fluoranthene	207-08-9	29	0	7.1E-11	7.5E-11	4.3E-11	2.3E-10	2.9E-10
Benzothiazole	95-16-9	35	2.1E-08	9.2E-08	4.8E-08	7.7E-08	1.8E-07	2.1E-07



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzothiazole, 2-phenyl-	883-93-2	35	6.6E-10	2.2E-09	1.8E-09	1.6E-09	6.0E-09	8.8E-09
1,3-Benzothiazole-2-thiol	149-30-4	1	0	6.1E-10	3.6E-09	0	0	2.1E-08
Benzothiazolone	934-34-9	35	5.6E-08	2.6E-07	8.3E-08	2.9E-07	3.6E-07	3.7E-07
Benzyl butyl phthalate	85-68-7	35	7.2E-11	1.1E-09	7.5E-10	8.9E-10	2.6E-09	3.0E-09
Cyclohexyl isothiocyanate	1122-82-3	1	0	3.9E-12	2.3E-11	0	0	1.4E-10
Dibenz[a,h]anthracene	53-70-3	14	0	4.2E-11	7.7E-11	0	2.1E-10	3.2E-10
Dibenzothiophene	132-65-0	20	0	2.5E-11	3.1E-11	2.1E-11	7.0E-11	1.5E-10
Diethyl Phthalate	84-66-2	5	0	1.9E-10	5.7E-10	0	1.1E-09	2.9E-09
Diisobutyl Phthalate	84-69-5	28	0	1.6E-10	2.7E-10	1.1E-10	4.2E-10	1.6E-09
Diisooctylphthalate	27554-26-3	35	4.5E-11	5.3E-09	3.7E-09	4.5E-09	1.3E-08	1.5E-08
Di-n-octyl phthalate	117-84-0	33	0	1.1E-09	1.6E-09	6.3E-10	3.9E-09	7.6E-09
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	9.9E-13	2.9E-12	0	7.7E-12	1.3E-11
3,5-Di-tert-butyl-4-hydroxybenzaldehyde	1620-98-0	25	0	4.9E-09	1.3E-08	1.1E-09	1.3E-08	7.4E-08
Fluoranthene	206-44-0	35	1.4E-10	8.8E-10	6.9E-10	6.1E-10	2.3E-09	2.4E-09
Fluorene	86-73-7	6	0	3.7E-12	1.2E-11	0	1.6E-11	6.3E-11
Hexanoic Acid, 2-ethyl	149-57-5	1	0	4.2E-12	2.5E-11	0	0	1.5E-10
1-Hydroxypyrene	5315-79-7	1	0	2.5E-10	1.5E-09	0	0	8.6E-09
2-(Methylthio)benzothiazole	615-22-5	9	0	3.4E-09	7.5E-09	0	1.6E-08	3.4E-08
Naphthalene	91-20-3	2	0	5.4E-12	2.2E-11	0	2.7E-11	9.9E-11
Naphthalene, 1,2-dimethyl-	573-98-8	1	0	3.8E-13	2.3E-12	0	0	1.3E-11
Naphthalene, 1,6-dimethyl-	575-43-9	6	0	4.1E-12	9.6E-12	0	2.9E-11	3.0E-11
Naphthalene, 1-methyl-	90-12-0	1	0	8.6E-13	5.1E-12	0	0	3.0E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	33	0	1.6E-10	1.3E-10	9.8E-11	3.8E-10	5.0E-10
Naphthalene, 2,3-dimethyl-	581-40-8	33	0	4.7E-11	4.2E-11	2.8E-11	1.4E-10	1.9E-10
Naphthalene, 2-methyl	91-57-6	1	0	8.3E-13	4.9E-12	0	0	2.9E-11
1-Octadecene	112-88-9	27	0	2.7E-10	7.6E-10	7.5E-11	4.8E-10	4.5E-09
Phenanthrene	85-01-8	29	0	1.9E-10	2.5E-10	9.1E-11	6.2E-10	1.2E-09
Phenanthrene, 1-methyl	832-69-9	35	1.8E-11	9.8E-11	7.5E-11	7.7E-11	2.4E-10	3.0E-10
Phenanthrene, 2-methyl-	2531-84-2	34	0	1.2E-10	1.1E-10	9.1E-11	3.3E-10	4.5E-10
Phenanthrene, 3-methyl	832-71-3	3	0	4.3E-11	1.6E-10	0	3.1E-10	8.7E-10
N-Phenylbenzamide	93-98-1	8	0	2.7E-10	5.5E-10	0	1.3E-09	2.0E-09
Phthalimide	85-41-6	26	0	1.8E-09	4.1E-09	2.4E-11	1.0E-08	1.8E-08



Chemical	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Pyrene	129-00-0	35	4.4E-10	2.4E-09	1.5E-09	2.1E-09	5.6E-09	6.3E-09
Pyridine, 2-(4-methylphenyl)-	4467-06-5	14	0	8.7E-11	1.7E-10	0	3.7E-10	7.6E-10
Triethylene glycol monobutyl ether	143-22-6	7	0	4.5E-10	1.1E-09	0	2.9E-09	4.9E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	32	0	6.1E-11	7.4E-11	2.2E-11	1.8E-10	2.9E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

F.5.4. Average Dermal Daily Dose (ADD_{der}) for Lifetime Cancer Risk Assessment of Carcinogens

Table F-270. Average Dermal Daily Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Athletes**

Chemical Name	CASRN	ADD _{der}						
		2<6 Years	6<11 Years	11<16 Years	16<30 Years	30<40 Years	40<50 Years	50<70 Years
Aniline	62-53-3	2.6E-10	2.0E-10	2.1E-10	2.7E-10	1.9E-10	1.6E-10	1.6E-10
Benz[a]anthracene	56-55-3	3.5E-10	2.8E-10	2.9E-10	3.7E-10	2.6E-10	2.2E-10	2.2E-10
Benzo[a]pyrene	50-32-8	2.6E-10	2.1E-10	2.2E-10	2.8E-10	2.0E-10	1.7E-10	1.7E-10
Benzo[b]fluoranthene	205-99-2	5.0E-10	3.9E-10	4.2E-10	5.2E-10	3.7E-10	3.2E-10	3.2E-10
Benzo[k]fluoranthene	207-08-9	2.2E-10	1.7E-10	1.8E-10	2.3E-10	1.6E-10	1.4E-10	1.4E-10
1,3-Benzothiazole-2-thiol	149-30-4	1.9E-09	1.5E-09	1.6E-09	2.0E-09	1.4E-09	1.2E-09	1.2E-09
Chrysene	218-01-9	2.3E-09	1.8E-09	1.9E-09	2.4E-09	1.7E-09	1.4E-09	1.4E-09
Cyclopenta[cd]pyrene	27208-37-3	4.1E-10	3.2E-10	3.4E-10	4.3E-10	3.0E-10	2.6E-10	2.6E-10
Dibenz[a,h]anthracene	53-70-3	1.3E-10	1.0E-10	1.1E-10	1.4E-10	9.7E-11	8.2E-11	8.3E-11
Indeno[1,2,3-cd]pyrene	193-39-5	2.6E-10	2.1E-10	2.2E-10	2.7E-10	1.9E-10	1.6E-10	1.7E-10
Naphthalene	91-20-3	1.7E-11	1.3E-11	1.4E-11	1.8E-11	1.2E-11	1.1E-11	1.1E-11

^a ADD_{der} was calculated from the mean of 35 individual field average concentrations (C_{der-crumb rubber}) of a chemical detected in artificial sweat extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-271. Average Dermal Daily Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Coaches**

Chemical Name	CASRN	ADD _{der}			
		16<30 Years	30<40 Years	40<50 Years	50<70 Years
Aniline	62-53-3	1.3E-10	1.2E-10	1.3E-10	1.3E-10
Benz[a]anthracene	56-55-3	1.8E-10	1.7E-10	1.7E-10	1.7E-10
Benzo[a]pyrene	50-32-8	1.4E-10	1.3E-10	1.3E-10	1.3E-10
Benzo[b]fluoranthene	205-99-2	2.6E-10	2.4E-10	2.5E-10	2.5E-10
Benzo[k]fluoranthene	207-08-9	1.1E-10	1.1E-10	1.1E-10	1.1E-10
1,3-Benzothiazole-2-thiol	149-30-4	9.8E-10	9.2E-10	9.3E-10	9.3E-10
Chrysene	218-01-9	1.2E-09	1.1E-09	1.1E-09	1.1E-09
Cyclopenta[cd]pyrene	27208-37-3	2.1E-10	2.0E-10	2.0E-10	2.0E-10
Dibenz[a,h]anthracene	53-70-3	6.7E-11	6.3E-11	6.4E-11	6.4E-11
Indeno[1,2,3-cd]pyrene	193-39-5	1.4E-10	1.3E-10	1.3E-10	1.3E-10
Naphthalene	91-20-3	8.7E-12	8.1E-12	8.2E-12	8.3E-12

^a ADD_{der} was calculated from the mean of 35 individual field average concentrations (C_{der-crumb rubber}) of a chemical detected in artificial sweat extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-272. Average Dermal Daily Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Referees**

Chemical Name	CASRN	ADD _{der}			
		16<30 Years	30<40 Years	40<50 Years	50<70 Years
Aniline	62-53-3	5.4E-11	5.0E-11	5.1E-11	5.1E-11
Benz[a]anthracene	56-55-3	7.3E-11	6.9E-11	7.0E-11	7.0E-11
Benzo[a]pyrene	50-32-8	5.5E-11	5.2E-11	5.2E-11	5.3E-11
Benzo[b]fluoranthene	205-99-2	1.0E-10	9.8E-11	9.9E-11	9.9E-11
Benzo[k]fluoranthene	207-08-9	4.6E-11	4.3E-11	4.3E-11	4.3E-11
1,3-Benzothiazole-2-thiol	149-30-4	3.9E-10	3.7E-10	3.7E-10	3.7E-10
Chrysene	218-01-9	4.7E-10	4.4E-10	4.5E-10	4.5E-10
Cyclopenta[cd]pyrene	27208-37-3	8.5E-11	7.9E-11	8.0E-11	8.1E-11
Dibenz[a,h]anthracene	53-70-3	2.7E-11	2.5E-11	2.6E-11	2.6E-11
Indeno[1,2,3-cd]pyrene	193-39-5	5.4E-11	5.1E-11	5.2E-11	5.2E-11
Naphthalene	91-20-3	3.5E-12	3.3E-12	3.3E-12	3.3E-12

^a ADD_{der} was calculated from the mean of 35 individual field average concentrations (C_{der-crumb rubber}) of a chemical detected in artificial sweat extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number



Values are rounded to two significant figures.

Table F-273. Average Dermal Daily Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Spectators**

Chemical Name	CASRN	ADD _{der}								
		Third Trimester Fetus	0<2 Years	2<6 Years	6<11 Years	11<16 Years	16<30 Years	30<40 Years	40<50 Years	50<70 Years
Aniline	62-53-3	7.6E-11	4.1E-10	3.0E-10	2.7E-10	2.3E-10	1.0E-10	8.7E-11	8.8E-11	8.3E-11
Benz[a]anthracene	56-55-3	1.0E-10	5.6E-10	4.1E-10	3.7E-10	3.2E-10	1.4E-10	1.2E-10	1.2E-10	1.1E-10
Benzo[a]pyrene	50-32-8	7.8E-11	4.2E-10	3.1E-10	2.8E-10	2.4E-10	1.0E-10	9.0E-11	9.1E-11	8.6E-11
Benzo[b]fluoranthene	205-99-2	1.5E-10	7.9E-10	5.8E-10	5.3E-10	4.5E-10	2.0E-10	1.7E-10	1.7E-10	1.6E-10
Benzo[k]fluoranthene	207-08-9	6.4E-11	3.4E-10	2.5E-10	2.3E-10	2.0E-10	8.6E-11	7.4E-11	7.5E-11	7.1E-11
1,3-Benzothiazole-2-thiol	149-30-4	5.6E-10	3.0E-09	2.2E-09	2.0E-09	1.7E-09	7.4E-10	6.4E-10	6.5E-10	6.1E-10
Chrysene	218-01-9	6.7E-10	3.6E-09	2.6E-09	2.4E-09	2.0E-09	9.0E-10	7.7E-10	7.8E-10	7.4E-10
Cyclopenta[cd]pyrene	27208-37-3	1.2E-10	6.4E-10	4.7E-10	4.3E-10	3.7E-10	1.6E-10	1.4E-10	1.4E-10	1.3E-10
Dibenz[a,h]anthracene	53-70-3	3.8E-11	2.0E-10	1.5E-10	1.4E-10	1.2E-10	5.1E-11	4.4E-11	4.5E-11	4.2E-11
Indeno[1,2,3-cd]pyrene	193-39-5	7.7E-11	4.1E-10	3.0E-10	2.7E-10	2.3E-10	1.0E-10	8.9E-11	9.0E-11	8.5E-11
Naphthalene	91-20-3	4.9E-12	2.6E-11	1.9E-11	1.8E-11	1.5E-11	6.6E-12	5.7E-12	5.8E-12	5.4E-12

^a ADD_{der} was calculated from the mean of 35 individual field average concentrations (C_{der-crumb rubber}) of a chemical detected in artificial sweat extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

INDIVIDUAL FIELD ASSESSMENT (Table F-274 to Table F-297)

Table F-274. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Athletes 2<6 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	2.6E-10	8.0E-10	0	2.0E-09	3.8E-09
Benz[a]anthracene	56-55-3	31	0	3.5E-10	4.1E-10	1.9E-10	9.9E-10	1.9E-09
Benzo[a]pyrene	50-32-8	32	0	2.6E-10	2.6E-10	1.7E-10	6.9E-10	1.1E-09
Benzo[b]fluoranthene	205-99-2	34	0	5.0E-10	4.4E-10	3.9E-10	1.2E-09	1.9E-09
Benzo[k]fluoranthene	207-08-9	29	0	2.2E-10	2.3E-10	1.3E-10	7.1E-10	8.8E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	1.9E-09	1.1E-08	0	0	6.6E-08
Chrysene	218-01-9	35	2.5E-10	2.3E-09	1.5E-09	2.3E-09	4.6E-09	5.2E-09
Cyclopenta[cd]pyrene	27208-37-3	35	4.1E-11	4.1E-10	2.4E-10	3.9E-10	7.8E-10	9.3E-10
Dibenz[a,h]anthracene	53-70-3	14	0	1.3E-10	2.4E-10	0	6.3E-10	9.9E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.6E-10	2.2E-10	2.3E-10	6.1E-10	9.5E-10



Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene	91-20-3	2	0	1.7E-11	6.9E-11	0	8.4E-11	3.0E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-275. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Athletes 6<11 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	2.0E-10	6.3E-10	0	1.6E-09	3.0E-09
Benz[a]anthracene	56-55-3	31	0	2.8E-10	3.2E-10	1.5E-10	7.8E-10	1.5E-09
Benzo[a]pyrene	50-32-8	32	0	2.1E-10	2.1E-10	1.4E-10	5.5E-10	8.5E-10
Benzo[b]fluoranthene	205-99-2	34	0	3.9E-10	3.5E-10	3.1E-10	9.9E-10	1.5E-09
Benzo[k]fluoranthene	207-08-9	29	0	1.7E-10	1.8E-10	1.0E-10	5.6E-10	6.9E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	1.5E-09	8.8E-09	0	0	5.2E-08
Chrysene	218-01-9	35	2.0E-10	1.8E-09	1.2E-09	1.8E-09	3.6E-09	4.1E-09
Cyclopenta[cd]pyrene	27208-37-3	35	3.3E-11	3.2E-10	1.9E-10	3.1E-10	6.1E-10	7.3E-10
Dibenz[a,h]anthracene	53-70-3	14	0	1.0E-10	1.9E-10	0	5.0E-10	7.8E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.1E-10	1.7E-10	1.8E-10	4.8E-10	7.5E-10
Naphthalene	91-20-3	2	0	1.3E-11	5.5E-11	0	6.7E-11	2.4E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-276. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Athletes 11<16 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	2.1E-10	6.6E-10	0	1.6E-09	3.2E-09
Benz[a]anthracene	56-55-3	31	0	2.9E-10	3.4E-10	1.5E-10	8.2E-10	1.6E-09



Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	32	0	2.2E-10	2.2E-10	1.4E-10	5.8E-10	9.0E-10
Benzo[b]fluoranthene	205-99-2	34	0	4.2E-10	3.7E-10	3.3E-10	1.0E-09	1.6E-09
Benzo[k]fluoranthene	207-08-9	29	0	1.8E-10	1.9E-10	1.1E-10	5.9E-10	7.3E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	1.6E-09	9.3E-09	0	0	5.5E-08
Chrysene	218-01-9	35	2.1E-10	1.9E-09	1.2E-09	1.9E-09	3.8E-09	4.3E-09
Cyclopenta[cd]pyrene	27208-37-3	35	3.5E-11	3.4E-10	2.0E-10	3.3E-10	6.5E-10	7.7E-10
Dibenz[a,h]anthracene	53-70-3	14	0	1.1E-10	2.0E-10	0	5.3E-10	8.2E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.2E-10	1.8E-10	1.9E-10	5.1E-10	7.9E-10
Naphthalene	91-20-3	2	0	1.4E-11	5.8E-11	0	7.0E-11	2.5E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-277. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Athletes 16<30 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	2.7E-10	8.4E-10	0	2.1E-09	4.0E-09
Benz[a]anthracene	56-55-3	31	0	3.7E-10	4.3E-10	2.0E-10	1.0E-09	2.0E-09
Benzo[a]pyrene	50-32-8	32	0	2.8E-10	2.7E-10	1.8E-10	7.3E-10	1.1E-09
Benzo[b]fluoranthene	205-99-2	34	0	5.2E-10	4.6E-10	4.1E-10	1.3E-09	2.0E-09
Benzo[k]fluoranthene	207-08-9	29	0	2.3E-10	2.4E-10	1.4E-10	7.4E-10	9.2E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	2.0E-09	1.2E-08	0	0	6.9E-08
Chrysene	218-01-9	35	2.6E-10	2.4E-09	1.5E-09	2.4E-09	4.8E-09	5.4E-09
Cyclopenta[cd]pyrene	27208-37-3	35	4.4E-11	4.3E-10	2.5E-10	4.1E-10	8.2E-10	9.8E-10
Dibenz[a,h]anthracene	53-70-3	14	0	1.4E-10	2.5E-10	0	6.6E-10	1.0E-09
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.7E-10	2.3E-10	2.4E-10	6.4E-10	1.0E-09
Naphthalene	91-20-3	2	0	1.8E-11	7.3E-11	0	8.9E-11	3.2E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number



Values are rounded to two significant figures.

Table F-278. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Athletes 30<40 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	1.9E-10	5.9E-10	0	1.5E-09	2.8E-09
Benz[a]anthracene	56-55-3	31	0	2.6E-10	3.0E-10	1.4E-10	7.4E-10	1.4E-09
Benzo[a]pyrene	50-32-8	32	0	2.0E-10	1.9E-10	1.3E-10	5.2E-10	8.1E-10
Benzo[b]fluoranthene	205-99-2	34	0	3.7E-10	3.3E-10	2.9E-10	9.3E-10	1.4E-09
Benzo[k]fluoranthene	207-08-9	29	0	1.6E-10	1.7E-10	9.9E-11	5.3E-10	6.6E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	1.4E-09	8.3E-09	0	0	4.9E-08
Chrysene	218-01-9	35	1.8E-10	1.7E-09	1.1E-09	1.7E-09	3.4E-09	3.8E-09
Cyclopenta[cd]pyrene	27208-37-3	35	3.1E-11	3.0E-10	1.8E-10	2.9E-10	5.8E-10	6.9E-10
Dibenz[a,h]anthracene	53-70-3	14	0	9.7E-11	1.8E-10	0	4.7E-10	7.4E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	1.9E-10	1.6E-10	1.7E-10	4.5E-10	7.1E-10
Naphthalene	91-20-3	2	0	1.2E-11	5.1E-11	0	6.3E-11	2.3E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-279. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Athletes 40<50 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	1.6E-10	5.0E-10	0	1.2E-09	2.4E-09
Benz[a]anthracene	56-55-3	31	0	2.2E-10	2.6E-10	1.2E-10	6.3E-10	1.2E-09
Benzo[a]pyrene	50-32-8	32	0	1.7E-10	1.7E-10	1.1E-10	4.4E-10	6.8E-10
Benzo[b]fluoranthene	205-99-2	34	0	3.2E-10	2.8E-10	2.5E-10	7.9E-10	1.2E-09
Benzo[k]fluoranthene	207-08-9	29	0	1.4E-10	1.5E-10	8.4E-11	4.5E-10	5.6E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	1.2E-09	7.1E-09	0	0	4.2E-08
Chrysene	218-01-9	35	1.6E-10	1.4E-09	9.3E-10	1.4E-09	2.9E-09	3.3E-09
Cyclopenta[cd]pyrene	27208-37-3	35	2.6E-11	2.6E-10	1.5E-10	2.5E-10	4.9E-10	5.9E-10
Dibenz[a,h]anthracene	53-70-3	14	0	8.2E-11	1.5E-10	0	4.0E-10	6.3E-10



Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	1.6E-10	1.4E-10	1.5E-10	3.8E-10	6.0E-10
Naphthalene	91-20-3	2	0	1.1E-11	4.4E-11	0	5.3E-11	1.9E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-280. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Athletes 50<70 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	1.6E-10	5.1E-10	0	1.3E-09	2.4E-09
Benz[a]anthracene	56-55-3	31	0	2.2E-10	2.6E-10	1.2E-10	6.3E-10	1.2E-09
Benzo[a]pyrene	50-32-8	32	0	1.7E-10	1.7E-10	1.1E-10	4.4E-10	6.9E-10
Benzo[b]fluoranthene	205-99-2	34	0	3.2E-10	2.8E-10	2.5E-10	8.0E-10	1.2E-09
Benzo[k]fluoranthene	207-08-9	29	0	1.4E-10	1.5E-10	8.4E-11	4.5E-10	5.6E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	1.2E-09	7.1E-09	0	0	4.2E-08
Chrysene	218-01-9	35	1.6E-10	1.4E-09	9.3E-10	1.4E-09	2.9E-09	3.3E-09
Cyclopenta[cd]pyrene	27208-37-3	35	2.6E-11	2.6E-10	1.5E-10	2.5E-10	5.0E-10	5.9E-10
Dibenz[a,h]anthracene	53-70-3	14	0	8.3E-11	1.5E-10	0	4.0E-10	6.3E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	1.7E-10	1.4E-10	1.5E-10	3.9E-10	6.1E-10
Naphthalene	91-20-3	2	0	1.1E-11	4.4E-11	0	5.4E-11	1.9E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-281. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Coaches 16<30 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	1.3E-10	4.1E-10	0	1.0E-09	2.0E-09



Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benz[a]anthracene	56-55-3	31	0	1.8E-10	2.1E-10	9.6E-11	5.1E-10	9.7E-10
Benzo[a]pyrene	50-32-8	32	0	1.4E-10	1.4E-10	9.0E-11	3.6E-10	5.6E-10
Benzo[b]fluoranthene	205-99-2	34	0	2.6E-10	2.3E-10	2.0E-10	6.5E-10	1.0E-09
Benzo[k]fluoranthene	207-08-9	29	0	1.1E-10	1.2E-10	6.9E-11	3.7E-10	4.6E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	9.8E-10	5.8E-09	0	0	3.4E-08
Chrysene	218-01-9	35	1.3E-10	1.2E-09	7.6E-10	1.2E-09	2.4E-09	2.7E-09
Cyclopenta[cd]pyrene	27208-37-3	35	2.2E-11	2.1E-10	1.3E-10	2.0E-10	4.0E-10	4.8E-10
Dibenz[a,h]anthracene	53-70-3	14	0	6.7E-11	1.2E-10	0	3.3E-10	5.1E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	1.4E-10	1.1E-10	1.2E-10	3.1E-10	4.9E-10
Naphthalene	91-20-3	2	0	8.7E-12	3.6E-11	0	4.4E-11	1.6E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-282. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Coaches 30<40 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	1.2E-10	3.9E-10	0	9.6E-10	1.9E-09
Benz[a]anthracene	56-55-3	31	0	1.7E-10	2.0E-10	9.0E-11	4.8E-10	9.1E-10
Benzo[a]pyrene	50-32-8	32	0	1.3E-10	1.3E-10	8.5E-11	3.4E-10	5.3E-10
Benzo[b]fluoranthene	205-99-2	34	0	2.4E-10	2.1E-10	1.9E-10	6.1E-10	9.4E-10
Benzo[k]fluoranthene	207-08-9	29	0	1.1E-10	1.1E-10	6.4E-11	3.4E-10	4.3E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	9.2E-10	5.4E-09	0	0	3.2E-08
Chrysene	218-01-9	35	1.2E-10	1.1E-09	7.1E-10	1.1E-09	2.2E-09	2.5E-09
Cyclopenta[cd]pyrene	27208-37-3	35	2.0E-11	2.0E-10	1.2E-10	1.9E-10	3.8E-10	4.5E-10
Dibenz[a,h]anthracene	53-70-3	14	0	6.3E-11	1.2E-10	0	3.1E-10	4.8E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	1.3E-10	1.1E-10	1.1E-10	3.0E-10	4.6E-10
Naphthalene	91-20-3	2	0	8.1E-12	3.4E-11	0	4.1E-11	1.5E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.



CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures.

Table F-283. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Coaches 40<50 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	1.3E-10	3.9E-10	0	9.7E-10	1.9E-09
Benz[a]anthracene	56-55-3	31	0	1.7E-10	2.0E-10	9.1E-11	4.9E-10	9.2E-10
Benzo[a]pyrene	50-32-8	32	0	1.3E-10	1.3E-10	8.5E-11	3.4E-10	5.3E-10
Benzo[b]fluoranthene	205-99-2	34	0	2.5E-10	2.2E-10	1.9E-10	6.1E-10	9.5E-10
Benzo[k]fluoranthene	207-08-9	29	0	1.1E-10	1.1E-10	6.5E-11	3.5E-10	4.3E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	9.3E-10	5.5E-09	0	0	3.2E-08
Chrysene	218-01-9	35	1.2E-10	1.1E-09	7.2E-10	1.1E-09	2.2E-09	2.5E-09
Cyclopenta[cd]pyrene	27208-37-3	35	2.0E-11	2.0E-10	1.2E-10	1.9E-10	3.8E-10	4.6E-10
Dibenz[a,h]anthracene	53-70-3	14	0	6.4E-11	1.2E-10	0	3.1E-10	4.9E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	1.3E-10	1.1E-10	1.1E-10	3.0E-10	4.7E-10
Naphthalene	91-20-3	2	0	8.2E-12	3.4E-11	0	4.2E-11	1.5E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures.

Table F-284. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Coaches 50<70 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	1.3E-10	3.9E-10	0	9.7E-10	1.9E-09
Benz[a]anthracene	56-55-3	31	0	1.7E-10	2.0E-10	9.2E-11	4.9E-10	9.2E-10
Benzo[a]pyrene	50-32-8	32	0	1.3E-10	1.3E-10	8.6E-11	3.4E-10	5.3E-10
Benzo[b]fluoranthene	205-99-2	34	0	2.5E-10	2.2E-10	1.9E-10	6.2E-10	9.5E-10
Benzo[k]fluoranthene	207-08-9	29	0	1.1E-10	1.1E-10	6.5E-11	3.5E-10	4.3E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	9.3E-10	5.5E-09	0	0	3.3E-08
Chrysene	218-01-9	35	1.2E-10	1.1E-09	7.2E-10	1.1E-09	2.3E-09	2.5E-09
Cyclopenta[cd]pyrene	27208-37-3	35	2.0E-11	2.0E-10	1.2E-10	1.9E-10	3.8E-10	4.6E-10
Dibenz[a,h]anthracene	53-70-3	14	0	6.4E-11	1.2E-10	0	3.1E-10	4.9E-10



Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	1.3E-10	1.1E-10	1.1E-10	3.0E-10	4.7E-10
Naphthalene	91-20-3	2	0	8.3E-12	3.4E-11	0	4.2E-11	1.5E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-285. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Referees 16<30 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	5.4E-11	1.7E-10	0	4.1E-10	8.0E-10
Benz[a]anthracene	56-55-3	31	0	7.3E-11	8.5E-11	3.9E-11	2.1E-10	3.9E-10
Benzo[a]pyrene	50-32-8	32	0	5.5E-11	5.5E-11	3.6E-11	1.4E-10	2.3E-10
Benzo[b]fluoranthene	205-99-2	34	0	1.0E-10	9.2E-11	8.2E-11	2.6E-10	4.0E-10
Benzo[k]fluoranthene	207-08-9	29	0	4.6E-11	4.8E-11	2.8E-11	1.5E-10	1.8E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	3.9E-10	2.3E-09	0	0	1.4E-08
Chrysene	218-01-9	35	5.2E-11	4.7E-10	3.1E-10	4.7E-10	9.5E-10	1.1E-09
Cyclopenta[cd]pyrene	27208-37-3	35	8.7E-12	8.5E-11	5.1E-11	8.2E-11	1.6E-10	1.9E-10
Dibenz[a,h]anthracene	53-70-3	14	0	2.7E-11	4.9E-11	0	1.3E-10	2.1E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	5.4E-11	4.5E-11	4.8E-11	1.3E-10	2.0E-10
Naphthalene	91-20-3	2	0	3.5E-12	1.4E-11	0	1.8E-11	6.4E-11

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-286. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Referees 30<40 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	5.0E-11	1.6E-10	0	3.8E-10	7.5E-10



Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benz[a]anthracene	56-55-3	31	0	6.9E-11	8.0E-11	3.6E-11	1.9E-10	3.7E-10
Benzo[a]pyrene	50-32-8	32	0	5.2E-11	5.1E-11	3.4E-11	1.4E-10	2.1E-10
Benzo[b]fluoranthene	205-99-2	34	0	9.8E-11	8.6E-11	7.7E-11	2.4E-10	3.8E-10
Benzo[k]fluoranthene	207-08-9	29	0	4.3E-11	4.5E-11	2.6E-11	1.4E-10	1.7E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	3.7E-10	2.2E-09	0	0	1.3E-08
Chrysene	218-01-9	35	4.8E-11	4.4E-10	2.9E-10	4.4E-10	8.9E-10	1.0E-09
Cyclopenta[cd]pyrene	27208-37-3	35	8.1E-12	7.9E-11	4.7E-11	7.7E-11	1.5E-10	1.8E-10
Dibenz[a,h]anthracene	53-70-3	14	0	2.5E-11	4.6E-11	0	1.2E-10	1.9E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	5.1E-11	4.3E-11	4.5E-11	1.2E-10	1.9E-10
Naphthalene	91-20-3	2	0	3.3E-12	1.4E-11	0	1.7E-11	6.0E-11

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-287. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Referees 40<50 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	5.1E-11	1.6E-10	0	3.9E-10	7.6E-10
Benz[a]anthracene	56-55-3	31	0	7.0E-11	8.0E-11	3.7E-11	2.0E-10	3.7E-10
Benzo[a]pyrene	50-32-8	32	0	5.2E-11	5.2E-11	3.4E-11	1.4E-10	2.1E-10
Benzo[b]fluoranthene	205-99-2	34	0	9.9E-11	8.7E-11	7.8E-11	2.5E-10	3.8E-10
Benzo[k]fluoranthene	207-08-9	29	0	4.3E-11	4.6E-11	2.6E-11	1.4E-10	1.7E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	3.7E-10	2.2E-09	0	0	1.3E-08
Chrysene	218-01-9	35	4.9E-11	4.5E-10	2.9E-10	4.5E-10	9.0E-10	1.0E-09
Cyclopenta[cd]pyrene	27208-37-3	35	8.2E-12	8.0E-11	4.8E-11	7.7E-11	1.5E-10	1.8E-10
Dibenz[a,h]anthracene	53-70-3	14	0	2.6E-11	4.7E-11	0	1.2E-10	2.0E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	5.2E-11	4.3E-11	4.6E-11	1.2E-10	1.9E-10
Naphthalene	91-20-3	2	0	3.3E-12	1.4E-11	0	1.7E-11	6.0E-11

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.



CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures.

Table F-288. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Referees 50<70 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	5.1E-11	1.6E-10	0	3.9E-10	7.6E-10
Benz[a]anthracene	56-55-3	31	0	7.0E-11	8.1E-11	3.7E-11	2.0E-10	3.7E-10
Benzo[a]pyrene	50-32-8	32	0	5.3E-11	5.2E-11	3.4E-11	1.4E-10	2.1E-10
Benzo[b]fluoranthene	205-99-2	34	0	9.9E-11	8.7E-11	7.8E-11	2.5E-10	3.8E-10
Benzo[k]fluoranthene	207-08-9	29	0	4.3E-11	4.6E-11	2.6E-11	1.4E-10	1.7E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	3.7E-10	2.2E-09	0	0	1.3E-08
Chrysene	218-01-9	35	4.9E-11	4.5E-10	2.9E-10	4.5E-10	9.1E-10	1.0E-09
Cyclopenta[cd]pyrene	27208-37-3	35	8.2E-12	8.1E-11	4.8E-11	7.8E-11	1.5E-10	1.8E-10
Dibenz[a,h]anthracene	53-70-3	14	0	2.6E-11	4.7E-11	0	1.3E-10	2.0E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	5.2E-11	4.3E-11	4.6E-11	1.2E-10	1.9E-10
Naphthalene	91-20-3	2	0	3.3E-12	1.4E-11	0	1.7E-11	6.0E-11

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures.

Table F-289. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Spectators Third Trimester Fetus<0 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	7.6E-11	2.3E-10	0	5.8E-10	1.1E-09
Benz[a]anthracene	56-55-3	31	0	1.0E-10	1.2E-10	5.5E-11	2.9E-10	5.5E-10
Benzo[a]pyrene	50-32-8	32	0	7.8E-11	7.7E-11	5.1E-11	2.0E-10	3.2E-10
Benzo[b]fluoranthene	205-99-2	34	0	1.5E-10	1.3E-10	1.2E-10	3.7E-10	5.7E-10
Benzo[k]fluoranthene	207-08-9	29	0	6.4E-11	6.8E-11	3.9E-11	2.1E-10	2.6E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	5.6E-10	3.3E-09	0	0	1.9E-08
Chrysene	218-01-9	35	7.3E-11	6.7E-10	4.3E-10	6.7E-10	1.3E-09	1.5E-09
Cyclopenta[cd]pyrene	27208-37-3	35	1.2E-11	1.2E-10	7.1E-11	1.2E-10	2.3E-10	2.7E-10



Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dibenz[a,h]anthracene	53-70-3	14	0	3.8E-11	7.0E-11	0	1.9E-10	2.9E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	7.7E-11	6.4E-11	6.8E-11	1.8E-10	2.8E-10
Naphthalene	91-20-3	2	0	4.9E-12	2.0E-11	0	2.5E-11	9.0E-11

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-290. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 0<2 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	4.1E-10	1.3E-09	0	3.1E-09	6.0E-09
Benz[a]anthracene	56-55-3	31	0	5.6E-10	6.4E-10	2.9E-10	1.6E-09	3.0E-09
Benzo[a]pyrene	50-32-8	32	0	4.2E-10	4.1E-10	2.7E-10	1.1E-09	1.7E-09
Benzo[b]fluoranthene	205-99-2	34	0	7.9E-10	7.0E-10	6.2E-10	2.0E-09	3.1E-09
Benzo[k]fluoranthene	207-08-9	29	0	3.4E-10	3.6E-10	2.1E-10	1.1E-09	1.4E-09
1,3-Benzothiazole-2-thiol	149-30-4	1	0	3.0E-09	1.8E-08	0	0	1.0E-07
Chrysene	218-01-9	35	3.9E-10	3.6E-09	2.3E-09	3.6E-09	7.2E-09	8.1E-09
Cyclopenta[cd]pyrene	27208-37-3	35	6.5E-11	6.4E-10	3.8E-10	6.2E-10	1.2E-09	1.5E-09
Dibenz[a,h]anthracene	53-70-3	14	0	2.0E-10	3.7E-10	0	1.0E-09	1.6E-09
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	4.1E-10	3.4E-10	3.6E-10	9.6E-10	1.5E-09
Naphthalene	91-20-3	2	0	2.6E-11	1.1E-10	0	1.3E-10	4.8E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-291. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 2<6 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	3.0E-10	9.3E-10	0	2.3E-09	4.4E-09
Benz[a]anthracene	56-55-3	31	0	4.1E-10	4.7E-10	2.2E-10	1.1E-09	2.2E-09
Benzo[a]pyrene	50-32-8	32	0	3.1E-10	3.0E-10	2.0E-10	8.0E-10	1.3E-09
Benzo[b]fluoranthene	205-99-2	34	0	5.8E-10	5.1E-10	4.6E-10	1.4E-09	2.2E-09
Benzo[k]fluoranthene	207-08-9	29	0	2.5E-10	2.7E-10	1.5E-10	8.2E-10	1.0E-09
1,3-Benzothiazole-2-thiol	149-30-4	1	0	2.2E-09	1.3E-08	0	0	7.7E-08
Chrysene	218-01-9	35	2.9E-10	2.6E-09	1.7E-09	2.6E-09	5.3E-09	6.0E-09
Cyclopenta[cd]pyrene	27208-37-3	35	4.8E-11	4.7E-10	2.8E-10	4.5E-10	9.0E-10	1.1E-09
Dibenz[a,h]anthracene	53-70-3	14	0	1.5E-10	2.7E-10	0	7.3E-10	1.1E-09
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	3.0E-10	2.5E-10	2.7E-10	7.0E-10	1.1E-09
Naphthalene	91-20-3	2	0	1.9E-11	8.0E-11	0	9.8E-11	3.5E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-292. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 6<11 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	2.7E-10	8.4E-10	0	2.1E-09	4.0E-09
Benz[a]anthracene	56-55-3	31	0	3.7E-10	4.3E-10	2.0E-10	1.0E-09	2.0E-09
Benzo[a]pyrene	50-32-8	32	0	2.8E-10	2.7E-10	1.8E-10	7.3E-10	1.1E-09
Benzo[b]fluoranthene	205-99-2	34	0	5.3E-10	4.6E-10	4.1E-10	1.3E-09	2.0E-09
Benzo[k]fluoranthene	207-08-9	29	0	2.3E-10	2.4E-10	1.4E-10	7.5E-10	9.3E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	2.0E-09	1.2E-08	0	0	7.0E-08
Chrysene	218-01-9	35	2.6E-10	2.4E-09	1.5E-09	2.4E-09	4.8E-09	5.4E-09
Cyclopenta[cd]pyrene	27208-37-3	35	4.4E-11	4.3E-10	2.6E-10	4.1E-10	8.2E-10	9.8E-10
Dibenz[a,h]anthracene	53-70-3	14	0	1.4E-10	2.5E-10	0	6.7E-10	1.0E-09
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.7E-10	2.3E-10	2.4E-10	6.4E-10	1.0E-09
Naphthalene	91-20-3	2	0	1.8E-11	7.3E-11	0	8.9E-11	3.2E-10



^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-293. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 11<16 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	2.3E-10	7.2E-10	0	1.8E-09	3.4E-09
Benz[a]anthracene	56-55-3	31	0	3.2E-10	3.7E-10	1.7E-10	8.9E-10	1.7E-09
Benzo[a]pyrene	50-32-8	32	0	2.4E-10	2.4E-10	1.6E-10	6.2E-10	9.8E-10
Benzo[b]fluoranthene	205-99-2	34	0	4.5E-10	4.0E-10	3.5E-10	1.1E-09	1.7E-09
Benzo[k]fluoranthene	207-08-9	29	0	2.0E-10	2.1E-10	1.2E-10	6.4E-10	7.9E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	1.7E-09	1.0E-08	0	0	6.0E-08
Chrysene	218-01-9	35	2.2E-10	2.0E-09	1.3E-09	2.0E-09	4.1E-09	4.7E-09
Cyclopenta[cd]pyrene	27208-37-3	35	3.7E-11	3.7E-10	2.2E-10	3.5E-10	7.0E-10	8.4E-10
Dibenz[a,h]anthracene	53-70-3	14	0	1.2E-10	2.1E-10	0	5.7E-10	8.9E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	2.3E-10	2.0E-10	2.1E-10	5.5E-10	8.6E-10
Naphthalene	91-20-3	2	0	1.5E-11	6.2E-11	0	7.6E-11	2.7E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-294. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 16<30 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	1.0E-10	3.1E-10	0	7.8E-10	1.5E-09
Benz[a]anthracene	56-55-3	31	0	1.4E-10	1.6E-10	7.3E-11	3.9E-10	7.4E-10
Benzo[a]pyrene	50-32-8	32	0	1.0E-10	1.0E-10	6.9E-11	2.7E-10	4.3E-10
Benzo[b]fluoranthene	205-99-2	34	0	2.0E-10	1.7E-10	1.6E-10	4.9E-10	7.6E-10
Benzo[k]fluoranthene	207-08-9	29	0	8.6E-11	9.1E-11	5.2E-11	2.8E-10	3.5E-10



Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
1,3-Benzothiazole-2-thiol	149-30-4	1	0	7.4E-10	4.4E-09	0	0	2.6E-08
Chrysene	218-01-9	35	9.8E-11	9.0E-10	5.8E-10	9.0E-10	1.8E-09	2.0E-09
Cyclopenta[cd]pyrene	27208-37-3	35	1.6E-11	1.6E-10	9.6E-11	1.5E-10	3.1E-10	3.7E-10
Dibenz[a,h]anthracene	53-70-3	14	0	5.1E-11	9.3E-11	0	2.5E-10	3.9E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	1.0E-10	8.6E-11	9.1E-11	2.4E-10	3.7E-10
Naphthalene	91-20-3	2	0	6.6E-12	2.7E-11	0	3.3E-11	1.2E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-295. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 30<40 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	8.7E-11	2.7E-10	0	6.7E-10	1.3E-09
Benz[a]anthracene	56-55-3	31	0	1.2E-10	1.4E-10	6.3E-11	3.4E-10	6.4E-10
Benzo[a]pyrene	50-32-8	32	0	9.0E-11	8.9E-11	5.9E-11	2.4E-10	3.7E-10
Benzo[b]fluoranthene	205-99-2	34	0	1.7E-10	1.5E-10	1.3E-10	4.3E-10	6.6E-10
Benzo[k]fluoranthene	207-08-9	29	0	7.4E-11	7.9E-11	4.5E-11	2.4E-10	3.0E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	6.4E-10	3.8E-09	0	0	2.3E-08
Chrysene	218-01-9	35	8.4E-11	7.7E-10	5.0E-10	7.7E-10	1.6E-09	1.8E-09
Cyclopenta[cd]pyrene	27208-37-3	35	1.4E-11	1.4E-10	8.3E-11	1.3E-10	2.6E-10	3.2E-10
Dibenz[a,h]anthracene	53-70-3	14	0	4.4E-11	8.1E-11	0	2.2E-10	3.4E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	8.9E-11	7.4E-11	7.8E-11	2.1E-10	3.2E-10
Naphthalene	91-20-3	2	0	5.7E-12	2.4E-11	0	2.9E-11	1.0E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-296. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 40<50 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	8.8E-11	2.7E-10	0	6.8E-10	1.3E-09
Benz[a]anthracene	56-55-3	31	0	1.2E-10	1.4E-10	6.4E-11	3.4E-10	6.5E-10
Benzo[a]pyrene	50-32-8	32	0	9.1E-11	9.0E-11	6.0E-11	2.4E-10	3.7E-10
Benzo[b]fluoranthene	205-99-2	34	0	1.7E-10	1.5E-10	1.4E-10	4.3E-10	6.7E-10
Benzo[k]fluoranthene	207-08-9	29	0	7.5E-11	7.9E-11	4.6E-11	2.4E-10	3.0E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	6.5E-10	3.8E-09	0	0	2.3E-08
Chrysene	218-01-9	35	8.5E-11	7.8E-10	5.0E-10	7.8E-10	1.6E-09	1.8E-09
Cyclopenta[cd]pyrene	27208-37-3	35	1.4E-11	1.4E-10	8.4E-11	1.3E-10	2.7E-10	3.2E-10
Dibenz[a,h]anthracene	53-70-3	14	0	4.5E-11	8.1E-11	0	2.2E-10	3.4E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	9.0E-11	7.5E-11	7.9E-11	2.1E-10	3.3E-10
Naphthalene	91-20-3	2	0	5.8E-12	2.4E-11	0	2.9E-11	1.0E-10

^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-297. Field-Specific Average Dermal Dose^a (ADD_{der}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment (ADD_{der}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 50<70 Years**

Chemical Name	CASRN	Detection	ADD _{der-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	4	0	8.3E-11	2.6E-10	0	6.4E-10	1.2E-09
Benz[a]anthracene	56-55-3	31	0	1.1E-10	1.3E-10	6.0E-11	3.2E-10	6.1E-10
Benzo[a]pyrene	50-32-8	32	0	8.6E-11	8.5E-11	5.6E-11	2.2E-10	3.5E-10
Benzo[b]fluoranthene	205-99-2	34	0	1.6E-10	1.4E-10	1.3E-10	4.0E-10	6.3E-10
Benzo[k]fluoranthene	207-08-9	29	0	7.1E-11	7.5E-11	4.3E-11	2.3E-10	2.9E-10
1,3-Benzothiazole-2-thiol	149-30-4	1	0	6.1E-10	3.6E-09	0	0	2.1E-08
Chrysene	218-01-9	35	8.0E-11	7.4E-10	4.7E-10	7.4E-10	1.5E-09	1.7E-09
Cyclopenta[cd]pyrene	27208-37-3	35	1.3E-11	1.3E-10	7.9E-11	1.3E-10	2.5E-10	3.0E-10
Dibenz[a,h]anthracene	53-70-3	14	0	4.2E-11	7.7E-11	0	2.1E-10	3.2E-10
Indeno[1,2,3-cd]pyrene	193-39-5	28	0	8.5E-11	7.1E-11	7.5E-11	2.0E-10	3.1E-10
Naphthalene	91-20-3	2	0	5.4E-12	2.2E-11	0	2.7E-11	9.9E-11



^a 35 field-specific ADD_{der-field} are included in the table. ADD_{der-field} was calculated from field-specific average concentrations (C_{der-crumb rubber-field}) of a chemical detected in artificial sweat extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

F.5.5. Average Ingestion One-Day Dose (AD_{ing-DART}) for One-Day Non-Cancer Hazard Assessment of DART Chemicals

INDIVIDUAL FIELD ASSESSMENT (Table F-298 to Table F-321)

Table F-298. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Athletes 2<6 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	1.8E-08	1.5E-07	1.1E-07	1.4E-07	2.8E-07	5.9E-07
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	8.2E-08	3.3E-07	0	3.1E-07	1.9E-06
Benzo[a]pyrene	50-32-8	35	7.0E-09	3.5E-08	2.9E-08	2.5E-08	9.8E-08	1.0E-07
Benzo[e]pyrene	192-97-2	35	3.2E-08	1.1E-07	6.3E-08	8.2E-08	2.2E-07	2.4E-07
Benzo[g,h,i]perylene	191-24-2	35	4.9E-09	7.2E-08	4.5E-08	6.4E-08	1.4E-07	2.0E-07
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	2.3E-07	2.1E-07	1.6E-07	6.9E-07	8.4E-07
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	1.8E-07	5.3E-07	0	1.7E-06	2.0E-06
Boron		25	0	6.7E-06	7.5E-06	3.0E-06	2.0E-05	2.6E-05
n-Caproic acid vinyl ester	3050-69-9	1	0	2.0E-09	1.2E-08	0	0	7.0E-08
Chrysene	218-01-9	35	4.8E-08	1.9E-07	1.0E-07	1.9E-07	3.3E-07	5.3E-07
Coronene	191-07-1	7	0	2.6E-08	7.6E-08	0	1.1E-07	4.1E-07
Cyclopenta[cd]pyrene	27208-37-3	35	7.3E-09	4.1E-08	3.8E-08	3.6E-08	1.1E-07	1.9E-07
Dicyclohexylamine	101-83-7	25	0	6.1E-07	5.0E-07	5.8E-07	1.4E-06	1.7E-06
Dimethyl phthalate	131-11-3	9	0	1.4E-08	4.7E-08	0	1.2E-07	2.2E-07
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	7.2E-09	1.3E-08	0	3.1E-08	4.9E-08
Methyl stearate	112-61-8	35	3.7E-08	1.5E-07	1.3E-07	1.2E-07	3.5E-07	7.1E-07
Nickel		35	3.1E-07	1.4E-06	1.2E-06	1.0E-06	4.0E-06	5.5E-06
4-tert-Octylphenol	140-66-9	35	7.8E-08	1.0E-06	1.4E-06	4.5E-07	4.0E-06	6.4E-06
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	4.7E-08	3.8E-07	2.3E-07	3.4E-07	7.8E-07	1.0E-06
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	2.2E-07	2.8E-07	1.2E-07	7.4E-07	1.4E-06

^a 35 field-specific AD_{ing-DART-field} are included in the table

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not



included as the Study did not speciate the chemicals in the sample analysis
Values are rounded to two significant figures.

Table F-299. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Athletes 6<11 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	1.1E-08	9.7E-08	6.8E-08	8.7E-08	1.7E-07	3.7E-07
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	5.2E-08	2.1E-07	0	1.9E-07	1.2E-06
Benzo[a]pyrene	50-32-8	35	4.4E-09	2.2E-08	1.8E-08	1.6E-08	6.2E-08	6.6E-08
Benzo[e]pyrene	192-97-2	35	2.0E-08	6.7E-08	4.0E-08	5.2E-08	1.4E-07	1.5E-07
Benzo[g,h,i]perylene	191-24-2	35	3.1E-09	4.6E-08	2.9E-08	4.1E-08	9.1E-08	1.3E-07
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	1.5E-07	1.3E-07	1.0E-07	4.4E-07	5.4E-07
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	1.2E-07	3.4E-07	0	1.1E-06	1.2E-06
Boron		25	0	4.3E-06	4.8E-06	1.9E-06	1.3E-05	1.7E-05
n-Caproic acid vinyl ester	3050-69-9	1	0	1.3E-09	7.5E-09	0	0	4.5E-08
Chrysene	218-01-9	35	3.1E-08	1.2E-07	6.5E-08	1.2E-07	2.1E-07	3.3E-07
Coronene	191-07-1	7	0	1.7E-08	4.8E-08	0	7.1E-08	2.6E-07
Cyclopenta[cd]pyrene	27208-37-3	35	4.6E-09	2.6E-08	2.4E-08	2.3E-08	6.9E-08	1.2E-07
Dicyclohexylamine	101-83-7	25	0	3.9E-07	3.2E-07	3.7E-07	8.9E-07	1.1E-06
Dimethyl phthalate	131-11-3	9	0	8.8E-09	3.0E-08	0	7.9E-08	1.4E-07
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	4.6E-09	8.0E-09	0	1.9E-08	3.1E-08
Methyl stearate	112-61-8	35	2.4E-08	9.4E-08	8.3E-08	7.3E-08	2.2E-07	4.5E-07
Nickel		35	2.0E-07	9.1E-07	7.5E-07	6.3E-07	2.6E-06	3.5E-06
4-tert-Octylphenol	140-66-9	35	5.0E-08	6.5E-07	8.9E-07	2.9E-07	2.5E-06	4.1E-06
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	3.0E-08	2.4E-07	1.5E-07	2.2E-07	4.9E-07	6.6E-07
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	1.4E-07	1.8E-07	7.7E-08	4.7E-07	8.9E-07

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis
Values are rounded to two significant figures.



Table F-300. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Athletes 11<16 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	7.2E-09	6.1E-08	4.3E-08	5.5E-08	1.1E-07	2.4E-07
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	3.3E-08	1.3E-07	0	1.2E-07	7.5E-07
Benzo[a]pyrene	50-32-8	35	2.8E-09	1.4E-08	1.2E-08	1.0E-08	3.9E-08	4.2E-08
Benzo[e]pyrene	192-97-2	35	1.3E-08	4.3E-08	2.5E-08	3.3E-08	8.8E-08	9.6E-08
Benzo[g,h,i]perylene	191-24-2	35	2.0E-09	2.9E-08	1.8E-08	2.6E-08	5.8E-08	8.0E-08
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	9.3E-08	8.5E-08	6.5E-08	2.8E-07	3.4E-07
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	7.4E-08	2.1E-07	0	6.8E-07	7.9E-07
Boron		25	0	2.7E-06	3.0E-06	1.2E-06	7.9E-06	1.1E-05
n-Caproic acid vinyl ester	3050-69-9	1	0	8.1E-10	4.8E-09	0	0	2.8E-08
Chrysene	218-01-9	35	1.9E-08	7.7E-08	4.1E-08	7.7E-08	1.3E-07	2.1E-07
Coronene	191-07-1	7	0	1.1E-08	3.1E-08	0	4.5E-08	1.7E-07
Cyclopenta[cd]pyrene	27208-37-3	35	2.9E-09	1.7E-08	1.5E-08	1.4E-08	4.4E-08	7.8E-08
Dicyclohexylamine	101-83-7	25	0	2.5E-07	2.0E-07	2.3E-07	5.6E-07	6.8E-07
Dimethyl phthalate	131-11-3	9	0	5.6E-09	1.9E-08	0	5.0E-08	8.8E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	2.9E-09	5.1E-09	0	1.2E-08	2.0E-08
Methyl stearate	112-61-8	35	1.5E-08	6.0E-08	5.3E-08	4.6E-08	1.4E-07	2.8E-07
Nickel		35	1.2E-07	5.8E-07	4.7E-07	4.0E-07	1.6E-06	2.2E-06
4-tert-Octylphenol	140-66-9	35	3.2E-08	4.1E-07	5.6E-07	1.8E-07	1.6E-06	2.6E-06
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.9E-08	1.5E-07	9.3E-08	1.4E-07	3.1E-07	4.2E-07
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	8.7E-08	1.1E-07	4.9E-08	3.0E-07	5.6E-07

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis

Values are rounded to two significant figures.

Table F-301. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Athletes 16<30 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	5.8E-09	4.9E-08	3.5E-08	4.4E-08	8.8E-08	1.9E-07
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	2.6E-08	1.1E-07	0	9.9E-08	6.0E-07



Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	35	2.2E-09	1.1E-08	9.3E-09	8.1E-09	3.2E-08	3.3E-08
Benzo[e]pyrene	192-97-2	35	1.0E-08	3.4E-08	2.0E-08	2.7E-08	7.0E-08	7.6E-08
Benzo[g,h,i]perylene	191-24-2	35	1.6E-09	2.3E-08	1.4E-08	2.1E-08	4.6E-08	6.4E-08
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	7.4E-08	6.8E-08	5.2E-08	2.2E-07	2.7E-07
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	5.9E-08	1.7E-07	0	5.4E-07	6.3E-07
Boron		25	0	2.2E-06	2.4E-06	9.7E-07	6.3E-06	8.5E-06
n-Caproic acid vinyl ester	3050-69-9	1	0	6.5E-10	3.8E-09	0	0	2.3E-08
Chrysene	218-01-9	35	1.6E-08	6.2E-08	3.3E-08	6.2E-08	1.1E-07	1.7E-07
Coronene	191-07-1	7	0	8.5E-09	2.5E-08	0	3.6E-08	1.3E-07
Cyclopenta[cd]pyrene	27208-37-3	35	2.3E-09	1.3E-08	1.2E-08	1.1E-08	3.5E-08	6.2E-08
Dicyclohexylamine	101-83-7	25	0	2.0E-07	1.6E-07	1.9E-07	4.5E-07	5.4E-07
Dimethyl phthalate	131-11-3	9	0	4.4E-09	1.5E-08	0	4.0E-08	7.1E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	2.3E-09	4.1E-09	0	9.8E-09	1.6E-08
Methyl stearate	112-61-8	35	1.2E-08	4.8E-08	4.2E-08	3.7E-08	1.1E-07	2.3E-07
Nickel		35	9.9E-08	4.6E-07	3.8E-07	3.2E-07	1.3E-06	1.8E-06
4-tert-Octylphenol	140-66-9	35	2.5E-08	3.3E-07	4.5E-07	1.5E-07	1.3E-06	2.1E-06
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.5E-08	1.2E-07	7.4E-08	1.1E-07	2.5E-07	3.3E-07
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	6.9E-08	9.0E-08	3.9E-08	2.4E-07	4.5E-07

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis

Values are rounded to two significant figures.

Table F-302. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Athletes 30<40 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	4.9E-09	4.2E-08	3.0E-08	3.8E-08	7.6E-08	1.6E-07
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	2.3E-08	9.2E-08	0	8.5E-08	5.1E-07
Benzo[a]pyrene	50-32-8	35	1.9E-09	9.8E-09	8.0E-09	7.0E-09	2.7E-08	2.9E-08
Benzo[e]pyrene	192-97-2	35	8.8E-09	2.9E-08	1.7E-08	2.3E-08	6.0E-08	6.5E-08
Benzo[g,h,i]perylene	191-24-2	35	1.4E-09	2.0E-08	1.2E-08	1.8E-08	4.0E-08	5.5E-08
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	6.4E-08	5.8E-08	4.4E-08	1.9E-07	2.3E-07
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	5.1E-08	1.5E-07	0	4.7E-07	5.4E-07



Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Boron		25	0	1.9E-06	2.1E-06	8.3E-07	5.4E-06	7.3E-06
n-Caproic acid vinyl ester	3050-69-9	1	0	5.5E-10	3.3E-09	0	0	1.9E-08
Chrysene	218-01-9	35	1.3E-08	5.3E-08	2.8E-08	5.3E-08	9.1E-08	1.5E-07
Coronene	191-07-1	7	0	7.3E-09	2.1E-08	0	3.1E-08	1.1E-07
Cyclopenta[cd]pyrene	27208-37-3	35	2.0E-09	1.1E-08	1.1E-08	9.8E-09	3.0E-08	5.3E-08
Dicyclohexylamine	101-83-7	25	0	1.7E-07	1.4E-07	1.6E-07	3.9E-07	4.6E-07
Dimethyl phthalate	131-11-3	9	0	3.8E-09	1.3E-08	0	3.4E-08	6.0E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	2.0E-09	3.5E-09	0	8.4E-09	1.4E-08
Methyl stearate	112-61-8	35	1.0E-08	4.1E-08	3.6E-08	3.2E-08	9.7E-08	1.9E-07
Nickel		35	8.5E-08	4.0E-07	3.2E-07	2.8E-07	1.1E-06	1.5E-06
4-tert-Octylphenol	140-66-9	35	2.2E-08	2.8E-07	3.9E-07	1.2E-07	1.1E-06	1.8E-06
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.3E-08	1.1E-07	6.4E-08	9.4E-08	2.1E-07	2.9E-07
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	5.9E-08	7.8E-08	3.3E-08	2.0E-07	3.9E-07

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis
Values are rounded to two significant figures.

Table F-303. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Athletes 40<50 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	4.9E-09	4.1E-08	2.9E-08	3.7E-08	7.5E-08	1.6E-07
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	2.2E-08	9.0E-08	0	8.3E-08	5.0E-07
Benzo[a]pyrene	50-32-8	35	1.9E-09	9.6E-09	7.9E-09	6.8E-09	2.7E-08	2.8E-08
Benzo[e]pyrene	192-97-2	35	8.6E-09	2.9E-08	1.7E-08	2.2E-08	5.9E-08	6.4E-08
Benzo[g,h,i]perylene	191-24-2	35	1.3E-09	2.0E-08	1.2E-08	1.7E-08	3.9E-08	5.4E-08
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	6.3E-08	5.7E-08	4.4E-08	1.9E-07	2.3E-07
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	5.0E-08	1.4E-07	0	4.6E-07	5.3E-07
Boron		25	0	1.8E-06	2.0E-06	8.2E-07	5.3E-06	7.1E-06
n-Caproic acid vinyl ester	3050-69-9	1	0	5.4E-10	3.2E-09	0	0	1.9E-08
Chrysene	218-01-9	35	1.3E-08	5.2E-08	2.8E-08	5.2E-08	8.9E-08	1.4E-07
Coronene	191-07-1	7	0	7.1E-09	2.1E-08	0	3.0E-08	1.1E-07
Cyclopenta[cd]pyrene	27208-37-3	35	2.0E-09	1.1E-08	1.0E-08	9.6E-09	2.9E-08	5.2E-08
Dicyclohexylamine	101-83-7	25	0	1.6E-07	1.4E-07	1.6E-07	3.8E-07	4.6E-07



Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dimethyl phthalate	131-11-3	9	0	3.7E-09	1.3E-08	0	3.4E-08	5.9E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	2.0E-09	3.4E-09	0	8.3E-09	1.3E-08
Methyl stearate	112-61-8	35	1.0E-08	4.0E-08	3.5E-08	3.1E-08	9.6E-08	1.9E-07
Nickel		35	8.3E-08	3.9E-07	3.2E-07	2.7E-07	1.1E-06	1.5E-06
4-tert-Octylphenol	140-66-9	35	2.1E-08	2.8E-07	3.8E-07	1.2E-07	1.1E-06	1.7E-06
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.3E-08	1.0E-07	6.3E-08	9.2E-08	2.1E-07	2.8E-07
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	5.8E-08	7.6E-08	3.3E-08	2.0E-07	3.8E-07

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis
Values are rounded to two significant figures.

Table F-304. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Athletes 50<70 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	4.9E-09	4.1E-08	2.9E-08	3.7E-08	7.5E-08	1.6E-07
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	2.2E-08	9.0E-08	0	8.3E-08	5.0E-07
Benzo[a]pyrene	50-32-8	35	1.9E-09	9.6E-09	7.9E-09	6.8E-09	2.7E-08	2.8E-08
Benzo[e]pyrene	192-97-2	35	8.6E-09	2.9E-08	1.7E-08	2.2E-08	5.9E-08	6.4E-08
Benzo[g,h,i]perylene	191-24-2	35	1.3E-09	2.0E-08	1.2E-08	1.7E-08	3.9E-08	5.4E-08
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	6.3E-08	5.7E-08	4.4E-08	1.9E-07	2.3E-07
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	5.0E-08	1.4E-07	0	4.6E-07	5.3E-07
Boron		25	0	1.8E-06	2.0E-06	8.2E-07	5.4E-06	7.1E-06
n-Caproic acid vinyl ester	3050-69-9	1	0	5.5E-10	3.2E-09	0	0	1.9E-08
Chrysene	218-01-9	35	1.3E-08	5.2E-08	2.8E-08	5.2E-08	8.9E-08	1.4E-07
Coronene	191-07-1	7	0	7.1E-09	2.1E-08	0	3.0E-08	1.1E-07
Cyclopenta[cd]pyrene	27208-37-3	35	2.0E-09	1.1E-08	1.0E-08	9.7E-09	3.0E-08	5.3E-08
Dicyclohexylamine	101-83-7	25	0	1.7E-07	1.4E-07	1.6E-07	3.8E-07	4.6E-07
Dimethyl phthalate	131-11-3	9	0	3.8E-09	1.3E-08	0	3.4E-08	6.0E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	2.0E-09	3.4E-09	0	8.3E-09	1.3E-08
Methyl stearate	112-61-8	35	1.0E-08	4.0E-08	3.5E-08	3.1E-08	9.6E-08	1.9E-07
Nickel		35	8.3E-08	3.9E-07	3.2E-07	2.7E-07	1.1E-06	1.5E-06
4-tert-Octylphenol	140-66-9	35	2.1E-08	2.8E-07	3.8E-07	1.2E-07	1.1E-06	1.7E-06



Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.3E-08	1.0E-07	6.3E-08	9.3E-08	2.1E-07	2.8E-07
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	5.8E-08	7.6E-08	3.3E-08	2.0E-07	3.8E-07

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis
Values are rounded to two significant figures.

Table F-305. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Coaches 16<30 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	1.7E-09	1.4E-08	1.0E-08	1.3E-08	2.6E-08	5.5E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	7.7E-09	3.1E-08	0	2.9E-08	1.7E-07
Benzo[a]pyrene	50-32-8	35	6.5E-10	3.3E-09	2.7E-09	2.4E-09	9.2E-09	9.7E-09
Benzo[e]pyrene	192-97-2	35	3.0E-09	9.9E-09	5.9E-09	7.7E-09	2.0E-08	2.2E-08
Benzo[g,h,i]perylene	191-24-2	35	4.6E-10	6.7E-09	4.2E-09	6.0E-09	1.3E-08	1.9E-08
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	2.2E-08	2.0E-08	1.5E-08	6.5E-08	7.9E-08
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	1.7E-08	5.0E-08	0	1.6E-07	1.8E-07
Boron		25	0	6.3E-07	7.1E-07	2.8E-07	1.8E-06	2.5E-06
n-Caproic acid vinyl ester	3050-69-9	1	0	1.9E-10	1.1E-09	0	0	6.6E-09
Chrysene	218-01-9	35	4.5E-09	1.8E-08	9.6E-09	1.8E-08	3.1E-08	4.9E-08
Coronene	191-07-1	7	0	2.5E-09	7.1E-09	0	1.0E-08	3.9E-08
Cyclopenta[cd]pyrene	27208-37-3	35	6.8E-10	3.9E-09	3.6E-09	3.3E-09	1.0E-08	1.8E-08
Dicyclohexylamine	101-83-7	25	0	5.7E-08	4.7E-08	5.5E-08	1.3E-07	1.6E-07
Dimethyl phthalate	131-11-3	9	0	1.3E-09	4.4E-09	0	1.2E-08	2.1E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	6.8E-10	1.2E-09	0	2.9E-09	4.6E-09
Methyl stearate	112-61-8	35	3.5E-09	1.4E-08	1.2E-08	1.1E-08	3.3E-08	6.6E-08
Nickel		35	2.9E-08	1.3E-07	1.1E-07	9.4E-08	3.8E-07	5.2E-07
4-tert-Octylphenol	140-66-9	35	7.3E-09	9.6E-08	1.3E-07	4.2E-08	3.8E-07	6.0E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	4.4E-09	3.6E-08	2.2E-08	3.2E-08	7.3E-08	9.7E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	2.0E-08	2.6E-08	1.1E-08	6.9E-08	1.3E-07

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.



° CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis
Values are rounded to two significant figures.

Table F-306. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Coaches 30<40 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	1.6E-09	1.3E-08	9.5E-09	1.2E-08	2.4E-08	5.2E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	7.2E-09	2.9E-08	0	2.7E-08	1.6E-07
Benzo[a]pyrene	50-32-8	35	6.1E-10	3.1E-09	2.6E-09	2.2E-09	8.6E-09	9.1E-09
Benzo[e]pyrene	192-97-2	35	2.8E-09	9.3E-09	5.5E-09	7.2E-09	1.9E-08	2.1E-08
Benzo[g,h,i]perylene	191-24-2	35	4.3E-10	6.3E-09	4.0E-09	5.7E-09	1.3E-08	1.8E-08
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	2.0E-08	1.9E-08	1.4E-08	6.1E-08	7.4E-08
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	1.6E-08	4.7E-08	0	1.5E-07	1.7E-07
Boron		25	0	5.9E-07	6.6E-07	2.7E-07	1.7E-06	2.3E-06
n-Caproic acid vinyl ester	3050-69-9	1	0	1.8E-10	1.0E-09	0	0	6.2E-09
Chrysene	218-01-9	35	4.3E-09	1.7E-08	9.0E-09	1.7E-08	2.9E-08	4.6E-08
Coronene	191-07-1	7	0	2.3E-09	6.7E-09	0	9.8E-09	3.6E-08
Cyclopenta[cd]pyrene	27208-37-3	35	6.4E-10	3.6E-09	3.4E-09	3.1E-09	9.6E-09	1.7E-08
Dicyclohexylamine	101-83-7	25	0	5.3E-08	4.4E-08	5.1E-08	1.2E-07	1.5E-07
Dimethyl phthalate	131-11-3	9	0	1.2E-09	4.1E-09	0	1.1E-08	1.9E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	6.3E-10	1.1E-09	0	2.7E-09	4.3E-09
Methyl stearate	112-61-8	35	3.3E-09	1.3E-08	1.1E-08	1.0E-08	3.1E-08	6.2E-08
Nickel		35	2.7E-08	1.3E-07	1.0E-07	8.8E-08	3.6E-07	4.9E-07
4-tert-Octylphenol	140-66-9	35	6.9E-09	9.0E-08	1.2E-07	4.0E-08	3.5E-07	5.7E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	4.1E-09	3.4E-08	2.0E-08	3.0E-08	6.8E-08	9.1E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	1.9E-08	2.5E-08	1.1E-08	6.5E-08	1.2E-07

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis
Values are rounded to two significant figures.



Table F-307. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Coaches 40<50 Years**

Chemical	CASRN	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	1.5E-09	1.3E-08	9.3E-09	1.2E-08	2.4E-08	5.1E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	7.1E-09	2.9E-08	0	2.6E-08	1.6E-07
Benzo[a]pyrene	50-32-8	35	6.0E-10	3.0E-09	2.5E-09	2.2E-09	8.4E-09	9.0E-09
Benzo[e]pyrene	192-97-2	35	2.7E-09	9.1E-09	5.4E-09	7.1E-09	1.9E-08	2.0E-08
Benzo[g,h,i]perylene	191-24-2	35	4.2E-10	6.2E-09	3.9E-09	5.5E-09	1.2E-08	1.7E-08
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	2.0E-08	1.8E-08	1.4E-08	5.9E-08	7.3E-08
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	1.6E-08	4.6E-08	0	1.5E-07	1.7E-07
Boron		25	0	5.8E-07	6.5E-07	2.6E-07	1.7E-06	2.3E-06
n-Caproic acid vinyl ester	3050-69-9	1	0	1.7E-10	1.0E-09	0	0	6.1E-09
Chrysene	218-01-9	35	4.2E-09	1.7E-08	8.8E-09	1.7E-08	2.8E-08	4.5E-08
Coronene	191-07-1	7	0	2.3E-09	6.6E-09	0	9.6E-09	3.5E-08
Cyclopenta[cd]pyrene	27208-37-3	35	6.2E-10	3.5E-09	3.3E-09	3.1E-09	9.4E-09	1.7E-08
Dicyclohexylamine	101-83-7	25	0	5.2E-08	4.3E-08	5.0E-08	1.2E-07	1.4E-07
Dimethyl phthalate	131-11-3	9	0	1.2E-09	4.0E-09	0	1.1E-08	1.9E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	6.2E-10	1.1E-09	0	2.6E-09	4.2E-09
Methyl stearate	112-61-8	35	3.2E-09	1.3E-08	1.1E-08	9.9E-09	3.0E-08	6.1E-08
Nickel		35	2.6E-08	1.2E-07	1.0E-07	8.6E-08	3.5E-07	4.8E-07
4-tert-Octylphenol	140-66-9	35	6.7E-09	8.8E-08	1.2E-07	3.9E-08	3.5E-07	5.5E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	4.0E-09	3.3E-08	2.0E-08	2.9E-08	6.7E-08	8.9E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	1.9E-08	2.4E-08	1.0E-08	6.3E-08	1.2E-07

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis

Values are rounded to two significant figures.

Table F-308. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Coaches 50<70 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	1.5E-09	1.3E-08	9.3E-09	1.2E-08	2.4E-08	5.1E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	7.1E-09	2.9E-08	0	2.6E-08	1.6E-07



Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	35	6.0E-10	3.1E-09	2.5E-09	2.2E-09	8.5E-09	9.0E-09
Benzo[e]pyrene	192-97-2	35	2.7E-09	9.1E-09	5.4E-09	7.1E-09	1.9E-08	2.0E-08
Benzo[g,h,i]perylene	191-24-2	35	4.3E-10	6.2E-09	3.9E-09	5.6E-09	1.2E-08	1.7E-08
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	2.0E-08	1.8E-08	1.4E-08	6.0E-08	7.3E-08
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	1.6E-08	4.6E-08	0	1.5E-07	1.7E-07
Boron		25	0	5.8E-07	6.5E-07	2.6E-07	1.7E-06	2.3E-06
n-Caproic acid vinyl ester	3050-69-9	1	0	1.7E-10	1.0E-09	0	0	6.1E-09
Chrysene	218-01-9	35	4.2E-09	1.7E-08	8.8E-09	1.7E-08	2.8E-08	4.6E-08
Coronene	191-07-1	7	0	2.3E-09	6.6E-09	0	9.6E-09	3.6E-08
Cyclopenta[cd]pyrene	27208-37-3	35	6.3E-10	3.6E-09	3.3E-09	3.1E-09	9.4E-09	1.7E-08
Dicyclohexylamine	101-83-7	25	0	5.2E-08	4.3E-08	5.0E-08	1.2E-07	1.5E-07
Dimethyl phthalate	131-11-3	9	0	1.2E-09	4.0E-09	0	1.1E-08	1.9E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	6.2E-10	1.1E-09	0	2.6E-09	4.3E-09
Methyl stearate	112-61-8	35	3.2E-09	1.3E-08	1.1E-08	9.9E-09	3.0E-08	6.1E-08
Nickel		35	2.7E-08	1.2E-07	1.0E-07	8.6E-08	3.5E-07	4.8E-07
4-tert-Octylphenol	140-66-9	35	6.8E-09	8.8E-08	1.2E-07	3.9E-08	3.5E-07	5.6E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	4.0E-09	3.3E-08	2.0E-08	2.9E-08	6.7E-08	8.9E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	1.9E-08	2.4E-08	1.0E-08	6.4E-08	1.2E-07

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis

Values are rounded to two significant figures.

Table F-309. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Referees 16<30 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	1.7E-09	1.4E-08	1.0E-08	1.3E-08	2.6E-08	5.5E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	7.7E-09	3.1E-08	0	2.9E-08	1.7E-07
Benzo[a]pyrene	50-32-8	35	6.5E-10	3.3E-09	2.7E-09	2.4E-09	9.2E-09	9.7E-09
Benzo[e]pyrene	192-97-2	35	3.0E-09	9.9E-09	5.9E-09	7.7E-09	2.0E-08	2.2E-08
Benzo[g,h,i]perylene	191-24-2	35	4.6E-10	6.7E-09	4.2E-09	6.0E-09	1.3E-08	1.9E-08
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	2.2E-08	2.0E-08	1.5E-08	6.5E-08	7.9E-08
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	1.7E-08	5.0E-08	0	1.6E-07	1.8E-07



Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Boron		25	0	6.3E-07	7.1E-07	2.8E-07	1.8E-06	2.5E-06
n-Caproic acid vinyl ester	3050-69-9	1	0	1.9E-10	1.1E-09	0	0	6.6E-09
Chrysene	218-01-9	35	4.5E-09	1.8E-08	9.6E-09	1.8E-08	3.1E-08	4.9E-08
Coronene	191-07-1	7	0	2.5E-09	7.1E-09	0	1.0E-08	3.9E-08
Cyclopenta[cd]pyrene	27208-37-3	35	6.8E-10	3.9E-09	3.6E-09	3.3E-09	1.0E-08	1.8E-08
Dicyclohexylamine	101-83-7	25	0	5.7E-08	4.7E-08	5.5E-08	1.3E-07	1.6E-07
Dimethyl phthalate	131-11-3	9	0	1.3E-09	4.4E-09	0	1.2E-08	2.1E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	6.8E-10	1.2E-09	0	2.9E-09	4.6E-09
Methyl stearate	112-61-8	35	3.5E-09	1.4E-08	1.2E-08	1.1E-08	3.3E-08	6.6E-08
Nickel		35	2.9E-08	1.3E-07	1.1E-07	9.4E-08	3.8E-07	5.2E-07
4-tert-Octylphenol	140-66-9	35	7.3E-09	9.6E-08	1.3E-07	4.2E-08	3.8E-07	6.0E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	4.4E-09	3.6E-08	2.2E-08	3.2E-08	7.3E-08	9.7E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	2.0E-08	2.6E-08	1.1E-08	6.9E-08	1.3E-07

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis
Values are rounded to two significant figures.

Table F-310. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Referees 30<40 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	1.6E-09	1.3E-08	9.5E-09	1.2E-08	2.4E-08	5.2E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	7.2E-09	2.9E-08	0	2.7E-08	1.6E-07
Benzo[a]pyrene	50-32-8	35	6.1E-10	3.1E-09	2.6E-09	2.2E-09	8.6E-09	9.1E-09
Benzo[e]pyrene	192-97-2	35	2.8E-09	9.3E-09	5.5E-09	7.2E-09	1.9E-08	2.1E-08
Benzo[g,h,i]perylene	191-24-2	35	4.3E-10	6.3E-09	4.0E-09	5.7E-09	1.3E-08	1.8E-08
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	2.0E-08	1.9E-08	1.4E-08	6.1E-08	7.4E-08
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	1.6E-08	4.7E-08	0	1.5E-07	1.7E-07
Boron		25	0	5.9E-07	6.6E-07	2.7E-07	1.7E-06	2.3E-06
n-Caproic acid vinyl ester	3050-69-9	1	0	1.8E-10	1.0E-09	0	0	6.2E-09
Chrysene	218-01-9	35	4.3E-09	1.7E-08	9.0E-09	1.7E-08	2.9E-08	4.6E-08
Coronene	191-07-1	7	0	2.3E-09	6.7E-09	0	9.8E-09	3.6E-08
Cyclopenta[cd]pyrene	27208-37-3	35	6.4E-10	3.6E-09	3.4E-09	3.1E-09	9.6E-09	1.7E-08
Dicyclohexylamine	101-83-7	25	0	5.3E-08	4.4E-08	5.1E-08	1.2E-07	1.5E-07



Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dimethyl phthalate	131-11-3	9	0	1.2E-09	4.1E-09	0	1.1E-08	1.9E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	6.3E-10	1.1E-09	0	2.7E-09	4.3E-09
Methyl stearate	112-61-8	35	3.3E-09	1.3E-08	1.1E-08	1.0E-08	3.1E-08	6.2E-08
Nickel		35	2.7E-08	1.3E-07	1.0E-07	8.8E-08	3.6E-07	4.9E-07
4-tert-Octylphenol	140-66-9	35	6.9E-09	9.0E-08	1.2E-07	4.0E-08	3.5E-07	5.7E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	4.1E-09	3.4E-08	2.0E-08	3.0E-08	6.8E-08	9.1E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	1.9E-08	2.5E-08	1.1E-08	6.5E-08	1.2E-07

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis

Values are rounded to two significant figures.

Table F-311. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Referees 40<50 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	1.5E-09	1.3E-08	9.3E-09	1.2E-08	2.4E-08	5.1E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	7.1E-09	2.9E-08	0	2.6E-08	1.6E-07
Benzo[a]pyrene	50-32-8	35	6.0E-10	3.0E-09	2.5E-09	2.2E-09	8.4E-09	9.0E-09
Benzo[e]pyrene	192-97-2	35	2.7E-09	9.1E-09	5.4E-09	7.1E-09	1.9E-08	2.0E-08
Benzo[g,h,i]perylene	191-24-2	35	4.2E-10	6.2E-09	3.9E-09	5.5E-09	1.2E-08	1.7E-08
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	2.0E-08	1.8E-08	1.4E-08	5.9E-08	7.3E-08
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	1.6E-08	4.6E-08	0	1.5E-07	1.7E-07
Boron		25	0	5.8E-07	6.5E-07	2.6E-07	1.7E-06	2.3E-06
n-Caproic acid vinyl ester	3050-69-9	1	0	1.7E-10	1.0E-09	0	0	6.1E-09
Chrysene	218-01-9	35	4.2E-09	1.7E-08	8.8E-09	1.7E-08	2.8E-08	4.5E-08
Coronene	191-07-1	7	0	2.3E-09	6.6E-09	0	9.6E-09	3.5E-08
Cyclopenta[cd]pyrene	27208-37-3	35	6.2E-10	3.5E-09	3.3E-09	3.1E-09	9.4E-09	1.7E-08
Dicyclohexylamine	101-83-7	25	0	5.2E-08	4.3E-08	5.0E-08	1.2E-07	1.4E-07
Dimethyl phthalate	131-11-3	9	0	1.2E-09	4.0E-09	0	1.1E-08	1.9E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	6.2E-10	1.1E-09	0	2.6E-09	4.2E-09
Methyl stearate	112-61-8	35	3.2E-09	1.3E-08	1.1E-08	9.9E-09	3.0E-08	6.1E-08
Nickel		35	2.6E-08	1.2E-07	1.0E-07	8.6E-08	3.5E-07	4.8E-07
4-tert-Octylphenol	140-66-9	35	6.7E-09	8.8E-08	1.2E-07	3.9E-08	3.5E-07	5.5E-07



Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	4.0E-09	3.3E-08	2.0E-08	2.9E-08	6.7E-08	8.9E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	1.9E-08	2.4E-08	1.0E-08	6.3E-08	1.2E-07

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis
Values are rounded to two significant figures.

Table F-312. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Referees 50<70 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	1.5E-09	1.3E-08	9.3E-09	1.2E-08	2.4E-08	5.1E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	7.1E-09	2.9E-08	0	2.6E-08	1.6E-07
Benzo[a]pyrene	50-32-8	35	6.0E-10	3.1E-09	2.5E-09	2.2E-09	8.5E-09	9.0E-09
Benzo[e]pyrene	192-97-2	35	2.7E-09	9.1E-09	5.4E-09	7.1E-09	1.9E-08	2.0E-08
Benzo[g,h,i]perylene	191-24-2	35	4.3E-10	6.2E-09	3.9E-09	5.6E-09	1.2E-08	1.7E-08
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	2.0E-08	1.8E-08	1.4E-08	6.0E-08	7.3E-08
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	1.6E-08	4.6E-08	0	1.5E-07	1.7E-07
Boron		25	0	5.8E-07	6.5E-07	2.6E-07	1.7E-06	2.3E-06
n-Caproic acid vinyl ester	3050-69-9	1	0	1.7E-10	1.0E-09	0	0	6.1E-09
Chrysene	218-01-9	35	4.2E-09	1.7E-08	8.8E-09	1.7E-08	2.8E-08	4.6E-08
Coronene	191-07-1	7	0	2.3E-09	6.6E-09	0	9.6E-09	3.6E-08
Cyclopenta[cd]pyrene	27208-37-3	35	6.3E-10	3.6E-09	3.3E-09	3.1E-09	9.4E-09	1.7E-08
Dicyclohexylamine	101-83-7	25	0	5.2E-08	4.3E-08	5.0E-08	1.2E-07	1.5E-07
Dimethyl phthalate	131-11-3	9	0	1.2E-09	4.0E-09	0	1.1E-08	1.9E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	6.2E-10	1.1E-09	0	2.6E-09	4.3E-09
Methyl stearate	112-61-8	35	3.2E-09	1.3E-08	1.1E-08	9.9E-09	3.0E-08	6.1E-08
Nickel		35	2.7E-08	1.2E-07	1.0E-07	8.6E-08	3.5E-07	4.8E-07
4-tert-Octylphenol	140-66-9	35	6.8E-09	8.8E-08	1.2E-07	3.9E-08	3.5E-07	5.6E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	4.0E-09	3.3E-08	2.0E-08	2.9E-08	6.7E-08	8.9E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	1.9E-08	2.4E-08	1.0E-08	6.4E-08	1.2E-07

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.



° CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis
Values are rounded to two significant figures.

Table F-313. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Spectators Third Trimester Fetus<0 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	6.8E-10	5.7E-09	4.1E-09	5.2E-09	1.0E-08	2.2E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	3.1E-09	1.3E-08	0	1.2E-08	7.0E-08
Benzo[a]pyrene	50-32-8	35	2.6E-10	1.3E-09	1.1E-09	9.5E-10	3.7E-09	3.9E-09
Benzo[e]pyrene	192-97-2	35	1.2E-09	4.0E-09	2.4E-09	3.1E-09	8.2E-09	8.9E-09
Benzo[g,h,i]perylene	191-24-2	35	1.9E-10	2.7E-09	1.7E-09	2.4E-09	5.4E-09	7.5E-09
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	8.7E-09	7.9E-09	6.1E-09	2.6E-08	3.2E-08
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	6.9E-09	2.0E-08	0	6.4E-08	7.4E-08
Boron		25	0	2.5E-07	2.8E-07	1.1E-07	7.4E-07	9.9E-07
n-Caproic acid vinyl ester	3050-69-9	1	0	7.6E-11	4.5E-10	0	0	2.6E-09
Chrysene	218-01-9	35	1.8E-09	7.2E-09	3.8E-09	7.2E-09	1.2E-08	2.0E-08
Coronene	191-07-1	7	0	9.9E-10	2.9E-09	0	4.2E-09	1.6E-08
Cyclopenta[cd]pyrene	27208-37-3	35	2.7E-10	1.6E-09	1.4E-09	1.3E-09	4.1E-09	7.3E-09
Dicyclohexylamine	101-83-7	25	0	2.3E-08	1.9E-08	2.2E-08	5.3E-08	6.3E-08
Dimethyl phthalate	131-11-3	9	0	5.2E-10	1.8E-09	0	4.7E-09	8.3E-09
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	2.7E-10	4.7E-10	0	1.2E-09	1.9E-09
Methyl stearate	112-61-8	35	1.4E-09	5.6E-09	4.9E-09	4.3E-09	1.3E-08	2.7E-08
Nickel		35	1.2E-08	5.4E-08	4.4E-08	3.8E-08	1.5E-07	2.1E-07
4-tert-Octylphenol	140-66-9	35	2.9E-09	3.9E-08	5.3E-08	1.7E-08	1.5E-07	2.4E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.8E-09	1.4E-08	8.7E-09	1.3E-08	2.9E-08	3.9E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	8.1E-09	1.1E-08	4.5E-09	2.8E-08	5.3E-08

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis
Values are rounded to two significant figures.



Table F-314. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Spectators 0<2 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	4.5E-08	3.9E-07	2.7E-07	3.5E-07	7.0E-07	1.5E-06
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	2.1E-07	8.4E-07	0	7.8E-07	4.7E-06
Benzo[a]pyrene	50-32-8	35	1.8E-08	9.0E-08	7.4E-08	6.4E-08	2.5E-07	2.6E-07
Benzo[e]pyrene	192-97-2	35	8.1E-08	2.7E-07	1.6E-07	2.1E-07	5.5E-07	6.0E-07
Benzo[g,h,i]perylene	191-24-2	35	1.2E-08	1.8E-07	1.1E-07	1.6E-07	3.6E-07	5.1E-07
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	5.9E-07	5.3E-07	4.1E-07	1.7E-06	2.1E-06
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	4.6E-07	1.4E-06	0	4.3E-06	5.0E-06
Boron		25	0	1.7E-05	1.9E-05	7.7E-06	5.0E-05	6.7E-05
n-Caproic acid vinyl ester	3050-69-9	1	0	5.1E-09	3.0E-08	0	0	1.8E-07
Chrysene	218-01-9	35	1.2E-07	4.9E-07	2.6E-07	4.9E-07	8.3E-07	1.3E-06
Coronene	191-07-1	7	0	6.7E-08	1.9E-07	0	2.8E-07	1.0E-06
Cyclopenta[cd]pyrene	27208-37-3	35	1.8E-08	1.0E-07	9.7E-08	9.0E-08	2.8E-07	4.9E-07
Dicyclohexylamine	101-83-7	25	0	1.5E-06	1.3E-06	1.5E-06	3.5E-06	4.3E-06
Dimethyl phthalate	131-11-3	9	0	3.5E-08	1.2E-07	0	3.1E-07	5.6E-07
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	1.8E-08	3.2E-08	0	7.7E-08	1.3E-07
Methyl stearate	112-61-8	35	9.4E-08	3.8E-07	3.3E-07	2.9E-07	8.9E-07	1.8E-06
Nickel		35	7.8E-07	3.6E-06	3.0E-06	2.5E-06	1.0E-05	1.4E-05
4-tert-Octylphenol	140-66-9	35	2.0E-07	2.6E-06	3.6E-06	1.1E-06	1.0E-05	1.6E-05
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.2E-07	9.7E-07	5.9E-07	8.7E-07	2.0E-06	2.6E-06
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	5.4E-07	7.1E-07	3.1E-07	1.9E-06	3.5E-06

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis

Values are rounded to two significant figures.

Table F-315. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Spectators 2<6 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	2.2E-08	1.8E-07	1.3E-07	1.7E-07	3.3E-07	7.1E-07
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	9.9E-08	4.0E-07	0	3.7E-07	2.2E-06



Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	35	8.4E-09	4.3E-08	3.5E-08	3.0E-08	1.2E-07	1.3E-07
Benzo[e]pyrene	192-97-2	35	3.8E-08	1.3E-07	7.6E-08	9.9E-08	2.6E-07	2.9E-07
Benzo[g,h,i]perylene	191-24-2	35	5.9E-09	8.7E-08	5.4E-08	7.7E-08	1.7E-07	2.4E-07
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	2.8E-07	2.5E-07	1.9E-07	8.3E-07	1.0E-06
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	2.2E-07	6.4E-07	0	2.0E-06	2.3E-06
Boron		25	0	8.1E-06	9.1E-06	3.6E-06	2.4E-05	3.2E-05
n-Caproic acid vinyl ester	3050-69-9	1	0	2.4E-09	1.4E-08	0	0	8.5E-08
Chrysene	218-01-9	35	5.8E-08	2.3E-07	1.2E-07	2.3E-07	4.0E-07	6.3E-07
Coronene	191-07-1	7	0	3.2E-08	9.2E-08	0	1.3E-07	5.0E-07
Cyclopenta[cd]pyrene	27208-37-3	35	8.7E-09	5.0E-08	4.6E-08	4.3E-08	1.3E-07	2.3E-07
Dicyclohexylamine	101-83-7	25	0	7.3E-07	6.0E-07	7.0E-07	1.7E-06	2.0E-06
Dimethyl phthalate	131-11-3	9	0	1.7E-08	5.6E-08	0	1.5E-07	2.6E-07
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	8.7E-09	1.5E-08	0	3.7E-08	5.9E-08
Methyl stearate	112-61-8	35	4.5E-08	1.8E-07	1.6E-07	1.4E-07	4.2E-07	8.5E-07
Nickel		35	3.7E-07	1.7E-06	1.4E-06	1.2E-06	4.9E-06	6.6E-06
4-tert-Octylphenol	140-66-9	35	9.4E-08	1.2E-06	1.7E-06	5.4E-07	4.8E-06	7.7E-06
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	5.6E-08	4.6E-07	2.8E-07	4.1E-07	9.3E-07	1.2E-06
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	2.6E-07	3.4E-07	1.4E-07	8.9E-07	1.7E-06

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis

Values are rounded to two significant figures.

Table F-316. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Spectators 6<11 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	1.5E-08	1.3E-07	9.1E-08	1.2E-07	2.3E-07	5.0E-07
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	7.0E-08	2.8E-07	0	2.6E-07	1.6E-06
Benzo[a]pyrene	50-32-8	35	5.9E-09	3.0E-08	2.5E-08	2.1E-08	8.3E-08	8.8E-08
Benzo[e]pyrene	192-97-2	35	2.7E-08	8.9E-08	5.3E-08	7.0E-08	1.8E-07	2.0E-07
Benzo[g,h,i]perylene	191-24-2	35	4.2E-09	6.1E-08	3.8E-08	5.4E-08	1.2E-07	1.7E-07
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	2.0E-07	1.8E-07	1.4E-07	5.8E-07	7.1E-07
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	1.5E-07	4.5E-07	0	1.4E-06	1.7E-06



Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Boron		25	0	5.7E-06	6.4E-06	2.6E-06	1.7E-05	2.2E-05
n-Caproic acid vinyl ester	3050-69-9	1	0	1.7E-09	1.0E-08	0	0	5.9E-08
Chrysene	218-01-9	35	4.1E-08	1.6E-07	8.6E-08	1.6E-07	2.8E-07	4.5E-07
Coronene	191-07-1	7	0	2.2E-08	6.4E-08	0	9.4E-08	3.5E-07
Cyclopenta[cd]pyrene	27208-37-3	35	6.1E-09	3.5E-08	3.2E-08	3.0E-08	9.2E-08	1.6E-07
Dicyclohexylamine	101-83-7	25	0	5.1E-07	4.2E-07	4.9E-07	1.2E-06	1.4E-06
Dimethyl phthalate	131-11-3	9	0	1.2E-08	4.0E-08	0	1.0E-07	1.9E-07
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	6.1E-09	1.1E-08	0	2.6E-08	4.2E-08
Methyl stearate	112-61-8	35	3.1E-08	1.3E-07	1.1E-07	9.7E-08	3.0E-07	6.0E-07
Nickel		35	2.6E-07	1.2E-06	1.0E-06	8.4E-07	3.4E-06	4.7E-06
4-tert-Octylphenol	140-66-9	35	6.6E-08	8.7E-07	1.2E-06	3.8E-07	3.4E-06	5.4E-06
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	3.9E-08	3.2E-07	2.0E-07	2.9E-07	6.6E-07	8.8E-07
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	1.8E-07	2.4E-07	1.0E-07	6.2E-07	1.2E-06

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis
Values are rounded to two significant figures.

Table F-317. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Spectators 11<16 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	1.2E-09	1.0E-08	7.3E-09	9.4E-09	1.9E-08	4.0E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	5.6E-09	2.3E-08	0	2.1E-08	1.3E-07
Benzo[a]pyrene	50-32-8	35	4.7E-10	2.4E-09	2.0E-09	1.7E-09	6.7E-09	7.1E-09
Benzo[e]pyrene	192-97-2	35	2.2E-09	7.2E-09	4.3E-09	5.6E-09	1.5E-08	1.6E-08
Benzo[g,h,i]perylene	191-24-2	35	3.4E-10	4.9E-09	3.1E-09	4.4E-09	9.8E-09	1.4E-08
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	1.6E-08	1.4E-08	1.1E-08	4.7E-08	5.7E-08
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	1.2E-08	3.6E-08	0	1.2E-07	1.3E-07
Boron		25	0	4.6E-07	5.1E-07	2.1E-07	1.3E-06	1.8E-06
n-Caproic acid vinyl ester	3050-69-9	1	0	1.4E-10	8.1E-10	0	0	4.8E-09
Chrysene	218-01-9	35	3.3E-09	1.3E-08	7.0E-09	1.3E-08	2.2E-08	3.6E-08
Coronene	191-07-1	7	0	1.8E-09	5.2E-09	0	7.6E-09	2.8E-08
Cyclopenta[cd]pyrene	27208-37-3	35	4.9E-10	2.8E-09	2.6E-09	2.4E-09	7.4E-09	1.3E-08
Dicyclohexylamine	101-83-7	25	0	4.1E-08	3.4E-08	4.0E-08	9.5E-08	1.1E-07



Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dimethyl phthalate	131-11-3	9	0	9.4E-10	3.2E-09	0	8.4E-09	1.5E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	4.9E-10	8.6E-10	0	2.1E-09	3.4E-09
Methyl stearate	112-61-8	35	2.5E-09	1.0E-08	8.9E-09	7.8E-09	2.4E-08	4.8E-08
Nickel		35	2.1E-08	9.8E-08	8.0E-08	6.8E-08	2.8E-07	3.8E-07
4-tert-Octylphenol	140-66-9	35	5.3E-09	7.0E-08	9.5E-08	3.1E-08	2.7E-07	4.4E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	3.2E-09	2.6E-08	1.6E-08	2.3E-08	5.3E-08	7.1E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	1.5E-08	1.9E-08	8.2E-09	5.0E-08	9.5E-08

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis

Values are rounded to two significant figures.

Table F-318. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b**—Combined Gender **Spectators 16<30 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	7.0E-10	5.9E-09	4.2E-09	5.3E-09	1.1E-08	2.3E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	3.2E-09	1.3E-08	0	1.2E-08	7.2E-08
Benzo[a]pyrene	50-32-8	35	2.7E-10	1.4E-09	1.1E-09	9.8E-10	3.8E-09	4.0E-09
Benzo[e]pyrene	192-97-2	35	1.2E-09	4.1E-09	2.4E-09	3.2E-09	8.4E-09	9.2E-09
Benzo[g,h,i]perylene	191-24-2	35	1.9E-10	2.8E-09	1.7E-09	2.5E-09	5.6E-09	7.7E-09
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	8.9E-09	8.2E-09	6.3E-09	2.7E-08	3.3E-08
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	7.1E-09	2.1E-08	0	6.6E-08	7.6E-08
Boron		25	0	2.6E-07	2.9E-07	1.2E-07	7.6E-07	1.0E-06
n-Caproic acid vinyl ester	3050-69-9	1	0	7.8E-11	4.6E-10	0	0	2.7E-09
Chrysene	218-01-9	35	1.9E-09	7.4E-09	4.0E-09	7.4E-09	1.3E-08	2.0E-08
Coronene	191-07-1	7	0	1.0E-09	3.0E-09	0	4.3E-09	1.6E-08
Cyclopenta[cd]pyrene	27208-37-3	35	2.8E-10	1.6E-09	1.5E-09	1.4E-09	4.2E-09	7.5E-09
Dicyclohexylamine	101-83-7	25	0	2.4E-08	1.9E-08	2.3E-08	5.4E-08	6.5E-08
Dimethyl phthalate	131-11-3	9	0	5.4E-10	1.8E-09	0	4.8E-09	8.5E-09
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	2.8E-10	4.9E-10	0	1.2E-09	1.9E-09
Methyl stearate	112-61-8	35	1.4E-09	5.8E-09	5.1E-09	4.5E-09	1.4E-08	2.7E-08
Nickel		35	1.2E-08	5.6E-08	4.6E-08	3.9E-08	1.6E-07	2.1E-07
4-tert-Octylphenol	140-66-9	35	3.0E-09	4.0E-08	5.4E-08	1.7E-08	1.6E-07	2.5E-07



Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.8E-09	1.5E-08	8.9E-09	1.3E-08	3.0E-08	4.0E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	8.3E-09	1.1E-08	4.7E-09	2.9E-08	5.4E-08

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis
Values are rounded to two significant figures.

Table F-319. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Spectators 30<40 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	6.6E-10	5.6E-09	4.0E-09	5.0E-09	1.0E-08	2.2E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	3.0E-09	1.2E-08	0	1.1E-08	6.8E-08
Benzo[a]pyrene	50-32-8	35	2.5E-10	1.3E-09	1.1E-09	9.2E-10	3.6E-09	3.8E-09
Benzo[e]pyrene	192-97-2	35	1.2E-09	3.9E-09	2.3E-09	3.0E-09	8.0E-09	8.7E-09
Benzo[g,h,i]perylene	191-24-2	35	1.8E-10	2.6E-09	1.6E-09	2.4E-09	5.3E-09	7.3E-09
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	8.5E-09	7.7E-09	5.9E-09	2.5E-08	3.1E-08
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	6.7E-09	2.0E-08	0	6.2E-08	7.2E-08
Boron		25	0	2.5E-07	2.8E-07	1.1E-07	7.2E-07	9.6E-07
n-Caproic acid vinyl ester	3050-69-9	1	0	7.4E-11	4.4E-10	0	0	2.6E-09
Chrysene	218-01-9	35	1.8E-09	7.0E-09	3.7E-09	7.0E-09	1.2E-08	1.9E-08
Coronene	191-07-1	7	0	9.6E-10	2.8E-09	0	4.1E-09	1.5E-08
Cyclopenta[cd]pyrene	27208-37-3	35	2.7E-10	1.5E-09	1.4E-09	1.3E-09	4.0E-09	7.1E-09
Dicyclohexylamine	101-83-7	25	0	2.2E-08	1.8E-08	2.1E-08	5.1E-08	6.2E-08
Dimethyl phthalate	131-11-3	9	0	5.1E-10	1.7E-09	0	4.5E-09	8.0E-09
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	2.6E-10	4.6E-10	0	1.1E-09	1.8E-09
Methyl stearate	112-61-8	35	1.4E-09	5.4E-09	4.8E-09	4.2E-09	1.3E-08	2.6E-08
Nickel		35	1.1E-08	5.3E-08	4.3E-08	3.7E-08	1.5E-07	2.0E-07
4-tert-Octylphenol	140-66-9	35	2.9E-09	3.8E-08	5.1E-08	1.7E-08	1.5E-07	2.4E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.7E-09	1.4E-08	8.5E-09	1.3E-08	2.8E-08	3.8E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	7.9E-09	1.0E-08	4.4E-09	2.7E-08	5.1E-08

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.



° CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis
Values are rounded to two significant figures.

Table F-320. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Spectators 40<50 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	6.2E-10	5.3E-09	3.7E-09	4.8E-09	9.5E-09	2.0E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	2.9E-09	1.2E-08	0	1.1E-08	6.4E-08
Benzo[a]pyrene	50-32-8	35	2.4E-10	1.2E-09	1.0E-09	8.7E-10	3.4E-09	3.6E-09
Benzo[e]pyrene	192-97-2	35	1.1E-09	3.7E-09	2.2E-09	2.9E-09	7.5E-09	8.2E-09
Benzo[g,h,i]perylene	191-24-2	35	1.7E-10	2.5E-09	1.6E-09	2.2E-09	5.0E-09	6.9E-09
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	8.0E-09	7.3E-09	5.6E-09	2.4E-08	2.9E-08
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	6.4E-09	1.8E-08	0	5.9E-08	6.8E-08
Boron		25	0	2.3E-07	2.6E-07	1.0E-07	6.8E-07	9.1E-07
n-Caproic acid vinyl ester	3050-69-9	1	0	7.0E-11	4.1E-10	0	0	2.4E-09
Chrysene	218-01-9	35	1.7E-09	6.7E-09	3.5E-09	6.7E-09	1.1E-08	1.8E-08
Coronene	191-07-1	7	0	9.1E-10	2.6E-09	0	3.9E-09	1.4E-08
Cyclopenta[cd]pyrene	27208-37-3	35	2.5E-10	1.4E-09	1.3E-09	1.2E-09	3.8E-09	6.7E-09
Dicyclohexylamine	101-83-7	25	0	2.1E-08	1.7E-08	2.0E-08	4.9E-08	5.8E-08
Dimethyl phthalate	131-11-3	9	0	4.8E-10	1.6E-09	0	4.3E-09	7.6E-09
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	2.5E-10	4.4E-10	0	1.1E-09	1.7E-09
Methyl stearate	112-61-8	35	1.3E-09	5.1E-09	4.5E-09	4.0E-09	1.2E-08	2.4E-08
Nickel		35	1.1E-08	5.0E-08	4.1E-08	3.5E-08	1.4E-07	1.9E-07
4-tert-Octylphenol	140-66-9	35	2.7E-09	3.6E-08	4.9E-08	1.6E-08	1.4E-07	2.2E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.6E-09	1.3E-08	8.0E-09	1.2E-08	2.7E-08	3.6E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	7.5E-09	9.8E-09	4.2E-09	2.6E-08	4.9E-08

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.
Values are rounded to two significant figures.



Table F-321. Field-Specific Average Ingestion One-Day Dose^a (AD_{ing-DART-field}, milligrams per kilogram BW per day) for One-Day Non-Cancer Hazard Assessment of **Field-Related DARTs^b—Combined Gender Spectators 50<70 Years**

Chemical	CASRN ^c	Detection	AD _{ing-DART-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	6.2E-10	5.3E-09	3.8E-09	4.8E-09	9.6E-09	2.0E-08
1,4-Benzenediamine, N,N'-diphenyl-	74-31-7	7	0	2.9E-09	1.2E-08	0	1.1E-08	6.5E-08
Benzo[a]pyrene	50-32-8	35	2.4E-10	1.2E-09	1.0E-09	8.8E-10	3.4E-09	3.6E-09
Benzo[e]pyrene	192-97-2	35	1.1E-09	3.7E-09	2.2E-09	2.9E-09	7.6E-09	8.3E-09
Benzo[g,h,i]perylene	191-24-2	35	1.7E-10	2.5E-09	1.6E-09	2.2E-09	5.0E-09	6.9E-09
Bis(2-Ethylhexyl)adipate	103-23-1	33	0	8.0E-09	7.3E-09	5.6E-09	2.4E-08	2.9E-08
Bis(2,2,6,6-tetramethyl-4-piperidyl)sebacate	52829-07-9	4	0	6.4E-09	1.9E-08	0	5.9E-08	6.8E-08
Boron		25	0	2.3E-07	2.6E-07	1.1E-07	6.9E-07	9.1E-07
n-Caproic acid vinyl ester	3050-69-9	1	0	7.0E-11	4.1E-10	0	0	2.4E-09
Chrysene	218-01-9	35	1.7E-09	6.7E-09	3.5E-09	6.7E-09	1.1E-08	1.8E-08
Coronene	191-07-1	7	0	9.2E-10	2.7E-09	0	3.9E-09	1.4E-08
Cyclopenta[cd]pyrene	27208-37-3	35	2.5E-10	1.4E-09	1.3E-09	1.2E-09	3.8E-09	6.7E-09
Dicyclohexylamine	101-83-7	25	0	2.1E-08	1.7E-08	2.0E-08	4.9E-08	5.9E-08
Dimethyl phthalate	131-11-3	9	0	4.8E-10	1.6E-09	0	4.3E-09	7.6E-09
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	2.5E-10	4.4E-10	0	1.1E-09	1.7E-09
Methyl stearate	112-61-8	35	1.3E-09	5.2E-09	4.5E-09	4.0E-09	1.2E-08	2.5E-08
Nickel		35	1.1E-08	5.0E-08	4.1E-08	3.5E-08	1.4E-07	1.9E-07
4-tert-Octylphenol	140-66-9	35	2.7E-09	3.6E-08	4.9E-08	1.6E-08	1.4E-07	2.2E-07
Phenol, 2,4-bis(1-methyl-1-phenylethyl)-	2772-45-4	35	1.6E-09	1.3E-08	8.0E-09	1.2E-08	2.7E-08	3.6E-08
Phenol, 4-(1-phenylethyl)-	1988-89-2	34	0	7.5E-09	9.8E-09	4.2E-09	2.6E-08	4.9E-08

^a 35 field-specific AD_{ing-DART-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.



F.5.6. Average Ingestion Daily Dose (ADD_{ing}) for Chronic Non-Cancer Hazard of General Chemicals

Table F-322. Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—
Combined Gender **Athletes**

Chemical	CASRN ^c	ADD _{ing}						
		2<6 Years	6<11 Years	11<16 Years	16<30 Years	30<40 Years	40<50 Years	50<70 Years
Acenaphthylene	208-96-8	1.5E-09	9.8E-10	7.6E-10	8.5E-10	5.6E-10	4.8E-10	4.9E-10
Aluminum		1.0E-04	6.9E-05	5.3E-05	5.9E-05	4.0E-05	3.4E-05	3.4E-05
Aniline	62-53-3	8.2E-09	5.5E-09	4.2E-09	4.7E-09	3.1E-09	2.7E-09	2.7E-09
Anthracene	120-12-7	3.9E-09	2.6E-09	2.0E-09	2.2E-09	1.5E-09	1.3E-09	1.3E-09
Anthracene, 2-methyl-	613-12-7	1.5E-08	1.0E-08	8.0E-09	8.9E-09	5.9E-09	5.0E-09	5.1E-09
Anthracene, 9,10-dimethyl	781-43-1	1.2E-10	8.3E-11	6.4E-11	7.2E-11	4.8E-11	4.1E-11	4.1E-11
Anthracene, 9,10-diphenyl-	1499-10-1	9.6E-10	6.4E-10	5.0E-10	5.5E-10	3.7E-10	3.1E-10	3.2E-10
Anthracene, 9-phenyl	602-55-1	1.5E-09	9.7E-10	7.5E-10	8.4E-10	5.6E-10	4.7E-10	4.8E-10
Antimony		1.1E-07	7.5E-08	5.8E-08	6.4E-08	4.3E-08	3.6E-08	3.7E-08
Barium		7.1E-06	4.7E-06	3.7E-06	4.1E-06	2.7E-06	2.3E-06	2.4E-06
Benzene, n-butyl-	104-51-8	8.2E-10	5.5E-10	4.2E-10	4.7E-10	3.2E-10	2.7E-10	2.7E-10
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	2.2E-07	1.5E-07	1.2E-07	1.3E-07	8.6E-08	7.3E-08	7.5E-08
Benz[a]anthracene	56-55-3	1.3E-08	9.0E-09	7.0E-09	7.8E-09	5.2E-09	4.4E-09	4.5E-09
Benzo[b]fluoranthene	205-99-2	1.9E-08	1.2E-08	9.6E-09	1.1E-08	7.1E-09	6.0E-09	6.2E-09
7H-Benzo[c]fluorene	205-12-9	3.6E-09	2.4E-09	1.8E-09	2.1E-09	1.4E-09	1.2E-09	1.2E-09
Benzo[k]fluoranthene	207-08-9	6.0E-09	4.0E-09	3.1E-09	3.5E-09	2.3E-09	2.0E-09	2.0E-09
Benzothiazole	95-16-9	2.3E-06	1.5E-06	1.2E-06	1.3E-06	8.8E-07	7.4E-07	7.6E-07
Benzothiazole, 2-phenyl-	883-93-2	2.4E-07	1.6E-07	1.3E-07	1.4E-07	9.4E-08	8.0E-08	8.1E-08
Benzothiazolone	934-34-9	5.8E-06	3.9E-06	3.0E-06	3.3E-06	2.2E-06	1.9E-06	1.9E-06
Benzyl butyl phthalate	85-68-7	1.2E-07	7.8E-08	6.0E-08	6.7E-08	4.5E-08	3.8E-08	3.9E-08
Beryllium		2.9E-09	2.0E-09	1.5E-09	1.7E-09	1.1E-09	9.5E-10	9.7E-10
Butylated Hydroxytoluene	128-37-0	3.1E-09	2.1E-09	1.6E-09	1.8E-09	1.2E-09	1.0E-09	1.0E-09
Cadmium		7.9E-08	5.3E-08	4.1E-08	4.6E-08	3.0E-08	2.6E-08	2.6E-08
Chromium		2.2E-07	1.5E-07	1.1E-07	1.3E-07	8.5E-08	7.2E-08	7.3E-08
Cobalt		5.0E-06	3.4E-06	2.6E-06	2.9E-06	1.9E-06	1.6E-06	1.7E-06
Copper		9.8E-06	6.6E-06	5.1E-06	5.7E-06	3.8E-06	3.2E-06	3.3E-06
Cyclohexyl isothiocyanate	1122-82-3	7.3E-07	4.9E-07	3.8E-07	4.2E-07	2.8E-07	2.4E-07	2.4E-07
Dibenz[a,h]anthracene	53-70-3	8.0E-10	5.3E-10	4.1E-10	4.6E-10	3.1E-10	2.6E-10	2.7E-10
Dibenzothiophene	132-65-0	3.1E-09	2.1E-09	1.6E-09	1.8E-09	1.2E-09	1.0E-09	1.0E-09
Dibutyl phthalate	84-74-2	2.2E-07	1.5E-07	1.1E-07	1.3E-07	8.5E-08	7.2E-08	7.4E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	ADD _{ing}						
		2<6 Years	6<11 Years	11<16 Years	16<30 Years	30<40 Years	40<50 Years	50<70 Years
Diethyl Phthalate	84-66-2	1.9E-08	1.2E-08	9.6E-09	1.1E-08	7.2E-09	6.1E-09	6.2E-09
Diisobutyl Phthalate	84-69-5	2.2E-08	1.5E-08	1.1E-08	1.3E-08	8.5E-09	7.2E-09	7.4E-09
Diisooctylphthalate	27554-26-3	4.7E-07	3.1E-07	2.4E-07	2.7E-07	1.8E-07	1.5E-07	1.6E-07
Di-n-octyl phthalate	117-84-0	4.6E-08	3.1E-08	2.4E-08	2.7E-08	1.8E-08	1.5E-08	1.5E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	3.8E-10	2.6E-10	2.0E-10	2.2E-10	1.5E-10	1.2E-10	1.3E-10
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	8.2E-08	5.5E-08	4.2E-08	4.7E-08	3.2E-08	2.7E-08	2.7E-08
Fluoranthene	206-44-0	1.0E-07	7.0E-08	5.4E-08	6.0E-08	4.0E-08	3.4E-08	3.5E-08
Fluorene	86-73-7	1.4E-09	9.1E-10	7.0E-10	7.8E-10	5.2E-10	4.4E-10	4.5E-10
Hexadecane	544-76-3	9.8E-09	6.5E-09	5.1E-09	5.7E-09	3.8E-09	3.2E-09	3.3E-09
1-Hydroxypyrene	5315-79-7	5.3E-09	3.5E-09	2.8E-09	3.1E-09	2.0E-09	1.7E-09	1.8E-09
Lead		4.6E-06	3.1E-06	2.4E-06	2.7E-06	1.8E-06	1.5E-06	1.5E-06
Manganese		4.6E-06	3.1E-06	2.4E-06	2.6E-06	1.8E-06	1.5E-06	1.5E-06
2-(Methylthio)benzothiazole	615-22-5	7.0E-08	4.7E-08	3.6E-08	4.0E-08	2.7E-08	2.3E-08	2.3E-08
Molybdenum		8.5E-09	5.6E-09	4.4E-09	4.9E-09	3.3E-09	2.8E-09	2.8E-09
Naphthalene	91-20-3	2.4E-10	1.6E-10	1.3E-10	1.4E-10	9.4E-11	8.0E-11	8.1E-11
Naphthalene, 1-methyl-	90-12-0	6.3E-10	4.2E-10	3.3E-10	3.6E-10	2.4E-10	2.1E-10	2.1E-10
Naphthalene, 1,2-dimethyl-	573-98-8	1.5E-10	9.8E-11	7.6E-11	8.5E-11	5.7E-11	4.8E-11	4.9E-11
Naphthalene, 1,6-dimethyl-	575-43-9	9.7E-10	6.5E-10	5.0E-10	5.6E-10	3.7E-10	3.2E-10	3.2E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	2.6E-08	1.7E-08	1.3E-08	1.5E-08	9.8E-09	8.3E-09	8.5E-09
Naphthalene, 2,3-dimethyl-	581-40-8	7.5E-09	5.0E-09	3.9E-09	4.3E-09	2.9E-09	2.4E-09	2.5E-09
Naphthalene, 2-methyl	91-57-6	5.7E-10	3.8E-10	3.0E-10	3.3E-10	2.2E-10	1.9E-10	1.9E-10
1-Octadecene	112-88-9	2.2E-08	1.4E-08	1.1E-08	1.2E-08	8.3E-09	7.1E-09	7.2E-09
17-Pentatriacontene	6971-40-0	1.8E-08	1.2E-08	9.5E-09	1.1E-08	7.1E-09	6.0E-09	6.1E-09
Phenanthrene	85-01-8	4.6E-08	3.1E-08	2.4E-08	2.6E-08	1.8E-08	1.5E-08	1.5E-08
Phenanthrene, 1-methyl	832-69-9	1.2E-08	8.1E-09	6.3E-09	7.0E-09	4.7E-09	4.0E-09	4.0E-09
Phenanthrene, 2-methyl-	2531-84-2	1.6E-08	1.0E-08	8.1E-09	9.1E-09	6.0E-09	5.1E-09	5.2E-09
Phenanthrene, 3-methyl	832-71-3	2.6E-08	1.7E-08	1.3E-08	1.5E-08	1.0E-08	8.5E-09	8.6E-09
N-Phenylbenzamide	93-98-1	1.0E-07	6.8E-08	5.3E-08	5.8E-08	3.9E-08	3.3E-08	3.4E-08
Phthalimide	85-41-6	1.8E-08	1.2E-08	9.5E-09	1.1E-08	7.1E-09	6.0E-09	6.1E-09
Pyrene	129-00-0	2.2E-07	1.5E-07	1.1E-07	1.3E-07	8.4E-08	7.2E-08	7.3E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	4.7E-10	3.1E-10	2.4E-10	2.7E-10	1.8E-10	1.5E-10	1.6E-10
Selenium		2.8E-08	1.9E-08	1.4E-08	1.6E-08	1.1E-08	9.0E-09	9.2E-09
Strontium		3.9E-06	2.6E-06	2.0E-06	2.3E-06	1.5E-06	1.3E-06	1.3E-06
Thallium		1.1E-09	7.4E-10	5.7E-10	6.4E-10	4.2E-10	3.6E-10	3.7E-10
Tin		4.3E-08	2.9E-08	2.2E-08	2.5E-08	1.7E-08	1.4E-08	1.4E-08



Chemical	CASRN ^c	ADD _{ing}						
		2<6 Years	6<11 Years	11<16 Years	16<30 Years	30<40 Years	40<50 Years	50<70 Years
Triethylene glycol monobutyl ether	143-22-6	5.4E-09	3.6E-09	2.8E-09	3.1E-09	2.1E-09	1.7E-09	1.8E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	1.6E-08	1.1E-08	8.2E-09	9.1E-09	6.1E-09	5.2E-09	5.3E-09
Vanadium		2.2E-08	1.5E-08	1.2E-08	1.3E-08	8.6E-09	7.3E-09	7.5E-09
Zinc		9.7E-04	6.5E-04	5.0E-04	5.6E-04	3.7E-04	3.2E-04	3.2E-04

^a ADD_{ing} was calculated from the mean of 35 individual field average concentrations (C_{ing-crumb rubber}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-323. Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—
Combined Gender Coaches

Chemical	CASRN ^c	ADD _{ing}			
		16<30 Years	30<40 Years	40<50 Years	50<70 Years
Acenaphthylene	208-96-8	2.1E-10	2.0E-10	2.0E-10	2.0E-10
Aluminum		1.5E-05	1.4E-05	1.4E-05	1.4E-05
Aniline	62-53-3	1.2E-09	1.1E-09	1.1E-09	1.1E-09
Anthracene	120-12-7	5.6E-10	5.3E-10	5.2E-10	5.2E-10
Anthracene, 2-methyl-	613-12-7	2.2E-09	2.1E-09	2.1E-09	2.1E-09
Anthracene, 9,10-dimethyl	781-43-1	1.8E-11	1.7E-11	1.7E-11	1.7E-11
Anthracene, 9,10-diphenyl-	1499-10-1	1.4E-10	1.3E-10	1.3E-10	1.3E-10
Anthracene, 9-phenyl	602-55-1	2.1E-10	2.0E-10	1.9E-10	2.0E-10
Antimony		1.6E-08	1.5E-08	1.5E-08	1.5E-08
Barium		1.0E-06	9.7E-07	9.5E-07	9.5E-07
Benzene, n-butyl-	104-51-8	1.2E-10	1.1E-10	1.1E-10	1.1E-10
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	3.3E-08	3.1E-08	3.0E-08	3.0E-08
Benz[a]anthracene	56-55-3	2.0E-09	1.8E-09	1.8E-09	1.8E-09
Benzo[b]fluoranthene	205-99-2	2.7E-09	2.5E-09	2.5E-09	2.5E-09
7H-Benzo[c]fluorene	205-12-9	5.2E-10	4.9E-10	4.8E-10	4.8E-10
Benzo[k]fluoranthene	207-08-9	8.7E-10	8.2E-10	8.0E-10	8.0E-10
Benzothiazole	95-16-9	3.3E-07	3.1E-07	3.0E-07	3.1E-07
Benzothiazole, 2-phenyl-	883-93-2	3.6E-08	3.3E-08	3.3E-08	3.3E-08
Benzothiazolone	934-34-9	8.5E-07	7.9E-07	7.8E-07	7.8E-07
Benzyl butyl phthalate	85-68-7	1.7E-08	1.6E-08	1.6E-08	1.6E-08
Beryllium		4.3E-10	4.0E-10	3.9E-10	3.9E-10



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	ADD _{ing}			
		16<30 Years	30<40 Years	40<50 Years	50<70 Years
Butylated Hydroxytoluene	128-37-0	4.5E-10	4.2E-10	4.2E-10	4.2E-10
Cadmium		1.2E-08	1.1E-08	1.1E-08	1.1E-08
Chromium		3.2E-08	3.0E-08	3.0E-08	3.0E-08
Cobalt		7.3E-07	6.9E-07	6.7E-07	6.7E-07
Copper		1.4E-06	1.3E-06	1.3E-06	1.3E-06
Cyclohexyl isothiocyanate	1122-82-3	1.1E-07	9.9E-08	9.7E-08	9.8E-08
Dibenz[a,h]anthracene	53-70-3	1.2E-10	1.1E-10	1.1E-10	1.1E-10
Dibenzothiophene	132-65-0	4.5E-10	4.3E-10	4.2E-10	4.2E-10
Dibutyl phthalate	84-74-2	3.2E-08	3.0E-08	3.0E-08	3.0E-08
Diethyl Phthalate	84-66-2	2.7E-09	2.5E-09	2.5E-09	2.5E-09
Diisobutyl Phthalate	84-69-5	3.2E-09	3.0E-09	3.0E-09	3.0E-09
Diisooctylphthalate	27554-26-3	6.8E-08	6.4E-08	6.3E-08	6.3E-08
Di-n-octyl phthalate	117-84-0	6.7E-09	6.3E-09	6.2E-09	6.2E-09
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	5.6E-11	5.2E-11	5.1E-11	5.1E-11
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	1.2E-08	1.1E-08	1.1E-08	1.1E-08
Fluoranthene	206-44-0	1.5E-08	1.4E-08	1.4E-08	1.4E-08
Fluorene	86-73-7	2.0E-10	1.9E-10	1.8E-10	1.8E-10
Hexadecane	544-76-3	1.4E-09	1.3E-09	1.3E-09	1.3E-09
1-Hydroxypyrene	5315-79-7	7.7E-10	7.3E-10	7.1E-10	7.1E-10
Lead		6.7E-07	6.3E-07	6.2E-07	6.2E-07
Manganese		6.7E-07	6.3E-07	6.1E-07	6.1E-07
2-(Methylthio)benzothiazole	615-22-5	1.0E-08	9.5E-09	9.3E-09	9.4E-09
Molybdenum		1.2E-09	1.2E-09	1.1E-09	1.1E-09
Naphthalene	91-20-3	3.6E-11	3.3E-11	3.3E-11	3.3E-11
Naphthalene, 1-methyl-	90-12-0	9.2E-11	8.6E-11	8.4E-11	8.4E-11
Naphthalene, 1,2-dimethyl-	573-98-8	2.1E-11	2.0E-11	2.0E-11	2.0E-11
Naphthalene, 1,6-dimethyl-	575-43-9	1.4E-10	1.3E-10	1.3E-10	1.3E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	3.7E-09	3.5E-09	3.4E-09	3.4E-09
Naphthalene, 2,3-dimethyl-	581-40-8	1.1E-09	1.0E-09	1.0E-09	1.0E-09
Naphthalene, 2-methyl	91-57-6	8.3E-11	7.8E-11	7.6E-11	7.6E-11
1-Octadecene	112-88-9	3.2E-09	3.0E-09	2.9E-09	2.9E-09
17-Pentatriacontene	6971-40-0	2.7E-09	2.5E-09	2.5E-09	2.5E-09
Phenanthrene	85-01-8	6.7E-09	6.3E-09	6.1E-09	6.1E-09
Phenanthrene, 1-methyl	832-69-9	1.8E-09	1.7E-09	1.6E-09	1.6E-09
Phenanthrene, 2-methyl-	2531-84-2	2.3E-09	2.1E-09	2.1E-09	2.1E-09
Phenanthrene, 3-methyl	832-71-3	3.8E-09	3.5E-09	3.5E-09	3.5E-09
N-Phenylbenzamide	93-98-1	1.5E-08	1.4E-08	1.4E-08	1.4E-08
Phthalimide	85-41-6	2.7E-09	2.5E-09	2.5E-09	2.5E-09
Pyrene	129-00-0	3.2E-08	3.0E-08	2.9E-08	2.9E-08



Chemical	CASRN ^c	ADD _{ing}			
		16<30 Years	30<40 Years	40<50 Years	50<70 Years
Pyridine, 2-(4-methylphenyl)-	4467-06-5	6.8E-11	6.4E-11	6.3E-11	6.3E-11
Selenium		4.0E-09	3.8E-09	3.7E-09	3.7E-09
Strontium		5.7E-07	5.4E-07	5.3E-07	5.3E-07
Thallium		1.6E-10	1.5E-10	1.5E-10	1.5E-10
Tin		6.3E-09	5.9E-09	5.8E-09	5.8E-09
Triethylene glycol monobutyl ether	143-22-6	7.8E-10	7.3E-10	7.2E-10	7.2E-10
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	2.3E-09	2.2E-09	2.1E-09	2.1E-09
Vanadium		3.3E-09	3.1E-09	3.0E-09	3.0E-09
Zinc		1.4E-04	1.3E-04	1.3E-04	1.3E-04

^a ADD_{ing} was calculated from the mean of 35 individual field average concentrations (C_{ing-crumb rubber}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-324. Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—
Combined Gender **Referees**

Chemical	CASRN ^c	ADD _{ing}			
		16<30 Years	30<40 Years	40<50 Years	50<70 Years
Acenaphthylene	208-96-8	9.4E-11	8.8E-11	8.6E-11	8.6E-11
Aluminum		6.6E-06	6.2E-06	6.1E-06	6.1E-06
Aniline	62-53-3	5.2E-10	4.9E-10	4.8E-10	4.8E-10
Anthracene	120-12-7	2.5E-10	2.3E-10	2.3E-10	2.3E-10
Anthracene, 2-methyl-	613-12-7	9.8E-10	9.2E-10	9.0E-10	9.0E-10
Anthracene, 9,10-dimethyl	781-43-1	7.9E-12	7.5E-12	7.3E-12	7.3E-12
Anthracene, 9,10-diphenyl-	1499-10-1	6.1E-11	5.8E-11	5.6E-11	5.7E-11
Anthracene, 9-phenyl	602-55-1	9.3E-11	8.7E-11	8.5E-11	8.6E-11
Antimony		7.1E-09	6.7E-09	6.6E-09	6.6E-09
Barium		4.5E-07	4.3E-07	4.2E-07	4.2E-07
Benzene, n-butyl-	104-51-8	5.2E-11	4.9E-11	4.8E-11	4.8E-11
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	1.4E-08	1.3E-08	1.3E-08	1.3E-08
Benz[a]anthracene	56-55-3	8.6E-10	8.1E-10	7.9E-10	7.9E-10
Benzo[b]fluoranthene	205-99-2	1.2E-09	1.1E-09	1.1E-09	1.1E-09
7H-Benzo[c]fluorene	205-12-9	2.3E-10	2.1E-10	2.1E-10	2.1E-10
Benzo[k]fluoranthene	207-08-9	3.8E-10	3.6E-10	3.5E-10	3.5E-10
Benzothiazole	95-16-9	1.5E-07	1.4E-07	1.3E-07	1.3E-07



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	ADD _{ing}			
		16<30 Years	30<40 Years	40<50 Years	50<70 Years
Benzothiazole, 2-phenyl-	883-93-2	1.6E-08	1.5E-08	1.4E-08	1.4E-08
Benzothiazolone	934-34-9	3.7E-07	3.5E-07	3.4E-07	3.4E-07
Benzyl butyl phthalate	85-68-7	7.4E-09	7.0E-09	6.8E-09	6.9E-09
Beryllium		1.9E-10	1.8E-10	1.7E-10	1.7E-10
Butylated Hydroxytoluene	128-37-0	2.0E-10	1.9E-10	1.8E-10	1.8E-10
Cadmium		5.1E-09	4.8E-09	4.7E-09	4.7E-09
Chromium		1.4E-08	1.3E-08	1.3E-08	1.3E-08
Cobalt		3.2E-07	3.0E-07	3.0E-07	3.0E-07
Copper		6.3E-07	5.9E-07	5.8E-07	5.8E-07
Cyclohexyl isothiocyanate	1122-82-3	4.6E-08	4.4E-08	4.3E-08	4.3E-08
Dibenz[a,h]anthracene	53-70-3	5.1E-11	4.8E-11	4.7E-11	4.7E-11
Dibenzothiophene	132-65-0	2.0E-10	1.9E-10	1.8E-10	1.8E-10
Dibutyl phthalate	84-74-2	1.4E-08	1.3E-08	1.3E-08	1.3E-08
Diethyl Phthalate	84-66-2	1.2E-09	1.1E-09	1.1E-09	1.1E-09
Diisobutyl Phthalate	84-69-5	1.4E-09	1.3E-09	1.3E-09	1.3E-09
Diisooctylphthalate	27554-26-3	3.0E-08	2.8E-08	2.8E-08	2.8E-08
Di-n-octyl phthalate	117-84-0	3.0E-09	2.8E-09	2.7E-09	2.7E-09
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	2.4E-11	2.3E-11	2.3E-11	2.3E-11
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	5.2E-09	4.9E-09	4.8E-09	4.8E-09
Fluoranthene	206-44-0	6.7E-09	6.2E-09	6.1E-09	6.1E-09
Fluorene	86-73-7	8.7E-11	8.1E-11	8.0E-11	8.0E-11
Hexadecane	544-76-3	6.3E-10	5.9E-10	5.8E-10	5.8E-10
1-Hydroxypyrene	5315-79-7	3.4E-10	3.2E-10	3.1E-10	3.1E-10
Lead		3.0E-07	2.8E-07	2.7E-07	2.7E-07
Manganese		2.9E-07	2.8E-07	2.7E-07	2.7E-07
2-(Methylthio)benzothiazole	615-22-5	4.5E-09	4.2E-09	4.1E-09	4.1E-09
Molybdenum		5.4E-10	5.1E-10	5.0E-10	5.0E-10
Naphthalene	91-20-3	1.6E-11	1.5E-11	1.4E-11	1.4E-11
Naphthalene, 1-methyl-	90-12-0	4.0E-11	3.8E-11	3.7E-11	3.7E-11
Naphthalene, 1,2-dimethyl-	573-98-8	9.4E-12	8.8E-12	8.6E-12	8.7E-12
Naphthalene, 1,6-dimethyl-	575-43-9	6.2E-11	5.8E-11	5.7E-11	5.7E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	1.6E-09	1.5E-09	1.5E-09	1.5E-09
Naphthalene, 2,3-dimethyl-	581-40-8	4.8E-10	4.5E-10	4.4E-10	4.4E-10
Naphthalene, 2-methyl	91-57-6	3.6E-11	3.4E-11	3.4E-11	3.4E-11
1-Octadecene	112-88-9	1.4E-09	1.3E-09	1.3E-09	1.3E-09
17-Pentatriacontene	6971-40-0	1.2E-09	1.1E-09	1.1E-09	1.1E-09
Phenanthrene	85-01-8	2.9E-09	2.7E-09	2.7E-09	2.7E-09
Phenanthrene, 1-methyl	832-69-9	7.8E-10	7.3E-10	7.1E-10	7.2E-10
Phenanthrene, 2-methyl-	2531-84-2	1.0E-09	9.4E-10	9.2E-10	9.3E-10



Chemical	CASRN ^c	ADD _{ing}			
		16<30 Years	30<40 Years	40<50 Years	50<70 Years
Phenanthrene, 3-methyl	832-71-3	1.7E-09	1.6E-09	1.5E-09	1.5E-09
N-Phenylbenzamide	93-98-1	6.5E-09	6.1E-09	6.0E-09	6.0E-09
Phthalimide	85-41-6	1.2E-09	1.1E-09	1.1E-09	1.1E-09
Pyrene	129-00-0	1.4E-08	1.3E-08	1.3E-08	1.3E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	3.0E-11	2.8E-11	2.8E-11	2.8E-11
Selenium		1.8E-09	1.7E-09	1.6E-09	1.6E-09
Strontium		2.5E-07	2.4E-07	2.3E-07	2.3E-07
Thallium		7.0E-11	6.6E-11	6.5E-11	6.5E-11
Tin		2.8E-09	2.6E-09	2.5E-09	2.5E-09
Triethylene glycol monobutyl ether	143-22-6	3.4E-10	3.2E-10	3.1E-10	3.2E-10
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	1.0E-09	9.5E-10	9.3E-10	9.3E-10
Vanadium		1.4E-09	1.3E-09	1.3E-09	1.3E-09
Zinc		6.2E-05	5.8E-05	5.7E-05	5.7E-05

^a ADD_{ing} was calculated from the mean of 35 individual field average concentrations (C_{ing-crumb rubber}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-325. Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—
Combined Gender Spectators

Chemical Name	CASRN	ADD _{ing}								
		Third Trimester Fetus	0<2 Years	2<6 Years	6<11 Years	11<16 Years	16<30 Years	30<40 Years	40<50 Years	50<70 Years
Acenaphthylene	208-96-8	7.4E-11	5.1E-09	2.4E-09	1.7E-09	1.3E-10	7.6E-11	7.2E-11	6.8E-11	6.8E-11
Aluminum		5.2E-06	3.6E-04	1.7E-04	1.2E-04	9.4E-06	5.4E-06	5.1E-06	4.8E-06	4.8E-06
Aniline	62-53-3	4.1E-10	2.9E-08	1.4E-08	9.5E-09	7.5E-10	4.2E-10	4.0E-10	3.8E-10	3.8E-10
Anthracene	120-12-7	1.9E-10	1.4E-08	6.4E-09	4.5E-09	3.5E-10	2.0E-10	1.9E-10	1.8E-10	1.8E-10
Anthracene, 2-methyl-	613-12-7	7.7E-10	5.4E-08	2.5E-08	1.8E-08	1.4E-09	8.0E-10	7.5E-10	7.1E-10	7.2E-10
Anthracene, 9,10-dimethyl	781-43-1	6.3E-12	4.4E-10	2.1E-10	1.4E-10	1.1E-11	6.5E-12	6.1E-12	5.8E-12	5.8E-12
Anthracene, 9,10-diphenyl-	1499-10-1	4.8E-11	3.4E-09	1.6E-09	1.1E-09	8.8E-11	5.0E-11	4.7E-11	4.5E-11	4.5E-11
Anthracene, 9-phenyl	602-55-1	7.3E-11	5.1E-09	2.4E-09	1.7E-09	1.3E-10	7.5E-11	7.1E-11	6.8E-11	6.8E-11
Antimony		5.6E-09	3.9E-07	1.9E-07	1.3E-07	1.0E-08	5.8E-09	5.5E-09	5.2E-09	5.2E-09
Barium		3.6E-07	2.5E-05	1.2E-05	8.3E-06	6.5E-07	3.7E-07	3.5E-07	3.3E-07	3.3E-07
Benzene, n-butyl-	104-51-8	4.1E-11	2.9E-09	1.4E-09	9.6E-10	7.5E-11	4.3E-11	4.0E-11	3.8E-11	3.8E-11
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	1.1E-08	7.8E-07	3.7E-07	2.6E-07	2.0E-08	1.2E-08	1.1E-08	1.0E-08	1.0E-08
Benz[a]anthracene	56-55-3	6.8E-10	4.7E-08	2.2E-08	1.6E-08	1.2E-09	7.0E-10	6.6E-10	6.3E-10	6.3E-10



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical Name	CASRN	ADD _{ing}								
		Third Trimester Fetus	0<2 Years	2<6 Years	6<11 Years	11<16 Years	16<30 Years	30<40 Years	40<50 Years	50<70 Years
Benzo[b]fluoranthene	205-99-2	9.3E-10	6.5E-08	3.1E-08	2.2E-08	1.7E-09	9.6E-10	9.1E-10	8.6E-10	8.6E-10
7H-Benzo[c]fluorene	205-12-9	1.8E-10	1.3E-08	5.9E-09	4.2E-09	3.3E-10	1.9E-10	1.8E-10	1.7E-10	1.7E-10
Benzo[k]fluoranthene	207-08-9	3.0E-10	2.1E-08	1.0E-08	7.0E-09	5.5E-10	3.1E-10	2.9E-10	2.8E-10	2.8E-10
Benzothiazole	95-16-9	1.1E-07	8.0E-06	3.8E-06	2.7E-06	2.1E-07	1.2E-07	1.1E-07	1.1E-07	1.1E-07
Benzothiazole, 2-phenyl-	883-93-2	1.2E-08	8.6E-07	4.1E-07	2.8E-07	2.2E-08	1.3E-08	1.2E-08	1.1E-08	1.1E-08
Benzothiazolone	934-34-9	2.9E-07	2.0E-05	9.6E-06	6.8E-06	5.3E-07	3.0E-07	2.9E-07	2.7E-07	2.7E-07
Benzyl butyl phthalate	85-68-7	5.9E-09	4.1E-07	1.9E-07	1.4E-07	1.1E-08	6.0E-09	5.7E-09	5.4E-09	5.4E-09
Beryllium		1.5E-10	1.0E-08	4.9E-09	3.4E-09	2.7E-10	1.5E-10	1.4E-10	1.4E-10	1.4E-10
Butylated Hydroxytoluene	128-37-0	1.6E-10	1.1E-08	5.2E-09	3.6E-09	2.8E-10	1.6E-10	1.5E-10	1.4E-10	1.4E-10
Cadmium		4.0E-09	2.8E-07	1.3E-07	9.2E-08	7.2E-09	4.1E-09	3.9E-09	3.7E-09	3.7E-09
Chromium		1.1E-08	7.7E-07	3.7E-07	2.6E-07	2.0E-08	1.1E-08	1.1E-08	1.0E-08	1.0E-08
Cobalt		2.5E-07	1.8E-05	8.3E-06	5.8E-06	4.6E-07	2.6E-07	2.5E-07	2.3E-07	2.3E-07
Copper		5.0E-07	3.4E-05	1.6E-05	1.1E-05	9.0E-07	5.1E-07	4.8E-07	4.6E-07	4.6E-07
Cyclohexyl isothiocyanate	1122-82-3	3.7E-08	2.5E-06	1.2E-06	8.5E-07	6.6E-08	3.8E-08	3.6E-08	3.4E-08	3.4E-08
Dibenz[a,h]anthracene	53-70-3	4.0E-11	2.8E-09	1.3E-09	9.3E-10	7.3E-11	4.1E-11	3.9E-11	3.7E-11	3.7E-11
Dibenzothiophene	132-65-0	1.6E-10	1.1E-08	5.2E-09	3.6E-09	2.8E-10	1.6E-10	1.5E-10	1.4E-10	1.5E-10
Dibutyl phthalate	84-74-2	1.1E-08	7.8E-07	3.7E-07	2.6E-07	2.0E-08	1.1E-08	1.1E-08	1.0E-08	1.0E-08
Diethyl Phthalate	84-66-2	9.4E-10	6.5E-08	3.1E-08	2.2E-08	1.7E-09	9.7E-10	9.1E-10	8.7E-10	8.7E-10
Diisobutyl Phthalate	84-69-5	1.1E-09	7.8E-08	3.7E-08	2.6E-08	2.0E-09	1.2E-09	1.1E-09	1.0E-09	1.0E-09
Diisooctylphthalate	27554-26-3	2.4E-08	1.6E-06	7.8E-07	5.5E-07	4.3E-08	2.4E-08	2.3E-08	2.2E-08	2.2E-08
Di-n-octyl phthalate	117-84-0	2.3E-09	1.6E-07	7.7E-08	5.4E-08	4.2E-09	2.4E-09	2.3E-09	2.1E-09	2.2E-09
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	1.9E-11	1.3E-09	6.4E-10	4.5E-10	3.5E-11	2.0E-11	1.9E-11	1.8E-11	1.8E-11
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	4.1E-09	2.9E-07	1.4E-07	9.5E-08	7.5E-09	4.3E-09	4.0E-09	3.8E-09	3.8E-09
Fluoranthene	206-44-0	5.3E-09	3.6E-07	1.7E-07	1.2E-07	9.5E-09	5.4E-09	5.1E-09	4.8E-09	4.9E-09
Fluorene	86-73-7	6.8E-11	4.8E-09	2.3E-09	1.6E-09	1.2E-10	7.0E-11	6.7E-11	6.3E-11	6.3E-11
Hexadecane	544-76-3	4.9E-10	3.4E-08	1.6E-08	1.1E-08	8.9E-10	5.1E-10	4.8E-10	4.6E-10	4.6E-10
1-Hydroxypyrene	5315-79-7	2.7E-10	1.9E-08	8.8E-09	6.2E-09	4.9E-10	2.8E-10	2.6E-10	2.5E-10	2.5E-10
Lead		2.3E-07	1.6E-05	7.7E-06	5.4E-06	4.2E-07	2.4E-07	2.3E-07	2.2E-07	2.2E-07
Manganese		2.3E-07	1.6E-05	7.6E-06	5.3E-06	4.2E-07	2.4E-07	2.3E-07	2.1E-07	2.1E-07
2-(Methylthio)benzothiazole	615-22-5	3.5E-09	2.4E-07	1.2E-07	8.1E-08	6.4E-09	3.6E-09	3.4E-09	3.2E-09	3.3E-09
Molybdenum		4.3E-10	3.0E-08	1.4E-08	9.9E-09	7.7E-10	4.4E-10	4.2E-10	3.9E-10	3.9E-10
Naphthalene	91-20-3	1.2E-11	8.5E-10	4.1E-10	2.8E-10	2.2E-11	1.3E-11	1.2E-11	1.1E-11	1.1E-11
Naphthalene, 1-methyl-	90-12-0	3.2E-11	2.2E-09	1.0E-09	7.3E-10	5.7E-11	3.3E-11	3.1E-11	2.9E-11	2.9E-11
Naphthalene, 1,2-dimethyl-	573-98-8	7.4E-12	5.1E-10	2.4E-10	1.7E-10	1.3E-11	7.6E-12	7.2E-12	6.8E-12	6.8E-12
Naphthalene, 1,6-dimethyl-	575-43-9	4.9E-11	3.4E-09	1.6E-09	1.1E-09	8.8E-11	5.0E-11	4.8E-11	4.5E-11	4.5E-11
Naphthalene, 2-(bromomethyl)-	939-26-4	1.3E-09	8.9E-08	4.2E-08	3.0E-08	2.3E-09	1.3E-09	1.3E-09	1.2E-09	1.2E-09
Naphthalene, 2,3-dimethyl-	581-40-8	3.8E-10	2.6E-08	1.2E-08	8.7E-09	6.9E-10	3.9E-10	3.7E-10	3.5E-10	3.5E-10
Naphthalene, 2-methyl	91-57-6	2.9E-11	2.0E-09	9.5E-10	6.6E-10	5.2E-11	3.0E-11	2.8E-11	2.7E-11	2.7E-11
1-Octadecene	112-88-9	1.1E-09	7.6E-08	3.6E-08	2.5E-08	2.0E-09	1.1E-09	1.1E-09	1.0E-09	1.0E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical Name	CASRN	ADD _{ing}								
		Third Trimester Fetus	0<2 Years	2<6 Years	6<11 Years	11<16 Years	16<30 Years	30<40 Years	40<50 Years	50<70 Years
17-Pentatriacontene	6971-40-0	9.3E-10	6.5E-08	3.1E-08	2.1E-08	1.7E-09	9.6E-10	9.0E-10	8.6E-10	8.6E-10
Phenanthrene	85-01-8	2.3E-09	1.6E-07	7.6E-08	5.3E-08	4.2E-09	2.4E-09	2.2E-09	2.1E-09	2.1E-09
Phenanthrene, 1-methyl	832-69-9	6.1E-10	4.3E-08	2.0E-08	1.4E-08	1.1E-09	6.3E-10	6.0E-10	5.6E-10	5.7E-10
Phenanthrene, 2-methyl-	2531-84-2	7.9E-10	5.5E-08	2.6E-08	1.8E-08	1.4E-09	8.2E-10	7.7E-10	7.3E-10	7.3E-10
Phenanthrene, 3-methyl	832-71-3	1.3E-09	9.1E-08	4.3E-08	3.0E-08	2.4E-09	1.3E-09	1.3E-09	1.2E-09	1.2E-09
N-Phenylbenzamide	93-98-1	5.1E-09	3.6E-07	1.7E-07	1.2E-07	9.3E-09	5.3E-09	5.0E-09	4.7E-09	4.7E-09
Phthalimide	85-41-6	9.2E-10	6.4E-08	3.0E-08	2.1E-08	1.7E-09	9.5E-10	9.0E-10	8.5E-10	8.5E-10
Pyrene	129-00-0	1.1E-08	7.7E-07	3.6E-07	2.6E-07	2.0E-08	1.1E-08	1.1E-08	1.0E-08	1.0E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2.4E-11	1.6E-09	7.8E-10	5.5E-10	4.3E-11	2.4E-11	2.3E-11	2.2E-11	2.2E-11
Selenium		1.4E-09	9.7E-08	4.6E-08	3.2E-08	2.5E-09	1.4E-09	1.4E-09	1.3E-09	1.3E-09
Strontium		2.0E-07	1.4E-05	6.5E-06	4.6E-06	3.6E-07	2.0E-07	1.9E-07	1.8E-07	1.8E-07
Thallium		5.6E-11	3.9E-09	1.8E-09	1.3E-09	1.0E-10	5.7E-11	5.4E-11	5.1E-11	5.1E-11
Tin		2.2E-09	1.5E-07	7.2E-08	5.0E-08	3.9E-09	2.2E-09	2.1E-09	2.0E-09	2.0E-09
Triethylene glycol monobutyl ether	143-22-6	2.7E-10	1.9E-08	8.9E-09	6.2E-09	4.9E-10	2.8E-10	2.6E-10	2.5E-10	2.5E-10
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	8.0E-10	5.6E-08	2.6E-08	1.8E-08	1.4E-09	8.2E-10	7.8E-10	7.4E-10	7.4E-10
Vanadium		1.1E-09	7.9E-08	3.7E-08	2.6E-08	2.1E-09	1.2E-09	1.1E-09	1.0E-09	1.0E-09
Zinc		4.9E-05	3.4E-03	1.6E-03	1.1E-03	8.8E-05	5.0E-05	4.7E-05	4.5E-05	4.5E-05

^a ADD_{ing} was calculated from the mean of 35 individual field average concentrations (C_{ing-crumb rubber}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

INDIVIDUAL FIELD ASSESSMENT (Table F-326 to Table F-349)

Table F-326. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Athletes 2<6 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	1.9E-10	1.5E-09	1.4E-09	8.9E-10	4.3E-09	5.4E-09
Aluminum		35	3.3E-05	1.0E-04	5.3E-05	9.4E-05	1.9E-04	2.4E-04
Aniline	62-53-3	12	0	8.2E-09	1.9E-08	0	4.6E-08	8.2E-08
Anthracene	120-12-7	28	0	3.9E-09	6.8E-09	1.5E-09	1.5E-08	3.2E-08
Anthracene, 2-methyl-	613-12-7	35	2.2E-09	1.5E-08	1.3E-08	1.1E-08	3.5E-08	6.8E-08
Anthracene, 9,10-dimethyl	781-43-1	2	0	1.2E-10	6.9E-10	0	8.1E-11	4.1E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	9.6E-10	4.4E-09	0	2.7E-09	2.5E-08
Anthracene, 9-phenyl	602-55-1	30	0	1.5E-09	1.5E-09	8.9E-10	4.6E-09	6.5E-09
Antimony		35	6.8E-09	1.1E-07	9.9E-08	9.5E-08	2.6E-07	4.7E-07
Barium		35	1.8E-06	7.1E-06	4.8E-06	6.9E-06	1.3E-05	2.7E-05
Benzene, n-butyl-	104-51-8	29	0	8.2E-10	2.1E-09	4.6E-10	1.3E-09	1.3E-08
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	2.2E-07	3.1E-07	8.4E-08	8.3E-07	1.1E-06
Benz[a]anthracene	56-55-3	35	8.8E-10	1.3E-08	1.5E-08	6.4E-09	4.2E-08	5.9E-08
Benzo[b]fluoranthene	205-99-2	35	3.8E-09	1.9E-08	1.4E-08	1.2E-08	4.5E-08	5.4E-08
7H-Benzo[c]fluorene	205-12-9	31	0	3.6E-09	5.9E-09	1.2E-09	1.5E-08	2.9E-08
Benzo[k]fluoranthene	207-08-9	35	5.1E-10	6.0E-09	5.7E-09	3.8E-09	1.8E-08	2.3E-08
Benzothiazole	95-16-9	35	5.2E-07	2.3E-06	1.6E-06	1.7E-06	4.7E-06	5.4E-06
Benzothiazole, 2-phenyl-	883-93-2	35	7.0E-08	2.4E-07	2.3E-07	1.8E-07	5.6E-07	1.3E-06
Benzothiazolone	934-34-9	35	3.7E-06	5.8E-06	1.1E-06	5.7E-06	7.4E-06	7.7E-06
Benzyl butyl phthalate	85-68-7	35	6.0E-09	1.2E-07	9.7E-08	1.0E-07	2.6E-07	4.6E-07
Beryllium		11	0	2.9E-09	4.7E-09	0	1.0E-08	1.6E-08
Butylated Hydroxytoluene	128-37-0	19	0	3.1E-09	5.8E-09	1.1E-09	1.6E-08	2.6E-08
Cadmium		34	0	7.9E-08	4.9E-08	7.1E-08	1.7E-07	1.8E-07
Chromium		35	4.1E-08	2.2E-07	2.2E-07	1.5E-07	4.8E-07	1.2E-06
Cobalt		35	9.4E-07	5.0E-06	3.4E-06	4.8E-06	9.6E-06	1.7E-05
Copper		35	3.9E-06	9.8E-06	5.1E-06	9.0E-06	1.6E-05	3.1E-05
Cyclohexyl isothiocyanate	1122-82-3	29	0	7.3E-07	5.3E-07	7.2E-07	1.7E-06	1.9E-06
Dibenz[a,h]anthracene	53-70-3	7	0	8.0E-10	2.0E-09	0	3.6E-09	9.9E-09
Dibenzothiophene	132-65-0	27	0	3.1E-09	4.7E-09	9.1E-10	1.3E-08	1.9E-08
Dibutyl phthalate	84-74-2	31	0	2.2E-07	1.7E-07	1.8E-07	5.4E-07	6.3E-07
Diethyl Phthalate	84-66-2	23	0	1.9E-08	1.6E-08	2.0E-08	4.4E-08	5.8E-08
Diisobutyl Phthalate	84-69-5	32	0	2.2E-08	5.8E-08	5.0E-09	5.9E-08	3.4E-07
Diisooctylphthalate	27554-26-3	33	0	4.7E-07	3.5E-07	3.5E-07	1.2E-06	1.2E-06
Di-n-octyl phthalate	117-84-0	26	0	4.6E-08	1.1E-07	1.2E-08	2.3E-07	4.9E-07
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	3.8E-10	1.2E-09	0	3.0E-09	5.4E-09
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	2.3E-10	8.2E-08	1.7E-07	5.1E-09	3.4E-07	8.2E-07
Fluoranthene	206-44-0	35	1.4E-08	1.0E-07	1.0E-07	6.2E-08	2.6E-07	5.0E-07
Fluorene	86-73-7	11	0	1.4E-09	3.2E-09	0	6.7E-09	1.6E-08
Hexadecane	544-76-3	28	0	9.8E-09	8.2E-09	1.0E-08	2.3E-08	3.8E-08
1-Hydroxypyrene	5315-79-7	1	0	5.3E-09	3.1E-08	0	0	1.9E-07



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Lead		35	9.0E-07	4.6E-06	4.0E-06	3.5E-06	1.2E-05	2.1E-05
Manganese		35	8.4E-07	4.6E-06	2.7E-06	4.8E-06	7.5E-06	1.5E-05
2-(Methylthio)benzothiazole	615-22-5	5	0	7.0E-08	2.5E-07	0	3.9E-07	1.4E-06
Molybdenum		14	0	8.5E-09	1.2E-08	0	2.6E-08	4.2E-08
Naphthalene	91-20-3	1	0	2.4E-10	1.4E-09	0	0	8.5E-09
Naphthalene, 1-methyl-	90-12-0	31	0	6.3E-10	2.2E-09	1.1E-10	1.3E-09	1.3E-08
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	1.5E-10	3.3E-10	0	7.4E-10	1.7E-09
Naphthalene, 1,6-dimethyl-	575-43-9	35	3.9E-10	9.7E-10	4.6E-10	8.5E-10	2.0E-09	2.7E-09
Naphthalene, 2-(bromomethyl)-	939-26-4	35	9.1E-10	2.6E-08	1.2E-08	2.5E-08	4.5E-08	5.8E-08
Naphthalene, 2,3-dimethyl-	581-40-8	35	1.9E-10	7.5E-09	7.4E-09	5.2E-09	2.2E-08	3.3E-08
Naphthalene, 2-methyl	91-57-6	19	0	5.7E-10	2.4E-09	1.0E-10	1.3E-09	1.4E-08
1-Octadecene	112-88-9	32	0	2.2E-08	1.9E-08	1.7E-08	5.9E-08	6.7E-08
17-Pentatriacontene	6971-40-0	9	0	1.8E-08	4.3E-08	0	1.0E-07	2.0E-07
Phenanthrene	85-01-8	35	7.9E-09	4.6E-08	5.7E-08	2.6E-08	1.3E-07	2.9E-07
Phenanthrene, 1-methyl	832-69-9	35	1.1E-09	1.2E-08	1.2E-08	8.6E-09	3.0E-08	5.3E-08
Phenanthrene, 2-methyl-	2531-84-2	35	3.3E-10	1.6E-08	1.8E-08	9.4E-09	4.4E-08	8.6E-08
Phenanthrene, 3-methyl	832-71-3	35	2.0E-09	2.6E-08	2.8E-08	1.7E-08	7.1E-08	1.3E-07
N-Phenylbenzamide	93-98-1	22	0	1.0E-07	2.0E-07	1.6E-08	6.2E-07	7.2E-07
Phthalimide	85-41-6	16	0	1.8E-08	4.2E-08	0	9.2E-08	1.9E-07
Pyrene	129-00-0	35	4.1E-08	2.2E-07	1.4E-07	1.8E-07	4.6E-07	6.3E-07
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	4.7E-10	2.7E-09	0	1.6E-10	1.6E-08
Selenium		4	0	2.8E-08	8.5E-08	0	1.9E-07	4.1E-07
Strontium		35	3.8E-07	3.9E-06	7.6E-06	2.1E-06	1.2E-05	4.5E-05
Thallium		21	0	1.1E-09	1.9E-09	8.9E-10	3.4E-09	1.0E-08
Tin		30	0	4.3E-08	4.3E-08	3.2E-08	1.2E-07	1.7E-07
Triethylene glycol monobutyl ether	143-22-6	3	0	5.4E-09	2.2E-08	0	2.8E-08	1.2E-07
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	9.1E-09	1.6E-08	5.1E-09	1.5E-08	2.4E-08	3.0E-08
Vanadium		2	0	2.2E-08	9.3E-08	0	1.2E-07	4.0E-07
Zinc		35	2.1E-04	9.7E-04	5.1E-04	9.3E-04	1.7E-03	2.8E-03

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not



included as the Study did not speciate the chemicals in the sample analysis.
Values are rounded to two significant figures.

Table F-327. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Athletes 6<11 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	1.3E-10	9.8E-10	9.1E-10	5.9E-10	2.9E-09	3.6E-09
Aluminum		35	2.2E-05	6.9E-05	3.5E-05	6.3E-05	1.3E-04	1.6E-04
Aniline	62-53-3	12	0	5.5E-09	1.2E-08	0	3.0E-08	5.5E-08
Anthracene	120-12-7	28	0	2.6E-09	4.5E-09	1.0E-09	9.7E-09	2.1E-08
Anthracene, 2-methyl-	613-12-7	35	1.5E-09	1.0E-08	9.0E-09	7.4E-09	2.4E-08	4.6E-08
Anthracene, 9,10-dimethyl	781-43-1	2	0	8.3E-11	4.6E-10	0	5.4E-11	2.7E-09
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	6.4E-10	2.9E-09	0	1.8E-09	1.6E-08
Anthracene, 9-phenyl	602-55-1	30	0	9.7E-10	1.0E-09	5.9E-10	3.0E-09	4.4E-09
Antimony		35	4.5E-09	7.5E-08	6.6E-08	6.3E-08	1.7E-07	3.1E-07
Barium		35	1.2E-06	4.7E-06	3.2E-06	4.6E-06	9.0E-06	1.8E-05
Benzene, n-butyl-	104-51-8	29	0	5.5E-10	1.4E-09	3.1E-10	8.4E-10	8.5E-09
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	1.5E-07	2.0E-07	5.6E-08	5.5E-07	7.3E-07
Benz[a]anthracene	56-55-3	35	5.9E-10	9.0E-09	9.9E-09	4.3E-09	2.8E-08	3.9E-08
Benzo[b]fluoranthene	205-99-2	35	2.5E-09	1.2E-08	9.1E-09	8.3E-09	3.0E-08	3.6E-08
7H-Benzo[c]fluorene	205-12-9	31	0	2.4E-09	3.9E-09	7.8E-10	9.7E-09	1.9E-08
Benzo[k]fluoranthene	207-08-9	35	3.4E-10	4.0E-09	3.8E-09	2.5E-09	1.2E-08	1.5E-08
Benzothiazole	95-16-9	35	3.5E-07	1.5E-06	1.0E-06	1.1E-06	3.1E-06	3.6E-06
Benzothiazole, 2-phenyl-	883-93-2	35	4.7E-08	1.6E-07	1.5E-07	1.2E-07	3.8E-07	8.7E-07
Benzothiazolone	934-34-9	35	2.4E-06	3.9E-06	7.5E-07	3.8E-06	5.0E-06	5.1E-06
Benzyl butyl phthalate	85-68-7	35	4.0E-09	7.8E-08	6.4E-08	6.9E-08	1.7E-07	3.1E-07
Beryllium		11	0	2.0E-09	3.2E-09	0	6.8E-09	1.1E-08
Butylated Hydroxytoluene	128-37-0	19	0	2.1E-09	3.9E-09	7.2E-10	1.0E-08	1.7E-08
Cadmium		34	0	5.3E-08	3.3E-08	4.8E-08	1.1E-07	1.2E-07
Chromium		35	2.7E-08	1.5E-07	1.4E-07	9.7E-08	3.2E-07	8.2E-07
Cobalt		35	6.2E-07	3.4E-06	2.3E-06	3.2E-06	6.4E-06	1.2E-05
Copper		35	2.6E-06	6.6E-06	3.4E-06	6.0E-06	1.1E-05	2.1E-05
Cyclohexyl isothiocyanate	1122-82-3	29	0	4.9E-07	3.6E-07	4.8E-07	1.1E-06	1.3E-06
Dibenz[a,h]anthracene	53-70-3	7	0	5.3E-10	1.3E-09	0	2.4E-09	6.6E-09
Dibenzothiophene	132-65-0	27	0	2.1E-09	3.1E-09	6.1E-10	8.4E-09	1.3E-08
Dibutyl phthalate	84-74-2	31	0	1.5E-07	1.1E-07	1.2E-07	3.6E-07	4.2E-07



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Diethyl Phthalate	84-66-2	23	0	1.2E-08	1.1E-08	1.4E-08	2.9E-08	3.9E-08
Diisobutyl Phthalate	84-69-5	32	0	1.5E-08	3.8E-08	3.4E-09	4.0E-08	2.3E-07
Diisooctylphthalate	27554-26-3	33	0	3.1E-07	2.3E-07	2.3E-07	7.9E-07	8.3E-07
Di-n-octyl phthalate	117-84-0	26	0	3.1E-08	7.5E-08	8.1E-09	1.5E-07	3.3E-07
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	2.6E-10	8.1E-10	0	2.0E-09	3.6E-09
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	1.5E-10	5.5E-08	1.1E-07	3.4E-09	2.3E-07	5.5E-07
Fluoranthene	206-44-0	35	9.5E-09	7.0E-08	6.8E-08	4.2E-08	1.8E-07	3.3E-07
Fluorene	86-73-7	11	0	9.1E-10	2.2E-09	0	4.5E-09	1.1E-08
Hexadecane	544-76-3	28	0	6.5E-09	5.5E-09	6.9E-09	1.6E-08	2.5E-08
1-Hydroxypyrene	5315-79-7	1	0	3.5E-09	2.1E-08	0	0	1.2E-07
Lead		35	6.0E-07	3.1E-06	2.7E-06	2.3E-06	7.8E-06	1.4E-05
Manganese		35	5.6E-07	3.1E-06	1.8E-06	3.2E-06	5.0E-06	9.8E-06
2-(Methylthio)benzothiazole	615-22-5	5	0	4.7E-08	1.7E-07	0	2.6E-07	9.4E-07
Molybdenum		14	0	5.6E-09	7.9E-09	0	1.7E-08	2.8E-08
Naphthalene	91-20-3	1	0	1.6E-10	9.6E-10	0	0	5.7E-09
Naphthalene, 1-methyl-	90-12-0	31	0	4.2E-10	1.5E-09	7.0E-11	8.8E-10	8.7E-09
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	9.8E-11	2.2E-10	0	4.9E-10	1.1E-09
Naphthalene, 1,6-dimethyl-	575-43-9	35	2.6E-10	6.5E-10	3.1E-10	5.7E-10	1.3E-09	1.8E-09
Naphthalene, 2-(bromomethyl)-	939-26-4	35	6.0E-10	1.7E-08	8.0E-09	1.7E-08	3.0E-08	3.9E-08
Naphthalene, 2,3-dimethyl-	581-40-8	35	1.3E-10	5.0E-09	5.0E-09	3.5E-09	1.5E-08	2.2E-08
Naphthalene, 2-methyl	91-57-6	19	0	3.8E-10	1.6E-09	6.8E-11	8.6E-10	9.4E-09
1-Octadecene	112-88-9	32	0	1.4E-08	1.3E-08	1.1E-08	3.9E-08	4.5E-08
17-Pentatriacontene	6971-40-0	9	0	1.2E-08	2.9E-08	0	6.7E-08	1.3E-07
Phenanthrene	85-01-8	35	5.3E-09	3.1E-08	3.8E-08	1.7E-08	8.7E-08	1.9E-07
Phenanthrene, 1-methyl	832-69-9	35	7.0E-10	8.1E-09	7.8E-09	5.8E-09	2.0E-08	3.5E-08
Phenanthrene, 2-methyl-	2531-84-2	35	2.2E-10	1.0E-08	1.2E-08	6.2E-09	3.0E-08	5.7E-08
Phenanthrene, 3-methyl	832-71-3	35	1.3E-09	1.7E-08	1.9E-08	1.1E-08	4.8E-08	8.7E-08
N-Phenylbenzamide	93-98-1	22	0	6.8E-08	1.3E-07	1.0E-08	4.1E-07	4.8E-07
Phthalimide	85-41-6	16	0	1.2E-08	2.8E-08	0	6.2E-08	1.3E-07
Pyrene	129-00-0	35	2.7E-08	1.5E-07	9.5E-08	1.2E-07	3.1E-07	4.2E-07
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	3.1E-10	1.8E-09	0	1.1E-10	1.1E-08
Selenium		4	0	1.9E-08	5.7E-08	0	1.3E-07	2.7E-07
Strontium		35	2.5E-07	2.6E-06	5.1E-06	1.4E-06	7.8E-06	3.0E-05



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Thallium		21	0	7.4E-10	1.3E-09	5.9E-10	2.3E-09	6.7E-09
Tin		30	0	2.9E-08	2.9E-08	2.1E-08	7.7E-08	1.1E-07
Triethylene glycol monobutyl ether	143-22-6	3	0	3.6E-09	1.5E-08	0	1.9E-08	8.0E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	6.1E-09	1.1E-08	3.4E-09	9.8E-09	1.6E-08	2.0E-08
Vanadium		2	0	1.5E-08	6.2E-08	0	7.8E-08	2.6E-07
Zinc		35	1.4E-04	6.5E-04	3.4E-04	6.2E-04	1.2E-03	1.9E-03

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-328. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Athletes 11<16 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	9.9E-11	7.6E-10	7.1E-10	4.6E-10	2.2E-09	2.8E-09
Aluminum		35	1.7E-05	5.3E-05	2.7E-05	4.9E-05	9.7E-05	1.3E-04
Aniline	62-53-3	12	0	4.2E-09	9.7E-09	0	2.4E-08	4.3E-08
Anthracene	120-12-7	28	0	2.0E-09	3.5E-09	7.7E-10	7.5E-09	1.6E-08
Anthracene, 2-methyl-	613-12-7	35	1.2E-09	8.0E-09	7.0E-09	5.7E-09	1.8E-08	3.5E-08
Anthracene, 9,10-dimethyl	781-43-1	2	0	6.4E-11	3.6E-10	0	4.2E-11	2.1E-09
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	5.0E-10	2.3E-09	0	1.4E-09	1.3E-08
Anthracene, 9-phenyl	602-55-1	30	0	7.5E-10	7.9E-10	4.6E-10	2.4E-09	3.4E-09
Antimony		35	3.5E-09	5.8E-08	5.1E-08	4.9E-08	1.3E-07	2.4E-07
Barium		35	9.6E-07	3.7E-06	2.5E-06	3.6E-06	7.0E-06	1.4E-05
Benzene, n-butyl-	104-51-8	29	0	4.2E-10	1.1E-09	2.4E-10	6.5E-10	6.6E-09
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	1.2E-07	1.6E-07	4.4E-08	4.3E-07	5.6E-07
Benz[a]anthracene	56-55-3	35	4.5E-10	7.0E-09	7.6E-09	3.3E-09	2.2E-08	3.0E-08
Benzo[b]fluoranthene	205-99-2	35	2.0E-09	9.6E-09	7.1E-09	6.4E-09	2.3E-08	2.8E-08
7H-Benzo[c]fluorene	205-12-9	31	0	1.8E-09	3.0E-09	6.1E-10	7.5E-09	1.5E-08
Benzo[k]fluoranthene	207-08-9	35	2.6E-10	3.1E-09	2.9E-09	2.0E-09	9.1E-09	1.2E-08
Benzothiazole	95-16-9	35	2.7E-07	1.2E-06	8.1E-07	8.8E-07	2.4E-06	2.8E-06



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzothiazole, 2-phenyl-	883-93-2	35	3.6E-08	1.3E-07	1.2E-07	9.2E-08	2.9E-07	6.8E-07
Benzothiazolone	934-34-9	35	1.9E-06	3.0E-06	5.8E-07	3.0E-06	3.9E-06	4.0E-06
Benzyl butyl phthalate	85-68-7	35	3.1E-09	6.0E-08	5.0E-08	5.3E-08	1.3E-07	2.4E-07
Beryllium		11	0	1.5E-09	2.4E-09	0	5.3E-09	8.4E-09
Butylated Hydroxytoluene	128-37-0	19	0	1.6E-09	3.0E-09	5.6E-10	8.1E-09	1.4E-08
Cadmium		34	0	4.1E-08	2.5E-08	3.7E-08	8.7E-08	9.5E-08
Chromium		35	2.1E-08	1.1E-07	1.1E-07	7.5E-08	2.5E-07	6.3E-07
Cobalt		35	4.8E-07	2.6E-06	1.8E-06	2.5E-06	5.0E-06	9.0E-06
Copper		35	2.0E-06	5.1E-06	2.6E-06	4.7E-06	8.4E-06	1.6E-05
Cyclohexyl isothiocyanate	1122-82-3	29	0	3.8E-07	2.8E-07	3.7E-07	8.8E-07	9.8E-07
Dibenz[a,h]anthracene	53-70-3	7	0	4.1E-10	1.0E-09	0	1.9E-09	5.1E-09
Dibenzothiophene	132-65-0	27	0	1.6E-09	2.4E-09	4.7E-10	6.5E-09	9.8E-09
Dibutyl phthalate	84-74-2	31	0	1.1E-07	8.7E-08	9.2E-08	2.8E-07	3.3E-07
Diethyl Phthalate	84-66-2	23	0	9.6E-09	8.5E-09	1.0E-08	2.3E-08	3.0E-08
Diisobutyl Phthalate	84-69-5	32	0	1.1E-08	3.0E-08	2.6E-09	3.1E-08	1.8E-07
Diisooctylphthalate	27554-26-3	33	0	2.4E-07	1.8E-07	1.8E-07	6.1E-07	6.4E-07
Di-n-octyl phthalate	117-84-0	26	0	2.4E-08	5.8E-08	6.3E-09	1.2E-07	2.5E-07
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	2.0E-10	6.3E-10	0	1.5E-09	2.8E-09
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	1.2E-10	4.2E-08	8.5E-08	2.6E-09	1.7E-07	4.2E-07
Fluoranthene	206-44-0	35	7.3E-09	5.4E-08	5.2E-08	3.2E-08	1.4E-07	2.6E-07
Fluorene	86-73-7	11	0	7.0E-10	1.7E-09	0	3.5E-09	8.4E-09
Hexadecane	544-76-3	28	0	5.1E-09	4.3E-09	5.4E-09	1.2E-08	2.0E-08
1-Hydroxypyrene	5315-79-7	1	0	2.8E-09	1.6E-08	0	0	9.6E-08
Lead		35	4.6E-07	2.4E-06	2.1E-06	1.8E-06	6.0E-06	1.1E-05
Manganese		35	4.3E-07	2.4E-06	1.4E-06	2.5E-06	3.9E-06	7.6E-06
2-(Methylthio)benzothiazole	615-22-5	5	0	3.6E-08	1.3E-07	0	2.0E-07	7.3E-07
Molybdenum		14	0	4.4E-09	6.1E-09	0	1.4E-08	2.2E-08
Naphthalene	91-20-3	1	0	1.3E-10	7.5E-10	0	0	4.4E-09
Naphthalene, 1-methyl-	90-12-0	31	0	3.3E-10	1.1E-09	5.4E-11	6.8E-10	6.7E-09
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	7.6E-11	1.7E-10	0	3.8E-10	8.7E-10
Naphthalene, 1,6-dimethyl-	575-43-9	35	2.0E-10	5.0E-10	2.4E-10	4.4E-10	1.0E-09	1.4E-09
Naphthalene, 2-(bromomethyl)-	939-26-4	35	4.7E-10	1.3E-08	6.2E-09	1.3E-08	2.3E-08	3.0E-08
Naphthalene, 2,3-dimethyl-	581-40-8	35	9.9E-11	3.9E-09	3.8E-09	2.7E-09	1.2E-08	1.7E-08



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene, 2-methyl	91-57-6	19	0	3.0E-10	1.2E-09	5.3E-11	6.6E-10	7.3E-09
1-Octadecene	112-88-9	32	0	1.1E-08	9.8E-09	8.8E-09	3.0E-08	3.5E-08
17-Pentatriacontene	6971-40-0	9	0	9.5E-09	2.2E-08	0	5.2E-08	1.0E-07
Phenanthrene	85-01-8	35	4.1E-09	2.4E-08	2.9E-08	1.3E-08	6.8E-08	1.5E-07
Phenanthrene, 1-methyl	832-69-9	35	5.4E-10	6.3E-09	6.0E-09	4.5E-09	1.6E-08	2.7E-08
Phenanthrene, 2-methyl-	2531-84-2	35	1.7E-10	8.1E-09	9.4E-09	4.8E-09	2.3E-08	4.5E-08
Phenanthrene, 3-methyl	832-71-3	35	1.0E-09	1.3E-08	1.4E-08	8.6E-09	3.7E-08	6.8E-08
N-Phenylbenzamide	93-98-1	22	0	5.3E-08	1.0E-07	8.1E-09	3.2E-07	3.7E-07
Phthalimide	85-41-6	16	0	9.5E-09	2.2E-08	0	4.8E-08	1.0E-07
Pyrene	129-00-0	35	2.1E-08	1.1E-07	7.4E-08	9.3E-08	2.4E-07	3.3E-07
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	2.4E-10	1.4E-09	0	8.3E-11	8.2E-09
Selenium		4	0	1.4E-08	4.4E-08	0	1.0E-07	2.1E-07
Strontium		35	1.9E-07	2.0E-06	3.9E-06	1.1E-06	6.1E-06	2.3E-05
Thallium		21	0	5.7E-10	9.7E-10	4.6E-10	1.8E-09	5.2E-09
Tin		30	0	2.2E-08	2.2E-08	1.7E-08	6.0E-08	8.8E-08
Triethylene glycol monobutyl ether	143-22-6	3	0	2.8E-09	1.1E-08	0	1.4E-08	6.2E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	4.7E-09	8.2E-09	2.6E-09	7.6E-09	1.3E-08	1.6E-08
Vanadium		2	0	1.2E-08	4.8E-08	0	6.1E-08	2.1E-07
Zinc		35	1.1E-04	5.0E-04	2.7E-04	4.8E-04	8.9E-04	1.4E-03

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-329. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Athletes 16<30 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	1.1E-10	8.5E-10	7.9E-10	5.1E-10	2.5E-09	3.1E-09
Aluminum		35	1.9E-05	5.9E-05	3.0E-05	5.4E-05	1.1E-04	1.4E-04
Aniline	62-53-3	12	0	4.7E-09	1.1E-08	0	2.6E-08	4.7E-08
Anthracene	120-12-7	28	0	2.2E-09	3.9E-09	8.6E-10	8.4E-09	1.8E-08
Anthracene, 2-methyl-	613-12-7	35	1.3E-09	8.9E-09	7.8E-09	6.4E-09	2.0E-08	3.9E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Anthracene, 9,10-dimethyl	781-43-1	2	0	7.2E-11	4.0E-10	0	4.6E-11	2.4E-09
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	5.5E-10	2.5E-09	0	1.6E-09	1.4E-08
Anthracene, 9-phenyl	602-55-1	30	0	8.4E-10	8.8E-10	5.1E-10	2.6E-09	3.8E-09
Antimony		35	3.9E-09	6.4E-08	5.7E-08	5.4E-08	1.5E-07	2.7E-07
Barium		35	1.1E-06	4.1E-06	2.8E-06	4.0E-06	7.7E-06	1.5E-05
Benzene, n-butyl-	104-51-8	29	0	4.7E-10	1.2E-09	2.7E-10	7.3E-10	7.3E-09
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	1.3E-07	1.8E-07	4.8E-08	4.8E-07	6.3E-07
Benz[a]anthracene	56-55-3	35	5.1E-10	7.8E-09	8.5E-09	3.7E-09	2.4E-08	3.4E-08
Benzo[b]fluoranthene	205-99-2	35	2.2E-09	1.1E-08	7.9E-09	7.2E-09	2.6E-08	3.1E-08
7H-Benzo[c]fluorene	205-12-9	31	0	2.1E-09	3.4E-09	6.7E-10	8.4E-09	1.7E-08
Benzo[k]fluoranthene	207-08-9	35	2.9E-10	3.5E-09	3.3E-09	2.2E-09	1.0E-08	1.3E-08
Benzothiazole	95-16-9	35	3.0E-07	1.3E-06	9.0E-07	9.8E-07	2.7E-06	3.1E-06
Benzothiazole, 2-phenyl-	883-93-2	35	4.0E-08	1.4E-07	1.3E-07	1.0E-07	3.2E-07	7.5E-07
Benzothiazolone	934-34-9	35	2.1E-06	3.3E-06	6.5E-07	3.3E-06	4.3E-06	4.4E-06
Benzyl butyl phthalate	85-68-7	35	3.4E-09	6.7E-08	5.6E-08	5.9E-08	1.5E-07	2.7E-07
Beryllium		11	0	1.7E-09	2.7E-09	0	5.9E-09	9.3E-09
Butylated Hydroxytoluene	128-37-0	19	0	1.8E-09	3.3E-09	6.2E-10	9.0E-09	1.5E-08
Cadmium		34	0	4.6E-08	2.8E-08	4.1E-08	9.6E-08	1.1E-07
Chromium		35	2.3E-08	1.3E-07	1.2E-07	8.4E-08	2.8E-07	7.0E-07
Cobalt		35	5.4E-07	2.9E-06	2.0E-06	2.8E-06	5.5E-06	1.0E-05
Copper		35	2.2E-06	5.7E-06	2.9E-06	5.2E-06	9.3E-06	1.8E-05
Cyclohexyl isothiocyanate	1122-82-3	29	0	4.2E-07	3.1E-07	4.2E-07	9.8E-07	1.1E-06
Dibenz[a,h]anthracene	53-70-3	7	0	4.6E-10	1.1E-09	0	2.1E-09	5.7E-09
Dibenzothiophene	132-65-0	27	0	1.8E-09	2.7E-09	5.2E-10	7.2E-09	1.1E-08
Dibutyl phthalate	84-74-2	31	0	1.3E-07	9.7E-08	1.0E-07	3.1E-07	3.7E-07
Diethyl Phthalate	84-66-2	23	0	1.1E-08	9.4E-09	1.2E-08	2.5E-08	3.3E-08
Diisobutyl Phthalate	84-69-5	32	0	1.3E-08	3.3E-08	2.9E-09	3.4E-08	2.0E-07
Diisooctylphthalate	27554-26-3	33	0	2.7E-07	2.0E-07	2.0E-07	6.8E-07	7.1E-07
Di-n-octyl phthalate	117-84-0	26	0	2.7E-08	6.4E-08	7.0E-09	1.3E-07	2.8E-07
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	2.2E-10	7.0E-10	0	1.7E-09	3.1E-09
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	1.3E-10	4.7E-08	9.5E-08	2.9E-09	1.9E-07	4.7E-07
Fluoranthene	206-44-0	35	8.2E-09	6.0E-08	5.8E-08	3.6E-08	1.5E-07	2.9E-07
Fluorene	86-73-7	11	0	7.8E-10	1.9E-09	0	3.9E-09	9.3E-09
Hexadecane	544-76-3	28	0	5.7E-09	4.7E-09	6.0E-09	1.3E-08	2.2E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
1-Hydroxypyrene	5315-79-7	1	0	3.1E-09	1.8E-08	0	0	1.1E-07
Lead		35	5.2E-07	2.7E-06	2.3E-06	2.0E-06	6.7E-06	1.2E-05
Manganese		35	4.8E-07	2.6E-06	1.6E-06	2.7E-06	4.3E-06	8.4E-06
2-(Methylthio)benzothiazole	615-22-5	5	0	4.0E-08	1.5E-07	0	2.3E-07	8.1E-07
Molybdenum		14	0	4.9E-09	6.8E-09	0	1.5E-08	2.4E-08
Naphthalene	91-20-3	1	0	1.4E-10	8.3E-10	0	0	4.9E-09
Naphthalene, 1-methyl-	90-12-0	31	0	3.6E-10	1.3E-09	6.0E-11	7.6E-10	7.5E-09
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	8.5E-11	1.9E-10	0	4.2E-10	9.7E-10
Naphthalene, 1,6-dimethyl-	575-43-9	35	2.2E-10	5.6E-10	2.7E-10	4.9E-10	1.1E-09	1.6E-09
Naphthalene, 2-(bromomethyl)-	939-26-4	35	5.2E-10	1.5E-08	6.9E-09	1.4E-08	2.6E-08	3.3E-08
Naphthalene, 2,3-dimethyl-	581-40-8	35	1.1E-10	4.3E-09	4.3E-09	3.0E-09	1.3E-08	1.9E-08
Naphthalene, 2-methyl	91-57-6	19	0	3.3E-10	1.4E-09	5.9E-11	7.4E-10	8.2E-09
1-Octadecene	112-88-9	32	0	1.2E-08	1.1E-08	9.8E-09	3.4E-08	3.9E-08
17-Pentatriacontene	6971-40-0	9	0	1.1E-08	2.5E-08	0	5.8E-08	1.1E-07
Phenanthrene	85-01-8	35	4.6E-09	2.6E-08	3.3E-08	1.5E-08	7.5E-08	1.7E-07
Phenanthrene, 1-methyl	832-69-9	35	6.0E-10	7.0E-09	6.7E-09	5.0E-09	1.7E-08	3.1E-08
Phenanthrene, 2-methyl-	2531-84-2	35	1.9E-10	9.1E-09	1.0E-08	5.4E-09	2.6E-08	5.0E-08
Phenanthrene, 3-methyl	832-71-3	35	1.1E-09	1.5E-08	1.6E-08	9.5E-09	4.1E-08	7.5E-08
N-Phenylbenzamide	93-98-1	22	0	5.8E-08	1.1E-07	9.0E-09	3.6E-07	4.1E-07
Phthalimide	85-41-6	16	0	1.1E-08	2.4E-08	0	5.3E-08	1.1E-07
Pyrene	129-00-0	35	2.3E-08	1.3E-07	8.2E-08	1.0E-07	2.7E-07	3.6E-07
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	2.7E-10	1.5E-09	0	9.2E-11	9.2E-09
Selenium		4	0	1.6E-08	4.9E-08	0	1.1E-07	2.4E-07
Strontium		35	2.2E-07	2.3E-06	4.4E-06	1.2E-06	6.7E-06	2.6E-05
Thallium		21	0	6.4E-10	1.1E-09	5.1E-10	2.0E-09	5.8E-09
Tin		30	0	2.5E-08	2.5E-08	1.9E-08	6.6E-08	9.8E-08
Triethylene glycol monobutyl ether	143-22-6	3	0	3.1E-09	1.3E-08	0	1.6E-08	6.9E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	5.3E-09	9.1E-09	2.9E-09	8.5E-09	1.4E-08	1.7E-08
Vanadium		2	0	1.3E-08	5.3E-08	0	6.7E-08	2.3E-07
Zinc		35	1.2E-04	5.6E-04	3.0E-04	5.3E-04	1.0E-03	1.6E-03

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.



° CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.
Values are rounded to two significant figures.

Table F-330. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Athletes 30<40 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	7.3E-11	5.6E-10	5.3E-10	3.4E-10	1.7E-09	2.1E-09
Aluminum		35	1.3E-05	4.0E-05	2.0E-05	3.6E-05	7.2E-05	9.3E-05
Aniline	62-53-3	12	0	3.1E-09	7.2E-09	0	1.8E-08	3.2E-08
Anthracene	120-12-7	28	0	1.5E-09	2.6E-09	5.7E-10	5.6E-09	1.2E-08
Anthracene, 2-methyl-	613-12-7	35	8.6E-10	5.9E-09	5.2E-09	4.3E-09	1.4E-08	2.6E-08
Anthracene, 9,10-dimethyl	781-43-1	2	0	4.8E-11	2.7E-10	0	3.1E-11	1.6E-09
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	3.7E-10	1.7E-09	0	1.0E-09	9.4E-09
Anthracene, 9-phenyl	602-55-1	30	0	5.6E-10	5.9E-10	3.4E-10	1.8E-09	2.5E-09
Antimony		35	2.6E-09	4.3E-08	3.8E-08	3.6E-08	9.9E-08	1.8E-07
Barium		35	7.1E-07	2.7E-06	1.9E-06	2.7E-06	5.2E-06	1.0E-05
Benzene, n-butyl-	104-51-8	29	0	3.2E-10	8.1E-10	1.8E-10	4.9E-10	4.9E-09
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	8.6E-08	1.2E-07	3.2E-08	3.2E-07	4.2E-07
Benz[a]anthracene	56-55-3	35	3.4E-10	5.2E-09	5.7E-09	2.5E-09	1.6E-08	2.3E-08
Benzo[b]fluoranthene	205-99-2	35	1.5E-09	7.1E-09	5.3E-09	4.8E-09	1.7E-08	2.1E-08
7H-Benzo[c]fluorene	205-12-9	31	0	1.4E-09	2.2E-09	4.5E-10	5.6E-09	1.1E-08
Benzo[k]fluoranthene	207-08-9	35	1.9E-10	2.3E-09	2.2E-09	1.5E-09	6.8E-09	8.7E-09
Benzothiazole	95-16-9	35	2.0E-07	8.8E-07	6.0E-07	6.5E-07	1.8E-06	2.1E-06
Benzothiazole, 2-phenyl-	883-93-2	35	2.7E-08	9.4E-08	8.8E-08	6.8E-08	2.2E-07	5.0E-07
Benzothiazolone	934-34-9	35	1.4E-06	2.2E-06	4.3E-07	2.2E-06	2.9E-06	3.0E-06
Benzyl butyl phthalate	85-68-7	35	2.3E-09	4.5E-08	3.7E-08	4.0E-08	1.0E-07	1.8E-07
Beryllium		11	0	1.1E-09	1.8E-09	0	3.9E-09	6.2E-09
Butylated Hydroxytoluene	128-37-0	19	0	1.2E-09	2.2E-09	4.2E-10	6.0E-09	1.0E-08
Cadmium		34	0	3.0E-08	1.9E-08	2.7E-08	6.4E-08	7.1E-08
Chromium		35	1.6E-08	8.5E-08	8.3E-08	5.6E-08	1.9E-07	4.7E-07
Cobalt		35	3.6E-07	1.9E-06	1.3E-06	1.8E-06	3.7E-06	6.7E-06
Copper		35	1.5E-06	3.8E-06	2.0E-06	3.5E-06	6.2E-06	1.2E-05
Cyclohexyl isothiocyanate	1122-82-3	29	0	2.8E-07	2.1E-07	2.8E-07	6.6E-07	7.3E-07
Dibenz[a,h]anthracene	53-70-3	7	0	3.1E-10	7.7E-10	0	1.4E-09	3.8E-09
Dibenzothiophene	132-65-0	27	0	1.2E-09	1.8E-09	3.5E-10	4.8E-09	7.3E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dibutyl phthalate	84-74-2	31	0	8.5E-08	6.4E-08	6.8E-08	2.1E-07	2.4E-07
Diethyl Phthalate	84-66-2	23	0	7.2E-09	6.3E-09	7.8E-09	1.7E-08	2.2E-08
Diisobutyl Phthalate	84-69-5	32	0	8.5E-09	2.2E-08	1.9E-09	2.3E-08	1.3E-07
Diisooctylphthalate	27554-26-3	33	0	1.8E-07	1.3E-07	1.3E-07	4.6E-07	4.8E-07
Di-n-octyl phthalate	117-84-0	26	0	1.8E-08	4.3E-08	4.7E-09	8.7E-08	1.9E-07
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.5E-10	4.7E-10	0	1.1E-09	2.1E-09
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	8.7E-11	3.2E-08	6.4E-08	2.0E-09	1.3E-07	3.2E-07
Fluoranthene	206-44-0	35	5.5E-09	4.0E-08	3.9E-08	2.4E-08	1.0E-07	1.9E-07
Fluorene	86-73-7	11	0	5.2E-10	1.2E-09	0	2.6E-09	6.2E-09
Hexadecane	544-76-3	28	0	3.8E-09	3.2E-09	4.0E-09	9.0E-09	1.5E-08
1-Hydroxypyrene	5315-79-7	1	0	2.0E-09	1.2E-08	0	0	7.2E-08
Lead		35	3.4E-07	1.8E-06	1.5E-06	1.3E-06	4.5E-06	8.1E-06
Manganese		35	3.2E-07	1.8E-06	1.0E-06	1.8E-06	2.9E-06	5.6E-06
2-(Methylthio)benzothiazole	615-22-5	5	0	2.7E-08	9.7E-08	0	1.5E-07	5.4E-07
Molybdenum		14	0	3.3E-09	4.5E-09	0	1.0E-08	1.6E-08
Naphthalene	91-20-3	1	0	9.4E-11	5.6E-10	0	0	3.3E-09
Naphthalene, 1-methyl-	90-12-0	31	0	2.4E-10	8.4E-10	4.0E-11	5.1E-10	5.0E-09
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	5.7E-11	1.3E-10	0	2.8E-10	6.5E-10
Naphthalene, 1,6-dimethyl-	575-43-9	35	1.5E-10	3.7E-10	1.8E-10	3.3E-10	7.6E-10	1.1E-09
Naphthalene, 2-(bromomethyl)-	939-26-4	35	3.5E-10	9.8E-09	4.6E-09	9.6E-09	1.7E-08	2.2E-08
Naphthalene, 2,3-dimethyl-	581-40-8	35	7.3E-11	2.9E-09	2.9E-09	2.0E-09	8.6E-09	1.3E-08
Naphthalene, 2-methyl	91-57-6	19	0	2.2E-10	9.2E-10	3.9E-11	4.9E-10	5.4E-09
1-Octadecene	112-88-9	32	0	8.3E-09	7.3E-09	6.5E-09	2.3E-08	2.6E-08
17-Pentatriacontene	6971-40-0	9	0	7.1E-09	1.7E-08	0	3.9E-08	7.5E-08
Phenanthrene	85-01-8	35	3.0E-09	1.8E-08	2.2E-08	9.9E-09	5.0E-08	1.1E-07
Phenanthrene, 1-methyl	832-69-9	35	4.0E-10	4.7E-09	4.5E-09	3.3E-09	1.2E-08	2.0E-08
Phenanthrene, 2-methyl-	2531-84-2	35	1.3E-10	6.0E-09	7.0E-09	3.6E-09	1.7E-08	3.3E-08
Phenanthrene, 3-methyl	832-71-3	35	7.7E-10	1.0E-08	1.1E-08	6.4E-09	2.7E-08	5.0E-08
N-Phenylbenzamide	93-98-1	22	0	3.9E-08	7.6E-08	6.0E-09	2.4E-07	2.8E-07
Phthalimide	85-41-6	16	0	7.1E-09	1.6E-08	0	3.6E-08	7.5E-08
Pyrene	129-00-0	35	1.6E-08	8.4E-08	5.5E-08	6.9E-08	1.8E-07	2.4E-07
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	1.8E-10	1.0E-09	0	6.1E-11	6.1E-09
Selenium		4	0	1.1E-08	3.3E-08	0	7.4E-08	1.6E-07



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Strontium		35	1.4E-07	1.5E-06	2.9E-06	8.0E-07	4.5E-06	1.7E-05
Thallium		21	0	4.2E-10	7.2E-10	3.4E-10	1.3E-09	3.9E-09
Tin		30	0	1.7E-08	1.7E-08	1.2E-08	4.4E-08	6.5E-08
Triethylene glycol monobutyl ether	143-22-6	3	0	2.1E-09	8.4E-09	0	1.1E-08	4.6E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	3.5E-09	6.1E-09	1.9E-09	5.6E-09	9.4E-09	1.2E-08
Vanadium		2	0	8.6E-09	3.6E-08	0	4.5E-08	1.5E-07
Zinc		35	8.2E-05	3.7E-04	2.0E-04	3.6E-04	6.6E-04	1.1E-03

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-331. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Athletes 40<50 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	6.2E-11	4.8E-10	4.5E-10	2.9E-10	1.4E-09	1.8E-09
Aluminum		35	1.1E-05	3.4E-05	1.7E-05	3.1E-05	6.1E-05	7.9E-05
Aniline	62-53-3	12	0	2.7E-09	6.1E-09	0	1.5E-08	2.7E-08
Anthracene	120-12-7	28	0	1.3E-09	2.2E-09	4.9E-10	4.7E-09	1.0E-08
Anthracene, 2-methyl-	613-12-7	35	7.3E-10	5.0E-09	4.4E-09	3.6E-09	1.2E-08	2.2E-08
Anthracene, 9,10-dimethyl	781-43-1	2	0	4.1E-11	2.3E-10	0	2.6E-11	1.3E-09
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	3.1E-10	1.4E-09	0	8.9E-10	8.0E-09
Anthracene, 9-phenyl	602-55-1	30	0	4.7E-10	5.0E-10	2.9E-10	1.5E-09	2.1E-09
Antimony		35	2.2E-09	3.6E-08	3.2E-08	3.1E-08	8.4E-08	1.5E-07
Barium		35	6.0E-07	2.3E-06	1.6E-06	2.3E-06	4.4E-06	8.7E-06
Benzene, n-butyl-	104-51-8	29	0	2.7E-10	6.9E-10	1.5E-10	4.1E-10	4.2E-09
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	7.3E-08	1.0E-07	2.7E-08	2.7E-07	3.6E-07
Benz[a]anthracene	56-55-3	35	2.9E-10	4.4E-09	4.8E-09	2.1E-09	1.4E-08	1.9E-08
Benzo[b]fluoranthene	205-99-2	35	1.2E-09	6.0E-09	4.5E-09	4.1E-09	1.5E-08	1.8E-08
7H-Benzo[c]fluorene	205-12-9	31	0	1.2E-09	1.9E-09	3.8E-10	4.7E-09	9.4E-09
Benzo[k]fluoranthene	207-08-9	35	1.6E-10	2.0E-09	1.8E-09	1.2E-09	5.7E-09	7.3E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzothiazole	95-16-9	35	1.7E-07	7.4E-07	5.1E-07	5.5E-07	1.5E-06	1.8E-06
Benzothiazole, 2-phenyl-	883-93-2	35	2.3E-08	8.0E-08	7.5E-08	5.8E-08	1.8E-07	4.3E-07
Benzothiazolone	934-34-9	35	1.2E-06	1.9E-06	3.7E-07	1.9E-06	2.4E-06	2.5E-06
Benzyl butyl phthalate	85-68-7	35	1.9E-09	3.8E-08	3.1E-08	3.4E-08	8.5E-08	1.5E-07
Beryllium		11	0	9.5E-10	1.5E-09	0	3.3E-09	5.3E-09
Butylated Hydroxytoluene	128-37-0	19	0	1.0E-09	1.9E-09	3.5E-10	5.1E-09	8.5E-09
Cadmium		34	0	2.6E-08	1.6E-08	2.3E-08	5.5E-08	6.0E-08
Chromium		35	1.3E-08	7.2E-08	7.0E-08	4.8E-08	1.6E-07	4.0E-07
Cobalt		35	3.0E-07	1.6E-06	1.1E-06	1.6E-06	3.1E-06	5.7E-06
Copper		35	1.3E-06	3.2E-06	1.7E-06	2.9E-06	5.3E-06	1.0E-05
Cyclohexyl isothiocyanate	1122-82-3	29	0	2.4E-07	1.7E-07	2.4E-07	5.6E-07	6.2E-07
Dibenz[a,h]anthracene	53-70-3	7	0	2.6E-10	6.5E-10	0	1.2E-09	3.2E-09
Dibenzothiophene	132-65-0	27	0	1.0E-09	1.5E-09	3.0E-10	4.1E-09	6.2E-09
Dibutyl phthalate	84-74-2	31	0	7.2E-08	5.5E-08	5.8E-08	1.8E-07	2.1E-07
Diethyl Phthalate	84-66-2	23	0	6.1E-09	5.3E-09	6.6E-09	1.4E-08	1.9E-08
Diisobutyl Phthalate	84-69-5	32	0	7.2E-09	1.9E-08	1.6E-09	1.9E-08	1.1E-07
Diisooctylphthalate	27554-26-3	33	0	1.5E-07	1.1E-07	1.1E-07	3.9E-07	4.0E-07
Di-n-octyl phthalate	117-84-0	26	0	1.5E-08	3.6E-08	4.0E-09	7.3E-08	1.6E-07
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.2E-10	4.0E-10	0	9.6E-10	1.7E-09
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	7.4E-11	2.7E-08	5.4E-08	1.7E-09	1.1E-07	2.7E-07
Fluoranthene	206-44-0	35	4.6E-09	3.4E-08	3.3E-08	2.0E-08	8.6E-08	1.6E-07
Fluorene	86-73-7	11	0	4.4E-10	1.1E-09	0	2.2E-09	5.3E-09
Hexadecane	544-76-3	28	0	3.2E-09	2.7E-09	3.4E-09	7.6E-09	1.2E-08
1-Hydroxypyrene	5315-79-7	1	0	1.7E-09	1.0E-08	0	0	6.1E-08
Lead		35	2.9E-07	1.5E-06	1.3E-06	1.1E-06	3.8E-06	6.8E-06
Manganese		35	2.7E-07	1.5E-06	8.8E-07	1.6E-06	2.5E-06	4.8E-06
2-(Methylthio)benzothiazole	615-22-5	5	0	2.3E-08	8.2E-08	0	1.3E-07	4.6E-07
Molybdenum		14	0	2.8E-09	3.8E-09	0	8.5E-09	1.4E-08
Naphthalene	91-20-3	1	0	8.0E-11	4.7E-10	0	0	2.8E-09
Naphthalene, 1-methyl-	90-12-0	31	0	2.1E-10	7.1E-10	3.4E-11	4.3E-10	4.2E-09
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	4.8E-11	1.1E-10	0	2.4E-10	5.5E-10
Naphthalene, 1,6-dimethyl-	575-43-9	35	1.3E-10	3.2E-10	1.5E-10	2.8E-10	6.4E-10	8.9E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	35	2.9E-10	8.3E-09	3.9E-09	8.1E-09	1.5E-08	1.9E-08



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene, 2,3-dimethyl-	581-40-8	35	6.2E-11	2.4E-09	2.4E-09	1.7E-09	7.3E-09	1.1E-08
Naphthalene, 2-methyl	91-57-6	19	0	1.9E-10	7.8E-10	3.3E-11	4.2E-10	4.6E-09
1-Octadecene	112-88-9	32	0	7.1E-09	6.2E-09	5.5E-09	1.9E-08	2.2E-08
17-Pentatriacontene	6971-40-0	9	0	6.0E-09	1.4E-08	0	3.3E-08	6.4E-08
Phenanthrene	85-01-8	35	2.6E-09	1.5E-08	1.8E-08	8.4E-09	4.3E-08	9.5E-08
Phenanthrene, 1-methyl	832-69-9	35	3.4E-10	4.0E-09	3.8E-09	2.8E-09	9.9E-09	1.7E-08
Phenanthrene, 2-methyl-	2531-84-2	35	1.1E-10	5.1E-09	5.9E-09	3.0E-09	1.4E-08	2.8E-08
Phenanthrene, 3-methyl	832-71-3	35	6.5E-10	8.5E-09	9.1E-09	5.4E-09	2.3E-08	4.3E-08
N-Phenylbenzamide	93-98-1	22	0	3.3E-08	6.4E-08	5.1E-09	2.0E-07	2.3E-07
Phthalimide	85-41-6	16	0	6.0E-09	1.4E-08	0	3.0E-08	6.3E-08
Pyrene	129-00-0	35	1.3E-08	7.2E-08	4.6E-08	5.9E-08	1.5E-07	2.1E-07
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	1.5E-10	8.8E-10	0	5.2E-11	5.2E-09
Selenium		4	0	9.0E-09	2.8E-08	0	6.3E-08	1.3E-07
Strontium		35	1.2E-07	1.3E-06	2.5E-06	6.8E-07	3.8E-06	1.4E-05
Thallium		21	0	3.6E-10	6.1E-10	2.9E-10	1.1E-09	3.3E-09
Tin		30	0	1.4E-08	1.4E-08	1.0E-08	3.8E-08	5.5E-08
Triethylene glycol monobutyl ether	143-22-6	3	0	1.7E-09	7.1E-09	0	9.1E-09	3.9E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	3.0E-09	5.2E-09	1.6E-09	4.8E-09	8.0E-09	9.9E-09
Vanadium		2	0	7.3E-09	3.0E-08	0	3.8E-08	1.3E-07
Zinc		35	7.0E-05	3.2E-04	1.7E-04	3.0E-04	5.6E-04	9.0E-04

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-332. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Athletes 50<70 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	6.3E-11	4.9E-10	4.6E-10	3.0E-10	1.4E-09	1.8E-09
Aluminum		35	1.1E-05	3.4E-05	1.8E-05	3.1E-05	6.3E-05	8.1E-05
Aniline	62-53-3	12	0	2.7E-09	6.2E-09	0	1.5E-08	2.7E-08
Anthracene	120-12-7	28	0	1.3E-09	2.3E-09	5.0E-10	4.8E-09	1.1E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Anthracene, 2-methyl-	613-12-7	35	7.5E-10	5.1E-09	4.5E-09	3.7E-09	1.2E-08	2.3E-08
Anthracene, 9,10-dimethyl	781-43-1	2	0	4.1E-11	2.3E-10	0	2.7E-11	1.4E-09
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	3.2E-10	1.5E-09	0	9.1E-10	8.2E-09
Anthracene, 9-phenyl	602-55-1	30	0	4.8E-10	5.1E-10	2.9E-10	1.5E-09	2.2E-09
Antimony		35	2.3E-09	3.7E-08	3.3E-08	3.1E-08	8.6E-08	1.6E-07
Barium		35	6.1E-07	2.4E-06	1.6E-06	2.3E-06	4.5E-06	8.9E-06
Benzene, n-butyl-	104-51-8	29	0	2.7E-10	7.0E-10	1.5E-10	4.2E-10	4.2E-09
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	7.5E-08	1.0E-07	2.8E-08	2.7E-07	3.6E-07
Benz[a]anthracene	56-55-3	35	2.9E-10	4.5E-09	4.9E-09	2.1E-09	1.4E-08	2.0E-08
Benzo[b]fluoranthene	205-99-2	35	1.3E-09	6.2E-09	4.6E-09	4.1E-09	1.5E-08	1.8E-08
7H-Benzo[c]fluorene	205-12-9	31	0	1.2E-09	1.9E-09	3.9E-10	4.8E-09	9.6E-09
Benzo[k]fluoranthene	207-08-9	35	1.7E-10	2.0E-09	1.9E-09	1.3E-09	5.8E-09	7.5E-09
Benzothiazole	95-16-9	35	1.7E-07	7.6E-07	5.2E-07	5.6E-07	1.6E-06	1.8E-06
Benzothiazole, 2-phenyl-	883-93-2	35	2.3E-08	8.1E-08	7.6E-08	5.9E-08	1.9E-07	4.4E-07
Benzothiazolone	934-34-9	35	1.2E-06	1.9E-06	3.8E-07	1.9E-06	2.5E-06	2.6E-06
Benzyl butyl phthalate	85-68-7	35	2.0E-09	3.9E-08	3.2E-08	3.4E-08	8.7E-08	1.5E-07
Beryllium		11	0	9.7E-10	1.6E-09	0	3.4E-09	5.4E-09
Butylated Hydroxytoluene	128-37-0	19	0	1.0E-09	1.9E-09	3.6E-10	5.2E-09	8.7E-09
Cadmium		34	0	2.6E-08	1.6E-08	2.4E-08	5.6E-08	6.1E-08
Chromium		35	1.4E-08	7.3E-08	7.2E-08	4.9E-08	1.6E-07	4.1E-07
Cobalt		35	3.1E-07	1.7E-06	1.1E-06	1.6E-06	3.2E-06	5.8E-06
Copper		35	1.3E-06	3.3E-06	1.7E-06	3.0E-06	5.4E-06	1.0E-05
Cyclohexyl isothiocyanate	1122-82-3	29	0	2.4E-07	1.8E-07	2.4E-07	5.7E-07	6.3E-07
Dibenz[a,h]anthracene	53-70-3	7	0	2.7E-10	6.6E-10	0	1.2E-09	3.3E-09
Dibenzothiophene	132-65-0	27	0	1.0E-09	1.6E-09	3.0E-10	4.2E-09	6.3E-09
Dibutyl phthalate	84-74-2	31	0	7.4E-08	5.6E-08	5.9E-08	1.8E-07	2.1E-07
Diethyl Phthalate	84-66-2	23	0	6.2E-09	5.4E-09	6.7E-09	1.5E-08	1.9E-08
Diisobutyl Phthalate	84-69-5	32	0	7.4E-09	1.9E-08	1.7E-09	2.0E-08	1.1E-07
Diisooctylphthalate	27554-26-3	33	0	1.6E-07	1.2E-07	1.2E-07	3.9E-07	4.1E-07
Di-n-octyl phthalate	117-84-0	26	0	1.5E-08	3.7E-08	4.1E-09	7.5E-08	1.6E-07
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.3E-10	4.1E-10	0	9.8E-10	1.8E-09
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	7.6E-11	2.7E-08	5.5E-08	1.7E-09	1.1E-07	2.7E-07
Fluoranthene	206-44-0	35	4.7E-09	3.5E-08	3.4E-08	2.1E-08	8.8E-08	1.6E-07
Fluorene	86-73-7	11	0	4.5E-10	1.1E-09	0	2.2E-09	5.4E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Hexadecane	544-76-3	28	0	3.3E-09	2.7E-09	3.5E-09	7.7E-09	1.3E-08
1-Hydroxypyrene	5315-79-7	1	0	1.8E-09	1.0E-08	0	0	6.2E-08
Lead		35	3.0E-07	1.5E-06	1.3E-06	1.2E-06	3.9E-06	7.0E-06
Manganese		35	2.8E-07	1.5E-06	9.0E-07	1.6E-06	2.5E-06	4.9E-06
2-(Methylthio)benzothiazole	615-22-5	5	0	2.3E-08	8.4E-08	0	1.3E-07	4.7E-07
Molybdenum		14	0	2.8E-09	3.9E-09	0	8.7E-09	1.4E-08
Naphthalene	91-20-3	1	0	8.1E-11	4.8E-10	0	0	2.8E-09
Naphthalene, 1-methyl-	90-12-0	31	0	2.1E-10	7.3E-10	3.5E-11	4.4E-10	4.3E-09
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	4.9E-11	1.1E-10	0	2.5E-10	5.6E-10
Naphthalene, 1,6-dimethyl-	575-43-9	35	1.3E-10	3.2E-10	1.5E-10	2.8E-10	6.6E-10	9.1E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	35	3.0E-10	8.5E-09	4.0E-09	8.3E-09	1.5E-08	1.9E-08
Naphthalene, 2,3-dimethyl-	581-40-8	35	6.3E-11	2.5E-09	2.5E-09	1.7E-09	7.5E-09	1.1E-08
Naphthalene, 2-methyl	91-57-6	19	0	1.9E-10	7.9E-10	3.4E-11	4.3E-10	4.7E-09
1-Octadecene	112-88-9	32	0	7.2E-09	6.3E-09	5.6E-09	1.9E-08	2.2E-08
17-Pentatriacontene	6971-40-0	9	0	6.1E-09	1.4E-08	0	3.4E-08	6.5E-08
Phenanthrene	85-01-8	35	2.6E-09	1.5E-08	1.9E-08	8.5E-09	4.3E-08	9.7E-08
Phenanthrene, 1-methyl	832-69-9	35	3.5E-10	4.0E-09	3.9E-09	2.9E-09	1.0E-08	1.8E-08
Phenanthrene, 2-methyl-	2531-84-2	35	1.1E-10	5.2E-09	6.0E-09	3.1E-09	1.5E-08	2.9E-08
Phenanthrene, 3-methyl	832-71-3	35	6.6E-10	8.6E-09	9.3E-09	5.5E-09	2.4E-08	4.4E-08
N-Phenylbenzamide	93-98-1	22	0	3.4E-08	6.6E-08	5.2E-09	2.1E-07	2.4E-07
Phthalimide	85-41-6	16	0	6.1E-09	1.4E-08	0	3.1E-08	6.5E-08
Pyrene	129-00-0	35	1.4E-08	7.3E-08	4.7E-08	6.0E-08	1.5E-07	2.1E-07
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	1.6E-10	9.0E-10	0	5.3E-11	5.3E-09
Selenium		4	0	9.2E-09	2.8E-08	0	6.4E-08	1.4E-07
Strontium		35	1.3E-07	1.3E-06	2.5E-06	6.9E-07	3.9E-06	1.5E-05
Thallium		21	0	3.7E-10	6.2E-10	2.9E-10	1.1E-09	3.4E-09
Tin		30	0	1.4E-08	1.4E-08	1.1E-08	3.8E-08	5.7E-08
Triethylene glycol monobutyl ether	143-22-6	3	0	1.8E-09	7.3E-09	0	9.3E-09	4.0E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	3.0E-09	5.3E-09	1.7E-09	4.9E-09	8.1E-09	1.0E-08
Vanadium		2	0	7.5E-09	3.1E-08	0	3.9E-08	1.3E-07
Zinc		35	7.1E-05	3.2E-04	1.7E-04	3.1E-04	5.7E-04	9.2E-04

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.



^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis. Values are rounded to two significant figures.

Table F-333. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Coaches 16<30 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	2.8E-11	2.1E-10	2.0E-10	1.3E-10	6.2E-10	7.9E-10
Aluminum		35	4.7E-06	1.5E-05	7.7E-06	1.4E-05	2.7E-05	3.5E-05
Aniline	62-53-3	12	0	1.2E-09	2.7E-09	0	6.6E-09	1.2E-08
Anthracene	120-12-7	28	0	5.6E-10	9.8E-10	2.2E-10	2.1E-09	4.6E-09
Anthracene, 2-methyl-	613-12-7	35	3.3E-10	2.2E-09	2.0E-09	1.6E-09	5.1E-09	9.9E-09
Anthracene, 9,10-dimethyl	781-43-1	2	0	1.8E-11	1.0E-10	0	1.2E-11	5.9E-10
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	1.4E-10	6.4E-10	0	4.0E-10	3.6E-09
Anthracene, 9-phenyl	602-55-1	30	0	2.1E-10	2.2E-10	1.3E-10	6.6E-10	9.5E-10
Antimony		35	9.9E-10	1.6E-08	1.4E-08	1.4E-08	3.7E-08	6.8E-08
Barium		35	2.7E-07	1.0E-06	7.0E-07	1.0E-06	2.0E-06	3.9E-06
Benzene, n-butyl-	104-51-8	29	0	1.2E-10	3.1E-10	6.7E-11	1.8E-10	1.9E-09
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	3.3E-08	4.5E-08	1.2E-08	1.2E-07	1.6E-07
Benz[a]anthracene	56-55-3	35	1.3E-10	2.0E-09	2.1E-09	9.4E-10	6.1E-09	8.5E-09
Benzo[b]fluoranthene	205-99-2	35	5.5E-10	2.7E-09	2.0E-09	1.8E-09	6.5E-09	7.9E-09
7H-Benzo[c]fluorene	205-12-9	31	0	5.2E-10	8.5E-10	1.7E-10	2.1E-09	4.2E-09
Benzo[k]fluoranthene	207-08-9	35	7.3E-11	8.7E-10	8.2E-10	5.5E-10	2.6E-09	3.3E-09
Benzothiazole	95-16-9	35	7.6E-08	3.3E-07	2.3E-07	2.5E-07	6.9E-07	7.9E-07
Benzothiazole, 2-phenyl-	883-93-2	35	1.0E-08	3.6E-08	3.3E-08	2.6E-08	8.2E-08	1.9E-07
Benzothiazolone	934-34-9	35	5.3E-07	8.5E-07	1.6E-07	8.3E-07	1.1E-06	1.1E-06
Benzyl butyl phthalate	85-68-7	35	8.6E-10	1.7E-08	1.4E-08	1.5E-08	3.8E-08	6.7E-08
Beryllium		11	0	4.3E-10	6.9E-10	0	1.5E-09	2.4E-09
Butylated Hydroxytoluene	128-37-0	19	0	4.5E-10	8.4E-10	1.6E-10	2.3E-09	3.8E-09
Cadmium		34	0	1.2E-08	7.1E-09	1.0E-08	2.4E-08	2.7E-08
Chromium		35	5.9E-09	3.2E-08	3.1E-08	2.1E-08	7.0E-08	1.8E-07
Cobalt		35	1.4E-07	7.3E-07	5.0E-07	7.0E-07	1.4E-06	2.5E-06
Copper		35	5.7E-07	1.4E-06	7.4E-07	1.3E-06	2.4E-06	4.5E-06
Cyclohexyl isothiocyanate	1122-82-3	29	0	1.1E-07	7.8E-08	1.0E-07	2.5E-07	2.8E-07



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dibenz[a,h]anthracene	53-70-3	7	0	1.2E-10	2.9E-10	0	5.2E-10	1.4E-09
Dibenzothiophene	132-65-0	27	0	4.5E-10	6.8E-10	1.3E-10	1.8E-09	2.8E-09
Dibutyl phthalate	84-74-2	31	0	3.2E-08	2.4E-08	2.6E-08	7.8E-08	9.2E-08
Diethyl Phthalate	84-66-2	23	0	2.7E-09	2.4E-09	2.9E-09	6.4E-09	8.4E-09
Diisobutyl Phthalate	84-69-5	32	0	3.2E-09	8.4E-09	7.3E-10	8.6E-09	4.9E-08
Diisooctylphthalate	27554-26-3	33	0	6.8E-08	5.1E-08	5.0E-08	1.7E-07	1.8E-07
Di-n-octyl phthalate	117-84-0	26	0	6.7E-09	1.6E-08	1.8E-09	3.3E-08	7.1E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	5.6E-11	1.8E-10	0	4.3E-10	7.8E-10
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	3.3E-11	1.2E-08	2.4E-08	7.4E-10	4.9E-08	1.2E-07
Fluoranthene	206-44-0	35	2.1E-09	1.5E-08	1.5E-08	9.0E-09	3.8E-08	7.2E-08
Fluorene	86-73-7	11	0	2.0E-10	4.7E-10	0	9.8E-10	2.3E-09
Hexadecane	544-76-3	28	0	1.4E-09	1.2E-09	1.5E-09	3.4E-09	5.5E-09
1-Hydroxypyrene	5315-79-7	1	0	7.7E-10	4.6E-09	0	0	2.7E-08
Lead		35	1.3E-07	6.7E-07	5.8E-07	5.1E-07	1.7E-06	3.0E-06
Manganese		35	1.2E-07	6.7E-07	3.9E-07	6.9E-07	1.1E-06	2.1E-06
2-(Methylthio)benzothiazole	615-22-5	5	0	1.0E-08	3.7E-08	0	5.7E-08	2.0E-07
Molybdenum		14	0	1.2E-09	1.7E-09	0	3.8E-09	6.1E-09
Naphthalene	91-20-3	1	0	3.6E-11	2.1E-10	0	0	1.2E-09
Naphthalene, 1-methyl-	90-12-0	31	0	9.2E-11	3.2E-10	1.5E-11	1.9E-10	1.9E-09
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	2.1E-11	4.8E-11	0	1.1E-10	2.5E-10
Naphthalene, 1,6-dimethyl-	575-43-9	35	5.6E-11	1.4E-10	6.7E-11	1.2E-10	2.9E-10	4.0E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	35	1.3E-10	3.7E-09	1.8E-09	3.6E-09	6.5E-09	8.4E-09
Naphthalene, 2,3-dimethyl-	581-40-8	35	2.8E-11	1.1E-09	1.1E-09	7.5E-10	3.3E-09	4.7E-09
Naphthalene, 2-methyl	91-57-6	19	0	8.3E-11	3.5E-10	1.5E-11	1.9E-10	2.1E-09
1-Octadecene	112-88-9	32	0	3.2E-09	2.8E-09	2.5E-09	8.5E-09	9.8E-09
17-Pentatriacontene	6971-40-0	9	0	2.7E-09	6.3E-09	0	1.5E-08	2.8E-08
Phenanthrene	85-01-8	35	1.1E-09	6.7E-09	8.2E-09	3.7E-09	1.9E-08	4.2E-08
Phenanthrene, 1-methyl	832-69-9	35	1.5E-10	1.8E-09	1.7E-09	1.3E-09	4.4E-09	7.7E-09
Phenanthrene, 2-methyl-	2531-84-2	35	4.8E-11	2.3E-09	2.6E-09	1.4E-09	6.5E-09	1.3E-08
Phenanthrene, 3-methyl	832-71-3	35	2.9E-10	3.8E-09	4.1E-09	2.4E-09	1.0E-08	1.9E-08
N-Phenylbenzamide	93-98-1	22	0	1.5E-08	2.9E-08	2.3E-09	9.0E-08	1.0E-07
Phthalimide	85-41-6	16	0	2.7E-09	6.1E-09	0	1.3E-08	2.8E-08
Pyrene	129-00-0	35	5.9E-09	3.2E-08	2.1E-08	2.6E-08	6.7E-08	9.2E-08



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	6.8E-11	3.9E-10	0	2.3E-11	2.3E-09
Selenium		4	0	4.0E-09	1.2E-08	0	2.8E-08	6.0E-08
Strontium		35	5.5E-08	5.7E-07	1.1E-06	3.0E-07	1.7E-06	6.5E-06
Thallium		21	0	1.6E-10	2.7E-10	1.3E-10	5.0E-10	1.5E-09
Tin		30	0	6.3E-09	6.3E-09	4.7E-09	1.7E-08	2.5E-08
Triethylene glycol monobutyl ether	143-22-6	3	0	7.8E-10	3.2E-09	0	4.1E-09	1.8E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	1.3E-09	2.3E-09	7.3E-10	2.1E-09	3.6E-09	4.4E-09
Vanadium		2	0	3.3E-09	1.3E-08	0	1.7E-08	5.8E-08
Zinc		35	3.1E-05	1.4E-04	7.5E-05	1.4E-04	2.5E-04	4.0E-04

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-334. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Coaches 30<40 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	2.6E-11	2.0E-10	1.9E-10	1.2E-10	5.9E-10	7.4E-10
Aluminum		35	4.5E-06	1.4E-05	7.2E-06	1.3E-05	2.6E-05	3.3E-05
Aniline	62-53-3	12	0	1.1E-09	2.6E-09	0	6.2E-09	1.1E-08
Anthracene	120-12-7	28	0	5.3E-10	9.2E-10	2.0E-10	2.0E-09	4.3E-09
Anthracene, 2-methyl-	613-12-7	35	3.1E-10	2.1E-09	1.8E-09	1.5E-09	4.8E-09	9.3E-09
Anthracene, 9,10-dimethyl	781-43-1	2	0	1.7E-11	9.4E-11	0	1.1E-11	5.6E-10
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	1.3E-10	6.0E-10	0	3.7E-10	3.4E-09
Anthracene, 9-phenyl	602-55-1	30	0	2.0E-10	2.1E-10	1.2E-10	6.2E-10	8.9E-10
Antimony		35	9.3E-10	1.5E-08	1.3E-08	1.3E-08	3.5E-08	6.4E-08
Barium		35	2.5E-07	9.7E-07	6.6E-07	9.5E-07	1.8E-06	3.6E-06
Benzene, n-butyl-	104-51-8	29	0	1.1E-10	2.9E-10	6.3E-11	1.7E-10	1.7E-09
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	3.1E-08	4.2E-08	1.1E-08	1.1E-07	1.5E-07
Benz[a]anthracene	56-55-3	35	1.2E-10	1.8E-09	2.0E-09	8.8E-10	5.7E-09	8.0E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[b]fluoranthene	205-99-2	35	5.2E-10	2.5E-09	1.9E-09	1.7E-09	6.1E-09	7.4E-09
7H-Benzo[c]fluorene	205-12-9	31	0	4.9E-10	8.0E-10	1.6E-10	2.0E-09	4.0E-09
Benzo[k]fluoranthene	207-08-9	35	6.9E-11	8.2E-10	7.7E-10	5.2E-10	2.4E-09	3.1E-09
Benzothiazole	95-16-9	35	7.1E-08	3.1E-07	2.1E-07	2.3E-07	6.4E-07	7.4E-07
Benzothiazole, 2-phenyl-	883-93-2	35	9.5E-09	3.3E-08	3.1E-08	2.4E-08	7.7E-08	1.8E-07
Benzothiazolone	934-34-9	35	5.0E-07	7.9E-07	1.5E-07	7.8E-07	1.0E-06	1.0E-06
Benzyl butyl phthalate	85-68-7	35	8.1E-10	1.6E-08	1.3E-08	1.4E-08	3.6E-08	6.3E-08
Beryllium		11	0	4.0E-10	6.5E-10	0	1.4E-09	2.2E-09
Butylated Hydroxytoluene	128-37-0	19	0	4.2E-10	7.9E-10	1.5E-10	2.1E-09	3.6E-09
Cadmium		34	0	1.1E-08	6.7E-09	9.7E-09	2.3E-08	2.5E-08
Chromium		35	5.6E-09	3.0E-08	2.9E-08	2.0E-08	6.6E-08	1.7E-07
Cobalt		35	1.3E-07	6.9E-07	4.7E-07	6.5E-07	1.3E-06	2.4E-06
Copper		35	5.3E-07	1.3E-06	7.0E-07	1.2E-06	2.2E-06	4.3E-06
Cyclohexyl isothiocyanate	1122-82-3	29	0	9.9E-08	7.3E-08	9.9E-08	2.3E-07	2.6E-07
Dibenz[a,h]anthracene	53-70-3	7	0	1.1E-10	2.7E-10	0	4.9E-10	1.4E-09
Dibenzothiophene	132-65-0	27	0	4.3E-10	6.4E-10	1.2E-10	1.7E-09	2.6E-09
Dibutyl phthalate	84-74-2	31	0	3.0E-08	2.3E-08	2.4E-08	7.4E-08	8.7E-08
Diethyl Phthalate	84-66-2	23	0	2.5E-09	2.2E-09	2.8E-09	6.0E-09	7.9E-09
Diisobutyl Phthalate	84-69-5	32	0	3.0E-09	7.9E-09	6.9E-10	8.1E-09	4.6E-08
Diisooctylphthalate	27554-26-3	33	0	6.4E-08	4.8E-08	4.7E-08	1.6E-07	1.7E-07
Di-n-octyl phthalate	117-84-0	26	0	6.3E-09	1.5E-08	1.7E-09	3.1E-08	6.7E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	5.2E-11	1.7E-10	0	4.0E-10	7.3E-10
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	3.1E-11	1.1E-08	2.3E-08	6.9E-10	4.6E-08	1.1E-07
Fluoranthene	206-44-0	35	1.9E-09	1.4E-08	1.4E-08	8.5E-09	3.6E-08	6.8E-08
Fluorene	86-73-7	11	0	1.9E-10	4.4E-10	0	9.2E-10	2.2E-09
Hexadecane	544-76-3	28	0	1.3E-09	1.1E-09	1.4E-09	3.2E-09	5.2E-09
1-Hydroxypyrene	5315-79-7	1	0	7.3E-10	4.3E-09	0	0	2.5E-08
Lead		35	1.2E-07	6.3E-07	5.5E-07	4.8E-07	1.6E-06	2.9E-06
Manganese		35	1.1E-07	6.3E-07	3.7E-07	6.5E-07	1.0E-06	2.0E-06
2-(Methylthio)benzothiazole	615-22-5	5	0	9.5E-09	3.4E-08	0	5.3E-08	1.9E-07
Molybdenum		14	0	1.2E-09	1.6E-09	0	3.6E-09	5.7E-09
Naphthalene	91-20-3	1	0	3.3E-11	2.0E-10	0	0	1.2E-09
Naphthalene, 1-methyl-	90-12-0	31	0	8.6E-11	3.0E-10	1.4E-11	1.8E-10	1.8E-09
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	2.0E-11	4.5E-11	0	1.0E-10	2.3E-10



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene, 1,6-dimethyl-	575-43-9	35	5.3E-11	1.3E-10	6.3E-11	1.2E-10	2.7E-10	3.7E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	35	1.2E-10	3.5E-09	1.6E-09	3.4E-09	6.1E-09	7.9E-09
Naphthalene, 2,3-dimethyl-	581-40-8	35	2.6E-11	1.0E-09	1.0E-09	7.1E-10	3.1E-09	4.5E-09
Naphthalene, 2-methyl	91-57-6	19	0	7.8E-11	3.3E-10	1.4E-11	1.7E-10	1.9E-09
1-Octadecene	112-88-9	32	0	3.0E-09	2.6E-09	2.3E-09	8.0E-09	9.2E-09
17-Pentatriacontene	6971-40-0	9	0	2.5E-09	5.9E-09	0	1.4E-08	2.7E-08
Phenanthrene	85-01-8	35	1.1E-09	6.3E-09	7.7E-09	3.5E-09	1.8E-08	4.0E-08
Phenanthrene, 1-methyl	832-69-9	35	1.4E-10	1.7E-09	1.6E-09	1.2E-09	4.1E-09	7.2E-09
Phenanthrene, 2-methyl-	2531-84-2	35	4.5E-11	2.1E-09	2.5E-09	1.3E-09	6.1E-09	1.2E-08
Phenanthrene, 3-methyl	832-71-3	35	2.7E-10	3.5E-09	3.8E-09	2.3E-09	9.7E-09	1.8E-08
N-Phenylbenzamide	93-98-1	22	0	1.4E-08	2.7E-08	2.1E-09	8.5E-08	9.8E-08
Phthalimide	85-41-6	16	0	2.5E-09	5.8E-09	0	1.3E-08	2.7E-08
Pyrene	129-00-0	35	5.6E-09	3.0E-08	1.9E-08	2.5E-08	6.3E-08	8.6E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	6.4E-11	3.7E-10	0	2.2E-11	2.2E-09
Selenium		4	0	3.8E-09	1.2E-08	0	2.6E-08	5.6E-08
Strontium		35	5.1E-08	5.4E-07	1.0E-06	2.9E-07	1.6E-06	6.1E-06
Thallium		21	0	1.5E-10	2.6E-10	1.2E-10	4.7E-10	1.4E-09
Tin		30	0	5.9E-09	5.9E-09	4.4E-09	1.6E-08	2.3E-08
Triethylene glycol monobutyl ether	143-22-6	3	0	7.3E-10	3.0E-09	0	3.8E-09	1.6E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	1.2E-09	2.2E-09	6.9E-10	2.0E-09	3.3E-09	4.1E-09
Vanadium		2	0	3.1E-09	1.3E-08	0	1.6E-08	5.4E-08
Zinc		35	2.9E-05	1.3E-04	7.0E-05	1.3E-04	2.4E-04	3.8E-04

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-335. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Coaches 40<50 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	2.5E-11	2.0E-10	1.8E-10	1.2E-10	5.7E-10	7.3E-10



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aluminum		35	4.4E-06	1.4E-05	7.0E-06	1.3E-05	2.5E-05	3.2E-05
Aniline	62-53-3	12	0	1.1E-09	2.5E-09	0	6.1E-09	1.1E-08
Anthracene	120-12-7	28	0	5.2E-10	9.0E-10	2.0E-10	1.9E-09	4.2E-09
Anthracene, 2-methyl-	613-12-7	35	3.0E-10	2.1E-09	1.8E-09	1.5E-09	4.7E-09	9.1E-09
Anthracene, 9,10-dimethyl	781-43-1	2	0	1.7E-11	9.2E-11	0	1.1E-11	5.5E-10
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	1.3E-10	5.9E-10	0	3.6E-10	3.3E-09
Anthracene, 9-phenyl	602-55-1	30	0	1.9E-10	2.1E-10	1.2E-10	6.1E-10	8.7E-10
Antimony		35	9.1E-10	1.5E-08	1.3E-08	1.3E-08	3.4E-08	6.2E-08
Barium		35	2.5E-07	9.5E-07	6.5E-07	9.3E-07	1.8E-06	3.6E-06
Benzene, n-butyl-	104-51-8	29	0	1.1E-10	2.8E-10	6.2E-11	1.7E-10	1.7E-09
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	3.0E-08	4.1E-08	1.1E-08	1.1E-07	1.5E-07
Benz[a]anthracene	56-55-3	35	1.2E-10	1.8E-09	2.0E-09	8.6E-10	5.6E-09	7.8E-09
Benzo[b]fluoranthene	205-99-2	35	5.0E-10	2.5E-09	1.8E-09	1.7E-09	6.0E-09	7.3E-09
7H-Benzo[c]fluorene	205-12-9	31	0	4.8E-10	7.8E-10	1.6E-10	1.9E-09	3.9E-09
Benzo[k]fluoranthene	207-08-9	35	6.7E-11	8.0E-10	7.6E-10	5.1E-10	2.4E-09	3.0E-09
Benzothiazole	95-16-9	35	7.0E-08	3.0E-07	2.1E-07	2.3E-07	6.3E-07	7.2E-07
Benzothiazole, 2-phenyl-	883-93-2	35	9.3E-09	3.3E-08	3.1E-08	2.4E-08	7.5E-08	1.8E-07
Benzothiazolone	934-34-9	35	4.9E-07	7.8E-07	1.5E-07	7.6E-07	1.0E-06	1.0E-06
Benzyl butyl phthalate	85-68-7	35	7.9E-10	1.6E-08	1.3E-08	1.4E-08	3.5E-08	6.2E-08
Beryllium		11	0	3.9E-10	6.3E-10	0	1.4E-09	2.2E-09
Butylated Hydroxytoluene	128-37-0	19	0	4.2E-10	7.8E-10	1.4E-10	2.1E-09	3.5E-09
Cadmium		34	0	1.1E-08	6.5E-09	9.5E-09	2.2E-08	2.5E-08
Chromium		35	5.5E-09	3.0E-08	2.9E-08	1.9E-08	6.4E-08	1.6E-07
Cobalt		35	1.2E-07	6.7E-07	4.6E-07	6.4E-07	1.3E-06	2.3E-06
Copper		35	5.2E-07	1.3E-06	6.8E-07	1.2E-06	2.2E-06	4.2E-06
Cyclohexyl isothiocyanate	1122-82-3	29	0	9.7E-08	7.1E-08	9.6E-08	2.3E-07	2.5E-07
Dibenz[a,h]anthracene	53-70-3	7	0	1.1E-10	2.7E-10	0	4.8E-10	1.3E-09
Dibenzothiophene	132-65-0	27	0	4.2E-10	6.3E-10	1.2E-10	1.7E-09	2.5E-09
Dibutyl phthalate	84-74-2	31	0	3.0E-08	2.2E-08	2.4E-08	7.2E-08	8.5E-08
Diethyl Phthalate	84-66-2	23	0	2.5E-09	2.2E-09	2.7E-09	5.9E-09	7.7E-09
Diisobutyl Phthalate	84-69-5	32	0	3.0E-09	7.7E-09	6.7E-10	7.9E-09	4.5E-08
Diisooctylphthalate	27554-26-3	33	0	6.3E-08	4.7E-08	4.6E-08	1.6E-07	1.7E-07
Di-n-octyl phthalate	117-84-0	26	0	6.2E-09	1.5E-08	1.6E-09	3.0E-08	6.5E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	5.1E-11	1.6E-10	0	3.9E-10	7.2E-10



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	3.0E-11	1.1E-08	2.2E-08	6.8E-10	4.5E-08	1.1E-07
Fluoranthene	206-44-0	35	1.9E-09	1.4E-08	1.4E-08	8.3E-09	3.5E-08	6.6E-08
Fluorene	86-73-7	11	0	1.8E-10	4.3E-10	0	9.0E-10	2.2E-09
Hexadecane	544-76-3	28	0	1.3E-09	1.1E-09	1.4E-09	3.1E-09	5.1E-09
1-Hydroxypyrene	5315-79-7	1	0	7.1E-10	4.2E-09	0	0	2.5E-08
Lead		35	1.2E-07	6.2E-07	5.3E-07	4.7E-07	1.6E-06	2.8E-06
Manganese		35	1.1E-07	6.1E-07	3.6E-07	6.4E-07	1.0E-06	2.0E-06
2-(Methylthio)benzothiazole	615-22-5	5	0	9.3E-09	3.4E-08	0	5.2E-08	1.9E-07
Molybdenum		14	0	1.1E-09	1.6E-09	0	3.5E-09	5.6E-09
Naphthalene	91-20-3	1	0	3.3E-11	1.9E-10	0	0	1.1E-09
Naphthalene, 1-methyl-	90-12-0	31	0	8.4E-11	2.9E-10	1.4E-11	1.8E-10	1.7E-09
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	2.0E-11	4.4E-11	0	9.8E-11	2.3E-10
Naphthalene, 1,6-dimethyl-	575-43-9	35	5.2E-11	1.3E-10	6.2E-11	1.1E-10	2.6E-10	3.7E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	35	1.2E-10	3.4E-09	1.6E-09	3.3E-09	6.0E-09	7.7E-09
Naphthalene, 2,3-dimethyl-	581-40-8	35	2.5E-11	1.0E-09	9.9E-10	6.9E-10	3.0E-09	4.4E-09
Naphthalene, 2-methyl	91-57-6	19	0	7.6E-11	3.2E-10	1.4E-11	1.7E-10	1.9E-09
1-Octadecene	112-88-9	32	0	2.9E-09	2.5E-09	2.3E-09	7.8E-09	9.0E-09
17-Pentatriacontene	6971-40-0	9	0	2.5E-09	5.8E-09	0	1.3E-08	2.6E-08
Phenanthrene	85-01-8	35	1.1E-09	6.1E-09	7.6E-09	3.4E-09	1.7E-08	3.9E-08
Phenanthrene, 1-methyl	832-69-9	35	1.4E-10	1.6E-09	1.6E-09	1.2E-09	4.0E-09	7.1E-09
Phenanthrene, 2-methyl-	2531-84-2	35	4.4E-11	2.1E-09	2.4E-09	1.2E-09	5.9E-09	1.1E-08
Phenanthrene, 3-methyl	832-71-3	35	2.7E-10	3.5E-09	3.7E-09	2.2E-09	9.5E-09	1.8E-08
N-Phenylbenzamide	93-98-1	22	0	1.4E-08	2.6E-08	2.1E-09	8.3E-08	9.6E-08
Phthalimide	85-41-6	16	0	2.5E-09	5.6E-09	0	1.2E-08	2.6E-08
Pyrene	129-00-0	35	5.4E-09	2.9E-08	1.9E-08	2.4E-08	6.2E-08	8.4E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	6.3E-11	3.6E-10	0	2.1E-11	2.1E-09
Selenium		4	0	3.7E-09	1.1E-08	0	2.6E-08	5.5E-08
Strontium		35	5.0E-08	5.3E-07	1.0E-06	2.8E-07	1.6E-06	5.9E-06
Thallium		21	0	1.5E-10	2.5E-10	1.2E-10	4.6E-10	1.4E-09
Tin		30	0	5.8E-09	5.8E-09	4.3E-09	1.5E-08	2.3E-08
Triethylene glycol monobutyl ether	143-22-6	3	0	7.2E-10	2.9E-09	0	3.7E-09	1.6E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	1.2E-09	2.1E-09	6.7E-10	2.0E-09	3.3E-09	4.0E-09
Vanadium		2	0	3.0E-09	1.2E-08	0	1.6E-08	5.3E-08



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Zinc		35	2.9E-05	1.3E-04	6.9E-05	1.2E-04	2.3E-04	3.7E-04

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-336. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Coaches 50<70 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	2.6E-11	2.0E-10	1.8E-10	1.2E-10	5.8E-10	7.3E-10
Aluminum		35	4.4E-06	1.4E-05	7.1E-06	1.3E-05	2.5E-05	3.3E-05
Aniline	62-53-3	12	0	1.1E-09	2.5E-09	0	6.1E-09	1.1E-08
Anthracene	120-12-7	28	0	5.2E-10	9.1E-10	2.0E-10	1.9E-09	4.2E-09
Anthracene, 2-methyl-	613-12-7	35	3.0E-10	2.1E-09	1.8E-09	1.5E-09	4.7E-09	9.2E-09
Anthracene, 9,10-dimethyl	781-43-1	2	0	1.7E-11	9.3E-11	0	1.1E-11	5.5E-10
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	1.3E-10	5.9E-10	0	3.7E-10	3.3E-09
Anthracene, 9-phenyl	602-55-1	30	0	2.0E-10	2.1E-10	1.2E-10	6.1E-10	8.7E-10
Antimony		35	9.1E-10	1.5E-08	1.3E-08	1.3E-08	3.4E-08	6.3E-08
Barium		35	2.5E-07	9.5E-07	6.5E-07	9.3E-07	1.8E-06	3.6E-06
Benzene, n-butyl-	104-51-8	29	0	1.1E-10	2.8E-10	6.2E-11	1.7E-10	1.7E-09
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	3.0E-08	4.1E-08	1.1E-08	1.1E-07	1.5E-07
Benz[a]anthracene	56-55-3	35	1.2E-10	1.8E-09	2.0E-09	8.6E-10	5.6E-09	7.9E-09
Benzo[b]fluoranthene	205-99-2	35	5.1E-10	2.5E-09	1.8E-09	1.7E-09	6.0E-09	7.3E-09
7H-Benzo[c]fluorene	205-12-9	31	0	4.8E-10	7.8E-10	1.6E-10	1.9E-09	3.9E-09
Benzo[k]fluoranthene	207-08-9	35	6.8E-11	8.0E-10	7.6E-10	5.1E-10	2.4E-09	3.0E-09
Benzothiazole	95-16-9	35	7.0E-08	3.1E-07	2.1E-07	2.3E-07	6.3E-07	7.2E-07
Benzothiazole, 2-phenyl-	883-93-2	35	9.3E-09	3.3E-08	3.1E-08	2.4E-08	7.5E-08	1.8E-07
Benzothiazolone	934-34-9	35	4.9E-07	7.8E-07	1.5E-07	7.7E-07	1.0E-06	1.0E-06
Benzyl butyl phthalate	85-68-7	35	8.0E-10	1.6E-08	1.3E-08	1.4E-08	3.5E-08	6.2E-08
Beryllium		11	0	3.9E-10	6.3E-10	0	1.4E-09	2.2E-09
Butylated Hydroxytoluene	128-37-0	19	0	4.2E-10	7.8E-10	1.4E-10	2.1E-09	3.5E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Cadmium		34	0	1.1E-08	6.5E-09	9.6E-09	2.2E-08	2.5E-08
Chromium		35	5.5E-09	3.0E-08	2.9E-08	2.0E-08	6.5E-08	1.6E-07
Cobalt		35	1.3E-07	6.7E-07	4.6E-07	6.4E-07	1.3E-06	2.3E-06
Copper		35	5.2E-07	1.3E-06	6.8E-07	1.2E-06	2.2E-06	4.2E-06
Cyclohexyl isothiocyanate	1122-82-3	29	0	9.8E-08	7.2E-08	9.7E-08	2.3E-07	2.5E-07
Dibenz[a,h]anthracene	53-70-3	7	0	1.1E-10	2.7E-10	0	4.8E-10	1.3E-09
Dibenzothiophene	132-65-0	27	0	4.2E-10	6.3E-10	1.2E-10	1.7E-09	2.5E-09
Dibutyl phthalate	84-74-2	31	0	3.0E-08	2.2E-08	2.4E-08	7.2E-08	8.5E-08
Diethyl Phthalate	84-66-2	23	0	2.5E-09	2.2E-09	2.7E-09	5.9E-09	7.8E-09
Diisobutyl Phthalate	84-69-5	32	0	3.0E-09	7.7E-09	6.7E-10	7.9E-09	4.5E-08
Diisooctylphthalate	27554-26-3	33	0	6.3E-08	4.7E-08	4.6E-08	1.6E-07	1.7E-07
Di-n-octyl phthalate	117-84-0	26	0	6.2E-09	1.5E-08	1.6E-09	3.0E-08	6.6E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	5.1E-11	1.6E-10	0	4.0E-10	7.2E-10
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	3.0E-11	1.1E-08	2.2E-08	6.8E-10	4.5E-08	1.1E-07
Fluoranthene	206-44-0	35	1.9E-09	1.4E-08	1.4E-08	8.3E-09	3.5E-08	6.6E-08
Fluorene	86-73-7	11	0	1.8E-10	4.3E-10	0	9.0E-10	2.2E-09
Hexadecane	544-76-3	28	0	1.3E-09	1.1E-09	1.4E-09	3.1E-09	5.1E-09
1-Hydroxypyrene	5315-79-7	1	0	7.1E-10	4.2E-09	0	0	2.5E-08
Lead		35	1.2E-07	6.2E-07	5.4E-07	4.7E-07	1.6E-06	2.8E-06
Manganese		35	1.1E-07	6.1E-07	3.6E-07	6.4E-07	1.0E-06	2.0E-06
2-(Methylthio)benzothiazole	615-22-5	5	0	9.4E-09	3.4E-08	0	5.2E-08	1.9E-07
Molybdenum		14	0	1.1E-09	1.6E-09	0	3.5E-09	5.6E-09
Naphthalene	91-20-3	1	0	3.3E-11	1.9E-10	0	0	1.1E-09
Naphthalene, 1-methyl-	90-12-0	31	0	8.4E-11	2.9E-10	1.4E-11	1.8E-10	1.7E-09
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	2.0E-11	4.4E-11	0	9.9E-11	2.3E-10
Naphthalene, 1,6-dimethyl-	575-43-9	35	5.2E-11	1.3E-10	6.2E-11	1.1E-10	2.6E-10	3.7E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	35	1.2E-10	3.4E-09	1.6E-09	3.3E-09	6.0E-09	7.8E-09
Naphthalene, 2,3-dimethyl-	581-40-8	35	2.6E-11	1.0E-09	1.0E-09	6.9E-10	3.0E-09	4.4E-09
Naphthalene, 2-methyl	91-57-6	19	0	7.6E-11	3.2E-10	1.4E-11	1.7E-10	1.9E-09
1-Octadecene	112-88-9	32	0	2.9E-09	2.5E-09	2.3E-09	7.8E-09	9.0E-09
17-Pentatriacontene	6971-40-0	9	0	2.5E-09	5.8E-09	0	1.4E-08	2.6E-08
Phenanthrene	85-01-8	35	1.1E-09	6.1E-09	7.6E-09	3.4E-09	1.8E-08	3.9E-08
Phenanthrene, 1-methyl	832-69-9	35	1.4E-10	1.6E-09	1.6E-09	1.2E-09	4.1E-09	7.1E-09



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Phenanthrene, 2-methyl-	2531-84-2	35	4.4E-11	2.1E-09	2.4E-09	1.3E-09	5.9E-09	1.2E-08
Phenanthrene, 3-methyl	832-71-3	35	2.7E-10	3.5E-09	3.7E-09	2.2E-09	9.6E-09	1.8E-08
N-Phenylbenzamide	93-98-1	22	0	1.4E-08	2.6E-08	2.1E-09	8.3E-08	9.6E-08
Phthalimide	85-41-6	16	0	2.5E-09	5.7E-09	0	1.2E-08	2.6E-08
Pyrene	129-00-0	35	5.5E-09	2.9E-08	1.9E-08	2.4E-08	6.2E-08	8.5E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	6.3E-11	3.6E-10	0	2.1E-11	2.1E-09
Selenium		4	0	3.7E-09	1.1E-08	0	2.6E-08	5.5E-08
Strontium		35	5.0E-08	5.3E-07	1.0E-06	2.8E-07	1.6E-06	6.0E-06
Thallium		21	0	1.5E-10	2.5E-10	1.2E-10	4.6E-10	1.4E-09
Tin		30	0	5.8E-09	5.8E-09	4.3E-09	1.5E-08	2.3E-08
Triethylene glycol monobutyl ether	143-22-6	3	0	7.2E-10	2.9E-09	0	3.7E-09	1.6E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	1.2E-09	2.1E-09	6.8E-10	2.0E-09	3.3E-09	4.1E-09
Vanadium		2	0	3.0E-09	1.2E-08	0	1.6E-08	5.3E-08
Zinc		35	2.9E-05	1.3E-04	6.9E-05	1.2E-04	2.3E-04	3.7E-04

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-337. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Referees 16<30 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	1.2E-11	9.4E-11	8.7E-11	5.7E-11	2.7E-10	3.5E-10
Aluminum		35	2.1E-06	6.6E-06	3.4E-06	6.0E-06	1.2E-05	1.6E-05
Aniline	62-53-3	12	0	5.2E-10	1.2E-09	0	2.9E-09	5.3E-09
Anthracene	120-12-7	28	0	2.5E-10	4.3E-10	9.5E-11	9.3E-10	2.0E-09
Anthracene, 2-methyl-	613-12-7	35	1.4E-10	9.8E-10	8.6E-10	7.1E-10	2.3E-09	4.4E-09
Anthracene, 9,10-dimethyl	781-43-1	2	0	7.9E-12	4.4E-11	0	5.2E-12	2.6E-10
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	6.1E-11	2.8E-10	0	1.7E-10	1.6E-09
Anthracene, 9-phenyl	602-55-1	30	0	9.3E-11	9.8E-11	5.7E-11	2.9E-10	4.2E-10
Antimony		35	4.3E-10	7.1E-09	6.3E-09	6.0E-09	1.6E-08	3.0E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Barium		35	1.2E-07	4.5E-07	3.1E-07	4.4E-07	8.6E-07	1.7E-06
Benzene, n-butyl-	104-51-8	29	0	5.2E-11	1.3E-10	2.9E-11	8.1E-11	8.1E-10
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	1.4E-08	2.0E-08	5.4E-09	5.3E-08	7.0E-08
Benz[a]anthracene	56-55-3	35	5.6E-11	8.6E-10	9.4E-10	4.1E-10	2.7E-09	3.7E-09
Benzo[b]fluoranthene	205-99-2	35	2.4E-10	1.2E-09	8.8E-10	7.9E-10	2.9E-09	3.5E-09
7H-Benzo[c]fluorene	205-12-9	31	0	2.3E-10	3.7E-10	7.5E-11	9.3E-10	1.9E-09
Benzo[k]fluoranthene	207-08-9	35	3.2E-11	3.8E-10	3.6E-10	2.4E-10	1.1E-09	1.4E-09
Benzothiazole	95-16-9	35	3.3E-08	1.5E-07	1.0E-07	1.1E-07	3.0E-07	3.5E-07
Benzothiazole, 2-phenyl-	883-93-2	35	4.5E-09	1.6E-08	1.5E-08	1.1E-08	3.6E-08	8.4E-08
Benzothiazolone	934-34-9	35	2.3E-07	3.7E-07	7.2E-08	3.6E-07	4.8E-07	4.9E-07
Benzyl butyl phthalate	85-68-7	35	3.8E-10	7.4E-09	6.2E-09	6.6E-09	1.7E-08	3.0E-08
Beryllium		11	0	1.9E-10	3.0E-10	0	6.5E-10	1.0E-09
Butylated Hydroxytoluene	128-37-0	19	0	2.0E-10	3.7E-10	6.9E-11	1.0E-09	1.7E-09
Cadmium		34	0	5.1E-09	3.1E-09	4.6E-09	1.1E-08	1.2E-08
Chromium		35	2.6E-09	1.4E-08	1.4E-08	9.3E-09	3.1E-08	7.8E-08
Cobalt		35	6.0E-08	3.2E-07	2.2E-07	3.1E-07	6.1E-07	1.1E-06
Copper		35	2.5E-07	6.3E-07	3.3E-07	5.8E-07	1.0E-06	2.0E-06
Cyclohexyl isothiocyanate	1122-82-3	29	0	4.6E-08	3.4E-08	4.6E-08	1.1E-07	1.2E-07
Dibenz[a,h]anthracene	53-70-3	7	0	5.1E-11	1.3E-10	0	2.3E-10	6.3E-10
Dibenzothiophene	132-65-0	27	0	2.0E-10	3.0E-10	5.8E-11	8.0E-10	1.2E-09
Dibutyl phthalate	84-74-2	31	0	1.4E-08	1.1E-08	1.1E-08	3.4E-08	4.1E-08
Diethyl Phthalate	84-66-2	23	0	1.2E-09	1.0E-09	1.3E-09	2.8E-09	3.7E-09
Diisobutyl Phthalate	84-69-5	32	0	1.4E-09	3.7E-09	3.2E-10	3.8E-09	2.2E-08
Diisooctylphthalate	27554-26-3	33	0	3.0E-08	2.2E-08	2.2E-08	7.6E-08	7.9E-08
Di-n-octyl phthalate	117-84-0	26	0	3.0E-09	7.1E-09	7.8E-10	1.4E-08	3.1E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	2.4E-11	7.8E-11	0	1.9E-10	3.4E-10
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	1.5E-11	5.2E-09	1.1E-08	3.2E-10	2.2E-08	5.2E-08
Fluoranthene	206-44-0	35	9.1E-10	6.7E-09	6.5E-09	4.0E-09	1.7E-08	3.2E-08
Fluorene	86-73-7	11	0	8.7E-11	2.1E-10	0	4.3E-10	1.0E-09
Hexadecane	544-76-3	28	0	6.3E-10	5.3E-10	6.6E-10	1.5E-09	2.4E-09
1-Hydroxypyrene	5315-79-7	1	0	3.4E-10	2.0E-09	0	0	1.2E-08
Lead		35	5.7E-08	3.0E-07	2.6E-07	2.2E-07	7.4E-07	1.3E-06
Manganese		35	5.3E-08	2.9E-07	1.7E-07	3.0E-07	4.8E-07	9.3E-07



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
2-(Methylthio)benzothiazole	615-22-5	5	0	4.5E-09	1.6E-08	0	2.5E-08	9.0E-08
Molybdenum		14	0	5.4E-10	7.5E-10	0	1.7E-09	2.7E-09
Naphthalene	91-20-3	1	0	1.6E-11	9.2E-11	0	0	5.5E-10
Naphthalene, 1-methyl-	90-12-0	31	0	4.0E-11	1.4E-10	6.7E-12	8.4E-11	8.3E-10
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	9.4E-12	2.1E-11	0	4.7E-11	1.1E-10
Naphthalene, 1,6-dimethyl-	575-43-9	35	2.5E-11	6.2E-11	3.0E-11	5.4E-11	1.3E-10	1.7E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	35	5.8E-11	1.6E-09	7.7E-10	1.6E-09	2.9E-09	3.7E-09
Naphthalene, 2,3-dimethyl-	581-40-8	35	1.2E-11	4.8E-10	4.7E-10	3.3E-10	1.4E-09	2.1E-09
Naphthalene, 2-methyl	91-57-6	19	0	3.6E-11	1.5E-10	6.5E-12	8.2E-11	9.0E-10
1-Octadecene	112-88-9	32	0	1.4E-09	1.2E-09	1.1E-09	3.7E-09	4.3E-09
17-Pentatriacontene	6971-40-0	9	0	1.2E-09	2.8E-09	0	6.4E-09	1.2E-08
Phenanthrene	85-01-8	35	5.0E-10	2.9E-09	3.6E-09	1.6E-09	8.3E-09	1.9E-08
Phenanthrene, 1-methyl	832-69-9	35	6.7E-11	7.8E-10	7.5E-10	5.5E-10	1.9E-09	3.4E-09
Phenanthrene, 2-methyl-	2531-84-2	35	2.1E-11	1.0E-09	1.2E-09	6.0E-10	2.8E-09	5.5E-09
Phenanthrene, 3-methyl	832-71-3	35	1.3E-10	1.7E-09	1.8E-09	1.1E-09	4.6E-09	8.4E-09
N-Phenylbenzamide	93-98-1	22	0	6.5E-09	1.3E-08	1.0E-09	4.0E-08	4.6E-08
Phthalimide	85-41-6	16	0	1.2E-09	2.7E-09	0	5.9E-09	1.2E-08
Pyrene	129-00-0	35	2.6E-09	1.4E-08	9.1E-09	1.2E-08	2.9E-08	4.0E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	3.0E-11	1.7E-10	0	1.0E-11	1.0E-09
Selenium		4	0	1.8E-09	5.4E-09	0	1.2E-08	2.6E-08
Strontium		35	2.4E-08	2.5E-07	4.9E-07	1.3E-07	7.5E-07	2.8E-06
Thallium		21	0	7.0E-11	1.2E-10	5.7E-11	2.2E-10	6.5E-10
Tin		30	0	2.8E-09	2.8E-09	2.1E-09	7.4E-09	1.1E-08
Triethylene glycol monobutyl ether	143-22-6	3	0	3.4E-10	1.4E-09	0	1.8E-09	7.7E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	5.8E-10	1.0E-09	3.2E-10	9.4E-10	1.6E-09	1.9E-09
Vanadium		2	0	1.4E-09	5.9E-09	0	7.5E-09	2.5E-08
Zinc		35	1.4E-05	6.2E-05	3.3E-05	5.9E-05	1.1E-04	1.8E-04

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.



Table F-338. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Referees 30<40 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	1.1E-11	8.8E-11	8.2E-11	5.3E-11	2.6E-10	3.3E-10
Aluminum		35	2.0E-06	6.2E-06	3.2E-06	5.6E-06	1.1E-05	1.5E-05
Aniline	62-53-3	12	0	4.9E-10	1.1E-09	0	2.7E-09	4.9E-09
Anthracene	120-12-7	28	0	2.3E-10	4.1E-10	8.9E-11	8.7E-10	1.9E-09
Anthracene, 2-methyl-	613-12-7	35	1.3E-10	9.2E-10	8.1E-10	6.6E-10	2.1E-09	4.1E-09
Anthracene, 9,10-dimethyl	781-43-1	2	0	7.5E-12	4.1E-11	0	4.8E-12	2.4E-10
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	5.8E-11	2.6E-10	0	1.6E-10	1.5E-09
Anthracene, 9-phenyl	602-55-1	30	0	8.7E-11	9.2E-11	5.3E-11	2.7E-10	3.9E-10
Antimony		35	4.1E-10	6.7E-09	5.9E-09	5.7E-09	1.5E-08	2.8E-08
Barium		35	1.1E-07	4.3E-07	2.9E-07	4.2E-07	8.1E-07	1.6E-06
Benzene, n-butyl-	104-51-8	29	0	4.9E-11	1.3E-10	2.8E-11	7.6E-11	7.6E-10
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	1.3E-08	1.8E-08	5.0E-09	5.0E-08	6.5E-08
Benz[a]anthracene	56-55-3	35	5.3E-11	8.1E-10	8.9E-10	3.9E-10	2.5E-09	3.5E-09
Benzo[b]fluoranthene	205-99-2	35	2.3E-10	1.1E-09	8.2E-10	7.5E-10	2.7E-09	3.3E-09
7H-Benzo[c]fluorene	205-12-9	31	0	2.1E-10	3.5E-10	7.0E-11	8.7E-10	1.7E-09
Benzo[k]fluoranthene	207-08-9	35	3.0E-11	3.6E-10	3.4E-10	2.3E-10	1.1E-09	1.4E-09
Benzothiazole	95-16-9	35	3.1E-08	1.4E-07	9.4E-08	1.0E-07	2.8E-07	3.2E-07
Benzothiazole, 2-phenyl-	883-93-2	35	4.2E-09	1.5E-08	1.4E-08	1.1E-08	3.4E-08	7.9E-08
Benzothiazolone	934-34-9	35	2.2E-07	3.5E-07	6.8E-08	3.4E-07	4.5E-07	4.6E-07
Benzyl butyl phthalate	85-68-7	35	3.6E-10	7.0E-09	5.8E-09	6.2E-09	1.6E-08	2.8E-08
Beryllium		11	0	1.8E-10	2.8E-10	0	6.1E-10	9.7E-10
Butylated Hydroxytoluene	128-37-0	19	0	1.9E-10	3.5E-10	6.5E-11	9.4E-10	1.6E-09
Cadmium		34	0	4.8E-09	2.9E-09	4.3E-09	1.0E-08	1.1E-08
Chromium		35	2.4E-09	1.3E-08	1.3E-08	8.7E-09	2.9E-08	7.3E-08
Cobalt		35	5.6E-08	3.0E-07	2.1E-07	2.9E-07	5.8E-07	1.0E-06
Copper		35	2.3E-07	5.9E-07	3.1E-07	5.4E-07	9.7E-07	1.9E-06
Cyclohexyl isothiocyanate	1122-82-3	29	0	4.4E-08	3.2E-08	4.3E-08	1.0E-07	1.1E-07
Dibenz[a,h]anthracene	53-70-3	7	0	4.8E-11	1.2E-10	0	2.2E-10	6.0E-10
Dibenzothiophene	132-65-0	27	0	1.9E-10	2.8E-10	5.5E-11	7.5E-10	1.1E-09
Dibutyl phthalate	84-74-2	31	0	1.3E-08	1.0E-08	1.1E-08	3.2E-08	3.8E-08
Diethyl Phthalate	84-66-2	23	0	1.1E-09	9.8E-10	1.2E-09	2.6E-09	3.5E-09
Diisobutyl Phthalate	84-69-5	32	0	1.3E-09	3.4E-09	3.0E-10	3.6E-09	2.0E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Diisooctylphthalate	27554-26-3	33	0	2.8E-08	2.1E-08	2.1E-08	7.1E-08	7.4E-08
Di-n-octyl phthalate	117-84-0	26	0	2.8E-09	6.7E-09	7.3E-10	1.3E-08	2.9E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	2.3E-11	7.3E-11	0	1.8E-10	3.2E-10
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	1.4E-11	4.9E-09	9.9E-09	3.0E-10	2.0E-08	4.9E-08
Fluoranthene	206-44-0	35	8.5E-10	6.2E-09	6.1E-09	3.7E-09	1.6E-08	3.0E-08
Fluorene	86-73-7	11	0	8.1E-11	1.9E-10	0	4.0E-10	9.7E-10
Hexadecane	544-76-3	28	0	5.9E-10	4.9E-10	6.2E-10	1.4E-09	2.3E-09
1-Hydroxypyrene	5315-79-7	1	0	3.2E-10	1.9E-09	0	0	1.1E-08
Lead		35	5.4E-08	2.8E-07	2.4E-07	2.1E-07	7.0E-07	1.3E-06
Manganese		35	5.0E-08	2.8E-07	1.6E-07	2.9E-07	4.5E-07	8.8E-07
2-(Methylthio)benzothiazole	615-22-5	5	0	4.2E-09	1.5E-08	0	2.3E-08	8.4E-08
Molybdenum		14	0	5.1E-10	7.1E-10	0	1.6E-09	2.5E-09
Naphthalene	91-20-3	1	0	1.5E-11	8.7E-11	0	0	5.1E-10
Naphthalene, 1-methyl-	90-12-0	31	0	3.8E-11	1.3E-10	6.3E-12	7.9E-11	7.8E-10
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	8.8E-12	2.0E-11	0	4.4E-11	1.0E-10
Naphthalene, 1,6-dimethyl-	575-43-9	35	2.3E-11	5.8E-11	2.8E-11	5.1E-11	1.2E-10	1.6E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	35	5.4E-11	1.5E-09	7.2E-10	1.5E-09	2.7E-09	3.5E-09
Naphthalene, 2,3-dimethyl-	581-40-8	35	1.1E-11	4.5E-10	4.5E-10	3.1E-10	1.3E-09	2.0E-09
Naphthalene, 2-methyl	91-57-6	19	0	3.4E-11	1.4E-10	6.1E-12	7.7E-11	8.5E-10
1-Octadecene	112-88-9	32	0	1.3E-09	1.1E-09	1.0E-09	3.5E-09	4.0E-09
17-Pentatriacontene	6971-40-0	9	0	1.1E-09	2.6E-09	0	6.1E-09	1.2E-08
Phenanthrene	85-01-8	35	4.7E-10	2.7E-09	3.4E-09	1.5E-09	7.8E-09	1.8E-08
Phenanthrene, 1-methyl	832-69-9	35	6.3E-11	7.3E-10	7.0E-10	5.2E-10	1.8E-09	3.2E-09
Phenanthrene, 2-methyl-	2531-84-2	35	2.0E-11	9.4E-10	1.1E-09	5.6E-10	2.7E-09	5.2E-09
Phenanthrene, 3-methyl	832-71-3	35	1.2E-10	1.6E-09	1.7E-09	9.9E-10	4.3E-09	7.9E-09
N-Phenylbenzamide	93-98-1	22	0	6.1E-09	1.2E-08	9.4E-10	3.7E-08	4.3E-08
Phthalimide	85-41-6	16	0	1.1E-09	2.5E-09	0	5.5E-09	1.2E-08
Pyrene	129-00-0	35	2.4E-09	1.3E-08	8.5E-09	1.1E-08	2.8E-08	3.8E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	2.8E-11	1.6E-10	0	9.6E-12	9.5E-10
Selenium		4	0	1.7E-09	5.1E-09	0	1.2E-08	2.5E-08
Strontium		35	2.3E-08	2.4E-07	4.6E-07	1.3E-07	7.0E-07	2.7E-06
Thallium		21	0	6.6E-11	1.1E-10	5.3E-11	2.0E-10	6.1E-10
Tin		30	0	2.6E-09	2.6E-09	1.9E-09	6.9E-09	1.0E-08



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Triethylene glycol monobutyl ether	143-22-6	3	0	3.2E-10	1.3E-09	0	1.7E-09	7.2E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	5.5E-10	9.5E-10	3.0E-10	8.8E-10	1.5E-09	1.8E-09
Vanadium		2	0	1.3E-09	5.6E-09	0	7.0E-09	2.4E-08
Zinc		35	1.3E-05	5.8E-05	3.1E-05	5.6E-05	1.0E-04	1.7E-04

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-339. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender Referees 40<50 Years

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	1.1E-11	8.6E-11	8.0E-11	5.2E-11	2.5E-10	3.2E-10
Aluminum		35	1.9E-06	6.1E-06	3.1E-06	5.5E-06	1.1E-05	1.4E-05
Aniline	62-53-3	12	0	4.8E-10	1.1E-09	0	2.7E-09	4.8E-09
Anthracene	120-12-7	28	0	2.3E-10	4.0E-10	8.8E-11	8.5E-10	1.9E-09
Anthracene, 2-methyl-	613-12-7	35	1.3E-10	9.0E-10	7.9E-10	6.5E-10	2.1E-09	4.0E-09
Anthracene, 9,10-dimethyl	781-43-1	2	0	7.3E-12	4.1E-11	0	4.7E-12	2.4E-10
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	5.6E-11	2.6E-10	0	1.6E-10	1.4E-09
Anthracene, 9-phenyl	602-55-1	30	0	8.5E-11	9.0E-11	5.2E-11	2.7E-10	3.8E-10
Antimony		35	4.0E-10	6.6E-09	5.8E-09	5.6E-09	1.5E-08	2.7E-08
Barium		35	1.1E-07	4.2E-07	2.8E-07	4.1E-07	7.9E-07	1.6E-06
Benzene, n-butyl-	104-51-8	29	0	4.8E-11	1.2E-10	2.7E-11	7.4E-11	7.5E-10
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	1.3E-08	1.8E-08	4.9E-09	4.9E-08	6.4E-08
Benz[a]anthracene	56-55-3	35	5.2E-11	7.9E-10	8.7E-10	3.8E-10	2.5E-09	3.4E-09
Benzo[b]fluoranthene	205-99-2	35	2.2E-10	1.1E-09	8.0E-10	7.3E-10	2.6E-09	3.2E-09
7H-Benzo[c]fluorene	205-12-9	31	0	2.1E-10	3.4E-10	6.9E-11	8.5E-10	1.7E-09
Benzo[k]fluoranthene	207-08-9	35	3.0E-11	3.5E-10	3.3E-10	2.2E-10	1.0E-09	1.3E-09
Benzothiazole	95-16-9	35	3.1E-08	1.3E-07	9.2E-08	1.0E-07	2.8E-07	3.2E-07
Benzothiazole, 2-phenyl-	883-93-2	35	4.1E-09	1.4E-08	1.3E-08	1.0E-08	3.3E-08	7.7E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzothiazolone	934-34-9	35	2.1E-07	3.4E-07	6.6E-08	3.4E-07	4.4E-07	4.5E-07
Benzyl butyl phthalate	85-68-7	35	3.5E-10	6.8E-09	5.7E-09	6.0E-09	1.5E-08	2.7E-08
Beryllium		11	0	1.7E-10	2.8E-10	0	6.0E-10	9.5E-10
Butylated Hydroxytoluene	128-37-0	19	0	1.8E-10	3.4E-10	6.3E-11	9.2E-10	1.5E-09
Cadmium		34	0	4.7E-09	2.9E-09	4.2E-09	9.8E-09	1.1E-08
Chromium		35	2.4E-09	1.3E-08	1.3E-08	8.6E-09	2.8E-08	7.2E-08
Cobalt		35	5.5E-08	3.0E-07	2.0E-07	2.8E-07	5.6E-07	1.0E-06
Copper		35	2.3E-07	5.8E-07	3.0E-07	5.3E-07	9.5E-07	1.8E-06
Cyclohexyl isothiocyanate	1122-82-3	29	0	4.3E-08	3.1E-08	4.2E-08	1.0E-07	1.1E-07
Dibenz[a,h]anthracene	53-70-3	7	0	4.7E-11	1.2E-10	0	2.1E-10	5.8E-10
Dibenzothiophene	132-65-0	27	0	1.8E-10	2.8E-10	5.3E-11	7.4E-10	1.1E-09
Dibutyl phthalate	84-74-2	31	0	1.3E-08	9.8E-09	1.0E-08	3.2E-08	3.7E-08
Diethyl Phthalate	84-66-2	23	0	1.1E-09	9.6E-10	1.2E-09	2.6E-09	3.4E-09
Diisobutyl Phthalate	84-69-5	32	0	1.3E-09	3.4E-09	3.0E-10	3.5E-09	2.0E-08
Diisooctylphthalate	27554-26-3	33	0	2.8E-08	2.1E-08	2.0E-08	7.0E-08	7.3E-08
Di-n-octyl phthalate	117-84-0	26	0	2.7E-09	6.6E-09	7.2E-10	1.3E-08	2.9E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	2.3E-11	7.2E-11	0	1.7E-10	3.1E-10
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	1.3E-11	4.8E-09	9.7E-09	3.0E-10	2.0E-08	4.8E-08
Fluoranthene	206-44-0	35	8.3E-10	6.1E-09	5.9E-09	3.7E-09	1.5E-08	2.9E-08
Fluorene	86-73-7	11	0	8.0E-11	1.9E-10	0	4.0E-10	9.5E-10
Hexadecane	544-76-3	28	0	5.8E-10	4.8E-10	6.1E-10	1.4E-09	2.2E-09
1-Hydroxypyrene	5315-79-7	1	0	3.1E-10	1.8E-09	0	0	1.1E-08
Lead		35	5.3E-08	2.7E-07	2.3E-07	2.1E-07	6.8E-07	1.2E-06
Manganese		35	4.9E-08	2.7E-07	1.6E-07	2.8E-07	4.4E-07	8.6E-07
2-(Methylthio)benzothiazole	615-22-5	5	0	4.1E-09	1.5E-08	0	2.3E-08	8.3E-08
Molybdenum		14	0	5.0E-10	6.9E-10	0	1.5E-09	2.5E-09
Naphthalene	91-20-3	1	0	1.4E-11	8.5E-11	0	0	5.0E-10
Naphthalene, 1-methyl-	90-12-0	31	0	3.7E-11	1.3E-10	6.2E-12	7.8E-11	7.6E-10
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	8.6E-12	1.9E-11	0	4.3E-11	9.9E-11
Naphthalene, 1,6-dimethyl-	575-43-9	35	2.3E-11	5.7E-11	2.7E-11	5.0E-11	1.2E-10	1.6E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	35	5.3E-11	1.5E-09	7.1E-10	1.5E-09	2.6E-09	3.4E-09
Naphthalene, 2,3-dimethyl-	581-40-8	35	1.1E-11	4.4E-10	4.4E-10	3.0E-10	1.3E-09	1.9E-09
Naphthalene, 2-methyl	91-57-6	19	0	3.4E-11	1.4E-10	6.0E-12	7.5E-11	8.3E-10



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
1-Octadecene	112-88-9	32	0	1.3E-09	1.1E-09	9.9E-10	3.4E-09	3.9E-09
17-Pentatriacontene	6971-40-0	9	0	1.1E-09	2.5E-09	0	5.9E-09	1.1E-08
Phenanthrene	85-01-8	35	4.6E-10	2.7E-09	3.3E-09	1.5E-09	7.7E-09	1.7E-08
Phenanthrene, 1-methyl	832-69-9	35	6.2E-11	7.1E-10	6.9E-10	5.1E-10	1.8E-09	3.1E-09
Phenanthrene, 2-methyl-	2531-84-2	35	1.9E-11	9.2E-10	1.1E-09	5.5E-10	2.6E-09	5.0E-09
Phenanthrene, 3-methyl	832-71-3	35	1.2E-10	1.5E-09	1.6E-09	9.7E-10	4.2E-09	7.7E-09
N-Phenylbenzamide	93-98-1	22	0	6.0E-09	1.2E-08	9.2E-10	3.6E-08	4.2E-08
Phthalimide	85-41-6	16	0	1.1E-09	2.5E-09	0	5.4E-09	1.1E-08
Pyrene	129-00-0	35	2.4E-09	1.3E-08	8.4E-09	1.1E-08	2.7E-08	3.7E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	2.8E-11	1.6E-10	0	9.4E-12	9.3E-10
Selenium		4	0	1.6E-09	5.0E-09	0	1.1E-08	2.4E-08
Strontium		35	2.2E-08	2.3E-07	4.5E-07	1.2E-07	6.9E-07	2.6E-06
Thallium		21	0	6.5E-11	1.1E-10	5.2E-11	2.0E-10	5.9E-10
Tin		30	0	2.5E-09	2.5E-09	1.9E-09	6.8E-09	1.0E-08
Triethylene glycol monobutyl ether	143-22-6	3	0	3.1E-10	1.3E-09	0	1.6E-09	7.1E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	5.4E-10	9.3E-10	3.0E-10	8.6E-10	1.4E-09	1.8E-09
Vanadium		2	0	1.3E-09	5.4E-09	0	6.9E-09	2.3E-08
Zinc		35	1.3E-05	5.7E-05	3.0E-05	5.5E-05	1.0E-04	1.6E-04

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-340. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Referees 50<70 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	1.1E-11	8.6E-11	8.1E-11	5.2E-11	2.5E-10	3.2E-10
Aluminum		35	1.9E-06	6.1E-06	3.1E-06	5.5E-06	1.1E-05	1.4E-05
Aniline	62-53-3	12	0	4.8E-10	1.1E-09	0	2.7E-09	4.8E-09
Anthracene	120-12-7	28	0	2.3E-10	4.0E-10	8.8E-11	8.6E-10	1.9E-09
Anthracene, 2-methyl-	613-12-7	35	1.3E-10	9.0E-10	7.9E-10	6.5E-10	2.1E-09	4.0E-09
Anthracene, 9,10-dimethyl	781-43-1	2	0	7.3E-12	4.1E-11	0	4.8E-12	2.4E-10



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	5.7E-11	2.6E-10	0	1.6E-10	1.4E-09
Anthracene, 9-phenyl	602-55-1	30	0	8.6E-11	9.0E-11	5.2E-11	2.7E-10	3.8E-10
Antimony		35	4.0E-10	6.6E-09	5.8E-09	5.6E-09	1.5E-08	2.7E-08
Barium		35	1.1E-07	4.2E-07	2.9E-07	4.1E-07	7.9E-07	1.6E-06
Benzene, n-butyl-	104-51-8	29	0	4.8E-11	1.2E-10	2.7E-11	7.4E-11	7.5E-10
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	1.3E-08	1.8E-08	5.0E-09	4.9E-08	6.4E-08
Benz[a]anthracene	56-55-3	35	5.2E-11	7.9E-10	8.7E-10	3.8E-10	2.5E-09	3.5E-09
Benzo[b]fluoranthene	205-99-2	35	2.2E-10	1.1E-09	8.1E-10	7.3E-10	2.6E-09	3.2E-09
7H-Benzo[c]fluorene	205-12-9	31	0	2.1E-10	3.4E-10	6.9E-11	8.6E-10	1.7E-09
Benzo[k]fluoranthene	207-08-9	35	3.0E-11	3.5E-10	3.3E-10	2.2E-10	1.0E-09	1.3E-09
Benzothiazole	95-16-9	35	3.1E-08	1.3E-07	9.2E-08	1.0E-07	2.8E-07	3.2E-07
Benzothiazole, 2-phenyl-	883-93-2	35	4.1E-09	1.4E-08	1.3E-08	1.0E-08	3.3E-08	7.7E-08
Benzothiazolone	934-34-9	35	2.2E-07	3.4E-07	6.6E-08	3.4E-07	4.4E-07	4.5E-07
Benzyl butyl phthalate	85-68-7	35	3.5E-10	6.9E-09	5.7E-09	6.1E-09	1.5E-08	2.7E-08
Beryllium		11	0	1.7E-10	2.8E-10	0	6.0E-10	9.5E-10
Butylated Hydroxytoluene	128-37-0	19	0	1.8E-10	3.4E-10	6.4E-11	9.2E-10	1.5E-09
Cadmium		34	0	4.7E-09	2.9E-09	4.2E-09	9.9E-09	1.1E-08
Chromium		35	2.4E-09	1.3E-08	1.3E-08	8.6E-09	2.8E-08	7.2E-08
Cobalt		35	5.5E-08	3.0E-07	2.0E-07	2.8E-07	5.7E-07	1.0E-06
Copper		35	2.3E-07	5.8E-07	3.0E-07	5.3E-07	9.5E-07	1.8E-06
Cyclohexyl isothiocyanate	1122-82-3	29	0	4.3E-08	3.1E-08	4.2E-08	1.0E-07	1.1E-07
Dibenz[a,h]anthracene	53-70-3	7	0	4.7E-11	1.2E-10	0	2.1E-10	5.8E-10
Dibenzothiophene	132-65-0	27	0	1.8E-10	2.8E-10	5.4E-11	7.4E-10	1.1E-09
Dibutyl phthalate	84-74-2	31	0	1.3E-08	9.9E-09	1.0E-08	3.2E-08	3.7E-08
Diethyl Phthalate	84-66-2	23	0	1.1E-09	9.6E-10	1.2E-09	2.6E-09	3.4E-09
Diisobutyl Phthalate	84-69-5	32	0	1.3E-09	3.4E-09	3.0E-10	3.5E-09	2.0E-08
Diisooctylphthalate	27554-26-3	33	0	2.8E-08	2.1E-08	2.0E-08	7.0E-08	7.3E-08
Di-n-octyl phthalate	117-84-0	26	0	2.7E-09	6.6E-09	7.2E-10	1.3E-08	2.9E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	2.3E-11	7.2E-11	0	1.7E-10	3.1E-10
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	1.3E-11	4.8E-09	9.7E-09	3.0E-10	2.0E-08	4.8E-08
Fluoranthene	206-44-0	35	8.4E-10	6.1E-09	6.0E-09	3.7E-09	1.6E-08	2.9E-08
Fluorene	86-73-7	11	0	8.0E-11	1.9E-10	0	4.0E-10	9.5E-10
Hexadecane	544-76-3	28	0	5.8E-10	4.8E-10	6.1E-10	1.4E-09	2.2E-09
1-Hydroxypyrene	5315-79-7	1	0	3.1E-10	1.9E-09	0	0	1.1E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Lead		35	5.3E-08	2.7E-07	2.4E-07	2.1E-07	6.9E-07	1.2E-06
Manganese		35	4.9E-08	2.7E-07	1.6E-07	2.8E-07	4.4E-07	8.6E-07
2-(Methylthio)benzothiazole	615-22-5	5	0	4.1E-09	1.5E-08	0	2.3E-08	8.3E-08
Molybdenum		14	0	5.0E-10	6.9E-10	0	1.5E-09	2.5E-09
Naphthalene	91-20-3	1	0	1.4E-11	8.5E-11	0	0	5.0E-10
Naphthalene, 1-methyl-	90-12-0	31	0	3.7E-11	1.3E-10	6.2E-12	7.8E-11	7.7E-10
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	8.7E-12	1.9E-11	0	4.3E-11	9.9E-11
Naphthalene, 1,6-dimethyl-	575-43-9	35	2.3E-11	5.7E-11	2.7E-11	5.0E-11	1.2E-10	1.6E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	35	5.3E-11	1.5E-09	7.1E-10	1.5E-09	2.6E-09	3.4E-09
Naphthalene, 2,3-dimethyl-	581-40-8	35	1.1E-11	4.4E-10	4.4E-10	3.0E-10	1.3E-09	1.9E-09
Naphthalene, 2-methyl	91-57-6	19	0	3.4E-11	1.4E-10	6.0E-12	7.5E-11	8.3E-10
1-Octadecene	112-88-9	32	0	1.3E-09	1.1E-09	1.0E-09	3.4E-09	4.0E-09
17-Pentatriacontene	6971-40-0	9	0	1.1E-09	2.6E-09	0	5.9E-09	1.1E-08
Phenanthrene	85-01-8	35	4.7E-10	2.7E-09	3.3E-09	1.5E-09	7.7E-09	1.7E-08
Phenanthrene, 1-methyl	832-69-9	35	6.2E-11	7.2E-10	6.9E-10	5.1E-10	1.8E-09	3.1E-09
Phenanthrene, 2-methyl-	2531-84-2	35	1.9E-11	9.3E-10	1.1E-09	5.5E-10	2.6E-09	5.1E-09
Phenanthrene, 3-methyl	832-71-3	35	1.2E-10	1.5E-09	1.6E-09	9.7E-10	4.2E-09	7.7E-09
N-Phenylbenzamide	93-98-1	22	0	6.0E-09	1.2E-08	9.2E-10	3.6E-08	4.2E-08
Phthalimide	85-41-6	16	0	1.1E-09	2.5E-09	0	5.4E-09	1.1E-08
Pyrene	129-00-0	35	2.4E-09	1.3E-08	8.4E-09	1.1E-08	2.7E-08	3.7E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	2.8E-11	1.6E-10	0	9.4E-12	9.4E-10
Selenium		4	0	1.6E-09	5.0E-09	0	1.1E-08	2.4E-08
Strontium		35	2.2E-08	2.3E-07	4.5E-07	1.2E-07	6.9E-07	2.6E-06
Thallium		21	0	6.5E-11	1.1E-10	5.2E-11	2.0E-10	5.9E-10
Tin		30	0	2.5E-09	2.5E-09	1.9E-09	6.8E-09	1.0E-08
Triethylene glycol monobutyl ether	143-22-6	3	0	3.2E-10	1.3E-09	0	1.6E-09	7.1E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	5.4E-10	9.3E-10	3.0E-10	8.6E-10	1.4E-09	1.8E-09
Vanadium		2	0	1.3E-09	5.5E-09	0	6.9E-09	2.3E-08
Zinc		35	1.3E-05	5.7E-05	3.0E-05	5.5E-05	1.0E-04	1.6E-04

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not



included as the Study did not speciate the chemicals in the sample analysis.
Values are rounded to two significant figures.

Table F-341. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Spectators Third Trimester Fetus<0 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	9.6E-12	7.4E-11	6.9E-11	4.5E-11	2.2E-10	2.7E-10
Aluminum		35	1.6E-06	5.2E-06	2.7E-06	4.7E-06	9.5E-06	1.2E-05
Aniline	62-53-3	12	0	4.1E-10	9.4E-10	0	2.3E-09	4.2E-09
Anthracene	120-12-7	28	0	1.9E-10	3.4E-10	7.5E-11	7.3E-10	1.6E-09
Anthracene, 2-methyl-	613-12-7	35	1.1E-10	7.7E-10	6.8E-10	5.6E-10	1.8E-09	3.4E-09
Anthracene, 9,10-dimethyl	781-43-1	2	0	6.3E-12	3.5E-11	0	4.1E-12	2.1E-10
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	4.8E-11	2.2E-10	0	1.4E-10	1.2E-09
Anthracene, 9-phenyl	602-55-1	30	0	7.3E-11	7.7E-11	4.5E-11	2.3E-10	3.3E-10
Antimony		35	3.4E-10	5.6E-09	5.0E-09	4.8E-09	1.3E-08	2.4E-08
Barium		35	9.3E-08	3.6E-07	2.4E-07	3.5E-07	6.8E-07	1.3E-06
Benzene, n-butyl-	104-51-8	29	0	4.1E-11	1.1E-10	2.3E-11	6.4E-11	6.4E-10
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	1.1E-08	1.5E-08	4.2E-09	4.2E-08	5.5E-08
Benz[a]anthracene	56-55-3	35	4.4E-11	6.8E-10	7.4E-10	3.2E-10	2.1E-09	3.0E-09
Benzo[b]fluoranthene	205-99-2	35	1.9E-10	9.3E-10	6.9E-10	6.3E-10	2.3E-09	2.7E-09
7H-Benzo[c]fluorene	205-12-9	31	0	1.8E-10	2.9E-10	5.9E-11	7.3E-10	1.5E-09
Benzo[k]fluoranthene	207-08-9	35	2.5E-11	3.0E-10	2.9E-10	1.9E-10	8.9E-10	1.1E-09
Benzothiazole	95-16-9	35	2.6E-08	1.1E-07	7.9E-08	8.6E-08	2.4E-07	2.7E-07
Benzothiazole, 2-phenyl-	883-93-2	35	3.5E-09	1.2E-08	1.2E-08	8.9E-09	2.8E-08	6.6E-08
Benzothiazolone	934-34-9	35	1.8E-07	2.9E-07	5.7E-08	2.9E-07	3.8E-07	3.9E-07
Benzyl butyl phthalate	85-68-7	35	3.0E-10	5.9E-09	4.9E-09	5.2E-09	1.3E-08	2.3E-08
Beryllium		11	0	1.5E-10	2.4E-10	0	5.2E-10	8.2E-10
Butylated Hydroxytoluene	128-37-0	19	0	1.6E-10	2.9E-10	5.4E-11	7.9E-10	1.3E-09
Cadmium		34	0	4.0E-09	2.5E-09	3.6E-09	8.4E-09	9.3E-09
Chromium		35	2.1E-09	1.1E-08	1.1E-08	7.4E-09	2.4E-08	6.2E-08
Cobalt		35	4.7E-08	2.5E-07	1.7E-07	2.4E-07	4.8E-07	8.8E-07
Copper		35	2.0E-07	5.0E-07	2.6E-07	4.5E-07	8.1E-07	1.6E-06
Cyclohexyl isothiocyanate	1122-82-3	29	0	3.7E-08	2.7E-08	3.6E-08	8.6E-08	9.6E-08
Dibenz[a,h]anthracene	53-70-3	7	0	4.0E-11	1.0E-10	0	1.8E-10	5.0E-10
Dibenzothiophene	132-65-0	27	0	1.6E-10	2.4E-10	4.6E-11	6.3E-10	9.5E-10
Dibutyl phthalate	84-74-2	31	0	1.1E-08	8.4E-09	8.9E-09	2.7E-08	3.2E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Diethyl Phthalate	84-66-2	23	0	9.4E-10	8.3E-10	1.0E-09	2.2E-09	2.9E-09
Diisobutyl Phthalate	84-69-5	32	0	1.1E-09	2.9E-09	2.5E-10	3.0E-09	1.7E-08
Diisooctylphthalate	27554-26-3	33	0	2.4E-08	1.8E-08	1.7E-08	6.0E-08	6.3E-08
Di-n-octyl phthalate	117-84-0	26	0	2.3E-09	5.6E-09	6.1E-10	1.1E-08	2.5E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.9E-11	6.1E-11	0	1.5E-10	2.7E-10
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	1.1E-11	4.1E-09	8.3E-09	2.6E-10	1.7E-08	4.1E-08
Fluoranthene	206-44-0	35	7.2E-10	5.3E-09	5.1E-09	3.1E-09	1.3E-08	2.5E-08
Fluorene	86-73-7	11	0	6.8E-11	1.6E-10	0	3.4E-10	8.1E-10
Hexadecane	544-76-3	28	0	4.9E-10	4.2E-10	5.2E-10	1.2E-09	1.9E-09
1-Hydroxypyrene	5315-79-7	1	0	2.7E-10	1.6E-09	0	0	9.4E-09
Lead		35	4.5E-08	2.3E-07	2.0E-07	1.8E-07	5.9E-07	1.1E-06
Manganese		35	4.2E-08	2.3E-07	1.4E-07	2.4E-07	3.8E-07	7.4E-07
2-(Methylthio)benzothiazole	615-22-5	5	0	3.5E-09	1.3E-08	0	2.0E-08	7.1E-08
Molybdenum		14	0	4.3E-10	5.9E-10	0	1.3E-09	2.1E-09
Naphthalene	91-20-3	1	0	1.2E-11	7.3E-11	0	0	4.3E-10
Naphthalene, 1-methyl-	90-12-0	31	0	3.2E-11	1.1E-10	5.3E-12	6.7E-11	6.6E-10
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	7.4E-12	1.7E-11	0	3.7E-11	8.5E-11
Naphthalene, 1,6-dimethyl-	575-43-9	35	1.9E-11	4.9E-11	2.3E-11	4.3E-11	9.9E-11	1.4E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	35	4.6E-11	1.3E-09	6.1E-10	1.3E-09	2.3E-09	2.9E-09
Naphthalene, 2,3-dimethyl-	581-40-8	35	9.6E-12	3.8E-10	3.7E-10	2.6E-10	1.1E-09	1.6E-09
Naphthalene, 2-methyl	91-57-6	19	0	2.9E-11	1.2E-10	5.2E-12	6.5E-11	7.1E-10
1-Octadecene	112-88-9	32	0	1.1E-09	9.6E-10	8.5E-10	3.0E-09	3.4E-09
17-Pentatriacontene	6971-40-0	9	0	9.3E-10	2.2E-09	0	5.1E-09	9.8E-09
Phenanthrene	85-01-8	35	4.0E-10	2.3E-09	2.8E-09	1.3E-09	6.6E-09	1.5E-08
Phenanthrene, 1-methyl	832-69-9	35	5.3E-11	6.1E-10	5.9E-10	4.4E-10	1.5E-09	2.7E-09
Phenanthrene, 2-methyl-	2531-84-2	35	1.7E-11	7.9E-10	9.1E-10	4.7E-10	2.2E-09	4.3E-09
Phenanthrene, 3-methyl	832-71-3	35	1.0E-10	1.3E-09	1.4E-09	8.3E-10	3.6E-09	6.6E-09
N-Phenylbenzamide	93-98-1	22	0	5.1E-09	9.9E-09	7.9E-10	3.1E-08	3.6E-08
Phthalimide	85-41-6	16	0	9.2E-10	2.1E-09	0	4.7E-09	9.8E-09
Pyrene	129-00-0	35	2.1E-09	1.1E-08	7.2E-09	9.1E-09	2.3E-08	3.2E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	2.4E-11	1.4E-10	0	8.0E-12	8.0E-10
Selenium		4	0	1.4E-09	4.3E-09	0	9.7E-09	2.1E-08
Strontium		35	1.9E-08	2.0E-07	3.8E-07	1.1E-07	5.9E-07	2.2E-06



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Thallium		21	0	5.6E-11	9.5E-11	4.5E-11	1.7E-10	5.1E-10
Tin		30	0	2.2E-09	2.2E-09	1.6E-09	5.8E-09	8.6E-09
Triethylene glycol monobutyl ether	143-22-6	3	0	2.7E-10	1.1E-09	0	1.4E-09	6.1E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	4.6E-10	8.0E-10	2.5E-10	7.4E-10	1.2E-09	1.5E-09
Vanadium		2	0	1.1E-09	4.7E-09	0	5.9E-09	2.0E-08
Zinc		35	1.1E-05	4.9E-05	2.6E-05	4.7E-05	8.7E-05	1.4E-04

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-342. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Spectators 0<2 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	6.7E-10	5.1E-09	4.8E-09	3.1E-09	1.5E-08	1.9E-08
Aluminum		35	1.1E-04	3.6E-04	1.8E-04	3.3E-04	6.6E-04	8.5E-04
Aniline	62-53-3	12	0	2.9E-08	6.6E-08	0	1.6E-07	2.9E-07
Anthracene	120-12-7	28	0	1.4E-08	2.4E-08	5.2E-09	5.1E-08	1.1E-07
Anthracene, 2-methyl-	613-12-7	35	7.9E-09	5.4E-08	4.7E-08	3.9E-08	1.2E-07	2.4E-07
Anthracene, 9,10-dimethyl	781-43-1	2	0	4.4E-10	2.4E-09	0	2.8E-10	1.4E-08
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	3.4E-09	1.5E-08	0	9.6E-09	8.6E-08
Anthracene, 9-phenyl	602-55-1	30	0	5.1E-09	5.4E-09	3.1E-09	1.6E-08	2.3E-08
Antimony		35	2.4E-08	3.9E-07	3.5E-07	3.3E-07	9.0E-07	1.6E-06
Barium		35	6.5E-06	2.5E-05	1.7E-05	2.4E-05	4.7E-05	9.3E-05
Benzene, n-butyl-	104-51-8	29	0	2.9E-09	7.4E-09	1.6E-09	4.4E-09	4.5E-08
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	7.8E-07	1.1E-06	2.9E-07	2.9E-06	3.8E-06
Benz[a]anthracene	56-55-3	35	3.1E-09	4.7E-08	5.2E-08	2.3E-08	1.5E-07	2.1E-07
Benzo[b]fluoranthene	205-99-2	35	1.3E-08	6.5E-08	4.8E-08	4.4E-08	1.6E-07	1.9E-07
7H-Benzo[c]fluorene	205-12-9	31	0	1.3E-08	2.0E-08	4.1E-09	5.1E-08	1.0E-07
Benzo[k]fluoranthene	207-08-9	35	1.8E-09	2.1E-08	2.0E-08	1.3E-08	6.2E-08	7.9E-08
Benzothiazole	95-16-9	35	1.8E-06	8.0E-06	5.5E-06	5.9E-06	1.7E-05	1.9E-05



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzothiazole, 2-phenyl-	883-93-2	35	2.4E-07	8.6E-07	8.0E-07	6.2E-07	2.0E-06	4.6E-06
Benzothiazolone	934-34-9	35	1.3E-05	2.0E-05	4.0E-06	2.0E-05	2.6E-05	2.7E-05
Benzyl butyl phthalate	85-68-7	35	2.1E-08	4.1E-07	3.4E-07	3.6E-07	9.1E-07	1.6E-06
Beryllium		11	0	1.0E-08	1.7E-08	0	3.6E-08	5.7E-08
Butylated Hydroxytoluene	128-37-0	19	0	1.1E-08	2.0E-08	3.8E-09	5.5E-08	9.2E-08
Cadmium		34	0	2.8E-07	1.7E-07	2.5E-07	5.9E-07	6.4E-07
Chromium		35	1.4E-07	7.7E-07	7.6E-07	5.1E-07	1.7E-06	4.3E-06
Cobalt		35	3.3E-06	1.8E-05	1.2E-05	1.7E-05	3.4E-05	6.1E-05
Copper		35	1.4E-05	3.4E-05	1.8E-05	3.2E-05	5.7E-05	1.1E-04
Cyclohexyl isothiocyanate	1122-82-3	29	0	2.5E-06	1.9E-06	2.5E-06	6.0E-06	6.6E-06
Dibenz[a,h]anthracene	53-70-3	7	0	2.8E-09	7.0E-09	0	1.3E-08	3.5E-08
Dibenzothiophene	132-65-0	27	0	1.1E-08	1.6E-08	3.2E-09	4.4E-08	6.6E-08
Dibutyl phthalate	84-74-2	31	0	7.8E-07	5.9E-07	6.2E-07	1.9E-06	2.2E-06
Diethyl Phthalate	84-66-2	23	0	6.5E-08	5.7E-08	7.1E-08	1.5E-07	2.0E-07
Diisobutyl Phthalate	84-69-5	32	0	7.8E-08	2.0E-07	1.8E-08	2.1E-07	1.2E-06
Diisooctylphthalate	27554-26-3	33	0	1.6E-06	1.2E-06	1.2E-06	4.2E-06	4.3E-06
Di-n-octyl phthalate	117-84-0	26	0	1.6E-07	3.9E-07	4.3E-08	7.9E-07	1.7E-06
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.3E-09	4.3E-09	0	1.0E-08	1.9E-08
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	8.0E-10	2.9E-07	5.8E-07	1.8E-08	1.2E-06	2.9E-06
Fluoranthene	206-44-0	35	5.0E-08	3.6E-07	3.5E-07	2.2E-07	9.2E-07	1.7E-06
Fluorene	86-73-7	11	0	4.8E-09	1.1E-08	0	2.4E-08	5.7E-08
Hexadecane	544-76-3	28	0	3.4E-08	2.9E-08	3.6E-08	8.2E-08	1.3E-07
1-Hydroxypyrene	5315-79-7	1	0	1.9E-08	1.1E-07	0	0	6.5E-07
Lead		35	3.1E-06	1.6E-05	1.4E-05	1.2E-05	4.1E-05	7.3E-05
Manganese		35	2.9E-06	1.6E-05	9.5E-06	1.7E-05	2.6E-05	5.1E-05
2-(Methylthio)benzothiazole	615-22-5	5	0	2.4E-07	8.8E-07	0	1.4E-06	4.9E-06
Molybdenum		14	0	3.0E-08	4.1E-08	0	9.2E-08	1.5E-07
Naphthalene	91-20-3	1	0	8.5E-10	5.1E-09	0	0	3.0E-08
Naphthalene, 1-methyl-	90-12-0	31	0	2.2E-09	7.7E-09	3.7E-10	4.6E-09	4.6E-08
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	5.1E-10	1.2E-09	0	2.6E-09	5.9E-09
Naphthalene, 1,6-dimethyl-	575-43-9	35	1.4E-09	3.4E-09	1.6E-09	3.0E-09	6.9E-09	9.6E-09
Naphthalene, 2-(bromomethyl)-	939-26-4	35	3.2E-09	8.9E-08	4.2E-08	8.7E-08	1.6E-07	2.0E-07
Naphthalene, 2,3-dimethyl-	581-40-8	35	6.7E-10	2.6E-08	2.6E-08	1.8E-08	7.9E-08	1.1E-07



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene, 2-methyl	91-57-6	19	0	2.0E-09	8.3E-09	3.6E-10	4.5E-09	5.0E-08
1-Octadecene	112-88-9	32	0	7.6E-08	6.7E-08	5.9E-08	2.1E-07	2.4E-07
17-Pentatriacontene	6971-40-0	9	0	6.5E-08	1.5E-07	0	3.5E-07	6.8E-07
Phenanthrene	85-01-8	35	2.8E-08	1.6E-07	2.0E-07	9.0E-08	4.6E-07	1.0E-06
Phenanthrene, 1-methyl	832-69-9	35	3.7E-09	4.3E-08	4.1E-08	3.0E-08	1.1E-07	1.9E-07
Phenanthrene, 2-methyl-	2531-84-2	35	1.2E-09	5.5E-08	6.3E-08	3.3E-08	1.6E-07	3.0E-07
Phenanthrene, 3-methyl	832-71-3	35	7.0E-09	9.1E-08	9.8E-08	5.8E-08	2.5E-07	4.6E-07
N-Phenylbenzamide	93-98-1	22	0	3.6E-07	6.9E-07	5.5E-08	2.2E-06	2.5E-06
Phthalimide	85-41-6	16	0	6.4E-08	1.5E-07	0	3.2E-07	6.8E-07
Pyrene	129-00-0	35	1.4E-07	7.7E-07	5.0E-07	6.3E-07	1.6E-06	2.2E-06
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	1.6E-09	9.4E-09	0	5.6E-10	5.6E-08
Selenium		4	0	9.7E-08	3.0E-07	0	6.8E-07	1.4E-06
Strontium		35	1.3E-06	1.4E-05	2.7E-05	7.3E-06	4.1E-05	1.6E-04
Thallium		21	0	3.9E-09	6.6E-09	3.1E-09	1.2E-08	3.5E-08
Tin		30	0	1.5E-07	1.5E-07	1.1E-07	4.0E-07	6.0E-07
Triethylene glycol monobutyl ether	143-22-6	3	0	1.9E-08	7.6E-08	0	9.8E-08	4.2E-07
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	3.2E-08	5.6E-08	1.8E-08	5.1E-08	8.6E-08	1.1E-07
Vanadium		2	0	7.9E-08	3.2E-07	0	4.1E-07	1.4E-06
Zinc		35	7.5E-04	3.4E-03	1.8E-03	3.3E-03	6.1E-03	9.7E-03

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-343. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Spectators 2<6 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	3.2E-10	2.4E-09	2.3E-09	1.5E-09	7.1E-09	9.0E-09
Aluminum		35	5.4E-05	1.7E-04	8.7E-05	1.6E-04	3.1E-04	4.0E-04
Aniline	62-53-3	12	0	1.4E-08	3.1E-08	0	7.6E-08	1.4E-07
Anthracene	120-12-7	28	0	6.4E-09	1.1E-08	2.5E-09	2.4E-08	5.2E-08
Anthracene, 2-methyl-	613-12-7	35	3.7E-09	2.5E-08	2.2E-08	1.8E-08	5.9E-08	1.1E-07



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Anthracene, 9,10-dimethyl	781-43-1	2	0	2.1E-10	1.1E-09	0	1.3E-10	6.8E-09
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	1.6E-09	7.3E-09	0	4.5E-09	4.1E-08
Anthracene, 9-phenyl	602-55-1	30	0	2.4E-09	2.5E-09	1.5E-09	7.6E-09	1.1E-08
Antimony		35	1.1E-08	1.9E-07	1.6E-07	1.6E-07	4.3E-07	7.8E-07
Barium		35	3.1E-06	1.2E-05	8.0E-06	1.1E-05	2.2E-05	4.4E-05
Benzene, n-butyl-	104-51-8	29	0	1.4E-09	3.5E-09	7.6E-10	2.1E-09	2.1E-08
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	3.7E-07	5.1E-07	1.4E-07	1.4E-06	1.8E-06
Benz[a]anthracene	56-55-3	35	1.5E-09	2.2E-08	2.5E-08	1.1E-08	6.9E-08	9.7E-08
Benzo[b]fluoranthene	205-99-2	35	6.3E-09	3.1E-08	2.3E-08	2.1E-08	7.5E-08	9.0E-08
7H-Benzo[c]fluorene	205-12-9	31	0	5.9E-09	9.7E-09	1.9E-09	2.4E-08	4.8E-08
Benzo[k]fluoranthene	207-08-9	35	8.4E-10	1.0E-08	9.4E-09	6.3E-09	2.9E-08	3.7E-08
Benzothiazole	95-16-9	35	8.7E-07	3.8E-06	2.6E-06	2.8E-06	7.8E-06	9.0E-06
Benzothiazole, 2-phenyl-	883-93-2	35	1.2E-07	4.1E-07	3.8E-07	2.9E-07	9.3E-07	2.2E-06
Benzothiazolone	934-34-9	35	6.1E-06	9.6E-06	1.9E-06	9.5E-06	1.2E-05	1.3E-05
Benzyl butyl phthalate	85-68-7	35	9.9E-09	1.9E-07	1.6E-07	1.7E-07	4.3E-07	7.7E-07
Beryllium		11	0	4.9E-09	7.8E-09	0	1.7E-08	2.7E-08
Butylated Hydroxytoluene	128-37-0	19	0	5.2E-09	9.6E-09	1.8E-09	2.6E-08	4.3E-08
Cadmium		34	0	1.3E-07	8.1E-08	1.2E-07	2.8E-07	3.1E-07
Chromium		35	6.8E-08	3.7E-07	3.6E-07	2.4E-07	8.0E-07	2.0E-06
Cobalt		35	1.6E-06	8.3E-06	5.7E-06	7.9E-06	1.6E-05	2.9E-05
Copper		35	6.5E-06	1.6E-05	8.5E-06	1.5E-05	2.7E-05	5.2E-05
Cyclohexyl isothiocyanate	1122-82-3	29	0	1.2E-06	8.9E-07	1.2E-06	2.8E-06	3.2E-06
Dibenz[a,h]anthracene	53-70-3	7	0	1.3E-09	3.3E-09	0	6.0E-09	1.6E-08
Dibenzothiophene	132-65-0	27	0	5.2E-09	7.8E-09	1.5E-09	2.1E-08	3.1E-08
Dibutyl phthalate	84-74-2	31	0	3.7E-07	2.8E-07	2.9E-07	8.9E-07	1.1E-06
Diethyl Phthalate	84-66-2	23	0	3.1E-08	2.7E-08	3.4E-08	7.3E-08	9.6E-08
Diisobutyl Phthalate	84-69-5	32	0	3.7E-08	9.5E-08	8.3E-09	9.8E-08	5.6E-07
Diisooctylphthalate	27554-26-3	33	0	7.8E-07	5.8E-07	5.7E-07	2.0E-06	2.1E-06
Di-n-octyl phthalate	117-84-0	26	0	7.7E-08	1.9E-07	2.0E-08	3.7E-07	8.1E-07
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	6.4E-10	2.0E-09	0	4.9E-09	8.9E-09
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	3.8E-10	1.4E-07	2.7E-07	8.4E-09	5.6E-07	1.4E-06
Fluoranthene	206-44-0	35	2.4E-08	1.7E-07	1.7E-07	1.0E-07	4.4E-07	8.2E-07
Fluorene	86-73-7	11	0	2.3E-09	5.4E-09	0	1.1E-08	2.7E-08
Hexadecane	544-76-3	28	0	1.6E-08	1.4E-08	1.7E-08	3.9E-08	6.3E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
1-Hydroxypyrene	5315-79-7	1	0	8.8E-09	5.2E-08	0	0	3.1E-07
Lead		35	1.5E-06	7.7E-06	6.6E-06	5.8E-06	1.9E-05	3.5E-05
Manganese		35	1.4E-06	7.6E-06	4.5E-06	7.9E-06	1.2E-05	2.4E-05
2-(Methylthio)benzothiazole	615-22-5	5	0	1.2E-07	4.2E-07	0	6.5E-07	2.3E-06
Molybdenum		14	0	1.4E-08	2.0E-08	0	4.3E-08	7.0E-08
Naphthalene	91-20-3	1	0	4.1E-10	2.4E-09	0	0	1.4E-08
Naphthalene, 1-methyl-	90-12-0	31	0	1.0E-09	3.6E-09	1.7E-10	2.2E-09	2.2E-08
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	2.4E-10	5.5E-10	0	1.2E-09	2.8E-09
Naphthalene, 1,6-dimethyl-	575-43-9	35	6.4E-10	1.6E-09	7.7E-10	1.4E-09	3.3E-09	4.5E-09
Naphthalene, 2-(bromomethyl)-	939-26-4	35	1.5E-09	4.2E-08	2.0E-08	4.1E-08	7.4E-08	9.6E-08
Naphthalene, 2,3-dimethyl-	581-40-8	35	3.2E-10	1.2E-08	1.2E-08	8.6E-09	3.7E-08	5.4E-08
Naphthalene, 2-methyl	91-57-6	19	0	9.5E-10	4.0E-09	1.7E-10	2.1E-09	2.3E-08
1-Octadecene	112-88-9	32	0	3.6E-08	3.2E-08	2.8E-08	9.7E-08	1.1E-07
17-Pentatriacontene	6971-40-0	9	0	3.1E-08	7.2E-08	0	1.7E-07	3.2E-07
Phenanthrene	85-01-8	35	1.3E-08	7.6E-08	9.4E-08	4.3E-08	2.2E-07	4.8E-07
Phenanthrene, 1-methyl	832-69-9	35	1.7E-09	2.0E-08	1.9E-08	1.4E-08	5.0E-08	8.8E-08
Phenanthrene, 2-methyl-	2531-84-2	35	5.5E-10	2.6E-08	3.0E-08	1.6E-08	7.4E-08	1.4E-07
Phenanthrene, 3-methyl	832-71-3	35	3.3E-09	4.3E-08	4.6E-08	2.7E-08	1.2E-07	2.2E-07
N-Phenylbenzamide	93-98-1	22	0	1.7E-07	3.3E-07	2.6E-08	1.0E-06	1.2E-06
Phthalimide	85-41-6	16	0	3.0E-08	7.0E-08	0	1.5E-07	3.2E-07
Pyrene	129-00-0	35	6.8E-08	3.6E-07	2.4E-07	3.0E-07	7.7E-07	1.0E-06
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	7.8E-10	4.5E-09	0	2.6E-10	2.6E-08
Selenium		4	0	4.6E-08	1.4E-07	0	3.2E-07	6.8E-07
Strontium		35	6.2E-07	6.5E-06	1.3E-05	3.5E-06	1.9E-05	7.4E-05
Thallium		21	0	1.8E-09	3.1E-09	1.5E-09	5.7E-09	1.7E-08
Tin		30	0	7.2E-08	7.2E-08	5.3E-08	1.9E-07	2.8E-07
Triethylene glycol monobutyl ether	143-22-6	3	0	8.9E-09	3.6E-08	0	4.6E-08	2.0E-07
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	1.5E-08	2.6E-08	8.4E-09	2.4E-08	4.1E-08	5.0E-08
Vanadium		2	0	3.7E-08	1.5E-07	0	1.9E-07	6.6E-07
Zinc		35	3.5E-04	1.6E-03	8.5E-04	1.5E-03	2.9E-03	4.6E-03

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.



° CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.
Values are rounded to two significant figures.

Table F-344. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Spectators 6<11 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	2.2E-10	1.7E-09	1.6E-09	1.0E-09	5.0E-09	6.3E-09
Aluminum		35	3.8E-05	1.2E-04	6.1E-05	1.1E-04	2.2E-04	2.8E-04
Aniline	62-53-3	12	0	9.5E-09	2.2E-08	0	5.3E-08	9.6E-08
Anthracene	120-12-7	28	0	4.5E-09	7.9E-09	1.7E-09	1.7E-08	3.7E-08
Anthracene, 2-methyl-	613-12-7	35	2.6E-09	1.8E-08	1.6E-08	1.3E-08	4.1E-08	8.0E-08
Anthracene, 9,10-dimethyl	781-43-1	2	0	1.4E-10	8.0E-10	0	9.4E-11	4.8E-09
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	1.1E-09	5.1E-09	0	3.2E-09	2.9E-08
Anthracene, 9-phenyl	602-55-1	30	0	1.7E-09	1.8E-09	1.0E-09	5.3E-09	7.6E-09
Antimony		35	7.9E-09	1.3E-07	1.1E-07	1.1E-07	3.0E-07	5.4E-07
Barium		35	2.1E-06	8.3E-06	5.6E-06	8.1E-06	1.6E-05	3.1E-05
Benzene, n-butyl-	104-51-8	29	0	9.6E-10	2.4E-09	5.4E-10	1.5E-09	1.5E-08
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	2.6E-07	3.6E-07	9.8E-08	9.6E-07	1.3E-06
Benz[a]anthracene	56-55-3	35	1.0E-09	1.6E-08	1.7E-08	7.5E-09	4.9E-08	6.8E-08
Benzo[b]fluoranthene	205-99-2	35	4.4E-09	2.2E-08	1.6E-08	1.4E-08	5.2E-08	6.3E-08
7H-Benzo[c]fluorene	205-12-9	31	0	4.2E-09	6.8E-09	1.4E-09	1.7E-08	3.4E-08
Benzo[k]fluoranthene	207-08-9	35	5.9E-10	7.0E-09	6.6E-09	4.4E-09	2.0E-08	2.6E-08
Benzothiazole	95-16-9	35	6.1E-07	2.7E-06	1.8E-06	2.0E-06	5.5E-06	6.3E-06
Benzothiazole, 2-phenyl-	883-93-2	35	8.1E-08	2.8E-07	2.7E-07	2.1E-07	6.5E-07	1.5E-06
Benzothiazolone	934-34-9	35	4.3E-06	6.8E-06	1.3E-06	6.6E-06	8.7E-06	8.9E-06
Benzyl butyl phthalate	85-68-7	35	6.9E-09	1.4E-07	1.1E-07	1.2E-07	3.0E-07	5.4E-07
Beryllium		11	0	3.4E-09	5.5E-09	0	1.2E-08	1.9E-08
Butylated Hydroxytoluene	128-37-0	19	0	3.6E-09	6.8E-09	1.3E-09	1.8E-08	3.0E-08
Cadmium		34	0	9.2E-08	5.7E-08	8.3E-08	1.9E-07	2.1E-07
Chromium		35	4.7E-08	2.6E-07	2.5E-07	1.7E-07	5.6E-07	1.4E-06
Cobalt		35	1.1E-06	5.8E-06	4.0E-06	5.6E-06	1.1E-05	2.0E-05
Copper		35	4.5E-06	1.1E-05	5.9E-06	1.0E-05	1.9E-05	3.6E-05
Cyclohexyl isothiocyanate	1122-82-3	29	0	8.5E-07	6.2E-07	8.4E-07	2.0E-06	2.2E-06
Dibenz[a,h]anthracene	53-70-3	7	0	9.3E-10	2.3E-09	0	4.2E-09	1.2E-08
Dibenzothiophene	132-65-0	27	0	3.6E-09	5.5E-09	1.1E-09	1.5E-08	2.2E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dibutyl phthalate	84-74-2	31	0	2.6E-07	2.0E-07	2.1E-07	6.3E-07	7.4E-07
Diethyl Phthalate	84-66-2	23	0	2.2E-08	1.9E-08	2.4E-08	5.1E-08	6.7E-08
Diisobutyl Phthalate	84-69-5	32	0	2.6E-08	6.7E-08	5.9E-09	6.9E-08	3.9E-07
Diisooctylphthalate	27554-26-3	33	0	5.5E-07	4.1E-07	4.0E-07	1.4E-06	1.4E-06
Di-n-octyl phthalate	117-84-0	26	0	5.4E-08	1.3E-07	1.4E-08	2.6E-07	5.7E-07
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	4.5E-10	1.4E-09	0	3.4E-09	6.2E-09
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	2.6E-10	9.5E-08	1.9E-07	5.9E-09	3.9E-07	9.6E-07
Fluoranthene	206-44-0	35	1.7E-08	1.2E-07	1.2E-07	7.2E-08	3.1E-07	5.8E-07
Fluorene	86-73-7	11	0	1.6E-09	3.8E-09	0	7.9E-09	1.9E-08
Hexadecane	544-76-3	28	0	1.1E-08	9.6E-09	1.2E-08	2.7E-08	4.4E-08
1-Hydroxypyrene	5315-79-7	1	0	6.2E-09	3.7E-08	0	0	2.2E-07
Lead		35	1.0E-06	5.4E-06	4.7E-06	4.1E-06	1.4E-05	2.4E-05
Manganese		35	9.7E-07	5.3E-06	3.2E-06	5.5E-06	8.8E-06	1.7E-05
2-(Methylthio)benzothiazole	615-22-5	5	0	8.1E-08	2.9E-07	0	4.6E-07	1.6E-06
Molybdenum		14	0	9.9E-09	1.4E-08	0	3.0E-08	4.9E-08
Naphthalene	91-20-3	1	0	2.8E-10	1.7E-09	0	0	9.9E-09
Naphthalene, 1-methyl-	90-12-0	31	0	7.3E-10	2.5E-09	1.2E-10	1.5E-09	1.5E-08
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	1.7E-10	3.8E-10	0	8.6E-10	2.0E-09
Naphthalene, 1,6-dimethyl-	575-43-9	35	4.5E-10	1.1E-09	5.4E-10	9.9E-10	2.3E-09	3.2E-09
Naphthalene, 2-(bromomethyl)-	939-26-4	35	1.1E-09	3.0E-08	1.4E-08	2.9E-08	5.2E-08	6.7E-08
Naphthalene, 2,3-dimethyl-	581-40-8	35	2.2E-10	8.7E-09	8.7E-09	6.0E-09	2.6E-08	3.8E-08
Naphthalene, 2-methyl	91-57-6	19	0	6.6E-10	2.8E-09	1.2E-10	1.5E-09	1.6E-08
1-Octadecene	112-88-9	32	0	2.5E-08	2.2E-08	2.0E-08	6.8E-08	7.8E-08
17-Pentatriacontene	6971-40-0	9	0	2.1E-08	5.1E-08	0	1.2E-07	2.3E-07
Phenanthrene	85-01-8	35	9.2E-09	5.3E-08	6.6E-08	3.0E-08	1.5E-07	3.4E-07
Phenanthrene, 1-methyl	832-69-9	35	1.2E-09	1.4E-08	1.4E-08	1.0E-08	3.5E-08	6.2E-08
Phenanthrene, 2-methyl-	2531-84-2	35	3.8E-10	1.8E-08	2.1E-08	1.1E-08	5.2E-08	1.0E-07
Phenanthrene, 3-methyl	832-71-3	35	2.3E-09	3.0E-08	3.2E-08	1.9E-08	8.3E-08	1.5E-07
N-Phenylbenzamide	93-98-1	22	0	1.2E-07	2.3E-07	1.8E-08	7.2E-07	8.4E-07
Phthalimide	85-41-6	16	0	2.1E-08	4.9E-08	0	1.1E-07	2.3E-07
Pyrene	129-00-0	35	4.7E-08	2.6E-07	1.7E-07	2.1E-07	5.4E-07	7.3E-07
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	5.5E-10	3.1E-09	0	1.9E-10	1.9E-08
Selenium		4	0	3.2E-08	9.9E-08	0	2.3E-07	4.8E-07



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Strontium		35	4.4E-07	4.6E-06	8.9E-06	2.4E-06	1.4E-05	5.2E-05
Thallium		21	0	1.3E-09	2.2E-09	1.0E-09	4.0E-09	1.2E-08
Tin		30	0	5.0E-08	5.0E-08	3.7E-08	1.3E-07	2.0E-07
Triethylene glycol monobutyl ether	143-22-6	3	0	6.2E-09	2.5E-08	0	3.3E-08	1.4E-07
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	1.1E-08	1.8E-08	5.9E-09	1.7E-08	2.8E-08	3.5E-08
Vanadium		2	0	2.6E-08	1.1E-07	0	1.4E-07	4.6E-07
Zinc		35	2.5E-04	1.1E-03	6.0E-04	1.1E-03	2.0E-03	3.2E-03

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis. Values are rounded to two significant figures.

Table F-345. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Spectators 11<16 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	1.7E-11	1.3E-10	1.2E-10	8.1E-11	3.9E-10	5.0E-10
Aluminum		35	3.0E-06	9.4E-06	4.8E-06	8.6E-06	1.7E-05	2.2E-05
Aniline	62-53-3	12	0	7.5E-10	1.7E-09	0	4.2E-09	7.5E-09
Anthracene	120-12-7	28	0	3.5E-10	6.2E-10	1.4E-10	1.3E-09	2.9E-09
Anthracene, 2-methyl-	613-12-7	35	2.1E-10	1.4E-09	1.2E-09	1.0E-09	3.2E-09	6.2E-09
Anthracene, 9,10-dimethyl	781-43-1	2	0	1.1E-11	6.3E-11	0	7.4E-12	3.7E-10
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	8.8E-11	4.0E-10	0	2.5E-10	2.2E-09
Anthracene, 9-phenyl	602-55-1	30	0	1.3E-10	1.4E-10	8.1E-11	4.2E-10	6.0E-10
Antimony		35	6.2E-10	1.0E-08	9.0E-09	8.6E-09	2.3E-08	4.3E-08
Barium		35	1.7E-07	6.5E-07	4.4E-07	6.3E-07	1.2E-06	2.4E-06
Benzene, n-butyl-	104-51-8	29	0	7.5E-11	1.9E-10	4.2E-11	1.2E-10	1.2E-09
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	2.0E-08	2.8E-08	7.7E-09	7.5E-08	1.0E-07
Benz[a]anthracene	56-55-3	35	8.0E-11	1.2E-09	1.3E-09	5.9E-10	3.8E-09	5.4E-09
Benzo[b]fluoranthene	205-99-2	35	3.4E-10	1.7E-09	1.2E-09	1.1E-09	4.1E-09	5.0E-09
7H-Benzo[c]fluorene	205-12-9	31	0	3.3E-10	5.3E-10	1.1E-10	1.3E-09	2.6E-09
Benzo[k]fluoranthene	207-08-9	35	4.6E-11	5.5E-10	5.2E-10	3.5E-10	1.6E-09	2.1E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzothiazole	95-16-9	35	4.8E-08	2.1E-07	1.4E-07	1.5E-07	4.3E-07	4.9E-07
Benzothiazole, 2-phenyl-	883-93-2	35	6.4E-09	2.2E-08	2.1E-08	1.6E-08	5.1E-08	1.2E-07
Benzothiazolone	934-34-9	35	3.3E-07	5.3E-07	1.0E-07	5.2E-07	6.8E-07	7.0E-07
Benzyl butyl phthalate	85-68-7	35	5.4E-10	1.1E-08	8.8E-09	9.4E-09	2.4E-08	4.2E-08
Beryllium		11	0	2.7E-10	4.3E-10	0	9.3E-10	1.5E-09
Butylated Hydroxytoluene	128-37-0	19	0	2.8E-10	5.3E-10	9.9E-11	1.4E-09	2.4E-09
Cadmium		34	0	7.2E-09	4.4E-09	6.5E-09	1.5E-08	1.7E-08
Chromium		35	3.7E-09	2.0E-08	2.0E-08	1.3E-08	4.4E-08	1.1E-07
Cobalt		35	8.5E-08	4.6E-07	3.1E-07	4.4E-07	8.8E-07	1.6E-06
Copper		35	3.6E-07	9.0E-07	4.7E-07	8.2E-07	1.5E-06	2.8E-06
Cyclohexyl isothiocyanate	1122-82-3	29	0	6.6E-08	4.9E-08	6.6E-08	1.6E-07	1.7E-07
Dibenz[a,h]anthracene	53-70-3	7	0	7.3E-11	1.8E-10	0	3.3E-10	9.1E-10
Dibenzothiophene	132-65-0	27	0	2.8E-10	4.3E-10	8.3E-11	1.1E-09	1.7E-09
Dibutyl phthalate	84-74-2	31	0	2.0E-08	1.5E-08	1.6E-08	4.9E-08	5.8E-08
Diethyl Phthalate	84-66-2	23	0	1.7E-09	1.5E-09	1.8E-09	4.0E-09	5.3E-09
Diisobutyl Phthalate	84-69-5	32	0	2.0E-09	5.2E-09	4.6E-10	5.4E-09	3.1E-08
Diisooctylphthalate	27554-26-3	33	0	4.3E-08	3.2E-08	3.2E-08	1.1E-07	1.1E-07
Di-n-octyl phthalate	117-84-0	26	0	4.2E-09	1.0E-08	1.1E-09	2.1E-08	4.5E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	3.5E-11	1.1E-10	0	2.7E-10	4.9E-10
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	2.1E-11	7.5E-09	1.5E-08	4.6E-10	3.1E-08	7.5E-08
Fluoranthene	206-44-0	35	1.3E-09	9.5E-09	9.2E-09	5.7E-09	2.4E-08	4.5E-08
Fluorene	86-73-7	11	0	1.2E-10	3.0E-10	0	6.2E-10	1.5E-09
Hexadecane	544-76-3	28	0	8.9E-10	7.5E-10	9.5E-10	2.1E-09	3.5E-09
1-Hydroxypyrene	5315-79-7	1	0	4.9E-10	2.9E-09	0	0	1.7E-08
Lead		35	8.2E-08	4.2E-07	3.6E-07	3.2E-07	1.1E-06	1.9E-06
Manganese		35	7.6E-08	4.2E-07	2.5E-07	4.3E-07	6.9E-07	1.3E-06
2-(Methylthio)benzothiazole	615-22-5	5	0	6.4E-09	2.3E-08	0	3.6E-08	1.3E-07
Molybdenum		14	0	7.7E-10	1.1E-09	0	2.4E-09	3.8E-09
Naphthalene	91-20-3	1	0	2.2E-11	1.3E-10	0	0	7.8E-10
Naphthalene, 1-methyl-	90-12-0	31	0	5.7E-11	2.0E-10	9.6E-12	1.2E-10	1.2E-09
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	1.3E-11	3.0E-11	0	6.7E-11	1.5E-10
Naphthalene, 1,6-dimethyl-	575-43-9	35	3.5E-11	8.8E-11	4.2E-11	7.7E-11	1.8E-10	2.5E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	35	8.3E-11	2.3E-09	1.1E-09	2.3E-09	4.1E-09	5.3E-09



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene, 2,3-dimethyl-	581-40-8	35	1.7E-11	6.9E-10	6.8E-10	4.7E-10	2.0E-09	3.0E-09
Naphthalene, 2-methyl	91-57-6	19	0	5.2E-11	2.2E-10	9.3E-12	1.2E-10	1.3E-09
1-Octadecene	112-88-9	32	0	2.0E-09	1.7E-09	1.5E-09	5.3E-09	6.1E-09
17-Pentatriacontene	6971-40-0	9	0	1.7E-09	4.0E-09	0	9.2E-09	1.8E-08
Phenanthrene	85-01-8	35	7.2E-10	4.2E-09	5.2E-09	2.3E-09	1.2E-08	2.7E-08
Phenanthrene, 1-methyl	832-69-9	35	9.6E-11	1.1E-09	1.1E-09	7.9E-10	2.8E-09	4.8E-09
Phenanthrene, 2-methyl-	2531-84-2	35	3.0E-11	1.4E-09	1.7E-09	8.5E-10	4.0E-09	7.8E-09
Phenanthrene, 3-methyl	832-71-3	35	1.8E-10	2.4E-09	2.5E-09	1.5E-09	6.5E-09	1.2E-08
N-Phenylbenzamide	93-98-1	22	0	9.3E-09	1.8E-08	1.4E-09	5.7E-08	6.6E-08
Phthalimide	85-41-6	16	0	1.7E-09	3.8E-09	0	8.4E-09	1.8E-08
Pyrene	129-00-0	35	3.7E-09	2.0E-08	1.3E-08	1.6E-08	4.2E-08	5.8E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	4.3E-11	2.5E-10	0	1.5E-11	1.5E-09
Selenium		4	0	2.5E-09	7.8E-09	0	1.8E-08	3.7E-08
Strontium		35	3.4E-08	3.6E-07	7.0E-07	1.9E-07	1.1E-06	4.1E-06
Thallium		21	0	1.0E-10	1.7E-10	8.1E-11	3.1E-10	9.2E-10
Tin		30	0	3.9E-09	3.9E-09	2.9E-09	1.1E-08	1.6E-08
Triethylene glycol monobutyl ether	143-22-6	3	0	4.9E-10	2.0E-09	0	2.5E-09	1.1E-08
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	8.3E-10	1.4E-09	4.6E-10	1.3E-09	2.2E-09	2.8E-09
Vanadium		2	0	2.1E-09	8.5E-09	0	1.1E-08	3.6E-08
Zinc		35	1.9E-05	8.8E-05	4.7E-05	8.5E-05	1.6E-04	2.5E-04

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-346. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Spectators 16<30 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	9.9E-12	7.6E-11	7.1E-11	4.6E-11	2.2E-10	2.8E-10
Aluminum		35	1.7E-06	5.4E-06	2.7E-06	4.9E-06	9.8E-06	1.3E-05
Aniline	62-53-3	12	0	4.2E-10	9.7E-10	0	2.4E-09	4.3E-09
Anthracene	120-12-7	28	0	2.0E-10	3.5E-10	7.7E-11	7.5E-10	1.6E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Anthracene, 2-methyl-	613-12-7	35	1.2E-10	8.0E-10	7.0E-10	5.7E-10	1.8E-09	3.5E-09
Anthracene, 9,10-dimethyl	781-43-1	2	0	6.5E-12	3.6E-11	0	4.2E-12	2.1E-10
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	5.0E-11	2.3E-10	0	1.4E-10	1.3E-09
Anthracene, 9-phenyl	602-55-1	30	0	7.5E-11	8.0E-11	4.6E-11	2.4E-10	3.4E-10
Antimony		35	3.5E-10	5.8E-09	5.1E-09	4.9E-09	1.3E-08	2.4E-08
Barium		35	9.6E-08	3.7E-07	2.5E-07	3.6E-07	7.0E-07	1.4E-06
Benzene, n-butyl-	104-51-8	29	0	4.3E-11	1.1E-10	2.4E-11	6.6E-11	6.6E-10
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	1.2E-08	1.6E-08	4.4E-09	4.3E-08	5.7E-08
Benz[a]anthracene	56-55-3	35	4.6E-11	7.0E-10	7.7E-10	3.3E-10	2.2E-09	3.0E-09
Benzo[b]fluoranthene	205-99-2	35	2.0E-10	9.6E-10	7.1E-10	6.5E-10	2.3E-09	2.8E-09
7H-Benzo[c]fluorene	205-12-9	31	0	1.9E-10	3.0E-10	6.1E-11	7.5E-10	1.5E-09
Benzo[k]fluoranthene	207-08-9	35	2.6E-11	3.1E-10	2.9E-10	2.0E-10	9.1E-10	1.2E-09
Benzothiazole	95-16-9	35	2.7E-08	1.2E-07	8.1E-08	8.8E-08	2.4E-07	2.8E-07
Benzothiazole, 2-phenyl-	883-93-2	35	3.6E-09	1.3E-08	1.2E-08	9.2E-09	2.9E-08	6.8E-08
Benzothiazolone	934-34-9	35	1.9E-07	3.0E-07	5.9E-08	3.0E-07	3.9E-07	4.0E-07
Benzyl butyl phthalate	85-68-7	35	3.1E-10	6.0E-09	5.0E-09	5.3E-09	1.3E-08	2.4E-08
Beryllium		11	0	1.5E-10	2.5E-10	0	5.3E-10	8.4E-10
Butylated Hydroxytoluene	128-37-0	19	0	1.6E-10	3.0E-10	5.6E-11	8.1E-10	1.4E-09
Cadmium		34	0	4.1E-09	2.5E-09	3.7E-09	8.7E-09	9.5E-09
Chromium		35	2.1E-09	1.1E-08	1.1E-08	7.6E-09	2.5E-08	6.3E-08
Cobalt		35	4.8E-08	2.6E-07	1.8E-07	2.5E-07	5.0E-07	9.1E-07
Copper		35	2.0E-07	5.1E-07	2.6E-07	4.7E-07	8.4E-07	1.6E-06
Cyclohexyl isothiocyanate	1122-82-3	29	0	3.8E-08	2.8E-08	3.7E-08	8.8E-08	9.8E-08
Dibenz[a,h]anthracene	53-70-3	7	0	4.1E-11	1.0E-10	0	1.9E-10	5.2E-10
Dibenzothiophene	132-65-0	27	0	1.6E-10	2.4E-10	4.7E-11	6.5E-10	9.8E-10
Dibutyl phthalate	84-74-2	31	0	1.1E-08	8.7E-09	9.2E-09	2.8E-08	3.3E-08
Diethyl Phthalate	84-66-2	23	0	9.7E-10	8.5E-10	1.1E-09	2.3E-09	3.0E-09
Diisobutyl Phthalate	84-69-5	32	0	1.2E-09	3.0E-09	2.6E-10	3.1E-09	1.8E-08
Diisooctylphthalate	27554-26-3	33	0	2.4E-08	1.8E-08	1.8E-08	6.2E-08	6.4E-08
Di-n-octyl phthalate	117-84-0	26	0	2.4E-09	5.8E-09	6.3E-10	1.2E-08	2.5E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	2.0E-11	6.3E-11	0	1.5E-10	2.8E-10
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	1.2E-11	4.3E-09	8.6E-09	2.6E-10	1.8E-08	4.3E-08
Fluoranthene	206-44-0	35	7.4E-10	5.4E-09	5.3E-09	3.2E-09	1.4E-08	2.6E-08
Fluorene	86-73-7	11	0	7.0E-11	1.7E-10	0	3.5E-10	8.4E-10



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Hexadecane	544-76-3	28	0	5.1E-10	4.3E-10	5.4E-10	1.2E-09	2.0E-09
1-Hydroxypyrene	5315-79-7	1	0	2.8E-10	1.6E-09	0	0	9.7E-09
Lead		35	4.7E-08	2.4E-07	2.1E-07	1.8E-07	6.0E-07	1.1E-06
Manganese		35	4.3E-08	2.4E-07	1.4E-07	2.5E-07	3.9E-07	7.6E-07
2-(Methylthio)benzothiazole	615-22-5	5	0	3.6E-09	1.3E-08	0	2.0E-08	7.3E-08
Molybdenum		14	0	4.4E-10	6.1E-10	0	1.4E-09	2.2E-09
Naphthalene	91-20-3	1	0	1.3E-11	7.5E-11	0	0	4.4E-10
Naphthalene, 1-methyl-	90-12-0	31	0	3.3E-11	1.1E-10	5.4E-12	6.9E-11	6.8E-10
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	7.6E-12	1.7E-11	0	3.8E-11	8.7E-11
Naphthalene, 1,6-dimethyl-	575-43-9	35	2.0E-11	5.0E-11	2.4E-11	4.4E-11	1.0E-10	1.4E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	35	4.7E-11	1.3E-09	6.2E-10	1.3E-09	2.3E-09	3.0E-09
Naphthalene, 2,3-dimethyl-	581-40-8	35	9.9E-12	3.9E-10	3.9E-10	2.7E-10	1.2E-09	1.7E-09
Naphthalene, 2-methyl	91-57-6	19	0	3.0E-11	1.2E-10	5.3E-12	6.6E-11	7.3E-10
1-Octadecene	112-88-9	32	0	1.1E-09	9.9E-10	8.8E-10	3.0E-09	3.5E-09
17-Pentatriacontene	6971-40-0	9	0	9.6E-10	2.3E-09	0	5.2E-09	1.0E-08
Phenanthrene	85-01-8	35	4.1E-10	2.4E-09	2.9E-09	1.3E-09	6.8E-09	1.5E-08
Phenanthrene, 1-methyl	832-69-9	35	5.4E-11	6.3E-10	6.1E-10	4.5E-10	1.6E-09	2.7E-09
Phenanthrene, 2-methyl-	2531-84-2	35	1.7E-11	8.2E-10	9.4E-10	4.9E-10	2.3E-09	4.5E-09
Phenanthrene, 3-methyl	832-71-3	35	1.0E-10	1.3E-09	1.4E-09	8.6E-10	3.7E-09	6.8E-09
N-Phenylbenzamide	93-98-1	22	0	5.3E-09	1.0E-08	8.1E-10	3.2E-08	3.7E-08
Phthalimide	85-41-6	16	0	9.5E-10	2.2E-09	0	4.8E-09	1.0E-08
Pyrene	129-00-0	35	2.1E-09	1.1E-08	7.4E-09	9.4E-09	2.4E-08	3.3E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	2.4E-11	1.4E-10	0	8.3E-12	8.3E-10
Selenium		4	0	1.4E-09	4.4E-09	0	1.0E-08	2.1E-08
Strontium		35	2.0E-08	2.0E-07	4.0E-07	1.1E-07	6.1E-07	2.3E-06
Thallium		21	0	5.7E-11	9.7E-11	4.6E-11	1.8E-10	5.2E-10
Tin		30	0	2.2E-09	2.2E-09	1.7E-09	6.0E-09	8.8E-09
Triethylene glycol monobutyl ether	143-22-6	3	0	2.8E-10	1.1E-09	0	1.4E-09	6.2E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	4.7E-10	8.2E-10	2.6E-10	7.6E-10	1.3E-09	1.6E-09
Vanadium		2	0	1.2E-09	4.8E-09	0	6.1E-09	2.1E-08
Zinc		35	1.1E-05	5.0E-05	2.7E-05	4.8E-05	9.0E-05	1.4E-04

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.



^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-347. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Spectators 30<40 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	9.3E-12	7.2E-11	6.7E-11	4.4E-11	2.1E-10	2.7E-10
Aluminum		35	1.6E-06	5.1E-06	2.6E-06	4.6E-06	9.2E-06	1.2E-05
Aniline	62-53-3	12	0	4.0E-10	9.2E-10	0	2.2E-09	4.0E-09
Anthracene	120-12-7	28	0	1.9E-10	3.3E-10	7.3E-11	7.1E-10	1.6E-09
Anthracene, 2-methyl-	613-12-7	35	1.1E-10	7.5E-10	6.6E-10	5.4E-10	1.7E-09	3.4E-09
Anthracene, 9,10-dimethyl	781-43-1	2	0	6.1E-12	3.4E-11	0	4.0E-12	2.0E-10
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	4.7E-11	2.2E-10	0	1.3E-10	1.2E-09
Anthracene, 9-phenyl	602-55-1	30	0	7.1E-11	7.5E-11	4.3E-11	2.2E-10	3.2E-10
Antimony		35	3.3E-10	5.5E-09	4.8E-09	4.6E-09	1.3E-08	2.3E-08
Barium		35	9.1E-08	3.5E-07	2.4E-07	3.4E-07	6.6E-07	1.3E-06
Benzene, n-butyl-	104-51-8	29	0	4.0E-11	1.0E-10	2.3E-11	6.2E-11	6.2E-10
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	1.1E-08	1.5E-08	4.1E-09	4.1E-08	5.4E-08
Benz[a]anthracene	56-55-3	35	4.3E-11	6.6E-10	7.2E-10	3.2E-10	2.1E-09	2.9E-09
Benzo[b]fluoranthene	205-99-2	35	1.9E-10	9.1E-10	6.7E-10	6.1E-10	2.2E-09	2.7E-09
7H-Benzo[c]fluorene	205-12-9	31	0	1.8E-10	2.9E-10	5.7E-11	7.1E-10	1.4E-09
Benzo[k]fluoranthene	207-08-9	35	2.5E-11	2.9E-10	2.8E-10	1.9E-10	8.6E-10	1.1E-09
Benzothiazole	95-16-9	35	2.6E-08	1.1E-07	7.7E-08	8.3E-08	2.3E-07	2.7E-07
Benzothiazole, 2-phenyl-	883-93-2	35	3.4E-09	1.2E-08	1.1E-08	8.7E-09	2.8E-08	6.4E-08
Benzothiazolone	934-34-9	35	1.8E-07	2.9E-07	5.5E-08	2.8E-07	3.7E-07	3.8E-07
Benzyl butyl phthalate	85-68-7	35	2.9E-10	5.7E-09	4.7E-09	5.1E-09	1.3E-08	2.3E-08
Beryllium		11	0	1.4E-10	2.3E-10	0	5.0E-10	8.0E-10
Butylated Hydroxytoluene	128-37-0	19	0	1.5E-10	2.8E-10	5.3E-11	7.7E-10	1.3E-09
Cadmium		34	0	3.9E-09	2.4E-09	3.5E-09	8.2E-09	9.0E-09
Chromium		35	2.0E-09	1.1E-08	1.1E-08	7.2E-09	2.4E-08	6.0E-08
Cobalt		35	4.6E-08	2.5E-07	1.7E-07	2.3E-07	4.7E-07	8.6E-07
Copper		35	1.9E-07	4.8E-07	2.5E-07	4.4E-07	7.9E-07	1.5E-06
Cyclohexyl isothiocyanate	1122-82-3	29	0	3.6E-08	2.6E-08	3.5E-08	8.4E-08	9.3E-08



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dibenz[a,h]anthracene	53-70-3	7	0	3.9E-11	9.8E-11	0	1.8E-10	4.9E-10
Dibenzothiophene	132-65-0	27	0	1.5E-10	2.3E-10	4.5E-11	6.2E-10	9.3E-10
Dibutyl phthalate	84-74-2	31	0	1.1E-08	8.2E-09	8.7E-09	2.6E-08	3.1E-08
Diethyl Phthalate	84-66-2	23	0	9.1E-10	8.0E-10	9.9E-10	2.2E-09	2.8E-09
Diisobutyl Phthalate	84-69-5	32	0	1.1E-09	2.8E-09	2.5E-10	2.9E-09	1.7E-08
Diisooctylphthalate	27554-26-3	33	0	2.3E-08	1.7E-08	1.7E-08	5.8E-08	6.1E-08
Di-n-octyl phthalate	117-84-0	26	0	2.3E-09	5.5E-09	6.0E-10	1.1E-08	2.4E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.9E-11	6.0E-11	0	1.4E-10	2.6E-10
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	1.1E-11	4.0E-09	8.1E-09	2.5E-10	1.7E-08	4.0E-08
Fluoranthene	206-44-0	35	7.0E-10	5.1E-09	5.0E-09	3.1E-09	1.3E-08	2.4E-08
Fluorene	86-73-7	11	0	6.7E-11	1.6E-10	0	3.3E-10	7.9E-10
Hexadecane	544-76-3	28	0	4.8E-10	4.0E-10	5.1E-10	1.1E-09	1.9E-09
1-Hydroxypyrene	5315-79-7	1	0	2.6E-10	1.5E-09	0	0	9.1E-09
Lead		35	4.4E-08	2.3E-07	2.0E-07	1.7E-07	5.7E-07	1.0E-06
Manganese		35	4.1E-08	2.3E-07	1.3E-07	2.3E-07	3.7E-07	7.2E-07
2-(Methylthio)benzothiazole	615-22-5	5	0	3.4E-09	1.2E-08	0	1.9E-08	6.9E-08
Molybdenum		14	0	4.2E-10	5.8E-10	0	1.3E-09	2.1E-09
Naphthalene	91-20-3	1	0	1.2E-11	7.1E-11	0	0	4.2E-10
Naphthalene, 1-methyl-	90-12-0	31	0	3.1E-11	1.1E-10	5.2E-12	6.5E-11	6.4E-10
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	7.2E-12	1.6E-11	0	3.6E-11	8.3E-11
Naphthalene, 1,6-dimethyl-	575-43-9	35	1.9E-11	4.8E-11	2.3E-11	4.2E-11	9.7E-11	1.3E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	35	4.4E-11	1.3E-09	5.9E-10	1.2E-09	2.2E-09	2.8E-09
Naphthalene, 2,3-dimethyl-	581-40-8	35	9.3E-12	3.7E-10	3.6E-10	2.5E-10	1.1E-09	1.6E-09
Naphthalene, 2-methyl	91-57-6	19	0	2.8E-11	1.2E-10	5.0E-12	6.3E-11	6.9E-10
1-Octadecene	112-88-9	32	0	1.1E-09	9.3E-10	8.3E-10	2.9E-09	3.3E-09
17-Pentatriacontene	6971-40-0	9	0	9.0E-10	2.1E-09	0	5.0E-09	9.6E-09
Phenanthrene	85-01-8	35	3.9E-10	2.2E-09	2.8E-09	1.3E-09	6.4E-09	1.4E-08
Phenanthrene, 1-methyl	832-69-9	35	5.2E-11	6.0E-10	5.7E-10	4.2E-10	1.5E-09	2.6E-09
Phenanthrene, 2-methyl-	2531-84-2	35	1.6E-11	7.7E-10	8.9E-10	4.6E-10	2.2E-09	4.2E-09
Phenanthrene, 3-methyl	832-71-3	35	9.8E-11	1.3E-09	1.4E-09	8.1E-10	3.5E-09	6.4E-09
N-Phenylbenzamide	93-98-1	22	0	5.0E-09	9.7E-09	7.6E-10	3.0E-08	3.5E-08
Phthalimide	85-41-6	16	0	9.0E-10	2.1E-09	0	4.5E-09	9.5E-09
Pyrene	129-00-0	35	2.0E-09	1.1E-08	7.0E-09	8.9E-09	2.3E-08	3.1E-08



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	2.3E-11	1.3E-10	0	7.8E-12	7.8E-10
Selenium		4	0	1.4E-09	4.2E-09	0	9.5E-09	2.0E-08
Strontium		35	1.8E-08	1.9E-07	3.7E-07	1.0E-07	5.7E-07	2.2E-06
Thallium		21	0	5.4E-11	9.2E-11	4.3E-11	1.7E-10	5.0E-10
Tin		30	0	2.1E-09	2.1E-09	1.6E-09	5.7E-09	8.3E-09
Triethylene glycol monobutyl ether	143-22-6	3	0	2.6E-10	1.1E-09	0	1.4E-09	5.9E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	4.5E-10	7.8E-10	2.5E-10	7.2E-10	1.2E-09	1.5E-09
Vanadium		2	0	1.1E-09	4.5E-09	0	5.7E-09	1.9E-08
Zinc		35	1.0E-05	4.7E-05	2.5E-05	4.6E-05	8.5E-05	1.4E-04

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-348. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b—Combined Gender Spectators 40<50 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	8.8E-12	6.8E-11	6.4E-11	4.1E-11	2.0E-10	2.5E-10
Aluminum		35	1.5E-06	4.8E-06	2.4E-06	4.4E-06	8.7E-06	1.1E-05
Aniline	62-53-3	12	0	3.8E-10	8.7E-10	0	2.1E-09	3.8E-09
Anthracene	120-12-7	28	0	1.8E-10	3.1E-10	6.9E-11	6.7E-10	1.5E-09
Anthracene, 2-methyl-	613-12-7	35	1.0E-10	7.1E-10	6.3E-10	5.1E-10	1.6E-09	3.2E-09
Anthracene, 9,10-dimethyl	781-43-1	2	0	5.8E-12	3.2E-11	0	3.7E-12	1.9E-10
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	4.5E-11	2.0E-10	0	1.3E-10	1.1E-09
Anthracene, 9-phenyl	602-55-1	30	0	6.8E-11	7.1E-11	4.1E-11	2.1E-10	3.0E-10
Antimony		35	3.2E-10	5.2E-09	4.6E-09	4.4E-09	1.2E-08	2.2E-08
Barium		35	8.6E-08	3.3E-07	2.3E-07	3.2E-07	6.2E-07	1.2E-06
Benzene, n-butyl-	104-51-8	29	0	3.8E-11	9.8E-11	2.1E-11	5.9E-11	5.9E-10
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	1.0E-08	1.4E-08	3.9E-09	3.8E-08	5.1E-08
Benz[a]anthracene	56-55-3	35	4.1E-11	6.3E-10	6.9E-10	3.0E-10	1.9E-09	2.7E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[b]fluoranthene	205-99-2	35	1.8E-10	8.6E-10	6.4E-10	5.8E-10	2.1E-09	2.5E-09
7H-Benzo[c]fluorene	205-12-9	31	0	1.7E-10	2.7E-10	5.4E-11	6.8E-10	1.3E-09
Benzo[k]fluoranthene	207-08-9	35	2.3E-11	2.8E-10	2.6E-10	1.8E-10	8.2E-10	1.0E-09
Benzothiazole	95-16-9	35	2.4E-08	1.1E-07	7.3E-08	7.9E-08	2.2E-07	2.5E-07
Benzothiazole, 2-phenyl-	883-93-2	35	3.2E-09	1.1E-08	1.1E-08	8.2E-09	2.6E-08	6.1E-08
Benzothiazolone	934-34-9	35	1.7E-07	2.7E-07	5.2E-08	2.7E-07	3.5E-07	3.6E-07
Benzyl butyl phthalate	85-68-7	35	2.8E-10	5.4E-09	4.5E-09	4.8E-09	1.2E-08	2.2E-08
Beryllium		11	0	1.4E-10	2.2E-10	0	4.8E-10	7.5E-10
Butylated Hydroxytoluene	128-37-0	19	0	1.4E-10	2.7E-10	5.0E-11	7.3E-10	1.2E-09
Cadmium		34	0	3.7E-09	2.3E-09	3.3E-09	7.8E-09	8.5E-09
Chromium		35	1.9E-09	1.0E-08	1.0E-08	6.8E-09	2.2E-08	5.7E-08
Cobalt		35	4.3E-08	2.3E-07	1.6E-07	2.2E-07	4.5E-07	8.1E-07
Copper		35	1.8E-07	4.6E-07	2.4E-07	4.2E-07	7.5E-07	1.4E-06
Cyclohexyl isothiocyanate	1122-82-3	29	0	3.4E-08	2.5E-08	3.4E-08	7.9E-08	8.8E-08
Dibenz[a,h]anthracene	53-70-3	7	0	3.7E-11	9.2E-11	0	1.7E-10	4.6E-10
Dibenzothiophene	132-65-0	27	0	1.4E-10	2.2E-10	4.2E-11	5.8E-10	8.8E-10
Dibutyl phthalate	84-74-2	31	0	1.0E-08	7.8E-09	8.2E-09	2.5E-08	2.9E-08
Diethyl Phthalate	84-66-2	23	0	8.7E-10	7.6E-10	9.4E-10	2.0E-09	2.7E-09
Diisobutyl Phthalate	84-69-5	32	0	1.0E-09	2.7E-09	2.3E-10	2.8E-09	1.6E-08
Diisooctylphthalate	27554-26-3	33	0	2.2E-08	1.6E-08	1.6E-08	5.5E-08	5.8E-08
Di-n-octyl phthalate	117-84-0	26	0	2.1E-09	5.2E-09	5.7E-10	1.0E-08	2.3E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.8E-11	5.7E-11	0	1.4E-10	2.5E-10
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	1.1E-11	3.8E-09	7.7E-09	2.4E-10	1.6E-08	3.8E-08
Fluoranthene	206-44-0	35	6.6E-10	4.8E-09	4.7E-09	2.9E-09	1.2E-08	2.3E-08
Fluorene	86-73-7	11	0	6.3E-11	1.5E-10	0	3.1E-10	7.5E-10
Hexadecane	544-76-3	28	0	4.6E-10	3.8E-10	4.8E-10	1.1E-09	1.8E-09
1-Hydroxypyrene	5315-79-7	1	0	2.5E-10	1.5E-09	0	0	8.6E-09
Lead		35	4.2E-08	2.2E-07	1.9E-07	1.6E-07	5.4E-07	9.7E-07
Manganese		35	3.9E-08	2.1E-07	1.3E-07	2.2E-07	3.5E-07	6.8E-07
2-(Methylthio)benzothiazole	615-22-5	5	0	3.2E-09	1.2E-08	0	1.8E-08	6.5E-08
Molybdenum		14	0	3.9E-10	5.5E-10	0	1.2E-09	2.0E-09
Naphthalene	91-20-3	1	0	1.1E-11	6.7E-11	0	0	4.0E-10
Naphthalene, 1-methyl-	90-12-0	31	0	2.9E-11	1.0E-10	4.9E-12	6.1E-11	6.0E-10
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	6.8E-12	1.5E-11	0	3.4E-11	7.8E-11



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene, 1,6-dimethyl-	575-43-9	35	1.8E-11	4.5E-11	2.2E-11	3.9E-11	9.1E-11	1.3E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	35	4.2E-11	1.2E-09	5.6E-10	1.2E-09	2.1E-09	2.7E-09
Naphthalene, 2,3-dimethyl-	581-40-8	35	8.9E-12	3.5E-10	3.5E-10	2.4E-10	1.0E-09	1.5E-09
Naphthalene, 2-methyl	91-57-6	19	0	2.7E-11	1.1E-10	4.7E-12	6.0E-11	6.6E-10
1-Octadecene	112-88-9	32	0	1.0E-09	8.8E-10	7.9E-10	2.7E-09	3.1E-09
17-Pentatriacontene	6971-40-0	9	0	8.6E-10	2.0E-09	0	4.7E-09	9.1E-09
Phenanthrene	85-01-8	35	3.7E-10	2.1E-09	2.6E-09	1.2E-09	6.1E-09	1.4E-08
Phenanthrene, 1-methyl	832-69-9	35	4.9E-11	5.6E-10	5.4E-10	4.0E-10	1.4E-09	2.5E-09
Phenanthrene, 2-methyl-	2531-84-2	35	1.5E-11	7.3E-10	8.4E-10	4.3E-10	2.1E-09	4.0E-09
Phenanthrene, 3-methyl	832-71-3	35	9.3E-11	1.2E-09	1.3E-09	7.7E-10	3.3E-09	6.1E-09
N-Phenylbenzamide	93-98-1	22	0	4.7E-09	9.2E-09	7.2E-10	2.9E-08	3.3E-08
Phthalimide	85-41-6	16	0	8.5E-10	2.0E-09	0	4.3E-09	9.0E-09
Pyrene	129-00-0	35	1.9E-09	1.0E-08	6.6E-09	8.4E-09	2.1E-08	2.9E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	2.2E-11	1.2E-10	0	7.4E-12	7.4E-10
Selenium		4	0	1.3E-09	4.0E-09	0	9.0E-09	1.9E-08
Strontium		35	1.7E-08	1.8E-07	3.5E-07	9.7E-08	5.4E-07	2.1E-06
Thallium		21	0	5.1E-11	8.7E-11	4.1E-11	1.6E-10	4.7E-10
Tin		30	0	2.0E-09	2.0E-09	1.5E-09	5.4E-09	7.9E-09
Triethylene glycol monobutyl ether	143-22-6	3	0	2.5E-10	1.0E-09	0	1.3E-09	5.6E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	4.2E-10	7.4E-10	2.3E-10	6.8E-10	1.1E-09	1.4E-09
Vanadium		2	0	1.0E-09	4.3E-09	0	5.4E-09	1.8E-08
Zinc		35	9.9E-06	4.5E-05	2.4E-05	4.3E-05	8.0E-05	1.3E-04

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-349. Field-Specific Average Ingestion Daily Dose^a for Chronic Non-Cancer Hazard (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related General Chemicals^b**—Combined Gender **Spectators 50<70 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Acenaphthylene	208-96-8	35	8.9E-12	6.8E-11	6.4E-11	4.1E-11	2.0E-10	2.5E-10



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aluminum		35	1.5E-06	4.8E-06	2.4E-06	4.4E-06	8.8E-06	1.1E-05
Aniline	62-53-3	12	0	3.8E-10	8.7E-10	0	2.1E-09	3.8E-09
Anthracene	120-12-7	28	0	1.8E-10	3.2E-10	6.9E-11	6.8E-10	1.5E-09
Anthracene, 2-methyl-	613-12-7	35	1.0E-10	7.2E-10	6.3E-10	5.2E-10	1.6E-09	3.2E-09
Anthracene, 9,10-dimethyl	781-43-1	2	0	5.8E-12	3.2E-11	0	3.8E-12	1.9E-10
Anthracene, 9,10-diphenyl-	1499-10-1	2	0	4.5E-11	2.0E-10	0	1.3E-10	1.1E-09
Anthracene, 9-phenyl	602-55-1	30	0	6.8E-11	7.1E-11	4.1E-11	2.1E-10	3.0E-10
Antimony		35	3.2E-10	5.2E-09	4.6E-09	4.4E-09	1.2E-08	2.2E-08
Barium		35	8.6E-08	3.3E-07	2.3E-07	3.2E-07	6.3E-07	1.2E-06
Benzene, n-butyl-	104-51-8	29	0	3.8E-11	9.8E-11	2.1E-11	5.9E-11	5.9E-10
1,4-Benzenediamine, N-(1,3-dimethylbutyl)-N'-phenyl-	793-24-8	20	0	1.0E-08	1.4E-08	3.9E-09	3.8E-08	5.1E-08
Benz[a]anthracene	56-55-3	35	4.1E-11	6.3E-10	6.9E-10	3.0E-10	1.9E-09	2.7E-09
Benzo[b]fluoranthene	205-99-2	35	1.8E-10	8.6E-10	6.4E-10	5.8E-10	2.1E-09	2.5E-09
7H-Benzo[c]fluorene	205-12-9	31	0	1.7E-10	2.7E-10	5.5E-11	6.8E-10	1.3E-09
Benzo[k]fluoranthene	207-08-9	35	2.4E-11	2.8E-10	2.6E-10	1.8E-10	8.2E-10	1.1E-09
Benzothiazole	95-16-9	35	2.4E-08	1.1E-07	7.3E-08	7.9E-08	2.2E-07	2.5E-07
Benzothiazole, 2-phenyl-	883-93-2	35	3.2E-09	1.1E-08	1.1E-08	8.3E-09	2.6E-08	6.1E-08
Benzothiazolone	934-34-9	35	1.7E-07	2.7E-07	5.3E-08	2.7E-07	3.5E-07	3.6E-07
Benzyl butyl phthalate	85-68-7	35	2.8E-10	5.4E-09	4.5E-09	4.8E-09	1.2E-08	2.2E-08
Beryllium		11	0	1.4E-10	2.2E-10	0	4.8E-10	7.6E-10
Butylated Hydroxytoluene	128-37-0	19	0	1.4E-10	2.7E-10	5.0E-11	7.3E-10	1.2E-09
Cadmium		34	0	3.7E-09	2.3E-09	3.3E-09	7.8E-09	8.6E-09
Chromium		35	1.9E-09	1.0E-08	1.0E-08	6.8E-09	2.2E-08	5.7E-08
Cobalt		35	4.4E-08	2.3E-07	1.6E-07	2.2E-07	4.5E-07	8.1E-07
Copper		35	1.8E-07	4.6E-07	2.4E-07	4.2E-07	7.5E-07	1.4E-06
Cyclohexyl isothiocyanate	1122-82-3	29	0	3.4E-08	2.5E-08	3.4E-08	7.9E-08	8.8E-08
Dibenz[a,h]anthracene	53-70-3	7	0	3.7E-11	9.3E-11	0	1.7E-10	4.6E-10
Dibenzothiophene	132-65-0	27	0	1.5E-10	2.2E-10	4.2E-11	5.9E-10	8.8E-10
Dibutyl phthalate	84-74-2	31	0	1.0E-08	7.8E-09	8.2E-09	2.5E-08	3.0E-08
Diethyl Phthalate	84-66-2	23	0	8.7E-10	7.6E-10	9.4E-10	2.1E-09	2.7E-09
Diisobutyl Phthalate	84-69-5	32	0	1.0E-09	2.7E-09	2.3E-10	2.8E-09	1.6E-08
Diisooctylphthalate	27554-26-3	33	0	2.2E-08	1.6E-08	1.6E-08	5.5E-08	5.8E-08
Di-n-octyl phthalate	117-84-0	26	0	2.2E-09	5.2E-09	5.7E-10	1.0E-08	2.3E-08
2,5-di-tert-Butyl-1,4-benzoquinone	2460-77-7	4	0	1.8E-11	5.7E-11	0	1.4E-10	2.5E-10



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
3,5-di-tert-Butyl-4-hydroxybenzaldehyde	1620-98-0	35	1.1E-11	3.8E-09	7.7E-09	2.4E-10	1.6E-08	3.8E-08
Fluoranthene	206-44-0	35	6.6E-10	4.9E-09	4.7E-09	2.9E-09	1.2E-08	2.3E-08
Fluorene	86-73-7	11	0	6.3E-11	1.5E-10	0	3.1E-10	7.5E-10
Hexadecane	544-76-3	28	0	4.6E-10	3.8E-10	4.8E-10	1.1E-09	1.8E-09
1-Hydroxypyrene	5315-79-7	1	0	2.5E-10	1.5E-09	0	0	8.7E-09
Lead		35	4.2E-08	2.2E-07	1.9E-07	1.6E-07	5.4E-07	9.8E-07
Manganese		35	3.9E-08	2.1E-07	1.3E-07	2.2E-07	3.5E-07	6.8E-07
2-(Methylthio)benzothiazole	615-22-5	5	0	3.3E-09	1.2E-08	0	1.8E-08	6.6E-08
Molybdenum		14	0	3.9E-10	5.5E-10	0	1.2E-09	2.0E-09
Naphthalene	91-20-3	1	0	1.1E-11	6.7E-11	0	0	4.0E-10
Naphthalene, 1-methyl-	90-12-0	31	0	2.9E-11	1.0E-10	4.9E-12	6.2E-11	6.1E-10
Naphthalene, 1,2-dimethyl-	573-98-8	12	0	6.8E-12	1.5E-11	0	3.4E-11	7.8E-11
Naphthalene, 1,6-dimethyl-	575-43-9	35	1.8E-11	4.5E-11	2.2E-11	3.9E-11	9.2E-11	1.3E-10
Naphthalene, 2-(bromomethyl)-	939-26-4	35	4.2E-11	1.2E-09	5.6E-10	1.2E-09	2.1E-09	2.7E-09
Naphthalene, 2,3-dimethyl-	581-40-8	35	8.9E-12	3.5E-10	3.5E-10	2.4E-10	1.0E-09	1.5E-09
Naphthalene, 2-methyl	91-57-6	19	0	2.7E-11	1.1E-10	4.8E-12	6.0E-11	6.6E-10
1-Octadecene	112-88-9	32	0	1.0E-09	8.8E-10	7.9E-10	2.7E-09	3.1E-09
17-Pentatriacontene	6971-40-0	9	0	8.6E-10	2.0E-09	0	4.7E-09	9.1E-09
Phenanthrene	85-01-8	35	3.7E-10	2.1E-09	2.6E-09	1.2E-09	6.1E-09	1.4E-08
Phenanthrene, 1-methyl	832-69-9	35	4.9E-11	5.7E-10	5.4E-10	4.0E-10	1.4E-09	2.5E-09
Phenanthrene, 2-methyl-	2531-84-2	35	1.5E-11	7.3E-10	8.4E-10	4.4E-10	2.1E-09	4.0E-09
Phenanthrene, 3-methyl	832-71-3	35	9.3E-11	1.2E-09	1.3E-09	7.7E-10	3.3E-09	6.1E-09
N-Phenylbenzamide	93-98-1	22	0	4.7E-09	9.2E-09	7.3E-10	2.9E-08	3.3E-08
Phthalimide	85-41-6	16	0	8.5E-10	2.0E-09	0	4.3E-09	9.0E-09
Pyrene	129-00-0	35	1.9E-09	1.0E-08	6.6E-09	8.4E-09	2.1E-08	2.9E-08
Pyridine, 2-(4-methylphenyl)-	4467-06-5	2	0	2.2E-11	1.3E-10	0	7.4E-12	7.4E-10
Selenium		4	0	1.3E-09	4.0E-09	0	9.0E-09	1.9E-08
Strontium		35	1.8E-08	1.8E-07	3.6E-07	9.7E-08	5.4E-07	2.1E-06
Thallium		21	0	5.1E-11	8.7E-11	4.1E-11	1.6E-10	4.7E-10
Tin		30	0	2.0E-09	2.0E-09	1.5E-09	5.4E-09	7.9E-09
Triethylene glycol monobutyl ether	143-22-6	3	0	2.5E-10	1.0E-09	0	1.3E-09	5.6E-09
5,9-Undecadien-2-one, 6,10-dimethyl-	689-67-8	35	4.3E-10	7.4E-10	2.4E-10	6.8E-10	1.1E-09	1.4E-09
Vanadium		2	0	1.0E-09	4.3E-09	0	5.5E-09	1.8E-08



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Zinc		35	9.9E-06	4.5E-05	2.4E-05	4.3E-05	8.0E-05	1.3E-04

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

F.5.7. Average Ingestion Daily Dose (ADD_{ing}) for Lifetime Cancer Risk Assessment of Carcinogens

Table F-350. Average Ingestion Daily Dose^a (ADD_{ing}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of **Field-Related Carcinogens**—Combined Gender **Athletes**

Chemical	CASRN ^c	ADD _{ing}						
		2<6 Years	6<11 Years	11<16 Years	16<30 Years	30<40 Years	40<50 Years	50<70 Years
Aniline	62-53-3	8.2E-09	5.5E-09	4.2E-09	4.7E-09	3.1E-09	2.7E-09	2.7E-09
Arsenic		4.8E-08	3.2E-08	2.5E-08	2.7E-08	1.8E-08	1.6E-08	1.6E-08
Benz[a]anthracene	56-55-3	1.3E-08	9.0E-09	7.0E-09	7.8E-09	5.2E-09	4.4E-09	4.5E-09
Benzo[a]pyrene	50-32-8	1.1E-08	7.4E-09	5.7E-09	6.4E-09	4.3E-09	3.6E-09	3.7E-09
Benzo[b]fluoranthene	205-99-2	1.9E-08	1.2E-08	9.6E-09	1.1E-08	7.1E-09	6.0E-09	6.2E-09
Benzo[k]fluoranthene	207-08-9	6.0E-09	4.0E-09	3.1E-09	3.5E-09	2.3E-09	2.0E-09	2.0E-09
Chromium		2.2E-07	1.5E-07	1.1E-07	1.3E-07	8.5E-08	7.2E-08	7.3E-08
Chrysene	218-01-9	6.0E-08	4.0E-08	3.1E-08	3.5E-08	2.3E-08	2.0E-08	2.0E-08
Cyclopenta[cd]pyrene	27208-37-3	1.3E-08	8.6E-09	6.7E-09	7.4E-09	4.9E-09	4.2E-09	4.3E-09
Dibenz[a,h]anthracene	53-70-3	8.0E-10	5.3E-10	4.1E-10	4.6E-10	3.1E-10	2.6E-10	2.7E-10
Indeno[1,2,3-cd]pyrene	193-39-5	2.3E-09	1.5E-09	1.2E-09	1.3E-09	8.7E-10	7.3E-10	7.5E-10
Lead		4.6E-06	3.1E-06	2.4E-06	2.7E-06	1.8E-06	1.5E-06	1.5E-06
Naphthalene	91-20-3	2.4E-10	1.6E-10	1.3E-10	1.4E-10	9.4E-11	8.0E-11	8.1E-11

^a ADD_{ing} was calculated from the mean of 35 individual field average concentrations (C_{ing-crumb rubber}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.



Table F-351. Average Ingestion Daily Dose^a (ADD_{ing}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of **Field-Related Carcinogens**—Combined Gender **Coaches**

Chemical	CASRN ^c	ADD _{ing}			
		16<30 Years	30<40 Years	40<50 Years	50<70 Years
Aniline	62-53-3	1.2E-09	1.1E-09	1.1E-09	1.1E-09
Arsenic		6.9E-09	6.5E-09	6.4E-09	6.4E-09
Benz[a]anthracene	56-55-3	2.0E-09	1.8E-09	1.8E-09	1.8E-09
Benzo[a]pyrene	50-32-8	1.6E-09	1.5E-09	1.5E-09	1.5E-09
Benzo[b]fluoranthene	205-99-2	2.7E-09	2.5E-09	2.5E-09	2.5E-09
Benzo[k]fluoranthene	207-08-9	8.7E-10	8.2E-10	8.0E-10	8.0E-10
Chromium		3.2E-08	3.0E-08	3.0E-08	3.0E-08
Chrysene	218-01-9	8.7E-09	8.2E-09	8.0E-09	8.0E-09
Cyclopenta[cd]pyrene	27208-37-3	1.9E-09	1.8E-09	1.7E-09	1.7E-09
Dibenz[a,h]anthracene	53-70-3	1.2E-10	1.1E-10	1.1E-10	1.1E-10
Indeno[1,2,3-cd]pyrene	193-39-5	3.3E-10	3.1E-10	3.0E-10	3.0E-10
Lead		6.7E-07	6.3E-07	6.2E-07	6.2E-07
Naphthalene	91-20-3	3.6E-11	3.3E-11	3.3E-11	3.3E-11

^a ADD_{ing} was calculated from the mean of 35 individual field average concentrations (C_{ing-crumb rubber}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-352. Average Ingestion Daily Dose^a (ADD_{ing}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of **Field-Related Carcinogens**—Combined Gender **Referees**

Chemical	CASRN ^c	ADD _{ing}			
		16<30 Years	30<40 Years	40<50 Years	50<70 Years
Aniline	62-53-3	5.2E-10	4.9E-10	4.8E-10	4.8E-10
Arsenic		3.0E-09	2.9E-09	2.8E-09	2.8E-09
Benz[a]anthracene	56-55-3	8.6E-10	8.1E-10	7.9E-10	7.9E-10
Benzo[a]pyrene	50-32-8	7.1E-10	6.6E-10	6.5E-10	6.5E-10
Benzo[b]fluoranthene	205-99-2	1.2E-09	1.1E-09	1.1E-09	1.1E-09
Benzo[k]fluoranthene	207-08-9	3.8E-10	3.6E-10	3.5E-10	3.5E-10
Chromium		1.4E-08	1.3E-08	1.3E-08	1.3E-08
Chrysene	218-01-9	3.8E-09	3.6E-09	3.5E-09	3.5E-09
Cyclopenta[cd]pyrene	27208-37-3	8.2E-10	7.7E-10	7.6E-10	7.6E-10
Dibenz[a,h]anthracene	53-70-3	5.1E-11	4.8E-11	4.7E-11	4.7E-11
Indeno[1,2,3-cd]pyrene	193-39-5	1.4E-10	1.4E-10	1.3E-10	1.3E-10
Lead		3.0E-07	2.8E-07	2.7E-07	2.7E-07



Chemical	CASRN ^c	ADD _{ing}			
		16<30 Years	30<40 Years	40<50 Years	50<70 Years
Naphthalene	91-20-3	1.6E-11	1.5E-11	1.4E-11	1.4E-11

^a ADD_{ing} was calculated from the mean of 35 individual field average concentrations (C_{ing-crumb rubber}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-353. Average Ingestion Daily Dose^a (ADD_{ing}, milligrams per kilogram BW per day) for Lifetime Cancer Risk Assessment of **Field-Related Carcinogens**—Combined Gender **Spectators**

Chemical	CASRN ^c	ADD _{ing}								
		Third trimester fetus	0<2 Years	2<6 Years	6<11 Years	11<16 Years	16<30 Years	30<40 Years	40<50 Years	50<70 Years
Aniline	62-53-3	4.1E-10	2.9E-08	1.4E-08	9.5E-09	7.5E-10	4.2E-10	4.0E-10	3.8E-10	3.8E-10
Arsenic		2.4E-09	1.7E-07	7.9E-08	5.5E-08	4.3E-09	2.5E-09	2.3E-09	2.2E-09	2.2E-09
Benz[a]anthracene	56-55-3	6.8E-10	4.7E-08	2.2E-08	1.6E-08	1.2E-09	7.0E-10	6.6E-10	6.3E-10	6.3E-10
Benzo[a]pyrene	50-32-8	5.6E-10	3.9E-08	1.8E-08	1.3E-08	1.0E-09	5.7E-10	5.4E-10	5.1E-10	5.2E-10
Benzo[b]fluoranthene	205-99-2	9.3E-10	6.5E-08	3.1E-08	2.2E-08	1.7E-09	9.6E-10	9.1E-10	8.6E-10	8.6E-10
Benzo[k]fluoranthene	207-08-9	3.0E-10	2.1E-08	1.0E-08	7.0E-09	5.5E-10	3.1E-10	2.9E-10	2.8E-10	2.8E-10
Chromium		1.1E-08	7.7E-07	3.7E-07	2.6E-07	2.0E-08	1.1E-08	1.1E-08	1.0E-08	1.0E-08
Chrysene	218-01-9	3.0E-09	2.1E-07	1.0E-07	7.0E-08	5.5E-09	3.1E-09	2.9E-09	2.8E-09	2.8E-09
Cyclopenta[cd]pyrene	27208-37-3	6.5E-10	4.5E-08	2.1E-08	1.5E-08	1.2E-09	6.7E-10	6.3E-10	6.0E-10	6.0E-10
Dibenz[a,h]anthracene	53-70-3	4.0E-11	2.8E-09	1.3E-09	9.3E-10	7.3E-11	4.1E-11	3.9E-11	3.7E-11	3.7E-11
Indeno[1,2,3-cd]pyrene	193-39-5	1.1E-10	7.9E-09	3.7E-09	2.6E-09	2.1E-10	1.2E-10	1.1E-10	1.0E-10	1.0E-10
Lead		2.3E-07	1.6E-05	7.7E-06	5.4E-06	4.2E-07	2.4E-07	2.3E-07	2.2E-07	2.2E-07
Naphthalene	91-20-3	1.2E-11	8.5E-10	4.1E-10	2.8E-10	2.2E-11	1.3E-11	1.2E-11	1.1E-11	1.1E-11

^a ADD_{ing} was calculated from the mean of 35 individual field average concentrations (C_{ing-crumb rubber}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

INDIVIDUAL FIELD ASSESSMENT (Table F-354 to Table F-377)

Table F-354. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens**^b—Combined Gender **Athletes 2<6 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	8.2E-09	1.9E-08	0	4.6E-08	8.2E-08



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		35	5.6E-09	4.8E-08	3.4E-08	4.3E-08	8.6E-08	1.8E-07
Benz[a]anthracene	56-55-3	35	8.8E-10	1.3E-08	1.5E-08	6.4E-09	4.2E-08	5.9E-08
Benzo[a]pyrene	50-32-8	35	2.2E-09	1.1E-08	9.1E-09	7.9E-09	3.1E-08	3.3E-08
Benzo[b]fluoranthene	205-99-2	35	3.8E-09	1.9E-08	1.4E-08	1.2E-08	4.5E-08	5.4E-08
Benzo[k]fluoranthene	207-08-9	35	5.1E-10	6.0E-09	5.7E-09	3.8E-09	1.8E-08	2.3E-08
Chromium		35	4.1E-08	2.2E-07	2.2E-07	1.5E-07	4.8E-07	1.2E-06
Chrysene	218-01-9	35	1.5E-08	6.0E-08	3.2E-08	6.0E-08	1.0E-07	1.6E-07
Cyclopenta[cd]pyrene	27208-37-3	35	2.3E-09	1.3E-08	1.2E-08	1.1E-08	3.4E-08	6.1E-08
Dibenz[a,h]anthracene	53-70-3	7	0	8.0E-10	2.0E-09	0	3.6E-09	9.9E-09
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	2.3E-09	3.9E-09	0	9.6E-09	1.5E-08
Lead		35	9.0E-07	4.6E-06	4.0E-06	3.5E-06	1.2E-05	2.1E-05
Naphthalene	91-20-3	1	0	2.4E-10	1.4E-09	0	0	8.5E-09

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis. Values are rounded to two significant figures.

Table F-355. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Athletes 6<11 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	5.5E-09	1.2E-08	0	3.0E-08	5.5E-08
Arsenic		35	3.7E-09	3.2E-08	2.2E-08	2.9E-08	5.7E-08	1.2E-07
Benz[a]anthracene	56-55-3	35	5.9E-10	9.0E-09	9.9E-09	4.3E-09	2.8E-08	3.9E-08
Benzo[a]pyrene	50-32-8	35	1.4E-09	7.4E-09	6.1E-09	5.3E-09	2.0E-08	2.2E-08
Benzo[b]fluoranthene	205-99-2	35	2.5E-09	1.2E-08	9.1E-09	8.3E-09	3.0E-08	3.6E-08
Benzo[k]fluoranthene	207-08-9	35	3.4E-10	4.0E-09	3.8E-09	2.5E-09	1.2E-08	1.5E-08
Chromium		35	2.7E-08	1.5E-07	1.4E-07	9.7E-08	3.2E-07	8.2E-07
Chrysene	218-01-9	35	1.0E-08	4.0E-08	2.1E-08	4.0E-08	6.9E-08	1.1E-07
Cyclopenta[cd]pyrene	27208-37-3	35	1.5E-09	8.6E-09	8.0E-09	7.4E-09	2.3E-08	4.0E-08
Dibenz[a,h]anthracene	53-70-3	7	0	5.3E-10	1.3E-09	0	2.4E-09	6.6E-09
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	1.5E-09	2.6E-09	0	6.4E-09	1.0E-08
Lead		35	6.0E-07	3.1E-06	2.7E-06	2.3E-06	7.8E-06	1.4E-05
Naphthalene	91-20-3	1	0	1.6E-10	9.6E-10	0	0	5.7E-09

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average



concentrations ($C_{\text{ing-crumb rubber-field}}$) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-356. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment ($ADD_{\text{ing-field}}$, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Athletes 11<16 Years**

Chemical	CASRN ^c	Detection	$ADD_{\text{ing-field}}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	4.2E-09	9.7E-09	0	2.4E-08	4.3E-08
Arsenic		35	2.9E-09	2.5E-08	1.7E-08	2.2E-08	4.4E-08	9.5E-08
Benz[a]anthracene	56-55-3	35	4.5E-10	7.0E-09	7.6E-09	3.3E-09	2.2E-08	3.0E-08
Benzo[a]pyrene	50-32-8	35	1.1E-09	5.7E-09	4.7E-09	4.1E-09	1.6E-08	1.7E-08
Benzo[b]fluoranthene	205-99-2	35	2.0E-09	9.6E-09	7.1E-09	6.4E-09	2.3E-08	2.8E-08
Benzo[k]fluoranthene	207-08-9	35	2.6E-10	3.1E-09	2.9E-09	2.0E-09	9.1E-09	1.2E-08
Chromium		35	2.1E-08	1.1E-07	1.1E-07	7.5E-08	2.5E-07	6.3E-07
Chrysene	218-01-9	35	7.8E-09	3.1E-08	1.6E-08	3.1E-08	5.3E-08	8.5E-08
Cyclopenta[cd]pyrene	27208-37-3	35	1.2E-09	6.7E-09	6.2E-09	5.8E-09	1.8E-08	3.1E-08
Dibenz[a,h]anthracene	53-70-3	7	0	4.1E-10	1.0E-09	0	1.9E-09	5.1E-09
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	1.2E-09	2.0E-09	0	4.9E-09	8.0E-09
Lead		35	4.6E-07	2.4E-06	2.1E-06	1.8E-06	6.0E-06	1.1E-05
Naphthalene	91-20-3	1	0	1.3E-10	7.5E-10	0	0	4.4E-09

^a 35 field-specific $ADD_{\text{ing-field}}$ are included in the table. $ADD_{\text{ing-field}}$ was calculated from field-specific average concentrations ($C_{\text{ing-crumb rubber-field}}$) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-357. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment ($ADD_{\text{ing-field}}$, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Athletes 16<30 Years**

Chemical	CASRN ^c	Detection	$ADD_{\text{ing-field}}$					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	4.7E-09	1.1E-08	0	2.6E-08	4.7E-08
Arsenic		35	3.2E-09	2.7E-08	1.9E-08	2.5E-08	4.9E-08	1.1E-07
Benz[a]anthracene	56-55-3	35	5.1E-10	7.8E-09	8.5E-09	3.7E-09	2.4E-08	3.4E-08



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	35	1.3E-09	6.4E-09	5.2E-09	4.5E-09	1.8E-08	1.9E-08
Benzo[b]fluoranthene	205-99-2	35	2.2E-09	1.1E-08	7.9E-09	7.2E-09	2.6E-08	3.1E-08
Benzo[k]fluoranthene	207-08-9	35	2.9E-10	3.5E-09	3.3E-09	2.2E-09	1.0E-08	1.3E-08
Chromium		35	2.3E-08	1.3E-07	1.2E-07	8.4E-08	2.8E-07	7.0E-07
Chrysene	218-01-9	35	8.7E-09	3.5E-08	1.8E-08	3.5E-08	5.9E-08	9.5E-08
Cyclopenta[cd]pyrene	27208-37-3	35	1.3E-09	7.4E-09	6.9E-09	6.4E-09	2.0E-08	3.5E-08
Dibenz[a,h]anthracene	53-70-3	7	0	4.6E-10	1.1E-09	0	2.1E-09	5.7E-09
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	1.3E-09	2.3E-09	0	5.5E-09	8.9E-09
Lead		35	5.2E-07	2.7E-06	2.3E-06	2.0E-06	6.7E-06	1.2E-05
Naphthalene	91-20-3	1	0	1.4E-10	8.3E-10	0	0	4.9E-09

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis. Values are rounded to two significant figures.

Table F-358. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Athletes 30<40 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	3.1E-09	7.2E-09	0	1.8E-08	3.2E-08
Arsenic		35	2.2E-09	1.8E-08	1.3E-08	1.7E-08	3.3E-08	7.0E-08
Benz[a]anthracene	56-55-3	35	3.4E-10	5.2E-09	5.7E-09	2.5E-09	1.6E-08	2.3E-08
Benzo[a]pyrene	50-32-8	35	8.4E-10	4.3E-09	3.5E-09	3.0E-09	1.2E-08	1.2E-08
Benzo[b]fluoranthene	205-99-2	35	1.5E-09	7.1E-09	5.3E-09	4.8E-09	1.7E-08	2.1E-08
Benzo[k]fluoranthene	207-08-9	35	1.9E-10	2.3E-09	2.2E-09	1.5E-09	6.8E-09	8.7E-09
Chromium		35	1.6E-08	8.5E-08	8.3E-08	5.6E-08	1.9E-07	4.7E-07
Chrysene	218-01-9	35	5.8E-09	2.3E-08	1.2E-08	2.3E-08	4.0E-08	6.3E-08
Cyclopenta[cd]pyrene	27208-37-3	35	8.7E-10	4.9E-09	4.6E-09	4.3E-09	1.3E-08	2.3E-08
Dibenz[a,h]anthracene	53-70-3	7	0	3.1E-10	7.7E-10	0	1.4E-09	3.8E-09
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	8.7E-10	1.5E-09	0	3.7E-09	5.9E-09
Lead		35	3.4E-07	1.8E-06	1.5E-06	1.3E-06	4.5E-06	8.1E-06
Naphthalene	91-20-3	1	0	9.4E-11	5.6E-10	0	0	3.3E-09

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.



^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-359. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Athletes 40<50 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	2.7E-09	6.1E-09	0	1.5E-08	2.7E-08
Arsenic		35	1.8E-09	1.6E-08	1.1E-08	1.4E-08	2.8E-08	6.0E-08
Benz[a]anthracene	56-55-3	35	2.9E-10	4.4E-09	4.8E-09	2.1E-09	1.4E-08	1.9E-08
Benzo[a]pyrene	50-32-8	35	7.1E-10	3.6E-09	3.0E-09	2.6E-09	1.0E-08	1.1E-08
Benzo[b]fluoranthene	205-99-2	35	1.2E-09	6.0E-09	4.5E-09	4.1E-09	1.5E-08	1.8E-08
Benzo[k]fluoranthene	207-08-9	35	1.6E-10	2.0E-09	1.8E-09	1.2E-09	5.7E-09	7.3E-09
Chromium		35	1.3E-08	7.2E-08	7.0E-08	4.8E-08	1.6E-07	4.0E-07
Chrysene	218-01-9	35	4.9E-09	2.0E-08	1.0E-08	2.0E-08	3.3E-08	5.4E-08
Cyclopenta[cd]pyrene	27208-37-3	35	7.4E-10	4.2E-09	3.9E-09	3.6E-09	1.1E-08	2.0E-08
Dibenz[a,h]anthracene	53-70-3	7	0	2.6E-10	6.5E-10	0	1.2E-09	3.2E-09
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	7.3E-10	1.3E-09	0	3.1E-09	5.0E-09
Lead		35	2.9E-07	1.5E-06	1.3E-06	1.1E-06	3.8E-06	6.8E-06
Naphthalene	91-20-3	1	0	8.0E-11	4.7E-10	0	0	2.8E-09

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-360. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Athletes 50<70 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	2.7E-09	6.2E-09	0	1.5E-08	2.7E-08
Arsenic		35	1.9E-09	1.6E-08	1.1E-08	1.4E-08	2.9E-08	6.1E-08
Benz[a]anthracene	56-55-3	35	2.9E-10	4.5E-09	4.9E-09	2.1E-09	1.4E-08	2.0E-08
Benzo[a]pyrene	50-32-8	35	7.2E-10	3.7E-09	3.0E-09	2.6E-09	1.0E-08	1.1E-08
Benzo[b]fluoranthene	205-99-2	35	1.3E-09	6.2E-09	4.6E-09	4.1E-09	1.5E-08	1.8E-08



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[k]fluoranthene	207-08-9	35	1.7E-10	2.0E-09	1.9E-09	1.3E-09	5.8E-09	7.5E-09
Chromium		35	1.4E-08	7.3E-08	7.2E-08	4.9E-08	1.6E-07	4.1E-07
Chrysene	218-01-9	35	5.0E-09	2.0E-08	1.1E-08	2.0E-08	3.4E-08	5.5E-08
Cyclopenta[cd]pyrene	27208-37-3	35	7.5E-10	4.3E-09	4.0E-09	3.7E-09	1.1E-08	2.0E-08
Dibenz[a,h]anthracene	53-70-3	7	0	2.7E-10	6.6E-10	0	1.2E-09	3.3E-09
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	7.5E-10	1.3E-09	0	3.2E-09	5.1E-09
Lead		35	3.0E-07	1.5E-06	1.3E-06	1.2E-06	3.9E-06	7.0E-06
Naphthalene	91-20-3	1	0	8.1E-11	4.8E-10	0	0	2.8E-09

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-361. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Coaches 16<30 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	1.2E-09	2.7E-09	0	6.6E-09	1.2E-08
Arsenic		35	8.2E-10	6.9E-09	4.9E-09	6.3E-09	1.2E-08	2.7E-08
Benz[a]anthracene	56-55-3	35	1.3E-10	2.0E-09	2.1E-09	9.4E-10	6.1E-09	8.5E-09
Benzo[a]pyrene	50-32-8	35	3.2E-10	1.6E-09	1.3E-09	1.1E-09	4.5E-09	4.7E-09
Benzo[b]fluoranthene	205-99-2	35	5.5E-10	2.7E-09	2.0E-09	1.8E-09	6.5E-09	7.9E-09
Benzo[k]fluoranthene	207-08-9	35	7.3E-11	8.7E-10	8.2E-10	5.5E-10	2.6E-09	3.3E-09
Chromium		35	5.9E-09	3.2E-08	3.1E-08	2.1E-08	7.0E-08	1.8E-07
Chrysene	218-01-9	35	2.2E-09	8.7E-09	4.6E-09	8.7E-09	1.5E-08	2.4E-08
Cyclopenta[cd]pyrene	27208-37-3	35	3.3E-10	1.9E-09	1.7E-09	1.6E-09	4.9E-09	8.8E-09
Dibenz[a,h]anthracene	53-70-3	7	0	1.2E-10	2.9E-10	0	5.2E-10	1.4E-09
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	3.3E-10	5.7E-10	0	1.4E-09	2.2E-09
Lead		35	1.3E-07	6.7E-07	5.8E-07	5.1E-07	1.7E-06	3.0E-06
Naphthalene	91-20-3	1	0	3.6E-11	2.1E-10	0	0	1.2E-09

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.



° CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.
Values are rounded to two significant figures.

Table F-362. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Coaches 30<40 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	1.1E-09	2.6E-09	0	6.2E-09	1.1E-08
Arsenic		35	7.7E-10	6.5E-09	4.6E-09	5.9E-09	1.2E-08	2.5E-08
Benz[a]anthracene	56-55-3	35	1.2E-10	1.8E-09	2.0E-09	8.8E-10	5.7E-09	8.0E-09
Benzo[a]pyrene	50-32-8	35	3.0E-10	1.5E-09	1.2E-09	1.1E-09	4.2E-09	4.4E-09
Benzo[b]fluoranthene	205-99-2	35	5.2E-10	2.5E-09	1.9E-09	1.7E-09	6.1E-09	7.4E-09
Benzo[k]fluoranthene	207-08-9	35	6.9E-11	8.2E-10	7.7E-10	5.2E-10	2.4E-09	3.1E-09
Chromium		35	5.6E-09	3.0E-08	2.9E-08	2.0E-08	6.6E-08	1.7E-07
Chrysene	218-01-9	35	2.1E-09	8.2E-09	4.4E-09	8.2E-09	1.4E-08	2.2E-08
Cyclopenta[cd]pyrene	27208-37-3	35	3.1E-10	1.8E-09	1.6E-09	1.5E-09	4.6E-09	8.3E-09
Dibenz[a,h]anthracene	53-70-3	7	0	1.1E-10	2.7E-10	0	4.9E-10	1.4E-09
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	3.1E-10	5.4E-10	0	1.3E-09	2.1E-09
Lead		35	1.2E-07	6.3E-07	5.5E-07	4.8E-07	1.6E-06	2.9E-06
Naphthalene	91-20-3	1	0	3.3E-11	2.0E-10	0	0	1.2E-09

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.
Values are rounded to two significant figures.

Table F-363. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Coaches 40<50 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	1.1E-09	2.5E-09	0	6.1E-09	1.1E-08
Arsenic		35	7.5E-10	6.4E-09	4.5E-09	5.7E-09	1.1E-08	2.5E-08
Benz[a]anthracene	56-55-3	35	1.2E-10	1.8E-09	2.0E-09	8.6E-10	5.6E-09	7.8E-09
Benzo[a]pyrene	50-32-8	35	2.9E-10	1.5E-09	1.2E-09	1.1E-09	4.1E-09	4.3E-09
Benzo[b]fluoranthene	205-99-2	35	5.0E-10	2.5E-09	1.8E-09	1.7E-09	6.0E-09	7.3E-09
Benzo[k]fluoranthene	207-08-9	35	6.7E-11	8.0E-10	7.6E-10	5.1E-10	2.4E-09	3.0E-09
Chromium		35	5.5E-09	3.0E-08	2.9E-08	1.9E-08	6.4E-08	1.6E-07



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Chrysene	218-01-9	35	2.0E-09	8.0E-09	4.3E-09	8.0E-09	1.4E-08	2.2E-08
Cyclopenta[cd]pyrene	27208-37-3	35	3.0E-10	1.7E-09	1.6E-09	1.5E-09	4.5E-09	8.1E-09
Dibenz[a,h]anthracene	53-70-3	7	0	1.1E-10	2.7E-10	0	4.8E-10	1.3E-09
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	3.0E-10	5.3E-10	0	1.3E-09	2.1E-09
Lead		35	1.2E-07	6.2E-07	5.3E-07	4.7E-07	1.6E-06	2.8E-06
Naphthalene	91-20-3	1	0	3.3E-11	1.9E-10	0	0	1.1E-09

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-364. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Coaches 50<70 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	1.1E-09	2.5E-09	0	6.1E-09	1.1E-08
Arsenic		35	7.5E-10	6.4E-09	4.5E-09	5.8E-09	1.2E-08	2.5E-08
Benz[a]anthracene	56-55-3	35	1.2E-10	1.8E-09	2.0E-09	8.6E-10	5.6E-09	7.9E-09
Benzo[a]pyrene	50-32-8	35	2.9E-10	1.5E-09	1.2E-09	1.1E-09	4.1E-09	4.4E-09
Benzo[b]fluoranthene	205-99-2	35	5.1E-10	2.5E-09	1.8E-09	1.7E-09	6.0E-09	7.3E-09
Benzo[k]fluoranthene	207-08-9	35	6.8E-11	8.0E-10	7.6E-10	5.1E-10	2.4E-09	3.0E-09
Chromium		35	5.5E-09	3.0E-08	2.9E-08	2.0E-08	6.5E-08	1.6E-07
Chrysene	218-01-9	35	2.0E-09	8.0E-09	4.3E-09	8.0E-09	1.4E-08	2.2E-08
Cyclopenta[cd]pyrene	27208-37-3	35	3.0E-10	1.7E-09	1.6E-09	1.5E-09	4.5E-09	8.1E-09
Dibenz[a,h]anthracene	53-70-3	7	0	1.1E-10	2.7E-10	0	4.8E-10	1.3E-09
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	3.0E-10	5.3E-10	0	1.3E-09	2.1E-09
Lead		35	1.2E-07	6.2E-07	5.4E-07	4.7E-07	1.6E-06	2.8E-06
Naphthalene	91-20-3	1	0	3.3E-11	1.9E-10	0	0	1.1E-09

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.



Table F-365. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Referees 16<30 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	5.2E-10	1.2E-09	0	2.9E-09	5.3E-09
Arsenic		35	3.6E-10	3.0E-09	2.2E-09	2.7E-09	5.5E-09	1.2E-08
Benz[a]anthracene	56-55-3	35	5.6E-11	8.6E-10	9.4E-10	4.1E-10	2.7E-09	3.7E-09
Benzo[a]pyrene	50-32-8	35	1.4E-10	7.1E-10	5.8E-10	5.0E-10	2.0E-09	2.1E-09
Benzo[b]fluoranthene	205-99-2	35	2.4E-10	1.2E-09	8.8E-10	7.9E-10	2.9E-09	3.5E-09
Benzo[k]fluoranthene	207-08-9	35	3.2E-11	3.8E-10	3.6E-10	2.4E-10	1.1E-09	1.4E-09
Chromium		35	2.6E-09	1.4E-08	1.4E-08	9.3E-09	3.1E-08	7.8E-08
Chrysene	218-01-9	35	9.6E-10	3.8E-09	2.0E-09	3.8E-09	6.6E-09	1.1E-08
Cyclopenta[cd]pyrene	27208-37-3	35	1.4E-10	8.2E-10	7.6E-10	7.1E-10	2.2E-09	3.9E-09
Dibenz[a,h]anthracene	53-70-3	7	0	5.1E-11	1.3E-10	0	2.3E-10	6.3E-10
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	1.4E-10	2.5E-10	0	6.1E-10	9.8E-10
Lead		35	5.7E-08	3.0E-07	2.6E-07	2.2E-07	7.4E-07	1.3E-06
Naphthalene	91-20-3	1	0	1.6E-11	9.2E-11	0	0	5.5E-10

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-366. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Referees 30<40 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	4.9E-10	1.1E-09	0	2.7E-09	4.9E-09
Arsenic		35	3.4E-10	2.9E-09	2.0E-09	2.6E-09	5.2E-09	1.1E-08
Benz[a]anthracene	56-55-3	35	5.3E-11	8.1E-10	8.9E-10	3.9E-10	2.5E-09	3.5E-09
Benzo[a]pyrene	50-32-8	35	1.3E-10	6.6E-10	5.4E-10	4.7E-10	1.8E-09	1.9E-09
Benzo[b]fluoranthene	205-99-2	35	2.3E-10	1.1E-09	8.2E-10	7.5E-10	2.7E-09	3.3E-09
Benzo[k]fluoranthene	207-08-9	35	3.0E-11	3.6E-10	3.4E-10	2.3E-10	1.1E-09	1.4E-09
Chromium		35	2.4E-09	1.3E-08	1.3E-08	8.7E-09	2.9E-08	7.3E-08
Chrysene	218-01-9	35	9.1E-10	3.6E-09	1.9E-09	3.6E-09	6.2E-09	9.9E-09
Cyclopenta[cd]pyrene	27208-37-3	35	1.4E-10	7.7E-10	7.2E-10	6.7E-10	2.0E-09	3.6E-09



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dibenz[a,h]anthracene	53-70-3	7	0	4.8E-11	1.2E-10	0	2.2E-10	6.0E-10
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	1.4E-10	2.4E-10	0	5.7E-10	9.2E-10
Lead		35	5.4E-08	2.8E-07	2.4E-07	2.1E-07	7.0E-07	1.3E-06
Naphthalene	91-20-3	1	0	1.5E-11	8.7E-11	0	0	5.1E-10

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-367. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Referees 40<50 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	4.8E-10	1.1E-09	0	2.7E-09	4.8E-09
Arsenic		35	3.3E-10	2.8E-09	2.0E-09	2.5E-09	5.0E-09	1.1E-08
Benz[a]anthracene	56-55-3	35	5.2E-11	7.9E-10	8.7E-10	3.8E-10	2.5E-09	3.4E-09
Benzo[a]pyrene	50-32-8	35	1.3E-10	6.5E-10	5.3E-10	4.6E-10	1.8E-09	1.9E-09
Benzo[b]fluoranthene	205-99-2	35	2.2E-10	1.1E-09	8.0E-10	7.3E-10	2.6E-09	3.2E-09
Benzo[k]fluoranthene	207-08-9	35	3.0E-11	3.5E-10	3.3E-10	2.2E-10	1.0E-09	1.3E-09
Chromium		35	2.4E-09	1.3E-08	1.3E-08	8.6E-09	2.8E-08	7.2E-08
Chrysene	218-01-9	35	8.9E-10	3.5E-09	1.9E-09	3.5E-09	6.0E-09	9.7E-09
Cyclopenta[cd]pyrene	27208-37-3	35	1.3E-10	7.6E-10	7.0E-10	6.5E-10	2.0E-09	3.6E-09
Dibenz[a,h]anthracene	53-70-3	7	0	4.7E-11	1.2E-10	0	2.1E-10	5.8E-10
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	1.3E-10	2.3E-10	0	5.6E-10	9.1E-10
Lead		35	5.3E-08	2.7E-07	2.3E-07	2.1E-07	6.8E-07	1.2E-06
Naphthalene	91-20-3	1	0	1.4E-11	8.5E-11	0	0	5.0E-10

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.



Table F-368. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Referees 50<70 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	4.8E-10	1.1E-09	0	2.7E-09	4.8E-09
Arsenic		35	3.3E-10	2.8E-09	2.0E-09	2.5E-09	5.1E-09	1.1E-08
Benz[a]anthracene	56-55-3	35	5.2E-11	7.9E-10	8.7E-10	3.8E-10	2.5E-09	3.5E-09
Benzo[a]pyrene	50-32-8	35	1.3E-10	6.5E-10	5.3E-10	4.6E-10	1.8E-09	1.9E-09
Benzo[b]fluoranthene	205-99-2	35	2.2E-10	1.1E-09	8.1E-10	7.3E-10	2.6E-09	3.2E-09
Benzo[k]fluoranthene	207-08-9	35	3.0E-11	3.5E-10	3.3E-10	2.2E-10	1.0E-09	1.3E-09
Chromium		35	2.4E-09	1.3E-08	1.3E-08	8.6E-09	2.8E-08	7.2E-08
Chrysene	218-01-9	35	8.9E-10	3.5E-09	1.9E-09	3.5E-09	6.0E-09	9.7E-09
Cyclopenta[cd]pyrene	27208-37-3	35	1.3E-10	7.6E-10	7.0E-10	6.5E-10	2.0E-09	3.6E-09
Dibenz[a,h]anthracene	53-70-3	7	0	4.7E-11	1.2E-10	0	2.1E-10	5.8E-10
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	1.3E-10	2.3E-10	0	5.6E-10	9.1E-10
Lead		35	5.3E-08	2.7E-07	2.4E-07	2.1E-07	6.9E-07	1.2E-06
Naphthalene	91-20-3	1	0	1.4E-11	8.5E-11	0	0	5.0E-10

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-369. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators Third Trimester Fetus<0 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	4.1E-10	9.4E-10	0	2.3E-09	4.2E-09
Arsenic		35	2.8E-10	2.4E-09	1.7E-09	2.2E-09	4.3E-09	9.2E-09
Benz[a]anthracene	56-55-3	35	4.4E-11	6.8E-10	7.4E-10	3.2E-10	2.1E-09	3.0E-09
Benzo[a]pyrene	50-32-8	35	1.1E-10	5.6E-10	4.6E-10	4.0E-10	1.5E-09	1.6E-09
Benzo[b]fluoranthene	205-99-2	35	1.9E-10	9.3E-10	6.9E-10	6.3E-10	2.3E-09	2.7E-09
Benzo[k]fluoranthene	207-08-9	35	2.5E-11	3.0E-10	2.9E-10	1.9E-10	8.9E-10	1.1E-09
Chromium		35	2.1E-09	1.1E-08	1.1E-08	7.4E-09	2.4E-08	6.2E-08
Chrysene	218-01-9	35	7.6E-10	3.0E-09	1.6E-09	3.0E-09	5.2E-09	8.3E-09
Cyclopenta[cd]pyrene	27208-37-3	35	1.1E-10	6.5E-10	6.0E-10	5.6E-10	1.7E-09	3.1E-09



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dibenz[a,h]anthracene	53-70-3	7	0	4.0E-11	1.0E-10	0	1.8E-10	5.0E-10
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	1.1E-10	2.0E-10	0	4.8E-10	7.8E-10
Lead		35	4.5E-08	2.3E-07	2.0E-07	1.8E-07	5.9E-07	1.1E-06
Naphthalene	91-20-3	1	0	1.2E-11	7.3E-11	0	0	4.3E-10

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-370. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 0<2 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	2.9E-08	6.6E-08	0	1.6E-07	2.9E-07
Arsenic		35	2.0E-08	1.7E-07	1.2E-07	1.5E-07	3.0E-07	6.4E-07
Benz[a]anthracene	56-55-3	35	3.1E-09	4.7E-08	5.2E-08	2.3E-08	1.5E-07	2.1E-07
Benzo[a]pyrene	50-32-8	35	7.6E-09	3.9E-08	3.2E-08	2.8E-08	1.1E-07	1.1E-07
Benzo[b]fluoranthene	205-99-2	35	1.3E-08	6.5E-08	4.8E-08	4.4E-08	1.6E-07	1.9E-07
Benzo[k]fluoranthene	207-08-9	35	1.8E-09	2.1E-08	2.0E-08	1.3E-08	6.2E-08	7.9E-08
Chromium		35	1.4E-07	7.7E-07	7.6E-07	5.1E-07	1.7E-06	4.3E-06
Chrysene	218-01-9	35	5.3E-08	2.1E-07	1.1E-07	2.1E-07	3.6E-07	5.8E-07
Cyclopenta[cd]pyrene	27208-37-3	35	7.9E-09	4.5E-08	4.2E-08	3.9E-08	1.2E-07	2.1E-07
Dibenz[a,h]anthracene	53-70-3	7	0	2.8E-09	7.0E-09	0	1.3E-08	3.5E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	7.9E-09	1.4E-08	0	3.3E-08	5.4E-08
Lead		35	3.1E-06	1.6E-05	1.4E-05	1.2E-05	4.1E-05	7.3E-05
Naphthalene	91-20-3	1	0	8.5E-10	5.1E-09	0	0	3.0E-08

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.



Table F-371. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 2<6 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	1.4E-08	3.1E-08	0	7.6E-08	1.4E-07
Arsenic		35	9.3E-09	7.9E-08	5.6E-08	7.1E-08	1.4E-07	3.0E-07
Benz[a]anthracene	56-55-3	35	1.5E-09	2.2E-08	2.5E-08	1.1E-08	6.9E-08	9.7E-08
Benzo[a]pyrene	50-32-8	35	3.6E-09	1.8E-08	1.5E-08	1.3E-08	5.1E-08	5.4E-08
Benzo[b]fluoranthene	205-99-2	35	6.3E-09	3.1E-08	2.3E-08	2.1E-08	7.5E-08	9.0E-08
Benzo[k]fluoranthene	207-08-9	35	8.4E-10	1.0E-08	9.4E-09	6.3E-09	2.9E-08	3.7E-08
Chromium		35	6.8E-08	3.7E-07	3.6E-07	2.4E-07	8.0E-07	2.0E-06
Chrysene	218-01-9	35	2.5E-08	1.0E-07	5.3E-08	9.9E-08	1.7E-07	2.7E-07
Cyclopenta[cd]pyrene	27208-37-3	35	3.8E-09	2.1E-08	2.0E-08	1.8E-08	5.6E-08	1.0E-07
Dibenz[a,h]anthracene	53-70-3	7	0	1.3E-09	3.3E-09	0	6.0E-09	1.6E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	3.7E-09	6.5E-09	0	1.6E-08	2.6E-08
Lead		35	1.5E-06	7.7E-06	6.6E-06	5.8E-06	1.9E-05	3.5E-05
Naphthalene	91-20-3	1	0	4.1E-10	2.4E-09	0	0	1.4E-08

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-372. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 6<11 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	9.5E-09	2.2E-08	0	5.3E-08	9.6E-08
Arsenic		35	6.5E-09	5.5E-08	3.9E-08	5.0E-08	1.0E-07	2.1E-07
Benz[a]anthracene	56-55-3	35	1.0E-09	1.6E-08	1.7E-08	7.5E-09	4.9E-08	6.8E-08
Benzo[a]pyrene	50-32-8	35	2.5E-09	1.3E-08	1.1E-08	9.2E-09	3.6E-08	3.8E-08
Benzo[b]fluoranthene	205-99-2	35	4.4E-09	2.2E-08	1.6E-08	1.4E-08	5.2E-08	6.3E-08
Benzo[k]fluoranthene	207-08-9	35	5.9E-10	7.0E-09	6.6E-09	4.4E-09	2.0E-08	2.6E-08
Chromium		35	4.7E-08	2.6E-07	2.5E-07	1.7E-07	5.6E-07	1.4E-06
Chrysene	218-01-9	35	1.8E-08	7.0E-08	3.7E-08	7.0E-08	1.2E-07	1.9E-07
Cyclopenta[cd]pyrene	27208-37-3	35	2.6E-09	1.5E-08	1.4E-08	1.3E-08	3.9E-08	7.0E-08



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dibenz[a,h]anthracene	53-70-3	7	0	9.3E-10	2.3E-09	0	4.2E-09	1.2E-08
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	2.6E-09	4.6E-09	0	1.1E-08	1.8E-08
Lead		35	1.0E-06	5.4E-06	4.7E-06	4.1E-06	1.4E-05	2.4E-05
Naphthalene	91-20-3	1	0	2.8E-10	1.7E-09	0	0	9.9E-09

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-373. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 11<16 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	7.5E-10	1.7E-09	0	4.2E-09	7.5E-09
Arsenic		35	5.1E-10	4.3E-09	3.1E-09	3.9E-09	7.8E-09	1.7E-08
Benz[a]anthracene	56-55-3	35	8.0E-11	1.2E-09	1.3E-09	5.9E-10	3.8E-09	5.4E-09
Benzo[a]pyrene	50-32-8	35	2.0E-10	1.0E-09	8.3E-10	7.2E-10	2.8E-09	3.0E-09
Benzo[b]fluoranthene	205-99-2	35	3.4E-10	1.7E-09	1.2E-09	1.1E-09	4.1E-09	5.0E-09
Benzo[k]fluoranthene	207-08-9	35	4.6E-11	5.5E-10	5.2E-10	3.5E-10	1.6E-09	2.1E-09
Chromium		35	3.7E-09	2.0E-08	2.0E-08	1.3E-08	4.4E-08	1.1E-07
Chrysene	218-01-9	35	1.4E-09	5.5E-09	2.9E-09	5.5E-09	9.4E-09	1.5E-08
Cyclopenta[cd]pyrene	27208-37-3	35	2.1E-10	1.2E-09	1.1E-09	1.0E-09	3.1E-09	5.5E-09
Dibenz[a,h]anthracene	53-70-3	7	0	7.3E-11	1.8E-10	0	3.3E-10	9.1E-10
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	2.1E-10	3.6E-10	0	8.7E-10	1.4E-09
Lead		35	8.2E-08	4.2E-07	3.6E-07	3.2E-07	1.1E-06	1.9E-06
Naphthalene	91-20-3	1	0	2.2E-11	1.3E-10	0	0	7.8E-10

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.



Table F-374. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 16<30 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	4.2E-10	9.7E-10	0	2.4E-09	4.3E-09
Arsenic		35	2.9E-10	2.5E-09	1.7E-09	2.2E-09	4.5E-09	9.5E-09
Benz[a]anthracene	56-55-3	35	4.6E-11	7.0E-10	7.7E-10	3.3E-10	2.2E-09	3.0E-09
Benzo[a]pyrene	50-32-8	35	1.1E-10	5.7E-10	4.7E-10	4.1E-10	1.6E-09	1.7E-09
Benzo[b]fluoranthene	205-99-2	35	2.0E-10	9.6E-10	7.1E-10	6.5E-10	2.3E-09	2.8E-09
Benzo[k]fluoranthene	207-08-9	35	2.6E-11	3.1E-10	2.9E-10	2.0E-10	9.1E-10	1.2E-09
Chromium		35	2.1E-09	1.1E-08	1.1E-08	7.6E-09	2.5E-08	6.3E-08
Chrysene	218-01-9	35	7.8E-10	3.1E-09	1.7E-09	3.1E-09	5.3E-09	8.5E-09
Cyclopenta[cd]pyrene	27208-37-3	35	1.2E-10	6.7E-10	6.2E-10	5.8E-10	1.8E-09	3.1E-09
Dibenz[a,h]anthracene	53-70-3	7	0	4.1E-11	1.0E-10	0	1.9E-10	5.2E-10
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	1.2E-10	2.0E-10	0	5.0E-10	8.0E-10
Lead		35	4.7E-08	2.4E-07	2.1E-07	1.8E-07	6.0E-07	1.1E-06
Naphthalene	91-20-3	1	0	1.3E-11	7.5E-11	0	0	4.4E-10

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-375. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 30<40 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	4.0E-10	9.2E-10	0	2.2E-09	4.0E-09
Arsenic		35	2.7E-10	2.3E-09	1.7E-09	2.1E-09	4.2E-09	9.0E-09
Benz[a]anthracene	56-55-3	35	4.3E-11	6.6E-10	7.2E-10	3.2E-10	2.1E-09	2.9E-09
Benzo[a]pyrene	50-32-8	35	1.1E-10	5.4E-10	4.5E-10	3.9E-10	1.5E-09	1.6E-09
Benzo[b]fluoranthene	205-99-2	35	1.9E-10	9.1E-10	6.7E-10	6.1E-10	2.2E-09	2.7E-09
Benzo[k]fluoranthene	207-08-9	35	2.5E-11	2.9E-10	2.8E-10	1.9E-10	8.6E-10	1.1E-09
Chromium		35	2.0E-09	1.1E-08	1.1E-08	7.2E-09	2.4E-08	6.0E-08
Chrysene	218-01-9	35	7.4E-10	2.9E-09	1.6E-09	2.9E-09	5.0E-09	8.1E-09
Cyclopenta[cd]pyrene	27208-37-3	35	1.1E-10	6.3E-10	5.9E-10	5.5E-10	1.7E-09	3.0E-09



Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Dibenz[a,h]anthracene	53-70-3	7	0	3.9E-11	9.8E-11	0	1.8E-10	4.9E-10
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	1.1E-10	1.9E-10	0	4.7E-10	7.6E-10
Lead		35	4.4E-08	2.3E-07	2.0E-07	1.7E-07	5.7E-07	1.0E-06
Naphthalene	91-20-3	1	0	1.2E-11	7.1E-11	0	0	4.2E-10

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-376. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 40<50 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	3.8E-10	8.7E-10	0	2.1E-09	3.8E-09
Arsenic		35	2.6E-10	2.2E-09	1.6E-09	2.0E-09	4.0E-09	8.5E-09
Benz[a]anthracene	56-55-3	35	4.1E-11	6.3E-10	6.9E-10	3.0E-10	1.9E-09	2.7E-09
Benzo[a]pyrene	50-32-8	35	1.0E-10	5.1E-10	4.2E-10	3.7E-10	1.4E-09	1.5E-09
Benzo[b]fluoranthene	205-99-2	35	1.8E-10	8.6E-10	6.4E-10	5.8E-10	2.1E-09	2.5E-09
Benzo[k]fluoranthene	207-08-9	35	2.3E-11	2.8E-10	2.6E-10	1.8E-10	8.2E-10	1.0E-09
Chromium		35	1.9E-09	1.0E-08	1.0E-08	6.8E-09	2.2E-08	5.7E-08
Chrysene	218-01-9	35	7.0E-10	2.8E-09	1.5E-09	2.8E-09	4.8E-09	7.7E-09
Cyclopenta[cd]pyrene	27208-37-3	35	1.1E-10	6.0E-10	5.5E-10	5.2E-10	1.6E-09	2.8E-09
Dibenz[a,h]anthracene	53-70-3	7	0	3.7E-11	9.2E-11	0	1.7E-10	4.6E-10
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	1.0E-10	1.8E-10	0	4.4E-10	7.2E-10
Lead		35	4.2E-08	2.2E-07	1.9E-07	1.6E-07	5.4E-07	9.7E-07
Naphthalene	91-20-3	1	0	1.1E-11	6.7E-11	0	0	4.0E-10

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.



Table F-377. Field-Specific Average Ingestion Daily Dose^a for Lifetime Cancer Risk Assessment (ADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 50<70 Years**

Chemical	CASRN ^c	Detection	ADD _{ing-field}					
			Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	12	0	3.8E-10	8.7E-10	0	2.1E-09	3.8E-09
Arsenic		35	2.6E-10	2.2E-09	1.6E-09	2.0E-09	4.0E-09	8.5E-09
Benz[a]anthracene	56-55-3	35	4.1E-11	6.3E-10	6.9E-10	3.0E-10	1.9E-09	2.7E-09
Benzo[a]pyrene	50-32-8	35	1.0E-10	5.2E-10	4.2E-10	3.7E-10	1.4E-09	1.5E-09
Benzo[b]fluoranthene	205-99-2	35	1.8E-10	8.6E-10	6.4E-10	5.8E-10	2.1E-09	2.5E-09
Benzo[k]fluoranthene	207-08-9	35	2.4E-11	2.8E-10	2.6E-10	1.8E-10	8.2E-10	1.1E-09
Chromium		35	1.9E-09	1.0E-08	1.0E-08	6.8E-09	2.2E-08	5.7E-08
Chrysene	218-01-9	35	7.0E-10	2.8E-09	1.5E-09	2.8E-09	4.8E-09	7.7E-09
Cyclopenta[cd]pyrene	27208-37-3	35	1.1E-10	6.0E-10	5.6E-10	5.2E-10	1.6E-09	2.8E-09
Dibenz[a,h]anthracene	53-70-3	7	0	3.7E-11	9.3E-11	0	1.7E-10	4.6E-10
Indeno[1,2,3-cd]pyrene	193-39-5	12	0	1.0E-10	1.8E-10	0	4.4E-10	7.2E-10
Lead		35	4.2E-08	2.2E-07	1.9E-07	1.6E-07	5.4E-07	9.8E-07
Naphthalene	91-20-3	1	0	1.1E-11	6.7E-11	0	0	4.0E-10

^a 35 field-specific ADD_{ing-field} are included in the table. ADD_{ing-field} was calculated from field-specific average concentrations (C_{ing-crumb rubber-field}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples from an individual field.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

F.6. Lifetime Average Daily Dose (LADD)

This section presents the lifetime average daily dose (LADD) of all detected chemicals on synthetic turf fields for the inhalation (LADD_{inh}), ingestion (LADD_{ing}), and dermal (LADD_{der}) pathways when toxicity criteria are available for the calculation of cancer risk. Details of each exposure pathway and how to calculate the LADD are presented in Main Report Sections 5.2.3 and 5.4.5. An example calculation for each pathway is presented in Appendix G.1.2.



F.6.1. Inhalation Lifetime Average Daily Dose (LADD_{inh}) for Lifetime Cancer Risk Assessment of Carcinogens

Table F-378. **On-Field** Inhalation Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Athletes**

Chemical	CASRN	LADD _{inh}						
		2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Field-Related Chemicals								
Aniline	62-53-3	4.9E-08	8.0E-08	7.8E-08	1.4E-07	6.4E-08	6.3E-08	1.2E-07
Benz[a]anthracene	56-55-3	3.7E-11	6.1E-11	5.9E-11	1.0E-10	4.9E-11	4.8E-11	9.4E-11
Benzo[a]pyrene	50-32-8	8.7E-09	1.4E-08	1.4E-08	2.4E-08	1.1E-08	1.1E-08	2.2E-08
Benzo[b]fluoranthene	205-99-2	1.9E-10	3.0E-10	3.0E-10	5.2E-10	2.4E-10	2.4E-10	4.7E-10
Benzo[k]fluoranthene	207-08-9	2.3E-10	3.8E-10	3.7E-10	6.5E-10	3.0E-10	3.0E-10	5.9E-10
Chrysene	218-01-9	1.6E-09	2.5E-09	2.5E-09	4.3E-09	2.0E-09	2.0E-09	3.9E-09
Cyclopenta[cd]pyrene	27208-37-3	5.5E-10	8.9E-10	8.7E-10	1.5E-09	7.2E-10	7.1E-10	1.4E-09
Dibenz[a,h]anthracene	53-70-3	1.1E-09	1.8E-09	1.7E-09	3.0E-09	1.4E-09	1.4E-09	2.7E-09
Indeno[1,2,3-cd]pyrene	193-39-5	8.2E-10	1.3E-09	1.3E-09	2.3E-09	1.1E-09	1.0E-09	2.1E-09
Methyl Isobutyl Ketone	108-10-1	1.2E-07	1.9E-07	1.9E-07	3.3E-07	1.5E-07	1.5E-07	3.0E-07
Naphthalene	91-20-3	2.0E-07	3.3E-07	3.2E-07	5.6E-07	2.6E-07	2.6E-07	5.1E-07
Styrene	100-42-5	4.5E-07	7.3E-07	7.1E-07	1.2E-06	5.8E-07	5.7E-07	1.1E-06
Non-Field-Related Chemicals								
Acetaldehyde	75-07-0	1.9E-05	3.1E-05	3.0E-05	5.3E-05	2.5E-05	2.5E-05	4.8E-05
Benzene	71-43-2	4.6E-06	7.5E-06	7.3E-06	1.3E-05	6.0E-06	5.9E-06	1.2E-05
Benzene, 1,4-dichloro	106-46-7	1.5E-07	2.4E-07	2.3E-07	4.1E-07	1.9E-07	1.9E-07	3.7E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	4.4E-06	7.1E-06	6.9E-06	1.2E-05	5.7E-06	5.6E-06	1.1E-05
Ethylbenzene	100-41-4	1.3E-06	2.1E-06	2.1E-06	3.6E-06	1.7E-06	1.7E-06	3.3E-06
Formaldehyde	50-00-0	2.9E-05	4.7E-05	4.6E-05	8.1E-05	3.8E-05	3.7E-05	7.3E-05
Tetrachloroethylene	127-18-4	3.6E-07	5.9E-07	5.8E-07	1.0E-06	4.7E-07	4.7E-07	9.1E-07

^a On-Field LADD_{inh} was calculated from the 35 individual field average concentrations (C_{air-avg}) of a chemical detected in air on the fields and is the sum across all age groups in a receptor category. CASRN: Chemical Abstracts Service Registry Number. Values are rounded to two significant figures.

Table F-379. **On-Field** Inhalation Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Coaches**

Chemical	CASRN	LADD _{inh}			
		16<30 years	30<40 years	40<50 years	50<70 years
Field-Related Chemicals					
Aniline	62-53-3	5.7E-08	3.5E-08	3.6E-08	7.2E-08



Chemical	CASRN	LADD _{inh}			
		16<30 years	30<40 years	40<50 years	50<70 years
Benz[a]anthracene	56-55-3	4.3E-11	2.7E-11	2.7E-11	5.5E-11
Benzo[a]pyrene	50-32-8	1.0E-08	6.2E-09	6.3E-09	1.3E-08
Benzo[b]fluoranthene	205-99-2	2.2E-10	1.3E-10	1.4E-10	2.7E-10
Benzo[k]fluoranthene	207-08-9	2.7E-10	1.7E-10	1.7E-10	3.4E-10
Chrysene	218-01-9	1.8E-09	1.1E-09	1.1E-09	2.3E-09
Cyclopenta[cd]pyrene	27208-37-3	6.4E-10	4.0E-10	4.0E-10	8.0E-10
Dibenz[a,h]anthracene	53-70-3	1.3E-09	7.8E-10	7.9E-10	1.6E-09
Indeno[1,2,3-cd]pyrene	193-39-5	9.4E-10	5.9E-10	5.9E-10	1.2E-09
Methyl Isobutyl Ketone	108-10-1	1.4E-07	8.5E-08	8.6E-08	1.7E-07
Naphthalene	91-20-3	2.3E-07	1.5E-07	1.5E-07	2.9E-07
Styrene	100-42-5	5.1E-07	3.2E-07	3.2E-07	6.5E-07
Non-Field-Related Chemicals					
Acetaldehyde	75-07-0	2.2E-05	1.4E-05	1.4E-05	2.8E-05
Benzene	71-43-2	5.3E-06	3.3E-06	3.3E-06	6.7E-06
Benzene, 1,4-dichloro	106-46-7	1.7E-07	1.0E-07	1.1E-07	2.1E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	5.0E-06	3.1E-06	3.2E-06	6.4E-06
Ethylbenzene	100-41-4	1.5E-06	9.4E-07	9.5E-07	1.9E-06
Formaldehyde	50-00-0	3.3E-05	2.1E-05	2.1E-05	4.2E-05
Tetrachloroethylene	127-18-4	4.2E-07	2.6E-07	2.6E-07	5.3E-07

^a On-Field LADD_{inh} was calculated from the 35 individual field average concentrations ($C_{air-avg}$) of a chemical detected in air on the fields and is the sum across all age groups in a receptor category.
CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures.

Table F-380. **On-Field** Inhalation Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender
Referees

Chemical	CASRN	LADD _{inh}			
		16<30 years	30<40 years	40<50 years	50<70 years
Field-Related Chemicals					
Aniline	62-53-3	2.1E-08	1.3E-08	1.3E-08	2.7E-08
Benz[a]anthracene	56-55-3	1.6E-11	1.0E-11	1.0E-11	2.0E-11
Benzo[a]pyrene	50-32-8	3.7E-09	2.3E-09	2.3E-09	4.7E-09
Benzo[b]fluoranthene	205-99-2	8.0E-11	5.0E-11	5.1E-11	1.0E-10
Benzo[k]fluoranthene	207-08-9	1.0E-10	6.2E-11	6.3E-11	1.3E-10
Chrysene	218-01-9	6.7E-10	4.2E-10	4.2E-10	8.4E-10
Cyclopenta[cd]pyrene	27208-37-3	2.4E-10	1.5E-10	1.5E-10	3.0E-10
Dibenz[a,h]anthracene	53-70-3	4.7E-10	2.9E-10	2.9E-10	5.9E-10
Indeno[1,2,3-cd]pyrene	193-39-5	3.5E-10	2.2E-10	2.2E-10	4.4E-10
Methyl Isobutyl Ketone	108-10-1	5.1E-08	3.2E-08	3.2E-08	6.4E-08
Naphthalene	91-20-3	8.6E-08	5.4E-08	5.4E-08	1.1E-07



Chemical	CASRN	LADD _{inh}			
		16<30 years	30<40 years	40<50 years	50<70 years
Styrene	100-42-5	1.9E-07	1.2E-07	1.2E-07	2.4E-07
Non-Field-Related Chemicals					
Acetaldehyde	75-07-0	8.2E-06	5.1E-06	5.2E-06	1.0E-05
Benzene	71-43-2	2.0E-06	1.2E-06	1.2E-06	2.5E-06
Benzene, 1,4-dichloro	106-46-7	6.2E-08	3.9E-08	3.9E-08	7.9E-08
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	1.9E-06	1.2E-06	1.2E-06	2.4E-06
Ethylbenzene	100-41-4	5.6E-07	3.5E-07	3.5E-07	7.1E-07
Formaldehyde	50-00-0	1.2E-05	7.8E-06	7.8E-06	1.6E-05
Tetrachloroethylene	127-18-4	1.6E-07	9.7E-08	9.8E-08	2.0E-07

^a On-Field LADD_{inh} was calculated from the 35 individual field average concentrations ($C_{air-avg}$) of a chemical detected in air on the fields and is the sum across all age groups in a receptor category.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-381. On-Field Inhalation Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender Spectators

Chemical	CASRN	LADD _{inh}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Field-Related Chemicals										
Aniline	62-53-3	1.4E-09	8.8E-08	2.8E-08	3.3E-08	1.4E-08	8.8E-09	5.5E-09	5.5E-09	1.1E-08
Benz[a]anthracene	56-55-3	1.0E-12	6.7E-11	2.1E-11	2.5E-11	1.1E-11	6.7E-12	4.2E-12	4.2E-12	8.4E-12
Benzo[a]pyrene	50-32-8	2.4E-10	1.5E-08	4.9E-09	5.8E-09	2.5E-09	1.5E-09	9.6E-10	9.7E-10	2.0E-09
Benzo[b]fluoranthene	205-99-2	5.2E-12	3.3E-10	1.1E-10	1.3E-10	5.4E-11	3.3E-11	2.1E-11	2.1E-11	4.2E-11
Benzo[k]fluoranthene	207-08-9	6.4E-12	4.1E-10	1.3E-10	1.6E-10	6.7E-11	4.1E-11	2.6E-11	2.6E-11	5.2E-11
Chrysene	218-01-9	4.3E-11	2.8E-09	8.7E-10	1.0E-09	4.5E-10	2.8E-10	1.7E-10	1.7E-10	3.5E-10
Cyclopenta[cd]pyrene	27208-37-3	1.5E-11	9.8E-10	3.1E-10	3.7E-10	1.6E-10	9.8E-11	6.1E-11	6.2E-11	1.2E-10
Dibenz[a,h]anthracene	53-70-3	3.0E-11	1.9E-09	6.1E-10	7.3E-10	3.1E-10	1.9E-10	1.2E-10	1.2E-10	2.5E-10
Indeno[1,2,3-cd]pyrene	193-39-5	2.3E-11	1.5E-09	4.6E-10	5.5E-10	2.4E-10	1.5E-10	9.1E-11	9.2E-11	1.8E-10
Methyl Isobutyl Ketone	108-10-1	3.3E-09	2.1E-07	6.7E-08	8.0E-08	3.4E-08	2.1E-08	1.3E-08	1.3E-08	2.7E-08
Naphthalene	91-20-3	5.6E-09	3.6E-07	1.1E-07	1.4E-07	5.8E-08	3.6E-08	2.2E-08	2.3E-08	4.6E-08
Styrene	100-42-5	1.2E-08	8.0E-07	2.5E-07	3.0E-07	1.3E-07	8.0E-08	5.0E-08	5.0E-08	1.0E-07
Non-Field-Related Chemicals										
Acetaldehyde	75-07-0	5.3E-07	3.4E-05	1.1E-05	1.3E-05	5.5E-06	3.4E-06	2.1E-06	2.2E-06	4.3E-06
Benzene	71-43-2	1.3E-07	8.2E-06	2.6E-06	3.1E-06	1.3E-06	8.2E-07	5.1E-07	5.2E-07	1.0E-06
Benzene, 1,4-dichloro	106-46-7	4.0E-09	2.6E-07	8.2E-08	9.8E-08	4.2E-08	2.6E-08	1.6E-08	1.6E-08	3.3E-08
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	1.2E-07	7.8E-06	2.5E-06	2.9E-06	1.3E-06	7.8E-07	4.9E-07	4.9E-07	9.8E-07
Ethylbenzene	100-41-4	3.6E-08	2.3E-06	7.3E-07	8.7E-07	3.8E-07	2.3E-07	1.5E-07	1.5E-07	2.9E-07
Formaldehyde	50-00-0	8.1E-07	5.2E-05	1.6E-05	1.9E-05	8.4E-06	5.2E-06	3.2E-06	3.3E-06	6.6E-06
Tetrachloroethylene	127-18-4	1.0E-08	6.5E-07	2.0E-07	2.4E-07	1.1E-07	6.5E-08	4.0E-08	4.1E-08	8.2E-08



^a On-Field LADD_{inh} was calculated from the 35 individual field average concentrations (C_{air-avg}) of a chemical detected in air on the fields and is the sum across all age groups in a receptor category.
CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures.

Table F-382. Off-Field Inhalation Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender Spectators

Chemical	CASRN	LADD _{inh}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Field-Related Chemicals										
Aniline	62-53-3	1.5E-09	9.7E-08	3.1E-08	3.6E-08	1.6E-08	9.7E-09	6.0E-09	6.1E-09	1.2E-08
Benzo[a]pyrene	50-32-8	2.9E-10	1.9E-08	6.0E-09	7.1E-09	3.1E-09	1.9E-09	1.2E-09	1.2E-09	2.4E-09
Benzo[b]fluoranthene	205-99-2	8.0E-12	5.2E-10	1.6E-10	1.9E-10	8.4E-11	5.2E-11	3.2E-11	3.3E-11	6.5E-11
Benzo[k]fluoranthene	207-08-9	6.6E-12	4.3E-10	1.3E-10	1.6E-10	6.9E-11	4.3E-11	2.7E-11	2.7E-11	5.4E-11
Chrysene	218-01-9	2.2E-11	1.4E-09	4.5E-10	5.3E-10	2.3E-10	1.4E-10	8.9E-11	9.0E-11	1.8E-10
Cyclopenta[cd]pyrene	27208-37-3	1.3E-11	8.6E-10	2.7E-10	3.2E-10	1.4E-10	8.6E-11	5.4E-11	5.4E-11	1.1E-10
Dibenz[a,h]anthracene	53-70-3	1.9E-11	1.2E-09	3.8E-10	4.6E-10	2.0E-10	1.2E-10	7.6E-11	7.7E-11	1.5E-10
Indeno[1,2,3-cd]pyrene	193-39-5	1.6E-11	1.0E-09	3.2E-10	3.8E-10	1.6E-10	1.0E-10	6.2E-11	6.3E-11	1.3E-10
Methyl Isobutyl Ketone	108-10-1	7.5E-10	4.8E-08	1.5E-08	1.8E-08	7.9E-09	4.8E-09	3.0E-09	3.1E-09	6.1E-09
Naphthalene	91-20-3	6.1E-09	3.9E-07	1.2E-07	1.5E-07	6.4E-08	3.9E-08	2.5E-08	2.5E-08	5.0E-08
Styrene	100-42-5	1.3E-08	8.1E-07	2.6E-07	3.1E-07	1.3E-07	8.1E-08	5.1E-08	5.1E-08	1.0E-07
Non-Field-Related Chemicals										
Benzene	71-43-2	1.3E-07	8.6E-06	2.7E-06	3.2E-06	1.4E-06	8.6E-07	5.4E-07	5.4E-07	1.1E-06
Benzene, 1,4-dichloro	106-46-7	3.7E-09	2.4E-07	7.4E-08	8.9E-08	3.8E-08	2.4E-08	1.5E-08	1.5E-08	3.0E-08
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	1.2E-07	7.8E-06	2.5E-06	2.9E-06	1.3E-06	7.8E-07	4.9E-07	4.9E-07	9.9E-07
Ethylbenzene	100-41-4	3.8E-08	2.5E-06	7.8E-07	9.3E-07	4.0E-07	2.5E-07	1.5E-07	1.6E-07	3.1E-07
Tetrachloroethylene	127-18-4	1.0E-08	6.7E-07	2.1E-07	2.5E-07	1.1E-07	6.7E-08	4.2E-08	4.2E-08	8.4E-08

^a Off-Field LADD_{inh} was calculated from the 35 individual field average concentrations (C_{air-avg}) of a chemical detected in air on the fields and is the sum across all age groups in a receptor category.
CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures.

INDIVIDUAL FIELD ASSESSMENT (Table F-383 to Table F-415)

Table F-383. Field-Specific Inhalation On-Field Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender Athletes 2<6 Years

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	4.8E-08	8.6E-08	0.0E+00	1.9E-07	3.3E-07



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benz[a]anthracene	56-55-3	0.0E+00	3.6E-11	2.2E-10	0.0E+00	0.0E+00	1.3E-09
Benzo[a]pyrene	50-32-8	0.0E+00	8.4E-09	9.1E-09	4.4E-09	2.3E-08	3.6E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	1.8E-10	7.5E-10	0.0E+00	9.5E-10	3.2E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	2.3E-10	6.8E-10	0.0E+00	1.7E-09	2.6E-09
Chrysene	218-01-9	0.0E+00	1.5E-09	2.4E-09	0.0E+00	5.7E-09	1.0E-08
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	5.4E-10	6.7E-10	2.1E-10	1.6E-09	3.1E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.1E-09	2.1E-09	0.0E+00	5.4E-09	9.2E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	7.9E-10	2.6E-09	0.0E+00	9.3E-09	9.3E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	1.2E-07	3.2E-07	0.0E+00	9.2E-07	1.4E-06
Naphthalene	91-20-3	0.0E+00	2.0E-07	4.2E-07	0.0E+00	8.7E-07	2.0E-06
Styrene	100-42-5	0.0E+00	4.5E-07	9.2E-07	0.0E+00	1.5E-06	5.0E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	1.9E-05	1.4E-05	1.3E-05	3.7E-05	7.3E-05
Benzene	71-43-2	6.9E-07	4.6E-06	3.7E-06	3.2E-06	1.1E-05	1.9E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	1.5E-07	3.2E-07	0.0E+00	9.0E-07	9.0E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	4.4E-06	4.5E-06	2.9E-06	1.3E-05	1.6E-05
Ethylbenzene	100-41-4	0.0E+00	1.3E-06	2.2E-06	0.0E+00	5.3E-06	8.8E-06
Formaldehyde	50-00-0	0.0E+00	2.8E-05	2.2E-05	2.4E-05	4.8E-05	1.2E-04
Tetrachloroethylene	127-18-4	0.0E+00	3.6E-07	8.6E-07	0.0E+00	2.4E-06	3.2E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-384. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Athletes 6<11 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	7.8E-08	1.4E-07	0.0E+00	3.1E-07	5.4E-07
Benz[a]anthracene	56-55-3	0.0E+00	5.9E-11	3.5E-10	0.0E+00	0.0E+00	2.1E-09
Benzo[a]pyrene	50-32-8	0.0E+00	1.4E-08	1.5E-08	7.2E-09	3.8E-08	5.8E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	3.0E-10	1.2E-09	0.0E+00	1.5E-09	5.2E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	3.7E-10	1.1E-09	0.0E+00	2.8E-09	4.3E-09
Chrysene	218-01-9	0.0E+00	2.5E-09	3.9E-09	0.0E+00	9.3E-09	1.6E-08



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	8.7E-10	1.1E-09	3.3E-10	2.6E-09	5.1E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.7E-09	3.5E-09	0.0E+00	8.8E-09	1.5E-08
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.3E-09	4.3E-09	0.0E+00	1.5E-08	1.5E-08
Methyl Isobutyl Ketone	108-10-1	0.0E+00	1.9E-07	5.3E-07	0.0E+00	1.5E-06	2.2E-06
Naphthalene	91-20-3	0.0E+00	3.3E-07	6.8E-07	0.0E+00	1.4E-06	3.2E-06
Styrene	100-42-5	0.0E+00	7.3E-07	1.5E-06	0.0E+00	2.4E-06	8.1E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	3.0E-05	2.3E-05	2.1E-05	5.9E-05	1.2E-04
Benzene	71-43-2	1.1E-06	7.5E-06	6.0E-06	5.2E-06	1.9E-05	3.0E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	2.4E-07	5.2E-07	0.0E+00	1.5E-06	1.5E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	7.1E-06	7.4E-06	4.7E-06	2.1E-05	2.6E-05
Ethylbenzene	100-41-4	0.0E+00	2.1E-06	3.6E-06	0.0E+00	8.6E-06	1.4E-05
Formaldehyde	50-00-0	0.0E+00	4.6E-05	3.6E-05	3.9E-05	7.9E-05	2.0E-04
Tetrachloroethylene	127-18-4	0.0E+00	5.9E-07	1.4E-06	0.0E+00	3.9E-06	5.2E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-385. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Athletes 11<16 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	7.6E-08	1.4E-07	0.0E+00	3.0E-07	5.3E-07
Benz[a]anthracene	56-55-3	0.0E+00	5.8E-11	3.4E-10	0.0E+00	0.0E+00	2.0E-09
Benzo[a]pyrene	50-32-8	0.0E+00	1.3E-08	1.4E-08	7.0E-09	3.7E-08	5.7E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	2.9E-10	1.2E-09	0.0E+00	1.5E-09	5.0E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	3.6E-10	1.1E-09	0.0E+00	2.7E-09	4.2E-09
Chrysene	218-01-9	0.0E+00	2.4E-09	3.8E-09	0.0E+00	9.1E-09	1.6E-08
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	8.5E-10	1.1E-09	3.3E-10	2.5E-09	5.0E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.7E-09	3.4E-09	0.0E+00	8.6E-09	1.5E-08
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.3E-09	4.2E-09	0.0E+00	1.5E-08	1.5E-08
Methyl Isobutyl Ketone	108-10-1	0.0E+00	1.9E-07	5.1E-07	0.0E+00	1.5E-06	2.2E-06
Naphthalene	91-20-3	0.0E+00	3.2E-07	6.7E-07	0.0E+00	1.4E-06	3.1E-06
Styrene	100-42-5	0.0E+00	7.1E-07	1.5E-06	0.0E+00	2.4E-06	7.9E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	2.9E-05	2.3E-05	2.1E-05	5.8E-05	1.2E-04



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene	71-43-2	1.1E-06	7.3E-06	5.9E-06	5.1E-06	1.8E-05	3.0E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	2.3E-07	5.1E-07	0.0E+00	1.4E-06	1.4E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	6.9E-06	7.2E-06	4.6E-06	2.1E-05	2.5E-05
Ethylbenzene	100-41-4	0.0E+00	2.1E-06	3.5E-06	0.0E+00	8.4E-06	1.4E-05
Formaldehyde	50-00-0	0.0E+00	4.5E-05	3.5E-05	3.8E-05	7.7E-05	2.0E-04
Tetrachloroethylene	127-18-4	0.0E+00	5.8E-07	1.4E-06	0.0E+00	3.9E-06	5.0E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration ($C_{air-field}$) of a chemical detected in air off an individual field
CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures.

Table F-386. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Athletes 16<30 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	1.3E-07	2.4E-07	0.0E+00	5.3E-07	9.2E-07
Benz[a]anthracene	56-55-3	0.0E+00	1.0E-10	6.0E-10	0.0E+00	0.0E+00	3.5E-09
Benzo[a]pyrene	50-32-8	0.0E+00	2.3E-08	2.5E-08	1.2E-08	6.5E-08	1.0E-07
Benzo[b]fluoranthene	205-99-2	0.0E+00	5.1E-10	2.1E-09	0.0E+00	2.7E-09	8.9E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	6.3E-10	1.9E-09	0.0E+00	4.8E-09	7.4E-09
Chrysene	218-01-9	0.0E+00	4.2E-09	6.7E-09	0.0E+00	1.6E-08	2.8E-08
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	1.5E-09	1.9E-09	5.7E-10	4.4E-09	8.7E-09
Dibenzo[a,h]anthracene	53-70-3	0.0E+00	2.9E-09	6.0E-09	0.0E+00	1.5E-08	2.6E-08
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.2E-09	7.3E-09	0.0E+00	2.6E-08	2.6E-08
Methyl Isobutyl Ketone	108-10-1	0.0E+00	3.3E-07	9.0E-07	0.0E+00	2.6E-06	3.9E-06
Naphthalene	91-20-3	0.0E+00	5.6E-07	1.2E-06	0.0E+00	2.4E-06	5.5E-06
Styrene	100-42-5	0.0E+00	1.2E-06	2.6E-06	0.0E+00	4.2E-06	1.4E-05
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	5.2E-05	4.0E-05	3.7E-05	1.0E-04	2.0E-04
Benzene	71-43-2	1.9E-06	1.3E-05	1.0E-05	9.0E-06	3.2E-05	5.2E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	4.1E-07	8.9E-07	0.0E+00	2.5E-06	2.5E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	1.2E-05	1.3E-05	8.1E-06	3.6E-05	4.5E-05
Ethylbenzene	100-41-4	0.0E+00	3.6E-06	6.1E-06	0.0E+00	1.5E-05	2.4E-05
Formaldehyde	50-00-0	0.0E+00	7.9E-05	6.2E-05	6.8E-05	1.4E-04	3.5E-04
Tetrachloroethylene	127-18-4	0.0E+00	1.0E-06	2.4E-06	0.0E+00	6.8E-06	8.9E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration ($C_{air-field}$) of a chemical detected in air off an individual field



CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures.

Table F-387. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Athletes 30<40 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	6.2E-08	1.1E-07	0.0E+00	2.5E-07	4.3E-07
Benz[a]anthracene	56-55-3	0.0E+00	4.7E-11	2.8E-10	0.0E+00	0.0E+00	1.7E-09
Benzo[a]pyrene	50-32-8	0.0E+00	1.1E-08	1.2E-08	5.8E-09	3.0E-08	4.7E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	2.4E-10	9.7E-10	0.0E+00	1.2E-09	4.1E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	2.9E-10	8.8E-10	0.0E+00	2.2E-09	3.4E-09
Chrysene	218-01-9	0.0E+00	2.0E-09	3.1E-09	0.0E+00	7.4E-09	1.3E-08
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	7.0E-10	8.7E-10	2.7E-10	2.1E-09	4.1E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.4E-09	2.8E-09	0.0E+00	7.1E-09	1.2E-08
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.0E-09	3.4E-09	0.0E+00	1.2E-08	1.2E-08
Methyl Isobutyl Ketone	108-10-1	0.0E+00	1.5E-07	4.2E-07	0.0E+00	1.2E-06	1.8E-06
Naphthalene	91-20-3	0.0E+00	2.6E-07	5.5E-07	0.0E+00	1.1E-06	2.6E-06
Styrene	100-42-5	0.0E+00	5.8E-07	1.2E-06	0.0E+00	2.0E-06	6.5E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	2.4E-05	1.9E-05	1.7E-05	4.8E-05	9.5E-05
Benzene	71-43-2	8.9E-07	6.0E-06	4.8E-06	4.2E-06	1.5E-05	2.4E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	1.9E-07	4.2E-07	0.0E+00	1.2E-06	1.2E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	5.7E-06	5.9E-06	3.8E-06	1.7E-05	2.1E-05
Ethylbenzene	100-41-4	0.0E+00	1.7E-06	2.9E-06	0.0E+00	6.9E-06	1.1E-05
Formaldehyde	50-00-0	0.0E+00	3.7E-05	2.9E-05	3.2E-05	6.3E-05	1.6E-04
Tetrachloroethylene	127-18-4	0.0E+00	4.7E-07	1.1E-06	0.0E+00	3.2E-06	4.1E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field

CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures.

Table F-388. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Athletes 40<50 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	6.1E-08	1.1E-07	0.0E+00	2.5E-07	4.2E-07
Benz[a]anthracene	56-55-3	0.0E+00	4.7E-11	2.8E-10	0.0E+00	0.0E+00	1.6E-09



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[a]pyrene	50-32-8	0.0E+00	1.1E-08	1.2E-08	5.7E-09	3.0E-08	4.6E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	2.3E-10	9.6E-10	0.0E+00	1.2E-09	4.1E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	2.9E-10	8.7E-10	0.0E+00	2.2E-09	3.4E-09
Chrysene	218-01-9	0.0E+00	1.9E-09	3.1E-09	0.0E+00	7.3E-09	1.3E-08
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	6.9E-10	8.6E-10	2.6E-10	2.0E-09	4.0E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.4E-09	2.8E-09	0.0E+00	7.0E-09	1.2E-08
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.0E-09	3.4E-09	0.0E+00	1.2E-08	1.2E-08
Methyl Isobutyl Ketone	108-10-1	0.0E+00	1.5E-07	4.2E-07	0.0E+00	1.2E-06	1.8E-06
Naphthalene	91-20-3	0.0E+00	2.6E-07	5.4E-07	0.0E+00	1.1E-06	2.5E-06
Styrene	100-42-5	0.0E+00	5.7E-07	1.2E-06	0.0E+00	1.9E-06	6.4E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	2.4E-05	1.8E-05	1.7E-05	4.7E-05	9.3E-05
Benzene	71-43-2	8.8E-07	5.9E-06	4.8E-06	4.1E-06	1.5E-05	2.4E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	1.9E-07	4.1E-07	0.0E+00	1.2E-06	1.2E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	5.6E-06	5.8E-06	3.7E-06	1.7E-05	2.1E-05
Ethylbenzene	100-41-4	0.0E+00	1.7E-06	2.8E-06	0.0E+00	6.8E-06	1.1E-05
Formaldehyde	50-00-0	0.0E+00	3.6E-05	2.8E-05	3.1E-05	6.2E-05	1.6E-04
Tetrachloroethylene	127-18-4	0.0E+00	4.7E-07	1.1E-06	0.0E+00	3.1E-06	4.1E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-389. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Athletes 50<70 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	1.2E-07	2.2E-07	0.0E+00	4.8E-07	8.3E-07
Benz[a]anthracene	56-55-3	0.0E+00	9.2E-11	5.4E-10	0.0E+00	0.0E+00	3.2E-09
Benzo[a]pyrene	50-32-8	0.0E+00	2.1E-08	2.3E-08	1.1E-08	5.9E-08	9.0E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	4.6E-10	1.9E-09	0.0E+00	2.4E-09	8.0E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	5.7E-10	1.7E-09	0.0E+00	4.3E-09	6.6E-09
Chrysene	218-01-9	0.0E+00	3.8E-09	6.0E-09	0.0E+00	1.4E-08	2.5E-08
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	1.3E-09	1.7E-09	5.2E-10	4.0E-09	7.9E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	2.7E-09	5.4E-09	0.0E+00	1.4E-08	2.3E-08
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.0E-09	6.6E-09	0.0E+00	2.3E-08	2.3E-08
Methyl Isobutyl Ketone	108-10-1	0.0E+00	3.0E-07	8.2E-07	0.0E+00	2.3E-06	3.5E-06



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Naphthalene	91-20-3	0.0E+00	5.1E-07	1.1E-06	0.0E+00	2.2E-06	5.0E-06
Styrene	100-42-5	0.0E+00	1.1E-06	2.3E-06	0.0E+00	3.8E-06	1.3E-05
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	4.7E-05	3.6E-05	3.3E-05	9.2E-05	1.8E-04
Benzene	71-43-2	1.7E-06	1.2E-05	9.4E-06	8.1E-06	2.9E-05	4.7E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	3.7E-07	8.1E-07	0.0E+00	2.3E-06	2.3E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	1.1E-05	1.1E-05	7.3E-06	3.3E-05	4.1E-05
Ethylbenzene	100-41-4	0.0E+00	3.3E-06	5.5E-06	0.0E+00	1.3E-05	2.2E-05
Formaldehyde	50-00-0	0.0E+00	7.1E-05	5.6E-05	6.1E-05	1.2E-04	3.1E-04
Tetrachloroethylene	127-18-4	0.0E+00	9.1E-07	2.2E-06	0.0E+00	6.1E-06	8.0E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-390. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Coaches 16<30 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	5.5E-08	9.9E-08	0.0E+00	2.2E-07	3.8E-07
Benz[a]anthracene	56-55-3	0.0E+00	4.2E-11	2.5E-10	0.0E+00	0.0E+00	1.5E-09
Benzo[a]pyrene	50-32-8	0.0E+00	9.7E-09	1.0E-08	5.1E-09	2.7E-08	4.1E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	2.1E-10	8.6E-10	0.0E+00	1.1E-09	3.7E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	2.6E-10	7.8E-10	0.0E+00	2.0E-09	3.0E-09
Chrysene	218-01-9	0.0E+00	1.7E-09	2.8E-09	0.0E+00	6.6E-09	1.2E-08
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	6.2E-10	7.7E-10	2.4E-10	1.8E-09	3.6E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.2E-09	2.5E-09	0.0E+00	6.3E-09	1.1E-08
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	9.1E-10	3.0E-09	0.0E+00	1.1E-08	1.1E-08
Methyl Isobutyl Ketone	108-10-1	0.0E+00	1.4E-07	3.7E-07	0.0E+00	1.1E-06	1.6E-06
Naphthalene	91-20-3	0.0E+00	2.3E-07	4.9E-07	0.0E+00	1.0E-06	2.3E-06
Styrene	100-42-5	0.0E+00	5.1E-07	1.1E-06	0.0E+00	1.7E-06	5.8E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	2.1E-05	1.6E-05	1.5E-05	4.2E-05	8.4E-05
Benzene	71-43-2	7.9E-07	5.3E-06	4.3E-06	3.7E-06	1.3E-05	2.1E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	1.7E-07	3.7E-07	0.0E+00	1.0E-06	1.0E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	5.0E-06	5.2E-06	3.3E-06	1.5E-05	1.9E-05
Ethylbenzene	100-41-4	0.0E+00	1.5E-06	2.5E-06	0.0E+00	6.1E-06	1.0E-05



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Formaldehyde	50-00-0	0.0E+00	3.3E-05	2.5E-05	2.8E-05	5.6E-05	1.4E-04
Tetrachloroethylene	127-18-4	0.0E+00	4.2E-07	9.9E-07	0.0E+00	2.8E-06	3.7E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-391. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Coaches 30<40 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	3.4E-08	6.2E-08	0.0E+00	1.4E-07	2.4E-07
Benz[a]anthracene	56-55-3	0.0E+00	2.6E-11	1.5E-10	0.0E+00	0.0E+00	9.2E-10
Benzo[a]pyrene	50-32-8	0.0E+00	6.1E-09	6.5E-09	3.2E-09	1.7E-08	2.6E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	1.3E-10	5.4E-10	0.0E+00	6.9E-10	2.3E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	1.6E-10	4.9E-10	0.0E+00	1.2E-09	1.9E-09
Chrysene	218-01-9	0.0E+00	1.1E-09	1.7E-09	0.0E+00	4.1E-09	7.2E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	3.9E-10	4.8E-10	1.5E-10	1.1E-09	2.3E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	7.6E-10	1.5E-09	0.0E+00	3.9E-09	6.6E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	5.7E-10	1.9E-09	0.0E+00	6.7E-09	6.7E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	8.5E-08	2.3E-07	0.0E+00	6.6E-07	1.0E-06
Naphthalene	91-20-3	0.0E+00	1.5E-07	3.0E-07	0.0E+00	6.3E-07	1.4E-06
Styrene	100-42-5	0.0E+00	3.2E-07	6.6E-07	0.0E+00	1.1E-06	3.6E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	1.3E-05	1.0E-05	9.5E-06	2.6E-05	5.2E-05
Benzene	71-43-2	4.9E-07	3.3E-06	2.7E-06	2.3E-06	8.3E-06	1.3E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	1.0E-07	2.3E-07	0.0E+00	6.5E-07	6.5E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	3.1E-06	3.3E-06	2.1E-06	9.3E-06	1.2E-05
Ethylbenzene	100-41-4	0.0E+00	9.4E-07	1.6E-06	0.0E+00	3.8E-06	6.3E-06
Formaldehyde	50-00-0	0.0E+00	2.0E-05	1.6E-05	1.7E-05	3.5E-05	8.9E-05
Tetrachloroethylene	127-18-4	0.0E+00	2.6E-07	6.2E-07	0.0E+00	1.8E-06	2.3E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-392. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Coaches 40<50 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	3.5E-08	6.2E-08	0.0E+00	1.4E-07	2.4E-07
Benz[a]anthracene	56-55-3	0.0E+00	2.6E-11	1.6E-10	0.0E+00	0.0E+00	9.2E-10
Benzo[a]pyrene	50-32-8	0.0E+00	6.1E-09	6.6E-09	3.2E-09	1.7E-08	2.6E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	1.3E-10	5.4E-10	0.0E+00	6.9E-10	2.3E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	1.6E-10	4.9E-10	0.0E+00	1.2E-09	1.9E-09
Chrysene	218-01-9	0.0E+00	1.1E-09	1.7E-09	0.0E+00	4.2E-09	7.3E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	3.9E-10	4.8E-10	1.5E-10	1.2E-09	2.3E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	7.7E-10	1.6E-09	0.0E+00	3.9E-09	6.7E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	5.8E-10	1.9E-09	0.0E+00	6.7E-09	6.7E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	8.6E-08	2.4E-07	0.0E+00	6.7E-07	1.0E-06
Naphthalene	91-20-3	0.0E+00	1.5E-07	3.1E-07	0.0E+00	6.3E-07	1.4E-06
Styrene	100-42-5	0.0E+00	3.2E-07	6.7E-07	0.0E+00	1.1E-06	3.6E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	1.4E-05	1.0E-05	9.6E-06	2.7E-05	5.3E-05
Benzene	71-43-2	5.0E-07	3.3E-06	2.7E-06	2.4E-06	8.3E-06	1.4E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	1.1E-07	2.3E-07	0.0E+00	6.5E-07	6.5E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	3.2E-06	3.3E-06	2.1E-06	9.4E-06	1.2E-05
Ethylbenzene	100-41-4	0.0E+00	9.5E-07	1.6E-06	0.0E+00	3.9E-06	6.4E-06
Formaldehyde	50-00-0	0.0E+00	2.1E-05	1.6E-05	1.8E-05	3.5E-05	9.0E-05
Tetrachloroethylene	127-18-4	0.0E+00	2.6E-07	6.2E-07	0.0E+00	1.8E-06	2.3E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-393. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Coaches 50<70 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	7.0E-08	1.2E-07	0.0E+00	2.8E-07	4.8E-07
Benz[a]anthracene	56-55-3	0.0E+00	5.3E-11	3.1E-10	0.0E+00	0.0E+00	1.9E-09
Benzo[a]pyrene	50-32-8	0.0E+00	1.2E-08	1.3E-08	6.4E-09	3.4E-08	5.2E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	2.7E-10	1.1E-09	0.0E+00	1.4E-09	4.6E-09



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[k]fluoranthene	207-08-9	0.0E+00	3.3E-10	9.9E-10	0.0E+00	2.5E-09	3.8E-09
Chrysene	218-01-9	0.0E+00	2.2E-09	3.5E-09	0.0E+00	8.3E-09	1.5E-08
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	7.8E-10	9.7E-10	3.0E-10	2.3E-09	4.6E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.5E-09	3.1E-09	0.0E+00	7.9E-09	1.3E-08
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.2E-09	3.8E-09	0.0E+00	1.3E-08	1.3E-08
Methyl Isobutyl Ketone	108-10-1	0.0E+00	1.7E-07	4.7E-07	0.0E+00	1.3E-06	2.0E-06
Naphthalene	91-20-3	0.0E+00	2.9E-07	6.1E-07	0.0E+00	1.3E-06	2.9E-06
Styrene	100-42-5	0.0E+00	6.5E-07	1.3E-06	0.0E+00	2.2E-06	7.3E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	2.7E-05	2.1E-05	1.9E-05	5.3E-05	1.1E-04
Benzene	71-43-2	1.0E-06	6.7E-06	5.4E-06	4.7E-06	1.7E-05	2.7E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	2.1E-07	4.7E-07	0.0E+00	1.3E-06	1.3E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	6.4E-06	6.6E-06	4.2E-06	1.9E-05	2.3E-05
Ethylbenzene	100-41-4	0.0E+00	1.9E-06	3.2E-06	0.0E+00	7.8E-06	1.3E-05
Formaldehyde	50-00-0	0.0E+00	4.1E-05	3.2E-05	3.5E-05	7.1E-05	1.8E-04
Tetrachloroethylene	127-18-4	0.0E+00	5.3E-07	1.3E-06	0.0E+00	3.5E-06	4.6E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration ($C_{air-field}$) of a chemical detected in air off an individual field

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-394. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Referees 16<30 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	2.0E-08	3.7E-08	0.0E+00	8.2E-08	1.4E-07
Benz[a]anthracene	56-55-3	0.0E+00	1.6E-11	9.2E-11	0.0E+00	0.0E+00	5.4E-10
Benzo[a]pyrene	50-32-8	0.0E+00	3.6E-09	3.9E-09	1.9E-09	1.0E-08	1.5E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	7.8E-11	3.2E-10	0.0E+00	4.1E-10	1.4E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	9.7E-11	2.9E-10	0.0E+00	7.3E-10	1.1E-09
Chrysene	218-01-9	0.0E+00	6.5E-10	1.0E-09	0.0E+00	2.4E-09	4.3E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	2.3E-10	2.9E-10	8.8E-11	6.8E-10	1.3E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	4.5E-10	9.2E-10	0.0E+00	2.3E-09	3.9E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	3.4E-10	1.1E-09	0.0E+00	4.0E-09	4.0E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	5.1E-08	1.4E-07	0.0E+00	4.0E-07	5.9E-07
Naphthalene	91-20-3	0.0E+00	8.6E-08	1.8E-07	0.0E+00	3.7E-07	8.4E-07
Styrene	100-42-5	0.0E+00	1.9E-07	3.9E-07	0.0E+00	6.4E-07	2.1E-06



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	8.0E-06	6.1E-06	5.6E-06	1.6E-05	3.1E-05
Benzene	71-43-2	2.9E-07	2.0E-06	1.6E-06	1.4E-06	4.9E-06	8.0E-06
Benzene, 1,4-dichloro	106-46-7	0.0E+00	6.2E-08	1.4E-07	0.0E+00	3.8E-07	3.8E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	1.9E-06	1.9E-06	1.2E-06	5.6E-06	6.9E-06
Ethylbenzene	100-41-4	0.0E+00	5.6E-07	9.4E-07	0.0E+00	2.3E-06	3.8E-06
Formaldehyde	50-00-0	0.0E+00	1.2E-05	9.4E-06	1.0E-05	2.1E-05	5.3E-05
Tetrachloroethylene	127-18-4	0.0E+00	1.6E-07	3.7E-07	0.0E+00	1.0E-06	1.4E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-395. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Referees 30<40 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	1.3E-08	2.3E-08	0.0E+00	5.1E-08	8.8E-08
Benz[a]anthracene	56-55-3	0.0E+00	9.7E-12	5.7E-11	0.0E+00	0.0E+00	3.4E-10
Benzo[a]pyrene	50-32-8	0.0E+00	2.2E-09	2.4E-09	1.2E-09	6.2E-09	9.6E-09
Benzo[b]fluoranthene	205-99-2	0.0E+00	4.9E-11	2.0E-10	0.0E+00	2.5E-10	8.5E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	6.0E-11	1.8E-10	0.0E+00	4.6E-10	7.0E-10
Chrysene	218-01-9	0.0E+00	4.0E-10	6.4E-10	0.0E+00	1.5E-09	2.7E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	1.4E-10	1.8E-10	5.5E-11	4.3E-10	8.4E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	2.8E-10	5.7E-10	0.0E+00	1.4E-09	2.4E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.1E-10	7.0E-10	0.0E+00	2.5E-09	2.5E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	3.2E-08	8.6E-08	0.0E+00	2.5E-07	3.7E-07
Naphthalene	91-20-3	0.0E+00	5.4E-08	1.1E-07	0.0E+00	2.3E-07	5.3E-07
Styrene	100-42-5	0.0E+00	1.2E-07	2.5E-07	0.0E+00	4.0E-07	1.3E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	5.0E-06	3.8E-06	3.5E-06	9.8E-06	1.9E-05
Benzene	71-43-2	1.8E-07	1.2E-06	9.9E-07	8.6E-07	3.1E-06	5.0E-06
Benzene, 1,4-dichloro	106-46-7	0.0E+00	3.9E-08	8.5E-08	0.0E+00	2.4E-07	2.4E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	1.2E-06	1.2E-06	7.7E-07	3.5E-06	4.3E-06
Ethylbenzene	100-41-4	0.0E+00	3.5E-07	5.9E-07	0.0E+00	1.4E-06	2.3E-06
Formaldehyde	50-00-0	0.0E+00	7.5E-06	5.9E-06	6.5E-06	1.3E-05	3.3E-05
Tetrachloroethylene	127-18-4	0.0E+00	9.7E-08	2.3E-07	0.0E+00	6.5E-07	8.5E-07



^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field
CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures.

Table F-396. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Referees 40<50 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	1.3E-08	2.3E-08	0.0E+00	5.2E-08	8.9E-08
Benz[a]anthracene	56-55-3	0.0E+00	9.8E-12	5.8E-11	0.0E+00	0.0E+00	3.4E-10
Benzo[a]pyrene	50-32-8	0.0E+00	2.3E-09	2.5E-09	1.2E-09	6.3E-09	9.7E-09
Benzo[b]fluoranthene	205-99-2	0.0E+00	4.9E-11	2.0E-10	0.0E+00	2.6E-10	8.6E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	6.1E-11	1.8E-10	0.0E+00	4.6E-10	7.1E-10
Chrysene	218-01-9	0.0E+00	4.1E-10	6.5E-10	0.0E+00	1.5E-09	2.7E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	1.4E-10	1.8E-10	5.6E-11	4.3E-10	8.5E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	2.9E-10	5.8E-10	0.0E+00	1.5E-09	2.5E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.1E-10	7.1E-10	0.0E+00	2.5E-09	2.5E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	3.2E-08	8.7E-08	0.0E+00	2.5E-07	3.7E-07
Naphthalene	91-20-3	0.0E+00	5.4E-08	1.1E-07	0.0E+00	2.4E-07	5.3E-07
Styrene	100-42-5	0.0E+00	1.2E-07	2.5E-07	0.0E+00	4.1E-07	1.3E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	5.0E-06	3.8E-06	3.6E-06	9.9E-06	2.0E-05
Benzene	71-43-2	1.9E-07	1.2E-06	1.0E-06	8.7E-07	3.1E-06	5.0E-06
Benzene, 1,4-dichloro	106-46-7	0.0E+00	3.9E-08	8.6E-08	0.0E+00	2.4E-07	2.4E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	1.2E-06	1.2E-06	7.8E-07	3.5E-06	4.3E-06
Ethylbenzene	100-41-4	0.0E+00	3.5E-07	5.9E-07	0.0E+00	1.4E-06	2.4E-06
Formaldehyde	50-00-0	0.0E+00	7.6E-06	6.0E-06	6.5E-06	1.3E-05	3.4E-05
Tetrachloroethylene	127-18-4	0.0E+00	9.8E-08	2.3E-07	0.0E+00	6.6E-07	8.6E-07

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field
CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures.

Table F-397. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Referees 50<70 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	2.6E-08	4.6E-08	0.0E+00	1.0E-07	1.8E-07
Benz[a]anthracene	56-55-3	0.0E+00	2.0E-11	1.2E-10	0.0E+00	0.0E+00	6.9E-10
Benzo[a]pyrene	50-32-8	0.0E+00	4.6E-09	4.9E-09	2.4E-09	1.3E-08	1.9E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	9.8E-11	4.1E-10	0.0E+00	5.2E-10	1.7E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	1.2E-10	3.7E-10	0.0E+00	9.3E-10	1.4E-09
Chrysene	218-01-9	0.0E+00	8.2E-10	1.3E-09	0.0E+00	3.1E-09	5.4E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	2.9E-10	3.6E-10	1.1E-10	8.6E-10	1.7E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	5.7E-10	1.2E-09	0.0E+00	2.9E-09	5.0E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	4.3E-10	1.4E-09	0.0E+00	5.0E-09	5.0E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	6.4E-08	1.8E-07	0.0E+00	5.0E-07	7.5E-07
Naphthalene	91-20-3	0.0E+00	1.1E-07	2.3E-07	0.0E+00	4.7E-07	1.1E-06
Styrene	100-42-5	0.0E+00	2.4E-07	5.0E-07	0.0E+00	8.1E-07	2.7E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	1.0E-05	7.7E-06	7.1E-06	2.0E-05	3.9E-05
Benzene	71-43-2	3.7E-07	2.5E-06	2.0E-06	1.8E-06	6.2E-06	1.0E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	7.9E-08	1.7E-07	0.0E+00	4.9E-07	4.9E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	2.4E-06	2.5E-06	1.6E-06	7.0E-06	8.7E-06
Ethylbenzene	100-41-4	0.0E+00	7.1E-07	1.2E-06	0.0E+00	2.9E-06	4.7E-06
Formaldehyde	50-00-0	0.0E+00	1.5E-05	1.2E-05	1.3E-05	2.6E-05	6.7E-05
Tetrachloroethylene	127-18-4	0.0E+00	2.0E-07	4.6E-07	0.0E+00	1.3E-06	1.7E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration ($C_{air-field}$) of a chemical detected in air off an individual field

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-398. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Spectators Thrid Trimester Fetus**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	1.3E-09	2.4E-09	0.0E+00	5.3E-09	9.2E-09
Benz[a]anthracene	56-55-3	0.0E+00	1.0E-12	6.0E-12	0.0E+00	0.0E+00	3.5E-11
Benzo[a]pyrene	50-32-8	0.0E+00	2.3E-10	2.5E-10	1.2E-10	6.4E-10	9.9E-10
Benzo[b]fluoranthene	205-99-2	0.0E+00	5.0E-12	2.1E-11	0.0E+00	2.6E-11	8.8E-11
Benzo[k]fluoranthene	207-08-9	0.0E+00	6.3E-12	1.9E-11	0.0E+00	4.7E-11	7.3E-11
Chrysene	218-01-9	0.0E+00	4.2E-11	6.6E-11	0.0E+00	1.6E-10	2.8E-10
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	1.5E-11	1.8E-11	5.7E-12	4.4E-11	8.7E-11
Dibenz[a,h]anthracene	53-70-3	0.0E+00	2.9E-11	6.0E-11	0.0E+00	1.5E-10	2.5E-10



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.2E-11	7.3E-11	0.0E+00	2.6E-10	2.6E-10
Methyl Isobutyl Ketone	108-10-1	0.0E+00	3.3E-09	9.0E-09	0.0E+00	2.6E-08	3.8E-08
Naphthalene	91-20-3	0.0E+00	5.6E-09	1.2E-08	0.0E+00	2.4E-08	5.5E-08
Styrene	100-42-5	0.0E+00	1.2E-08	2.6E-08	0.0E+00	4.2E-08	1.4E-07
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	5.1E-07	3.9E-07	3.6E-07	1.0E-06	2.0E-06
Benzene	71-43-2	1.9E-08	1.3E-07	1.0E-07	9.0E-08	3.2E-07	5.2E-07
Benzene, 1,4-dichloro	106-46-7	0.0E+00	4.0E-09	8.9E-09	0.0E+00	2.5E-08	2.5E-08
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	1.2E-07	1.3E-07	8.0E-08	3.6E-07	4.5E-07
Ethylbenzene	100-41-4	0.0E+00	3.6E-08	6.1E-08	0.0E+00	1.5E-07	2.4E-07
Formaldehyde	50-00-0	0.0E+00	7.8E-07	6.1E-07	6.7E-07	1.3E-06	3.4E-06
Tetrachloroethylene	127-18-4	0.0E+00	1.0E-08	2.4E-08	0.0E+00	6.7E-08	8.8E-08

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-399. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Spectators 0<2 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	8.5E-08	1.5E-07	0.0E+00	3.4E-07	5.9E-07
Benz[a]anthracene	56-55-3	0.0E+00	6.5E-11	3.8E-10	0.0E+00	0.0E+00	2.3E-09
Benzo[a]pyrene	50-32-8	0.0E+00	1.5E-08	1.6E-08	7.9E-09	4.1E-08	6.4E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	3.2E-10	1.3E-09	0.0E+00	1.7E-09	5.7E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	4.0E-10	1.2E-09	0.0E+00	3.1E-09	4.7E-09
Chrysene	218-01-9	0.0E+00	2.7E-09	4.3E-09	0.0E+00	1.0E-08	1.8E-08
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	9.5E-10	1.2E-09	3.7E-10	2.8E-09	5.6E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.9E-09	3.8E-09	0.0E+00	9.7E-09	1.6E-08
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.4E-09	4.7E-09	0.0E+00	1.7E-08	1.7E-08
Methyl Isobutyl Ketone	108-10-1	0.0E+00	2.1E-07	5.8E-07	0.0E+00	1.6E-06	2.5E-06
Naphthalene	91-20-3	0.0E+00	3.6E-07	7.5E-07	0.0E+00	1.6E-06	3.5E-06
Styrene	100-42-5	0.0E+00	8.0E-07	1.6E-06	0.0E+00	2.7E-06	8.9E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	3.3E-05	2.5E-05	2.3E-05	6.5E-05	1.3E-04
Benzene	71-43-2	1.2E-06	8.2E-06	6.6E-06	5.8E-06	2.0E-05	3.3E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	2.6E-07	5.7E-07	0.0E+00	1.6E-06	1.6E-06



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	7.8E-06	8.1E-06	5.2E-06	2.3E-05	2.9E-05
Ethylbenzene	100-41-4	0.0E+00	2.3E-06	3.9E-06	0.0E+00	9.5E-06	1.6E-05
Formaldehyde	50-00-0	0.0E+00	5.0E-05	3.9E-05	4.3E-05	8.6E-05	2.2E-04
Tetrachloroethylene	127-18-4	0.0E+00	6.5E-07	1.5E-06	0.0E+00	4.3E-06	5.7E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-400. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Spectators 2<6 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	2.7E-08	4.8E-08	0.0E+00	1.1E-07	1.9E-07
Benz[a]anthracene	56-55-3	0.0E+00	2.0E-11	1.2E-10	0.0E+00	0.0E+00	7.2E-10
Benzo[a]pyrene	50-32-8	0.0E+00	4.7E-09	5.1E-09	2.5E-09	1.3E-08	2.0E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	1.0E-10	4.2E-10	0.0E+00	5.4E-10	1.8E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	1.3E-10	3.8E-10	0.0E+00	9.6E-10	1.5E-09
Chrysene	218-01-9	0.0E+00	8.5E-10	1.3E-09	0.0E+00	3.2E-09	5.6E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	3.0E-10	3.7E-10	1.2E-10	9.0E-10	1.8E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	5.9E-10	1.2E-09	0.0E+00	3.1E-09	5.2E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	4.5E-10	1.5E-09	0.0E+00	5.2E-09	5.2E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	6.7E-08	1.8E-07	0.0E+00	5.2E-07	7.8E-07
Naphthalene	91-20-3	0.0E+00	1.1E-07	2.4E-07	0.0E+00	4.9E-07	1.1E-06
Styrene	100-42-5	0.0E+00	2.5E-07	5.2E-07	0.0E+00	8.5E-07	2.8E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	1.0E-05	8.0E-06	7.4E-06	2.1E-05	4.1E-05
Benzene	71-43-2	3.9E-07	2.6E-06	2.1E-06	1.8E-06	6.5E-06	1.0E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	8.2E-08	1.8E-07	0.0E+00	5.1E-07	5.1E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	2.5E-06	2.5E-06	1.6E-06	7.3E-06	9.0E-06
Ethylbenzene	100-41-4	0.0E+00	7.3E-07	1.2E-06	0.0E+00	3.0E-06	4.9E-06
Formaldehyde	50-00-0	0.0E+00	1.6E-05	1.2E-05	1.4E-05	2.7E-05	7.0E-05
Tetrachloroethylene	127-18-4	0.0E+00	2.0E-07	4.8E-07	0.0E+00	1.4E-06	1.8E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number



Values are rounded to two significant figures.

Table F-401. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Spectators 6<11 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	3.2E-08	5.7E-08	0.0E+00	1.3E-07	2.2E-07
Benz[a]anthracene	56-55-3	0.0E+00	2.4E-11	1.4E-10	0.0E+00	0.0E+00	8.5E-10
Benzo[a]pyrene	50-32-8	0.0E+00	5.6E-09	6.1E-09	3.0E-09	1.6E-08	2.4E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	1.2E-10	5.0E-10	0.0E+00	6.4E-10	2.1E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	1.5E-10	4.5E-10	0.0E+00	1.1E-09	1.8E-09
Chrysene	218-01-9	0.0E+00	1.0E-09	1.6E-09	0.0E+00	3.8E-09	6.7E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	3.6E-10	4.5E-10	1.4E-10	1.1E-09	2.1E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	7.1E-10	1.4E-09	0.0E+00	3.6E-09	6.1E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	5.3E-10	1.8E-09	0.0E+00	6.2E-09	6.2E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	8.0E-08	2.2E-07	0.0E+00	6.2E-07	9.3E-07
Naphthalene	91-20-3	0.0E+00	1.4E-07	2.8E-07	0.0E+00	5.8E-07	1.3E-06
Styrene	100-42-5	0.0E+00	3.0E-07	6.2E-07	0.0E+00	1.0E-06	3.3E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	1.2E-05	9.5E-06	8.8E-06	2.5E-05	4.9E-05
Benzene	71-43-2	4.6E-07	3.1E-06	2.5E-06	2.2E-06	7.7E-06	1.3E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	9.8E-08	2.1E-07	0.0E+00	6.0E-07	6.0E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	2.9E-06	3.0E-06	1.9E-06	8.7E-06	1.1E-05
Ethylbenzene	100-41-4	0.0E+00	8.7E-07	1.5E-06	0.0E+00	3.6E-06	5.9E-06
Formaldehyde	50-00-0	0.0E+00	1.9E-05	1.5E-05	1.6E-05	3.2E-05	8.3E-05
Tetrachloroethylene	127-18-4	0.0E+00	2.4E-07	5.8E-07	0.0E+00	1.6E-06	2.1E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field

^b On-Field ADD_{inh} was calculated from the individual field concentrations (C_{air-field}) of a chemical detected in air on the fields.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-402. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Spectators 11<16 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	1.4E-08	2.5E-08	0.0E+00	5.5E-08	9.6E-08



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benz[a]anthracene	56-55-3	0.0E+00	1.1E-11	6.2E-11	0.0E+00	0.0E+00	3.7E-10
Benzo[a]pyrene	50-32-8	0.0E+00	2.4E-09	2.6E-09	1.3E-09	6.7E-09	1.0E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	5.3E-11	2.2E-10	0.0E+00	2.8E-10	9.2E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	6.5E-11	2.0E-10	0.0E+00	5.0E-10	7.6E-10
Chrysene	218-01-9	0.0E+00	4.4E-10	6.9E-10	0.0E+00	1.7E-09	2.9E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	1.6E-10	1.9E-10	6.0E-11	4.6E-10	9.1E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	3.1E-10	6.2E-10	0.0E+00	1.6E-09	2.7E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.3E-10	7.6E-10	0.0E+00	2.7E-09	2.7E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	3.4E-08	9.4E-08	0.0E+00	2.7E-07	4.0E-07
Naphthalene	91-20-3	0.0E+00	5.8E-08	1.2E-07	0.0E+00	2.5E-07	5.7E-07
Styrene	100-42-5	0.0E+00	1.3E-07	2.7E-07	0.0E+00	4.4E-07	1.4E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	5.4E-06	4.1E-06	3.8E-06	1.1E-05	2.1E-05
Benzene	71-43-2	2.0E-07	1.3E-06	1.1E-06	9.4E-07	3.3E-06	5.4E-06
Benzene, 1,4-dichloro	106-46-7	0.0E+00	4.2E-08	9.3E-08	0.0E+00	2.6E-07	2.6E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	1.3E-06	1.3E-06	8.4E-07	3.8E-06	4.7E-06
Ethylbenzene	100-41-4	0.0E+00	3.8E-07	6.4E-07	0.0E+00	1.5E-06	2.5E-06
Formaldehyde	50-00-0	0.0E+00	8.2E-06	6.4E-06	7.0E-06	1.4E-05	3.6E-05
Tetrachloroethylene	127-18-4	0.0E+00	1.1E-07	2.5E-07	0.0E+00	7.0E-07	9.2E-07

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-403. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Spectators 16<30 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	8.5E-09	1.5E-08	0.0E+00	3.4E-08	5.9E-08
Benz[a]anthracene	56-55-3	0.0E+00	6.5E-12	3.8E-11	0.0E+00	0.0E+00	2.3E-10
Benzo[a]pyrene	50-32-8	0.0E+00	1.5E-09	1.6E-09	7.9E-10	4.1E-09	6.4E-09
Benzo[b]fluoranthene	205-99-2	0.0E+00	3.2E-11	1.3E-10	0.0E+00	1.7E-10	5.7E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	4.0E-11	1.2E-10	0.0E+00	3.1E-10	4.7E-10
Chrysene	218-01-9	0.0E+00	2.7E-10	4.3E-10	0.0E+00	1.0E-09	1.8E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	9.5E-11	1.2E-10	3.7E-11	2.8E-10	5.6E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.9E-10	3.8E-10	0.0E+00	9.7E-10	1.6E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.4E-10	4.7E-10	0.0E+00	1.7E-09	1.7E-09



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Methyl Isobutyl Ketone	108-10-1	0.0E+00	2.1E-08	5.8E-08	0.0E+00	1.6E-07	2.5E-07
Naphthalene	91-20-3	0.0E+00	3.6E-08	7.5E-08	0.0E+00	1.6E-07	3.5E-07
Styrene	100-42-5	0.0E+00	8.0E-08	1.6E-07	0.0E+00	2.7E-07	8.9E-07
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	3.3E-06	2.5E-06	2.3E-06	6.5E-06	1.3E-05
Benzene	71-43-2	1.2E-07	8.2E-07	6.6E-07	5.8E-07	2.0E-06	3.3E-06
Benzene, 1,4-dichloro	106-46-7	0.0E+00	2.6E-08	5.7E-08	0.0E+00	1.6E-07	1.6E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	7.8E-07	8.1E-07	5.2E-07	2.3E-06	2.9E-06
Ethylbenzene	100-41-4	0.0E+00	2.3E-07	3.9E-07	0.0E+00	9.5E-07	1.6E-06
Formaldehyde	50-00-0	0.0E+00	5.0E-06	3.9E-06	4.3E-06	8.6E-06	2.2E-05
Tetrachloroethylene	127-18-4	0.0E+00	6.5E-08	1.5E-07	0.0E+00	4.3E-07	5.7E-07

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration ($C_{air-field}$) of a chemical detected in air off an individual field.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-404. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Spectators 30<40 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	5.3E-09	9.5E-09	0.0E+00	2.1E-08	3.7E-08
Benz[a]anthracene	56-55-3	0.0E+00	4.0E-12	2.4E-11	0.0E+00	0.0E+00	1.4E-10
Benzo[a]pyrene	50-32-8	0.0E+00	9.4E-10	1.0E-09	4.9E-10	2.6E-09	4.0E-09
Benzo[b]fluoranthene	205-99-2	0.0E+00	2.0E-11	8.3E-11	0.0E+00	1.1E-10	3.5E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	2.5E-11	7.5E-11	0.0E+00	1.9E-10	2.9E-10
Chrysene	218-01-9	0.0E+00	1.7E-10	2.7E-10	0.0E+00	6.4E-10	1.1E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	6.0E-11	7.4E-11	2.3E-11	1.8E-10	3.5E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.2E-10	2.4E-10	0.0E+00	6.0E-10	1.0E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	8.8E-11	2.9E-10	0.0E+00	1.0E-09	1.0E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	1.3E-08	3.6E-08	0.0E+00	1.0E-07	1.5E-07
Naphthalene	91-20-3	0.0E+00	2.2E-08	4.7E-08	0.0E+00	9.7E-08	2.2E-07
Styrene	100-42-5	0.0E+00	5.0E-08	1.0E-07	0.0E+00	1.7E-07	5.6E-07
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	2.1E-06	1.6E-06	1.5E-06	4.1E-06	8.1E-06
Benzene	71-43-2	7.6E-08	5.1E-07	4.1E-07	3.6E-07	1.3E-06	2.1E-06
Benzene, 1,4-dichloro	106-46-7	0.0E+00	1.6E-08	3.6E-08	0.0E+00	1.0E-07	1.0E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	4.9E-07	5.0E-07	3.2E-07	1.4E-06	1.8E-06



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Ethylbenzene	100-41-4	0.0E+00	1.5E-07	2.4E-07	0.0E+00	5.9E-07	9.8E-07
Formaldehyde	50-00-0	0.0E+00	3.1E-06	2.5E-06	2.7E-06	5.4E-06	1.4E-05
Tetrachloroethylene	127-18-4	0.0E+00	4.0E-08	9.6E-08	0.0E+00	2.7E-07	3.5E-07

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration ($C_{air-field}$) of a chemical detected in air off an individual field.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-405. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Spectators 40<50 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	5.4E-09	9.6E-09	0.0E+00	2.2E-08	3.7E-08
Benz[a]anthracene	56-55-3	0.0E+00	4.1E-12	2.4E-11	0.0E+00	0.0E+00	1.4E-10
Benzo[a]pyrene	50-32-8	0.0E+00	9.5E-10	1.0E-09	5.0E-10	2.6E-09	4.0E-09
Benzo[b]fluoranthene	205-99-2	0.0E+00	2.0E-11	8.4E-11	0.0E+00	1.1E-10	3.6E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	2.5E-11	7.6E-11	0.0E+00	1.9E-10	3.0E-10
Chrysene	218-01-9	0.0E+00	1.7E-10	2.7E-10	0.0E+00	6.4E-10	1.1E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	6.0E-11	7.5E-11	2.3E-11	1.8E-10	3.5E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.2E-10	2.4E-10	0.0E+00	6.1E-10	1.0E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	8.9E-11	3.0E-10	0.0E+00	1.0E-09	1.0E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	1.3E-08	3.6E-08	0.0E+00	1.0E-07	1.6E-07
Naphthalene	91-20-3	0.0E+00	2.3E-08	4.7E-08	0.0E+00	9.8E-08	2.2E-07
Styrene	100-42-5	0.0E+00	5.0E-08	1.0E-07	0.0E+00	1.7E-07	5.6E-07
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	2.1E-06	1.6E-06	1.5E-06	4.1E-06	8.2E-06
Benzene	71-43-2	7.7E-08	5.2E-07	4.2E-07	3.6E-07	1.3E-06	2.1E-06
Benzene, 1,4-dichloro	106-46-7	0.0E+00	1.6E-08	3.6E-08	0.0E+00	1.0E-07	1.0E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	4.9E-07	5.1E-07	3.3E-07	1.5E-06	1.8E-06
Ethylbenzene	100-41-4	0.0E+00	1.5E-07	2.5E-07	0.0E+00	6.0E-07	9.9E-07
Formaldehyde	50-00-0	0.0E+00	3.2E-06	2.5E-06	2.7E-06	5.4E-06	1.4E-05
Tetrachloroethylene	127-18-4	0.0E+00	4.1E-08	9.7E-08	0.0E+00	2.7E-07	3.6E-07

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration ($C_{air-field}$) of a chemical detected in air off an individual field.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-406. Field-Specific Inhalation **On-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Spectators 50<70 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	1.1E-08	1.9E-08	0.0E+00	4.3E-08	7.5E-08
Benz[a]anthracene	56-55-3	0.0E+00	8.2E-12	4.8E-11	0.0E+00	0.0E+00	2.9E-10
Benzo[a]pyrene	50-32-8	0.0E+00	1.9E-09	2.1E-09	1.0E-09	5.2E-09	8.1E-09
Benzo[b]fluoranthene	205-99-2	0.0E+00	4.1E-11	1.7E-10	0.0E+00	2.2E-10	7.2E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	5.1E-11	1.5E-10	0.0E+00	3.9E-10	5.9E-10
Chrysene	218-01-9	0.0E+00	3.4E-10	5.4E-10	0.0E+00	1.3E-09	2.3E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	1.2E-10	1.5E-10	4.6E-11	3.6E-10	7.1E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	2.4E-10	4.8E-10	0.0E+00	1.2E-09	2.1E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.8E-10	5.9E-10	0.0E+00	2.1E-09	2.1E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	2.7E-08	7.3E-08	0.0E+00	2.1E-07	3.1E-07
Naphthalene	91-20-3	0.0E+00	4.6E-08	9.5E-08	0.0E+00	2.0E-07	4.4E-07
Styrene	100-42-5	0.0E+00	1.0E-07	2.1E-07	0.0E+00	3.4E-07	1.1E-06
Non-Field-Related Chemicals							
Acetaldehyde	75-07-0	0.0E+00	4.2E-06	3.2E-06	3.0E-06	8.2E-06	1.6E-05
Benzene	71-43-2	1.5E-07	1.0E-06	8.4E-07	7.3E-07	2.6E-06	4.2E-06
Benzene, 1,4-dichloro	106-46-7	0.0E+00	3.3E-08	7.2E-08	0.0E+00	2.0E-07	2.0E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	9.8E-07	1.0E-06	6.5E-07	2.9E-06	3.6E-06
Ethylbenzene	100-41-4	0.0E+00	2.9E-07	5.0E-07	0.0E+00	1.2E-06	2.0E-06
Formaldehyde	50-00-0	0.0E+00	6.4E-06	5.0E-06	5.5E-06	1.1E-05	2.8E-05
Tetrachloroethylene	127-18-4	0.0E+00	8.2E-08	1.9E-07	0.0E+00	5.5E-07	7.2E-07

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-407. Field-Specific Inhalation **Off-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Spectators Third Trimester Fetus**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	1.4E-09	2.9E-09	0.0E+00	5.3E-09	1.3E-08
Benzo[a]pyrene	50-32-8	0.0E+00	2.8E-10	4.5E-10	1.2E-10	9.5E-10	2.1E-09
Benzo[b]fluoranthene	205-99-2	0.0E+00	7.6E-12	2.5E-11	0.0E+00	8.8E-11	8.8E-11
Benzo[k]fluoranthene	207-08-9	0.0E+00	6.3E-12	2.1E-11	0.0E+00	7.3E-11	7.3E-11



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Chrysene	218-01-9	0.0E+00	2.1E-11	4.6E-11	0.0E+00	8.9E-11	2.0E-10
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	1.3E-11	1.7E-11	5.7E-12	4.7E-11	5.5E-11
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.8E-11	4.1E-11	0.0E+00	1.2E-10	1.5E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.5E-11	6.0E-11	0.0E+00	7.7E-11	2.6E-10
Methyl Isobutyl Ketone	108-10-1	0.0E+00	7.3E-10	4.3E-09	0.0E+00	0.0E+00	2.6E-08
Naphthalene	91-20-3	0.0E+00	6.0E-09	1.2E-08	0.0E+00	2.7E-08	5.4E-08
Styrene	100-42-5	0.0E+00	1.2E-08	2.7E-08	0.0E+00	4.7E-08	1.4E-07
Non-Field-Related Chemicals							
Benzene	71-43-2	0.0E+00	1.3E-07	1.1E-07	8.1E-08	3.2E-07	5.7E-07
Benzene, 1,4-dichloro	106-46-7	0.0E+00	3.6E-09	8.4E-09	0.0E+00	2.5E-08	2.5E-08
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	1.2E-07	1.3E-07	4.4E-08	4.0E-07	4.2E-07
Ethylbenzene	100-41-4	0.0E+00	3.7E-08	6.5E-08	0.0E+00	1.6E-07	2.7E-07
Tetrachloroethylene	127-18-4	0.0E+00	1.0E-08	2.4E-08	0.0E+00	6.7E-08	8.8E-08

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-408. Field-Specific Inhalation **Off-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Spectators 0<2 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	9.1E-08	1.9E-07	0.0E+00	3.4E-07	8.1E-07
Benzo[a]pyrene	50-32-8	0.0E+00	1.8E-08	2.9E-08	7.6E-09	6.1E-08	1.3E-07
Benzo[b]fluoranthene	205-99-2	0.0E+00	4.9E-10	1.6E-09	0.0E+00	5.7E-09	5.7E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	4.0E-10	1.3E-09	0.0E+00	4.7E-09	4.7E-09
Chrysene	218-01-9	0.0E+00	1.3E-09	2.9E-09	0.0E+00	5.7E-09	1.3E-08
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	8.1E-10	1.1E-09	3.7E-10	3.0E-09	3.5E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.1E-09	2.6E-09	0.0E+00	7.5E-09	9.9E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	9.4E-10	3.9E-09	0.0E+00	5.0E-09	1.7E-08
Methyl Isobutyl Ketone	108-10-1	0.0E+00	4.7E-08	2.8E-07	0.0E+00	0.0E+00	1.6E-06
Naphthalene	91-20-3	0.0E+00	3.8E-07	7.8E-07	0.0E+00	1.8E-06	3.5E-06
Styrene	100-42-5	0.0E+00	7.9E-07	1.7E-06	0.0E+00	3.0E-06	9.1E-06
Non-Field-Related Chemicals							
Benzene	71-43-2	0.0E+00	8.4E-06	7.2E-06	5.2E-06	2.0E-05	3.7E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	2.3E-07	5.4E-07	0.0E+00	1.6E-06	1.6E-06
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	7.6E-06	8.6E-06	2.8E-06	2.6E-05	2.7E-05



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Ethylbenzene	100-41-4	0.0E+00	2.4E-06	4.2E-06	0.0E+00	1.1E-05	1.8E-05
Tetrachloroethylene	127-18-4	0.0E+00	6.5E-07	1.5E-06	0.0E+00	4.3E-06	5.7E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration ($C_{air-field}$) of a chemical detected in air off an individual field

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-409. Field-Specific Inhalation Off-Field Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender Spectators 2<6 Years

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	2.9E-08	5.9E-08	0.0E+00	1.1E-07	2.5E-07
Benzo[a]pyrene	50-32-8	0.0E+00	5.6E-09	9.1E-09	2.4E-09	1.9E-08	4.2E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	1.5E-10	5.1E-10	0.0E+00	1.8E-09	1.8E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	1.3E-10	4.2E-10	0.0E+00	1.5E-09	1.5E-09
Chrysene	218-01-9	0.0E+00	4.2E-10	9.2E-10	0.0E+00	1.8E-09	4.0E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	2.6E-10	3.4E-10	1.2E-10	9.5E-10	1.1E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	3.6E-10	8.4E-10	0.0E+00	2.4E-09	3.1E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	3.0E-10	1.2E-09	0.0E+00	1.6E-09	5.2E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	1.5E-08	8.8E-08	0.0E+00	0.0E+00	5.2E-07
Naphthalene	91-20-3	0.0E+00	1.2E-07	2.5E-07	0.0E+00	5.5E-07	1.1E-06
Styrene	100-42-5	0.0E+00	2.5E-07	5.4E-07	0.0E+00	9.5E-07	2.9E-06
Non-Field-Related Chemicals							
Benzene	71-43-2	0.0E+00	2.6E-06	2.3E-06	1.6E-06	6.5E-06	1.2E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	7.2E-08	1.7E-07	0.0E+00	5.1E-07	5.1E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	2.4E-06	2.7E-06	9.0E-07	8.2E-06	8.5E-06
Ethylbenzene	100-41-4	0.0E+00	7.5E-07	1.3E-06	0.0E+00	3.3E-06	5.6E-06
Tetrachloroethylene	127-18-4	0.0E+00	2.0E-07	4.8E-07	0.0E+00	1.4E-06	1.8E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration ($C_{air-field}$) of a chemical detected in air off an individual field

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-410. Field-Specific Inhalation **Off-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Spectators 6<11 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	3.4E-08	7.0E-08	0.0E+00	1.3E-07	3.0E-07
Benzo[a]pyrene	50-32-8	0.0E+00	6.7E-09	1.1E-08	2.9E-09	2.3E-08	5.0E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	1.8E-10	6.1E-10	0.0E+00	2.1E-09	2.1E-09
Benzo[k]fluoranthene	207-08-9	0.0E+00	1.5E-10	5.0E-10	0.0E+00	1.8E-09	1.8E-09
Chrysene	218-01-9	0.0E+00	5.0E-10	1.1E-09	0.0E+00	2.2E-09	4.7E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	3.1E-10	4.1E-10	1.4E-10	1.1E-09	1.3E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	4.3E-10	1.0E-09	0.0E+00	2.8E-09	3.7E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	3.5E-10	1.5E-09	0.0E+00	1.9E-09	6.2E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	1.8E-08	1.0E-07	0.0E+00	0.0E+00	6.2E-07
Naphthalene	91-20-3	0.0E+00	1.4E-07	2.9E-07	0.0E+00	6.6E-07	1.3E-06
Styrene	100-42-5	0.0E+00	3.0E-07	6.4E-07	0.0E+00	1.1E-06	3.4E-06
Non-Field-Related Chemicals							
Benzene	71-43-2	0.0E+00	3.1E-06	2.7E-06	2.0E-06	7.7E-06	1.4E-05
Benzene, 1,4-dichloro	106-46-7	0.0E+00	8.6E-08	2.0E-07	0.0E+00	6.0E-07	6.0E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	2.9E-06	3.2E-06	1.1E-06	9.7E-06	1.0E-05
Ethylbenzene	100-41-4	0.0E+00	9.0E-07	1.6E-06	0.0E+00	4.0E-06	6.6E-06
Tetrachloroethylene	127-18-4	0.0E+00	2.4E-07	5.7E-07	0.0E+00	1.6E-06	2.1E-06

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-411. Field-Specific Inhalation **Off-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Spectators 11<16 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	1.5E-08	3.0E-08	0.0E+00	5.5E-08	1.3E-07
Benzo[a]pyrene	50-32-8	0.0E+00	2.9E-09	4.7E-09	1.2E-09	9.9E-09	2.2E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	7.9E-11	2.6E-10	0.0E+00	9.2E-10	9.2E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	6.5E-11	2.2E-10	0.0E+00	7.6E-10	7.6E-10
Chrysene	218-01-9	0.0E+00	2.2E-10	4.8E-10	0.0E+00	9.3E-10	2.0E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	1.3E-10	1.8E-10	6.0E-11	4.9E-10	5.7E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.9E-10	4.3E-10	0.0E+00	1.2E-09	1.6E-09



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.5E-10	6.3E-10	0.0E+00	8.0E-10	2.7E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	7.6E-09	4.5E-08	0.0E+00	0.0E+00	2.7E-07
Naphthalene	91-20-3	0.0E+00	6.2E-08	1.3E-07	0.0E+00	2.9E-07	5.6E-07
Styrene	100-42-5	0.0E+00	1.3E-07	2.8E-07	0.0E+00	4.9E-07	1.5E-06
Non-Field-Related Chemicals							
Benzene	71-43-2	0.0E+00	1.4E-06	1.2E-06	8.4E-07	3.3E-06	6.0E-06
Benzene, 1,4-dichloro	106-46-7	0.0E+00	3.7E-08	8.8E-08	0.0E+00	2.6E-07	2.6E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	1.2E-06	1.4E-06	4.6E-07	4.2E-06	4.4E-06
Ethylbenzene	100-41-4	0.0E+00	3.9E-07	6.8E-07	0.0E+00	1.7E-06	2.9E-06
Tetrachloroethylene	127-18-4	0.0E+00	1.1E-07	2.5E-07	0.0E+00	7.0E-07	9.2E-07

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration ($C_{air-field}$) of a chemical detected in air off an individual field.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-412. Field-Specific Inhalation **Off-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Spectators 16<30 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	9.1E-09	1.9E-08	0.0E+00	3.4E-08	8.1E-08
Benzo[a]pyrene	50-32-8	0.0E+00	1.8E-09	2.9E-09	7.6E-10	6.1E-09	1.3E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	4.9E-11	1.6E-10	0.0E+00	5.7E-10	5.7E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	4.0E-11	1.3E-10	0.0E+00	4.7E-10	4.7E-10
Chrysene	218-01-9	0.0E+00	1.3E-10	2.9E-10	0.0E+00	5.7E-10	1.3E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	8.1E-11	1.1E-10	3.7E-11	3.0E-10	3.5E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.1E-10	2.6E-10	0.0E+00	7.5E-10	9.9E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	9.4E-11	3.9E-10	0.0E+00	5.0E-10	1.7E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	4.7E-09	2.8E-08	0.0E+00	0.0E+00	1.6E-07
Naphthalene	91-20-3	0.0E+00	3.8E-08	7.8E-08	0.0E+00	1.8E-07	3.5E-07
Styrene	100-42-5	0.0E+00	7.9E-08	1.7E-07	0.0E+00	3.0E-07	9.1E-07
Non-Field-Related Chemicals							
Benzene	71-43-2	0.0E+00	8.4E-07	7.2E-07	5.2E-07	2.0E-06	3.7E-06
Benzene, 1,4-dichloro	106-46-7	0.0E+00	2.3E-08	5.4E-08	0.0E+00	1.6E-07	1.6E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	7.6E-07	8.6E-07	2.8E-07	2.6E-06	2.7E-06
Ethylbenzene	100-41-4	0.0E+00	2.4E-07	4.2E-07	0.0E+00	1.1E-06	1.8E-06
Tetrachloroethylene	127-18-4	0.0E+00	6.5E-08	1.5E-07	0.0E+00	4.3E-07	5.7E-07

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific



average concentration ($C_{\text{air-field}}$) of a chemical detected in air off an individual field.
CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures.

Table F-413. Field-Specific Inhalation **Off-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment ($LADD_{\text{inh}}$, mg per kg BW per day) of Chemicals—Combined Gender **Spectators 30<40 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	5.7E-09	1.2E-08	0.0E+00	2.1E-08	5.0E-08
Benzo[a]pyrene	50-32-8	0.0E+00	1.1E-09	1.8E-09	4.7E-10	3.8E-09	8.4E-09
Benzo[b]fluoranthene	205-99-2	0.0E+00	3.0E-11	1.0E-10	0.0E+00	3.5E-10	3.5E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	2.5E-11	8.3E-11	0.0E+00	2.9E-10	2.9E-10
Chrysene	218-01-9	0.0E+00	8.4E-11	1.8E-10	0.0E+00	3.6E-10	7.9E-10
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	5.1E-11	6.8E-11	2.3E-11	1.9E-10	2.2E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	7.2E-11	1.7E-10	0.0E+00	4.7E-10	6.2E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	5.9E-11	2.4E-10	0.0E+00	3.1E-10	1.0E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	2.9E-09	1.7E-08	0.0E+00	0.0E+00	1.0E-07
Naphthalene	91-20-3	0.0E+00	2.4E-08	4.9E-08	0.0E+00	1.1E-07	2.2E-07
Styrene	100-42-5	0.0E+00	4.9E-08	1.1E-07	0.0E+00	1.9E-07	5.7E-07
Non-Field-Related Chemicals							
Benzene	71-43-2	0.0E+00	5.2E-07	4.5E-07	3.2E-07	1.3E-06	2.3E-06
Benzene, 1,4-dichloro	106-46-7	0.0E+00	1.4E-08	3.4E-08	0.0E+00	1.0E-07	1.0E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	4.7E-07	5.3E-07	1.8E-07	1.6E-06	1.7E-06
Ethylbenzene	100-41-4	0.0E+00	1.5E-07	2.6E-07	0.0E+00	6.6E-07	1.1E-06
Tetrachloroethylene	127-18-4	0.0E+00	4.0E-08	9.4E-08	0.0E+00	2.7E-07	3.5E-07

^a 35 field-specific $LADD_{\text{inh-field}}$ are included in the table. $LADD_{\text{inh-field}}$ was calculated from field-specific average concentration ($C_{\text{air-field}}$) of a chemical detected in air off an individual field.
CASRN: Chemical Abstracts Service Registry Number
Values are rounded to two significant figures.

Table F-414. Field-Specific Inhalation **Off-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment ($LADD_{\text{inh}}$, mg per kg BW per day) of Chemicals—Combined Gender **Spectators 40<50 Years**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	5.8E-09	1.2E-08	0.0E+00	2.2E-08	5.1E-08
Benzo[a]pyrene	50-32-8	0.0E+00	1.1E-09	1.8E-09	4.8E-10	3.9E-09	8.5E-09
Benzo[b]fluoranthene	205-99-2	0.0E+00	3.1E-11	1.0E-10	0.0E+00	3.6E-10	3.6E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	2.5E-11	8.4E-11	0.0E+00	3.0E-10	3.0E-10



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Chrysene	218-01-9	0.0E+00	8.5E-11	1.8E-10	0.0E+00	3.6E-10	7.9E-10
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	5.1E-11	6.9E-11	2.3E-11	1.9E-10	2.2E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	7.2E-11	1.7E-10	0.0E+00	4.7E-10	6.2E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	5.9E-11	2.5E-10	0.0E+00	3.1E-10	1.0E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	3.0E-09	1.8E-08	0.0E+00	0.0E+00	1.0E-07
Naphthalene	91-20-3	0.0E+00	2.4E-08	4.9E-08	0.0E+00	1.1E-07	2.2E-07
Styrene	100-42-5	0.0E+00	5.0E-08	1.1E-07	0.0E+00	1.9E-07	5.7E-07
Non-Field-Related Chemicals							
Benzene	71-43-2	0.0E+00	5.3E-07	4.6E-07	3.3E-07	1.3E-06	2.3E-06
Benzene, 1,4-dichloro	106-46-7	0.0E+00	1.4E-08	3.4E-08	0.0E+00	1.0E-07	1.0E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	4.8E-07	5.4E-07	1.8E-07	1.6E-06	1.7E-06
Ethylbenzene	100-41-4	0.0E+00	1.5E-07	2.6E-07	0.0E+00	6.7E-07	1.1E-06
Tetrachloroethylene	127-18-4	0.0E+00	4.1E-08	9.5E-08	0.0E+00	2.7E-07	3.6E-07

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration (C_{air-field}) of a chemical detected in air off an individual field.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-415. Field-Specific Inhalation **Off-Field** Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{inh}, mg per kg BW per day) of Chemicals—Combined Gender **Spectators 50<70 Year**

Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Field-Related Chemicals							
Aniline	62-53-3	0.0E+00	1.2E-08	2.4E-08	0.0E+00	4.3E-08	1.0E-07
Benzo[a]pyrene	50-32-8	0.0E+00	2.3E-09	3.6E-09	9.6E-10	7.7E-09	1.7E-08
Benzo[b]fluoranthene	205-99-2	0.0E+00	6.1E-11	2.0E-10	0.0E+00	7.2E-10	7.2E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	5.1E-11	1.7E-10	0.0E+00	5.9E-10	5.9E-10
Chrysene	218-01-9	0.0E+00	1.7E-10	3.7E-10	0.0E+00	7.2E-10	1.6E-09
Cyclopenta[cd]pyrene	27208-37-3	0.0E+00	1.0E-10	1.4E-10	4.6E-11	3.8E-10	4.5E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.5E-10	3.4E-10	0.0E+00	9.5E-10	1.3E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.2E-10	4.9E-10	0.0E+00	6.3E-10	2.1E-09
Methyl Isobutyl Ketone	108-10-1	0.0E+00	6.0E-09	3.5E-08	0.0E+00	0.0E+00	2.1E-07
Naphthalene	91-20-3	0.0E+00	4.9E-08	9.9E-08	0.0E+00	2.2E-07	4.4E-07
Styrene	100-42-5	0.0E+00	1.0E-07	2.2E-07	0.0E+00	3.8E-07	1.1E-06
Non-Field-Related Chemicals							
Benzene	71-43-2	0.0E+00	1.1E-06	9.2E-07	6.6E-07	2.6E-06	4.7E-06
Benzene, 1,4-dichloro	106-46-7	0.0E+00	2.9E-08	6.8E-08	0.0E+00	2.0E-07	2.0E-07
Benzene, 1-chloro-4-(trifluoromethyl)-	98-56-6	0.0E+00	9.6E-07	1.1E-06	3.6E-07	3.3E-06	3.4E-06



Chemical	CASRN	LADD _{inh-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Ethylbenzene	100-41-4	0.0E+00	3.0E-07	5.3E-07	0.0E+00	1.3E-06	2.2E-06
Tetrachloroethylene	127-18-4	0.0E+00	8.2E-08	1.9E-07	0.0E+00	5.5E-07	7.2E-07

^a 35 field-specific LADD_{inh-field} are included in the table. LADD_{inh-field} was calculated from field-specific average concentration ($C_{air-field}$) of a chemical detected in air off an individual field.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

F.6.2. Dermal Lifetime Average Daily Dose (LADD_{der}) for Lifetime Cancer Risk Assessment of Carcinogens

Table F-416. Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der}, mg per kg BW per day) of **Field-Related Carcinogens^b**—
Combined Gender **Athletes**

Chemical	CASRN ^a	LADD _{der}						
		2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Aniline	62-53-3	3.2E-10	3.2E-10	3.4E-10	4.0E-10	2.0E-10	1.7E-10	3.4E-10
Benz[a]anthracene	56-55-3	4.4E-11	4.3E-11	4.6E-11	5.4E-11	2.7E-11	2.3E-11	4.7E-11
Benzo[a]pyrene	50-32-8	6.0E-11	6.0E-11	6.3E-11	7.4E-11	3.7E-11	3.2E-11	6.4E-11
Benzo[b]fluoranthene	205-99-2	4.5E-11	4.5E-11	4.7E-11	5.6E-11	2.8E-11	2.4E-11	4.8E-11
Benzo[k]fluoranthene	207-08-9	8.5E-11	8.4E-11	8.9E-11	1.0E-10	5.3E-11	4.5E-11	9.1E-11
1,3-Benzothiazole-2-thiol	149-30-4	3.7E-11	3.7E-11	3.9E-11	4.6E-11	2.3E-11	2.0E-11	4.0E-11
Chrysene	218-01-9	3.9E-10	3.8E-10	4.1E-10	4.8E-10	2.4E-10	2.1E-10	4.1E-10
Cyclopenta[cd]pyrene	27208-37-3	6.9E-11	6.9E-11	7.2E-11	8.5E-11	4.3E-11	3.7E-11	7.4E-11
Dibenz[a,h]anthracene	53-70-3	2.2E-11	2.2E-11	2.3E-11	2.7E-11	1.4E-11	1.2E-11	2.4E-11
Indeno[1,2,3-cd]pyrene	193-39-5	4.5E-11	4.4E-11	4.6E-11	5.5E-11	2.8E-11	2.4E-11	4.7E-11
Naphthalene	91-20-3	2.9E-12	2.8E-12	3.0E-12	3.5E-12	1.8E-12	1.5E-12	3.1E-12

^a LADD_{der} was calculated from the mean of 35 individual field average concentrations ($C_{der-crumb\ rubber}$) of a chemical detected in artificial sweat extracts of crumb rubber samples and is the sum across all age groups in a receptor category.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.



Table F-417. Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der}, mg per kg BW per day) of **Field-Related Carcinogens^b**—
Combined Gender **Coaches**

Chemical	CASRN ^a	LADD _{der}			
		16<30 years	30<40 years	40<50 years	50<70 years
Aniline	62-53-3	2.0E-10	1.3E-10	1.3E-10	2.7E-10
Benz[a]anthracene	56-55-3	2.7E-11	1.8E-11	1.8E-11	3.6E-11
Benzo[a]pyrene	50-32-8	3.7E-11	2.4E-11	2.5E-11	5.0E-11
Benzo[b]fluoranthene	205-99-2	2.8E-11	1.8E-11	1.9E-11	3.7E-11
Benzo[k]fluoranthene	207-08-9	5.2E-11	3.5E-11	3.5E-11	7.0E-11
1,3-Benzothiazole-2-thiol	149-30-4	2.3E-11	1.5E-11	1.5E-11	3.1E-11
Chrysene	218-01-9	2.4E-10	1.6E-10	1.6E-10	3.2E-10
Cyclopenta[cd]pyrene	27208-37-3	4.2E-11	2.8E-11	2.9E-11	5.7E-11
Dibenz[a,h]anthracene	53-70-3	1.3E-11	9.0E-12	9.1E-12	1.8E-11
Indeno[1,2,3-cd]pyrene	193-39-5	2.7E-11	1.8E-11	1.8E-11	3.7E-11
Naphthalene	91-20-3	1.7E-12	1.2E-12	1.2E-12	2.4E-12

^a LADD_{der} was calculated from the mean of 35 individual field average concentrations (C_{der-crumb rubber}) of a chemical detected in artificial sweat extracts of crumb rubber samples and is the sum across all age groups in a receptor category.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.
Values are rounded to two significant figures.

Table F-418. Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der}, mg per kg BW per day) of **Field-Related Carcinogens^b**—
Combined Gender **Referees**

Chemical	CASRN ^a	LADD _{der}			
		16<30 years	30<40 years	40<50 years	50<70 years
Aniline	62-53-3	5.3E-11	7.9E-11	5.3E-11	1.1E-10
Benz[a]anthracene	56-55-3	7.2E-12	1.1E-11	7.3E-12	1.5E-11
Benzo[a]pyrene	50-32-8	9.8E-12	1.5E-11	9.9E-12	2.0E-11
Benzo[b]fluoranthene	205-99-2	7.4E-12	1.1E-11	7.5E-12	1.5E-11
Benzo[k]fluoranthene	207-08-9	1.4E-11	2.1E-11	1.4E-11	2.8E-11
1,3-Benzothiazole-2-thiol	149-30-4	6.1E-12	9.1E-12	6.2E-12	1.2E-11
Chrysene	218-01-9	6.3E-11	9.5E-11	6.4E-11	1.3E-10
Cyclopenta[cd]pyrene	27208-37-3	1.1E-11	1.7E-11	1.1E-11	2.3E-11
Dibenz[a,h]anthracene	53-70-3	3.6E-12	5.4E-12	3.7E-12	7.3E-12
Indeno[1,2,3-cd]pyrene	193-39-5	7.3E-12	1.1E-11	7.4E-12	1.5E-11
Naphthalene	91-20-3	4.7E-13	7.0E-13	4.7E-13	9.5E-13



^a LADD_{der} was calculated from the mean of 35 individual field average concentrations (C_{der-crumb rubber}) of a chemical detected in artificial sweat extracts of crumb rubber samples and is the sum across all age groups in a receptor category.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis. Values are rounded to two significant figures.

Table F-419. Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der}, mg per kg BW per day) of **Field-Related Carcinogens^b**—**Combined Gender Spectators**

Chemical	CASRN ^a	LADD _{der}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Aniline	62-53-3	2.0E-11	8.5E-10	3.8E-10	4.3E-10	3.6E-10	1.5E-10	9.2E-11	9.3E-11	1.8E-10
Benz[a]anthracene	56-55-3	2.7E-12	1.2E-10	5.1E-11	5.8E-11	5.0E-11	2.0E-11	1.2E-11	1.3E-11	2.4E-11
Benzo[a]pyrene	50-32-8	3.7E-12	1.6E-10	7.0E-11	7.9E-11	6.8E-11	2.8E-11	1.7E-11	1.7E-11	3.3E-11
Benzo[b]fluoranthene	205-99-2	2.8E-12	1.2E-10	5.3E-11	6.0E-11	5.1E-11	2.1E-11	1.3E-11	1.3E-11	2.5E-11
Benzo[k]fluoranthene	207-08-9	5.2E-12	2.3E-10	9.9E-11	1.1E-10	9.6E-11	3.9E-11	2.4E-11	2.5E-11	4.6E-11
1,3-Benzothiazole-2-thiol	149-30-4	2.3E-12	9.8E-11	4.3E-11	4.9E-11	4.2E-11	1.7E-11	1.1E-11	1.1E-11	2.0E-11
Chrysene	218-01-9	2.4E-11	1.0E-09	4.5E-10	5.1E-10	4.4E-10	1.8E-10	1.1E-10	1.1E-10	2.1E-10
Cyclopenta[cd]pyrene	27208-37-3	4.3E-12	1.8E-10	8.1E-11	9.2E-11	7.8E-11	3.2E-11	2.0E-11	2.0E-11	3.8E-11
Dibenz[a,h]anthracene	53-70-3	1.4E-12	5.8E-11	2.6E-11	2.9E-11	2.5E-11	1.0E-11	6.3E-12	6.4E-12	1.2E-11
Indeno[1,2,3-cd]pyrene	193-39-5	2.7E-12	1.2E-10	5.2E-11	5.9E-11	5.0E-11	2.1E-11	1.3E-11	1.3E-11	2.4E-11
Naphthalene	91-20-3	1.8E-13	7.6E-12	3.3E-12	3.8E-12	3.2E-12	1.3E-12	8.1E-13	8.2E-13	1.6E-12

^a LADD_{der} was calculated from the mean of 35 individual field average concentrations (C_{der-crumb rubber}) of a chemical detected in artificial sweat extracts of crumb rubber samples and is the sum across all age groups in a receptor category.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis. Values are rounded to two significant figures.

INDIVIDUAL FIELD ASSESSMENT (Table F-420 to Table F-443)



Table F-420. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Athletes 2<6 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	4.4E-11	1.4E-10	0.0E+00	3.4E-10	6.5E-10
Benz[a]anthracene	56-55-3	0.0E+00	6.0E-11	7.0E-11	3.2E-11	1.7E-10	3.2E-10
Benzo[a]pyrene	50-32-8	0.0E+00	4.5E-11	4.5E-11	3.0E-11	1.2E-10	1.9E-10
Benzo[b]fluoranthene	205-99-2	0.0E+00	8.5E-11	7.5E-11	6.7E-11	2.1E-10	3.3E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	3.7E-11	3.9E-11	2.3E-11	1.2E-10	1.5E-10
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	3.2E-10	1.9E-09	0.0E+00	0.0E+00	1.1E-08
Chrysene	218-01-9	4.2E-11	3.9E-10	2.5E-10	3.9E-10	7.8E-10	8.8E-10
Cyclopenta[cd]pyrene	27208-37-3	7.1E-12	6.9E-11	4.2E-11	6.7E-11	1.3E-10	1.6E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	2.2E-11	4.1E-11	0.0E+00	1.1E-10	1.7E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	4.5E-11	3.7E-11	3.9E-11	1.0E-10	1.6E-10
Naphthalene	91-20-3	0.0E+00	2.9E-12	1.2E-11	0.0E+00	1.4E-11	5.2E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-421. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Athletes 6<11 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	4.3E-11	1.3E-10	0.0E+00	3.3E-10	6.5E-10
Benz[a]anthracene	56-55-3	0.0E+00	6.0E-11	6.9E-11	3.1E-11	1.7E-10	3.2E-10
Benzo[a]pyrene	50-32-8	0.0E+00	4.5E-11	4.4E-11	2.9E-11	1.2E-10	1.8E-10
Benzo[b]fluoranthene	205-99-2	0.0E+00	8.4E-11	7.5E-11	6.7E-11	2.1E-10	3.3E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	3.7E-11	3.9E-11	2.2E-11	1.2E-10	1.5E-10
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	3.2E-10	1.9E-09	0.0E+00	0.0E+00	1.1E-08
Chrysene	218-01-9	4.2E-11	3.8E-10	2.5E-10	3.8E-10	7.7E-10	8.7E-10
Cyclopenta[cd]pyrene	27208-37-3	7.0E-12	6.9E-11	4.1E-11	6.6E-11	1.3E-10	1.6E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	2.2E-11	4.0E-11	0.0E+00	1.1E-10	1.7E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	4.4E-11	3.7E-11	3.9E-11	1.0E-10	1.6E-10
Naphthalene	91-20-3	0.0E+00	2.8E-12	1.2E-11	0.0E+00	1.4E-11	5.1E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-



Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-422. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Athletes 11<16 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	4.6E-11	1.4E-10	0.0E+00	3.5E-10	6.8E-10
Benz[a]anthracene	56-55-3	0.0E+00	6.3E-11	7.3E-11	3.3E-11	1.8E-10	3.3E-10
Benzo[a]pyrene	50-32-8	0.0E+00	4.7E-11	4.7E-11	3.1E-11	1.2E-10	1.9E-10
Benzo[b]fluoranthene	205-99-2	0.0E+00	8.9E-11	7.9E-11	7.0E-11	2.2E-10	3.5E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	3.9E-11	4.1E-11	2.4E-11	1.3E-10	1.6E-10
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	3.4E-10	2.0E-09	0.0E+00	0.0E+00	1.2E-08
Chrysene	218-01-9	4.4E-11	4.1E-10	2.6E-10	4.1E-10	8.2E-10	9.2E-10
Cyclopenta[cd]pyrene	27208-37-3	7.4E-12	7.2E-11	4.3E-11	7.0E-11	1.4E-10	1.7E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	2.3E-11	4.2E-11	0.0E+00	1.1E-10	1.8E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	4.6E-11	3.9E-11	4.1E-11	1.1E-10	1.7E-10
Naphthalene	91-20-3	0.0E+00	3.0E-12	1.2E-11	0.0E+00	1.5E-11	5.4E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-423. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Athletes 16<30 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	5.4E-11	1.7E-10	0.0E+00	4.1E-10	8.0E-10
Benz[a]anthracene	56-55-3	0.0E+00	7.4E-11	8.5E-11	3.9E-11	2.1E-10	3.9E-10
Benzo[a]pyrene	50-32-8	0.0E+00	5.6E-11	5.5E-11	3.7E-11	1.5E-10	2.3E-10
Benzo[b]fluoranthene	205-99-2	0.0E+00	1.0E-10	9.3E-11	8.3E-11	2.6E-10	4.1E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	4.6E-11	4.8E-11	2.8E-11	1.5E-10	1.8E-10
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	4.0E-10	2.3E-09	0.0E+00	0.0E+00	1.4E-08
Chrysene	218-01-9	5.2E-11	4.8E-10	3.1E-10	4.8E-10	9.6E-10	1.1E-09
Cyclopenta[cd]pyrene	27208-37-3	8.7E-12	8.5E-11	5.1E-11	8.2E-11	1.6E-10	2.0E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	2.7E-11	5.0E-11	0.0E+00	1.3E-10	2.1E-10



Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	5.5E-11	4.6E-11	4.8E-11	1.3E-10	2.0E-10
Naphthalene	91-20-3	0.0E+00	3.5E-12	1.5E-11	0.0E+00	1.8E-11	6.4E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-424. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Athletes 30<40 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	2.7E-11	8.5E-11	0.0E+00	2.1E-10	4.1E-10
Benz[a]anthracene	56-55-3	0.0E+00	3.7E-11	4.3E-11	2.0E-11	1.1E-10	2.0E-10
Benzo[a]pyrene	50-32-8	0.0E+00	2.8E-11	2.8E-11	1.9E-11	7.4E-11	1.2E-10
Benzo[b]fluoranthene	205-99-2	0.0E+00	5.3E-11	4.7E-11	4.2E-11	1.3E-10	2.1E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	2.3E-11	2.5E-11	1.4E-11	7.6E-11	9.4E-11
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	2.0E-10	1.2E-09	0.0E+00	0.0E+00	7.0E-09
Chrysene	218-01-9	2.6E-11	2.4E-10	1.6E-10	2.4E-10	4.9E-10	5.5E-10
Cyclopenta[cd]pyrene	27208-37-3	4.4E-12	4.3E-11	2.6E-11	4.2E-11	8.3E-11	9.9E-11
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.4E-11	2.5E-11	0.0E+00	6.7E-11	1.1E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.8E-11	2.3E-11	2.5E-11	6.5E-11	1.0E-10
Naphthalene	91-20-3	0.0E+00	1.8E-12	7.4E-12	0.0E+00	9.0E-12	3.2E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-425. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Athletes 40<50 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	2.3E-11	7.2E-11	0.0E+00	1.8E-10	3.5E-10
Benz[a]anthracene	56-55-3	0.0E+00	3.2E-11	3.7E-11	1.7E-11	8.9E-11	1.7E-10
Benzo[a]pyrene	50-32-8	0.0E+00	2.4E-11	2.4E-11	1.6E-11	6.3E-11	9.8E-11
Benzo[b]fluoranthene	205-99-2	0.0E+00	4.5E-11	4.0E-11	3.6E-11	1.1E-10	1.7E-10



Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[k]fluoranthene	207-08-9	0.0E+00	2.0E-11	2.1E-11	1.2E-11	6.4E-11	8.0E-11
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	1.7E-10	1.0E-09	0.0E+00	0.0E+00	6.0E-09
Chrysene	218-01-9	2.2E-11	2.1E-10	1.3E-10	2.1E-10	4.1E-10	4.7E-10
Cyclopenta[cd]pyrene	27208-37-3	3.7E-12	3.7E-11	2.2E-11	3.5E-11	7.0E-11	8.4E-11
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.2E-11	2.1E-11	0.0E+00	5.7E-11	8.9E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.4E-11	2.0E-11	2.1E-11	5.5E-11	8.6E-11
Naphthalene	91-20-3	0.0E+00	1.5E-12	6.2E-12	0.0E+00	7.6E-12	2.8E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-426. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Athletes 50<70 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	4.7E-11	1.5E-10	0.0E+00	3.6E-10	7.0E-10
Benz[a]anthracene	56-55-3	0.0E+00	6.4E-11	7.4E-11	3.4E-11	1.8E-10	3.4E-10
Benzo[a]pyrene	50-32-8	0.0E+00	4.8E-11	4.8E-11	3.2E-11	1.3E-10	2.0E-10
Benzo[b]fluoranthene	205-99-2	0.0E+00	9.1E-11	8.0E-11	7.2E-11	2.3E-10	3.5E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	4.0E-11	4.2E-11	2.4E-11	1.3E-10	1.6E-10
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	3.4E-10	2.0E-09	0.0E+00	0.0E+00	1.2E-08
Chrysene	218-01-9	4.5E-11	4.1E-10	2.7E-10	4.1E-10	8.3E-10	9.4E-10
Cyclopenta[cd]pyrene	27208-37-3	7.6E-12	7.4E-11	4.4E-11	7.1E-11	1.4E-10	1.7E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	2.4E-11	4.3E-11	0.0E+00	1.2E-10	1.8E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	4.7E-11	4.0E-11	4.2E-11	1.1E-10	1.7E-10
Naphthalene	91-20-3	0.0E+00	3.1E-12	1.3E-11	0.0E+00	1.5E-11	5.5E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-427. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Coaches 16<30 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	2.7E-11	8.3E-11	0.0E+00	2.0E-10	4.0E-10
Benz[a]anthracene	56-55-3	0.0E+00	3.7E-11	4.2E-11	1.9E-11	1.0E-10	1.9E-10
Benzo[a]pyrene	50-32-8	0.0E+00	2.8E-11	2.7E-11	1.8E-11	7.2E-11	1.1E-10
Benzo[b]fluoranthene	205-99-2	0.0E+00	5.2E-11	4.6E-11	4.1E-11	1.3E-10	2.0E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	2.3E-11	2.4E-11	1.4E-11	7.4E-11	9.1E-11
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	2.0E-10	1.2E-09	0.0E+00	0.0E+00	6.9E-09
Chrysene	218-01-9	2.6E-11	2.4E-10	1.5E-10	2.4E-10	4.7E-10	5.4E-10
Cyclopenta[cd]pyrene	27208-37-3	4.3E-12	4.2E-11	2.5E-11	4.1E-11	8.1E-11	9.7E-11
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.3E-11	2.5E-11	0.0E+00	6.6E-11	1.0E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.7E-11	2.3E-11	2.4E-11	6.3E-11	9.9E-11
Naphthalene	91-20-3	0.0E+00	1.7E-12	7.2E-12	0.0E+00	8.8E-12	3.2E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-428. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Coaches 30<40 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	1.8E-11	5.5E-11	0.0E+00	1.4E-10	2.7E-10
Benz[a]anthracene	56-55-3	0.0E+00	2.4E-11	2.8E-11	1.3E-11	6.9E-11	1.3E-10
Benzo[a]pyrene	50-32-8	0.0E+00	1.8E-11	1.8E-11	1.2E-11	4.8E-11	7.5E-11
Benzo[b]fluoranthene	205-99-2	0.0E+00	3.5E-11	3.1E-11	2.7E-11	8.7E-11	1.3E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	1.5E-11	1.6E-11	9.2E-12	4.9E-11	6.1E-11
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	1.3E-10	7.8E-10	0.0E+00	0.0E+00	4.6E-09
Chrysene	218-01-9	1.7E-11	1.6E-10	1.0E-10	1.6E-10	3.2E-10	3.6E-10
Cyclopenta[cd]pyrene	27208-37-3	2.9E-12	2.8E-11	1.7E-11	2.7E-11	5.4E-11	6.5E-11
Dibenz[a,h]anthracene	53-70-3	0.0E+00	9.0E-12	1.6E-11	0.0E+00	4.4E-11	6.9E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.8E-11	1.5E-11	1.6E-11	4.2E-11	6.6E-11
Naphthalene	91-20-3	0.0E+00	1.2E-12	4.8E-12	0.0E+00	5.9E-12	2.1E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-



Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-429. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Coaches 40<50 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	1.8E-11	5.6E-11	0.0E+00	1.4E-10	2.7E-10
Benz[a]anthracene	56-55-3	0.0E+00	2.5E-11	2.9E-11	1.3E-11	6.9E-11	1.3E-10
Benzo[a]pyrene	50-32-8	0.0E+00	1.9E-11	1.8E-11	1.2E-11	4.9E-11	7.6E-11
Benzo[b]fluoranthene	205-99-2	0.0E+00	3.5E-11	3.1E-11	2.8E-11	8.8E-11	1.4E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	1.5E-11	1.6E-11	9.3E-12	5.0E-11	6.2E-11
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	1.3E-10	7.8E-10	0.0E+00	0.0E+00	4.6E-09
Chrysene	218-01-9	1.7E-11	1.6E-10	1.0E-10	1.6E-10	3.2E-10	3.6E-10
Cyclopenta[cd]pyrene	27208-37-3	2.9E-12	2.9E-11	1.7E-11	2.8E-11	5.5E-11	6.5E-11
Dibenz[a,h]anthracene	53-70-3	0.0E+00	9.1E-12	1.7E-11	0.0E+00	4.4E-11	6.9E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.8E-11	1.5E-11	1.6E-11	4.3E-11	6.7E-11
Naphthalene	91-20-3	0.0E+00	1.2E-12	4.8E-12	0.0E+00	5.9E-12	2.1E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-430. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Coaches 50<70 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	3.6E-11	1.1E-10	0.0E+00	2.8E-10	5.4E-10
Benz[a]anthracene	56-55-3	0.0E+00	5.0E-11	5.7E-11	2.6E-11	1.4E-10	2.6E-10
Benzo[a]pyrene	50-32-8	0.0E+00	3.7E-11	3.7E-11	2.4E-11	9.7E-11	1.5E-10
Benzo[b]fluoranthene	205-99-2	0.0E+00	7.0E-11	6.2E-11	5.5E-11	1.8E-10	2.7E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	3.1E-11	3.2E-11	1.9E-11	1.0E-10	1.2E-10
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	2.7E-10	1.6E-09	0.0E+00	0.0E+00	9.3E-09
Chrysene	218-01-9	3.5E-11	3.2E-10	2.1E-10	3.2E-10	6.4E-10	7.3E-10
Cyclopenta[cd]pyrene	27208-37-3	5.8E-12	5.7E-11	3.4E-11	5.5E-11	1.1E-10	1.3E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.8E-11	3.3E-11	0.0E+00	8.9E-11	1.4E-10



Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	3.7E-11	3.1E-11	3.2E-11	8.5E-11	1.3E-10
Naphthalene	91-20-3	0.0E+00	2.4E-12	9.7E-12	0.0E+00	1.2E-11	4.3E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-431. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Referees 16<30 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	1.1E-11	3.3E-11	0.0E+00	8.2E-11	1.6E-10
Benz[a]anthracene	56-55-3	0.0E+00	1.5E-11	1.7E-11	7.8E-12	4.1E-11	7.8E-11
Benzo[a]pyrene	50-32-8	0.0E+00	1.1E-11	1.1E-11	7.3E-12	2.9E-11	4.5E-11
Benzo[b]fluoranthene	205-99-2	0.0E+00	2.1E-11	1.8E-11	1.6E-11	5.2E-11	8.1E-11
Benzo[k]fluoranthene	207-08-9	0.0E+00	9.1E-12	9.6E-12	5.5E-12	3.0E-11	3.7E-11
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	7.9E-11	4.7E-10	0.0E+00	0.0E+00	2.8E-09
Chrysene	218-01-9	1.0E-11	9.5E-11	6.1E-11	9.5E-11	1.9E-10	2.2E-10
Cyclopenta[cd]pyrene	27208-37-3	1.7E-12	1.7E-11	1.0E-11	1.6E-11	3.2E-11	3.9E-11
Dibenz[a,h]anthracene	53-70-3	0.0E+00	5.4E-12	9.9E-12	0.0E+00	2.6E-11	4.1E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.1E-11	9.1E-12	9.6E-12	2.5E-11	4.0E-11
Naphthalene	91-20-3	0.0E+00	7.0E-13	2.9E-12	0.0E+00	3.5E-12	1.3E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-432. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Referees 30<40 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	7.2E-12	2.2E-11	0.0E+00	5.5E-11	1.1E-10
Benz[a]anthracene	56-55-3	0.0E+00	9.8E-12	1.1E-11	5.2E-12	2.8E-11	5.2E-11
Benzo[a]pyrene	50-32-8	0.0E+00	7.4E-12	7.3E-12	4.9E-12	1.9E-11	3.0E-11
Benzo[b]fluoranthene	205-99-2	0.0E+00	1.4E-11	1.2E-11	1.1E-11	3.5E-11	5.4E-11



Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[k]fluoranthene	207-08-9	0.0E+00	6.1E-12	6.4E-12	3.7E-12	2.0E-11	2.5E-11
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	5.3E-11	3.1E-10	0.0E+00	0.0E+00	1.8E-09
Chrysene	218-01-9	6.9E-12	6.3E-11	4.1E-11	6.3E-11	1.3E-10	1.4E-10
Cyclopenta[cd]pyrene	27208-37-3	1.2E-12	1.1E-11	6.8E-12	1.1E-11	2.2E-11	2.6E-11
Dibenz[a,h]anthracene	53-70-3	0.0E+00	3.6E-12	6.6E-12	0.0E+00	1.8E-11	2.8E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	7.3E-12	6.1E-12	6.4E-12	1.7E-11	2.7E-11
Naphthalene	91-20-3	0.0E+00	4.7E-13	1.9E-12	0.0E+00	2.4E-12	8.5E-12

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-433. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Referees 40<50 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	7.3E-12	2.3E-11	0.0E+00	5.5E-11	1.1E-10
Benz[a]anthracene	56-55-3	0.0E+00	9.9E-12	1.1E-11	5.3E-12	2.8E-11	5.3E-11
Benzo[a]pyrene	50-32-8	0.0E+00	7.5E-12	7.4E-12	4.9E-12	2.0E-11	3.1E-11
Benzo[b]fluoranthene	205-99-2	0.0E+00	1.4E-11	1.2E-11	1.1E-11	3.5E-11	5.5E-11
Benzo[k]fluoranthene	207-08-9	0.0E+00	6.2E-12	6.5E-12	3.7E-12	2.0E-11	2.5E-11
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	5.3E-11	3.2E-10	0.0E+00	0.0E+00	1.9E-09
Chrysene	218-01-9	7.0E-12	6.4E-11	4.1E-11	6.4E-11	1.3E-10	1.5E-10
Cyclopenta[cd]pyrene	27208-37-3	1.2E-12	1.1E-11	6.9E-12	1.1E-11	2.2E-11	2.6E-11
Dibenz[a,h]anthracene	53-70-3	0.0E+00	3.7E-12	6.7E-12	0.0E+00	1.8E-11	2.8E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	7.4E-12	6.1E-12	6.5E-12	1.7E-11	2.7E-11
Naphthalene	91-20-3	0.0E+00	4.7E-13	2.0E-12	0.0E+00	2.4E-12	8.6E-12

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-434. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Referees 50<70 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	1.5E-11	4.5E-11	0.0E+00	1.1E-10	2.2E-10
Benz[a]anthracene	56-55-3	0.0E+00	2.0E-11	2.3E-11	1.1E-11	5.6E-11	1.1E-10
Benzo[a]pyrene	50-32-8	0.0E+00	1.5E-11	1.5E-11	9.9E-12	3.9E-11	6.1E-11
Benzo[b]fluoranthene	205-99-2	0.0E+00	2.8E-11	2.5E-11	2.2E-11	7.1E-11	1.1E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	1.2E-11	1.3E-11	7.5E-12	4.0E-11	5.0E-11
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	1.1E-10	6.3E-10	0.0E+00	0.0E+00	3.7E-09
Chrysene	218-01-9	1.4E-11	1.3E-10	8.3E-11	1.3E-10	2.6E-10	2.9E-10
Cyclopenta[cd]pyrene	27208-37-3	2.3E-12	2.3E-11	1.4E-11	2.2E-11	4.4E-11	5.3E-11
Dibenz[a,h]anthracene	53-70-3	0.0E+00	7.3E-12	1.3E-11	0.0E+00	3.6E-11	5.6E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.5E-11	1.2E-11	1.3E-11	3.4E-11	5.4E-11
Naphthalene	91-20-3	0.0E+00	9.5E-13	3.9E-12	0.0E+00	4.8E-12	1.7E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-435. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Spectators Third Trimester Fetus**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	2.7E-12	8.4E-12	0.0E+00	2.1E-11	4.0E-11
Benz[a]anthracene	56-55-3	0.0E+00	3.7E-12	4.3E-12	2.0E-12	1.0E-11	2.0E-11
Benzo[a]pyrene	50-32-8	0.0E+00	2.8E-12	2.7E-12	1.8E-12	7.3E-12	1.1E-11
Benzo[b]fluoranthene	205-99-2	0.0E+00	5.2E-12	4.6E-12	4.1E-12	1.3E-11	2.0E-11
Benzo[k]fluoranthene	207-08-9	0.0E+00	2.3E-12	2.4E-12	1.4E-12	7.5E-12	9.3E-12
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	2.0E-11	1.2E-10	0.0E+00	0.0E+00	6.9E-10
Chrysene	218-01-9	2.6E-12	2.4E-11	1.5E-11	2.4E-11	4.8E-11	5.4E-11
Cyclopenta[cd]pyrene	27208-37-3	4.4E-13	4.3E-12	2.6E-12	4.1E-12	8.2E-12	9.8E-12
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.4E-12	2.5E-12	0.0E+00	6.6E-12	1.0E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.7E-12	2.3E-12	2.4E-12	6.4E-12	1.0E-11
Naphthalene	91-20-3	0.0E+00	1.8E-13	7.3E-13	0.0E+00	8.9E-13	3.2E-12

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-



Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-436. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 0<2 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	1.2E-10	3.6E-10	0.0E+00	8.9E-10	1.7E-09
Benz[a]anthracene	56-55-3	0.0E+00	1.6E-10	1.8E-10	8.4E-11	4.5E-10	8.5E-10
Benzo[a]pyrene	50-32-8	0.0E+00	1.2E-10	1.2E-10	7.8E-11	3.1E-10	4.9E-10
Benzo[b]fluoranthene	205-99-2	0.0E+00	2.3E-10	2.0E-10	1.8E-10	5.6E-10	8.7E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	9.8E-11	1.0E-10	6.0E-11	3.2E-10	4.0E-10
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	8.5E-10	5.0E-09	0.0E+00	0.0E+00	3.0E-08
Chrysene	218-01-9	1.1E-10	1.0E-09	6.6E-10	1.0E-09	2.1E-09	2.3E-09
Cyclopenta[cd]pyrene	27208-37-3	1.9E-11	1.8E-10	1.1E-10	1.8E-10	3.5E-10	4.2E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	5.8E-11	1.1E-10	0.0E+00	2.9E-10	4.5E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.2E-10	9.8E-11	1.0E-10	2.7E-10	4.3E-10
Naphthalene	91-20-3	0.0E+00	7.6E-12	3.1E-11	0.0E+00	3.8E-11	1.4E-10

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-437. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 2<6 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	5.1E-11	1.6E-10	0.0E+00	3.9E-10	7.6E-10
Benz[a]anthracene	56-55-3	0.0E+00	7.0E-11	8.1E-11	3.7E-11	2.0E-10	3.7E-10
Benzo[a]pyrene	50-32-8	0.0E+00	5.3E-11	5.2E-11	3.5E-11	1.4E-10	2.2E-10
Benzo[b]fluoranthene	205-99-2	0.0E+00	9.9E-11	8.8E-11	7.8E-11	2.5E-10	3.8E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	4.3E-11	4.6E-11	2.6E-11	1.4E-10	1.8E-10
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	3.8E-10	2.2E-09	0.0E+00	0.0E+00	1.3E-08
Chrysene	218-01-9	4.9E-11	4.5E-10	2.9E-10	4.5E-10	9.1E-10	1.0E-09
Cyclopenta[cd]pyrene	27208-37-3	8.2E-12	8.1E-11	4.8E-11	7.8E-11	1.5E-10	1.9E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	2.6E-11	4.7E-11	0.0E+00	1.3E-10	2.0E-10



Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	5.2E-11	4.3E-11	4.6E-11	1.2E-10	1.9E-10
Naphthalene	91-20-3	0.0E+00	3.3E-12	1.4E-11	0.0E+00	1.7E-11	6.1E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-438. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 6<11 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	5.8E-11	1.8E-10	0.0E+00	4.4E-10	8.6E-10
Benz[a]anthracene	56-55-3	0.0E+00	7.9E-11	9.2E-11	4.2E-11	2.2E-10	4.2E-10
Benzo[a]pyrene	50-32-8	0.0E+00	6.0E-11	5.9E-11	3.9E-11	1.6E-10	2.4E-10
Benzo[b]fluoranthene	205-99-2	0.0E+00	1.1E-10	9.9E-11	8.9E-11	2.8E-10	4.4E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	4.9E-11	5.2E-11	3.0E-11	1.6E-10	2.0E-10
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	4.3E-10	2.5E-09	0.0E+00	0.0E+00	1.5E-08
Chrysene	218-01-9	5.6E-11	5.1E-10	3.3E-10	5.1E-10	1.0E-09	1.2E-09
Cyclopenta[cd]pyrene	27208-37-3	9.3E-12	9.2E-11	5.5E-11	8.8E-11	1.8E-10	2.1E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	2.9E-11	5.3E-11	0.0E+00	1.4E-10	2.2E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	5.9E-11	4.9E-11	5.2E-11	1.4E-10	2.1E-10
Naphthalene	91-20-3	0.0E+00	3.8E-12	1.6E-11	0.0E+00	1.9E-11	6.9E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-439. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 11<16 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	5.0E-11	1.5E-10	0.0E+00	3.8E-10	7.4E-10
Benz[a]anthracene	56-55-3	0.0E+00	6.8E-11	7.9E-11	3.6E-11	1.9E-10	3.6E-10
Benzo[a]pyrene	50-32-8	0.0E+00	5.1E-11	5.0E-11	3.4E-11	1.3E-10	2.1E-10
Benzo[b]fluoranthene	205-99-2	0.0E+00	9.6E-11	8.5E-11	7.6E-11	2.4E-10	3.7E-10



Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Benzo[k]fluoranthene	207-08-9	0.0E+00	4.2E-11	4.5E-11	2.6E-11	1.4E-10	1.7E-10
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	3.6E-10	2.2E-09	0.0E+00	0.0E+00	1.3E-08
Chrysene	218-01-9	4.8E-11	4.4E-10	2.8E-10	4.4E-10	8.8E-10	1.0E-09
Cyclopenta[cd]pyrene	27208-37-3	8.0E-12	7.8E-11	4.7E-11	7.6E-11	1.5E-10	1.8E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	2.5E-11	4.6E-11	0.0E+00	1.2E-10	1.9E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	5.0E-11	4.2E-11	4.4E-11	1.2E-10	1.8E-10
Naphthalene	91-20-3	0.0E+00	3.2E-12	1.3E-11	0.0E+00	1.6E-11	5.9E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-440. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 16<30 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	2.0E-11	6.3E-11	0.0E+00	1.6E-10	3.0E-10
Benz[a]anthracene	56-55-3	0.0E+00	2.8E-11	3.2E-11	1.5E-11	7.8E-11	1.5E-10
Benzo[a]pyrene	50-32-8	0.0E+00	2.1E-11	2.1E-11	1.4E-11	5.5E-11	8.5E-11
Benzo[b]fluoranthene	205-99-2	0.0E+00	3.9E-11	3.5E-11	3.1E-11	9.8E-11	1.5E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	1.7E-11	1.8E-11	1.0E-11	5.6E-11	6.9E-11
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	1.5E-10	8.8E-10	0.0E+00	0.0E+00	5.2E-09
Chrysene	218-01-9	2.0E-11	1.8E-10	1.2E-10	1.8E-10	3.6E-10	4.1E-10
Cyclopenta[cd]pyrene	27208-37-3	3.3E-12	3.2E-11	1.9E-11	3.1E-11	6.1E-11	7.3E-11
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.0E-11	1.9E-11	0.0E+00	5.0E-11	7.8E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.1E-11	1.7E-11	1.8E-11	4.8E-11	7.5E-11
Naphthalene	91-20-3	0.0E+00	1.3E-12	5.4E-12	0.0E+00	6.7E-12	2.4E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.



Table F-441. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 30<40 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	1.2E-11	3.9E-11	0.0E+00	9.6E-11	1.9E-10
Benz[a]anthracene	56-55-3	0.0E+00	1.7E-11	2.0E-11	9.0E-12	4.8E-11	9.1E-11
Benzo[a]pyrene	50-32-8	0.0E+00	1.3E-11	1.3E-11	8.5E-12	3.4E-11	5.3E-11
Benzo[b]fluoranthene	205-99-2	0.0E+00	2.4E-11	2.1E-11	1.9E-11	6.1E-11	9.4E-11
Benzo[k]fluoranthene	207-08-9	0.0E+00	1.1E-11	1.1E-11	6.4E-12	3.5E-11	4.3E-11
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	9.2E-11	5.4E-10	0.0E+00	0.0E+00	3.2E-09
Chrysene	218-01-9	1.2E-11	1.1E-10	7.1E-11	1.1E-10	2.2E-10	2.5E-10
Cyclopenta[cd]pyrene	27208-37-3	2.0E-12	2.0E-11	1.2E-11	1.9E-11	3.8E-11	4.5E-11
Dibenz[a,h]anthracene	53-70-3	0.0E+00	6.3E-12	1.2E-11	0.0E+00	3.1E-11	4.8E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.3E-11	1.1E-11	1.1E-11	3.0E-11	4.6E-11
Naphthalene	91-20-3	0.0E+00	8.1E-13	3.4E-12	0.0E+00	4.1E-12	1.5E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-442. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 40<50 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	1.3E-11	3.9E-11	0.0E+00	9.7E-11	1.9E-10
Benz[a]anthracene	56-55-3	0.0E+00	1.7E-11	2.0E-11	9.1E-12	4.9E-11	9.2E-11
Benzo[a]pyrene	50-32-8	0.0E+00	1.3E-11	1.3E-11	8.6E-12	3.4E-11	5.3E-11
Benzo[b]fluoranthene	205-99-2	0.0E+00	2.5E-11	2.2E-11	1.9E-11	6.1E-11	9.5E-11
Benzo[k]fluoranthene	207-08-9	0.0E+00	1.1E-11	1.1E-11	6.5E-12	3.5E-11	4.3E-11
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	9.3E-11	5.5E-10	0.0E+00	0.0E+00	3.3E-09
Chrysene	218-01-9	1.2E-11	1.1E-10	7.2E-11	1.1E-10	2.2E-10	2.5E-10
Cyclopenta[cd]pyrene	27208-37-3	2.0E-12	2.0E-11	1.2E-11	1.9E-11	3.8E-11	4.6E-11
Dibenz[a,h]anthracene	53-70-3	0.0E+00	6.4E-12	1.2E-11	0.0E+00	3.1E-11	4.9E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.3E-11	1.1E-11	1.1E-11	3.0E-11	4.7E-11
Naphthalene	91-20-3	0.0E+00	8.2E-13	3.4E-12	0.0E+00	4.2E-12	1.5E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-



Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

Table F-443. Field-Specific Dermal Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{der-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Spectators 50<70 Years**

Chemical Name	CASRN	LADD _{der-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	2.4E-11	7.4E-11	0.0E+00	1.8E-10	3.5E-10
Benz[a]anthracene	56-55-3	0.0E+00	3.3E-11	3.8E-11	1.7E-11	9.2E-11	1.7E-10
Benzo[a]pyrene	50-32-8	0.0E+00	2.5E-11	2.4E-11	1.6E-11	6.4E-11	1.0E-10
Benzo[b]fluoranthene	205-99-2	0.0E+00	4.6E-11	4.1E-11	3.6E-11	1.2E-10	1.8E-10
Benzo[k]fluoranthene	207-08-9	0.0E+00	2.0E-11	2.1E-11	1.2E-11	6.6E-11	8.2E-11
1,3-Benzothiazole-2-thiol	149-30-4	0.0E+00	1.8E-10	1.0E-09	0.0E+00	0.0E+00	6.1E-09
Chrysene	218-01-9	2.3E-11	2.1E-10	1.4E-10	2.1E-10	4.2E-10	4.8E-10
Cyclopenta[cd]pyrene	27208-37-3	3.8E-12	3.8E-11	2.2E-11	3.6E-11	7.2E-11	8.6E-11
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.2E-11	2.2E-11	0.0E+00	5.9E-11	9.2E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.4E-11	2.0E-11	2.1E-11	5.6E-11	8.8E-11
Naphthalene	91-20-3	0.0E+00	1.6E-12	6.4E-12	0.0E+00	7.8E-12	2.8E-11

^a 35 field-specific LADD_{der-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

CASRN: Chemical Abstracts Service Registry Number

Values are rounded to two significant figures.

F.6.3. Ingestion Lifetime Average Daily Dose (LADD_{ing}) for Lifetime Cancer Risk Assessment of Carcinogens

Table F-444. Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing}, mg per kg BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Athletes**

Chemical	CASRN ^a	LADD _{ing}						
		2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Aniline	62-53-3	1.4E-09	1.2E-09	9.1E-10	9.4E-10	4.5E-10	3.8E-10	7.8E-10
Arsenic		8.2E-09	6.8E-09	5.3E-09	5.5E-09	2.6E-09	2.2E-09	4.5E-09
Benz[a]anthracene	56-55-3	2.3E-09	1.9E-09	1.5E-09	1.6E-09	7.4E-10	6.3E-10	1.3E-09
Benzo[a]pyrene	50-32-8	1.9E-09	1.6E-09	1.2E-09	1.3E-09	6.1E-10	5.2E-10	1.1E-09
Benzo[b]fluoranthene	205-99-2	3.2E-09	2.6E-09	2.1E-09	2.1E-09	1.0E-09	8.6E-10	1.8E-09
Benzo[k]fluoranthene	207-08-9	1.0E-09	8.6E-10	6.7E-10	6.9E-10	3.3E-10	2.8E-10	5.7E-10
Chromium		3.8E-08	3.2E-08	2.4E-08	2.5E-08	1.2E-08	1.0E-08	2.1E-08



Chemical	CASRN ^a	LADD _{ing}						
		2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Chrysene	218-01-9	1.0E-08	8.6E-09	6.7E-09	6.9E-09	3.3E-09	2.8E-09	5.7E-09
Cyclopenta[cd]pyrene	27208-37-3	2.2E-09	1.8E-09	1.4E-09	1.5E-09	7.1E-10	6.0E-10	1.2E-09
Dibenz[a,h]anthracene	53-70-3	1.4E-10	1.1E-10	8.8E-11	9.2E-11	4.4E-11	3.7E-11	7.6E-11
Indeno[1,2,3-cd]pyrene	193-39-5	3.9E-10	3.2E-10	2.5E-10	2.6E-10	1.2E-10	1.0E-10	2.1E-10
Lead		7.9E-07	6.6E-07	5.1E-07	5.3E-07	2.5E-07	2.2E-07	4.4E-07
Naphthalene	91-20-3	4.2E-11	3.5E-11	2.7E-11	2.8E-11	1.3E-11	1.1E-11	2.3E-11

^a LADD_{ing} was calculated from the mean of 35 individual field average concentrations (C_{GI-crumb rubber}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-445. Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing}, mg per kg BW per day) of **Field-Related Carcinogens^b**—**Combined Gender Coaches**

Chemical	CASRN ^a	LADD _{ing}			
		16<30 years	30<40 years	40<50 years	50<70 years
Aniline	62-53-3	2.4E-10	1.6E-10	1.6E-10	3.1E-10
Arsenic		1.4E-09	9.3E-10	9.1E-10	1.8E-09
Benz[a]anthracene	56-55-3	3.9E-10	2.6E-10	2.6E-10	5.2E-10
Benzo[a]pyrene	50-32-8	3.2E-10	2.2E-10	2.1E-10	4.2E-10
Benzo[b]fluoranthene	205-99-2	5.4E-10	3.6E-10	3.5E-10	7.1E-10
Benzo[k]fluoranthene	207-08-9	1.7E-10	1.2E-10	1.1E-10	2.3E-10
Chromium		6.4E-09	4.3E-09	4.2E-09	8.5E-09
Chrysene	218-01-9	1.7E-09	1.2E-09	1.1E-09	2.3E-09
Cyclopenta[cd]pyrene	27208-37-3	3.7E-10	2.5E-10	2.5E-10	4.9E-10
Dibenz[a,h]anthracene	53-70-3	2.3E-11	1.6E-11	1.5E-11	3.1E-11
Indeno[1,2,3-cd]pyrene	193-39-5	6.6E-11	4.4E-11	4.3E-11	8.6E-11
Lead		1.3E-07	9.0E-08	8.8E-08	1.8E-07
Naphthalene	91-20-3	7.1E-12	4.8E-12	4.7E-12	9.3E-12

^a LADD_{ing} was calculated from the mean of 35 individual field average concentrations (C_{GI-crumb rubber}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.



Table F-446. Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing}, mg per kg BW per day) of **Field-Related Carcinogens^b**—
Combined Gender **Referees**

Chemical	CASRN ^a	LADD _{ing}			
		16<30 years	30<40 years	40<50 years	50<70 years
Aniline	62-53-3	1.0E-10	7.0E-11	6.9E-11	1.4E-10
Arsenic		6.1E-10	4.1E-10	4.0E-10	8.0E-10
Benz[a]anthracene	56-55-3	1.7E-10	1.2E-10	1.1E-10	2.3E-10
Benzo[a]pyrene	50-32-8	1.4E-10	9.5E-11	9.3E-11	1.9E-10
Benzo[b]fluoranthene	205-99-2	2.4E-10	1.6E-10	1.6E-10	3.1E-10
Benzo[k]fluoranthene	207-08-9	7.7E-11	5.1E-11	5.0E-11	1.0E-10
Chromium		2.8E-09	1.9E-09	1.9E-09	3.7E-09
Chrysene	218-01-9	7.7E-10	5.1E-10	5.0E-10	1.0E-09
Cyclopenta[cd]pyrene	27208-37-3	1.6E-10	1.1E-10	1.1E-10	2.2E-10
Dibenz[a,h]anthracene	53-70-3	1.0E-11	6.8E-12	6.7E-12	1.3E-11
Indeno[1,2,3-cd]pyrene	193-39-5	2.9E-11	1.9E-11	1.9E-11	3.8E-11
Lead		5.9E-08	4.0E-08	3.9E-08	7.8E-08
Naphthalene	91-20-3	3.1E-12	2.1E-12	2.0E-12	4.1E-12

^a LADD_{ing} was calculated from the mean of 35 individual field average concentrations (C_{GI-crumb rubber}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.
Values are rounded to two significant figures.

Table F-447. Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing}, mg per kg BW per day) of **Field-Related Carcinogens^b**—
Combined Gender **Spectators**

Chemical	CASRN ^a	LADD _{ing}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Aniline	62-53-3	1.5E-11	8.2E-09	2.3E-09	2.0E-09	1.6E-10	8.5E-11	5.7E-11	5.4E-11	1.1E-10
Arsenic		8.6E-11	4.8E-08	1.4E-08	1.2E-08	9.3E-10	4.9E-10	3.3E-10	3.2E-10	6.3E-10
Benz[a]anthracene	56-55-3	2.4E-11	1.3E-08	3.8E-09	3.4E-09	2.6E-10	1.4E-10	9.4E-11	8.9E-11	1.8E-10
Benzo[a]pyrene	50-32-8	2.0E-11	1.1E-08	3.1E-09	2.8E-09	2.2E-10	1.1E-10	7.8E-11	7.3E-11	1.5E-10
Benzo[b]fluoranthene	205-99-2	3.3E-11	1.9E-08	5.3E-09	4.6E-09	3.6E-10	1.9E-10	1.3E-10	1.2E-10	2.5E-10
Benzo[k]fluoranthene	207-08-9	1.1E-11	6.0E-09	1.7E-09	1.5E-09	1.2E-10	6.2E-11	4.2E-11	4.0E-11	8.0E-11
Chromium		4.0E-10	2.2E-07	6.3E-08	5.5E-08	4.3E-09	2.3E-09	1.5E-09	1.5E-09	2.9E-09



Chemical	CASRN ^a	LADD _{ing}								
		Third trimester fetus	0<2 years	2<6 years	6<11 years	11<16 years	16<30 years	30<40 years	40<50 years	50<70 years
Chrysene	218-01-9	1.1E-10	6.0E-08	1.7E-08	1.5E-08	1.2E-09	6.2E-10	4.2E-10	4.0E-10	8.0E-10
Cyclopenta[cd]pyrene	27208-37-3	2.3E-11	1.3E-08	3.7E-09	3.2E-09	2.5E-10	1.3E-10	9.0E-11	8.5E-11	1.7E-10
Dibenz[a,h]anthracene	53-70-3	1.4E-12	8.0E-10	2.3E-10	2.0E-10	1.6E-11	8.3E-12	5.6E-12	5.3E-12	1.1E-11
Indeno[1,2,3-cd]pyrene	193-39-5	4.1E-12	2.3E-09	6.4E-10	5.6E-10	4.4E-11	2.3E-11	1.6E-11	1.5E-11	3.0E-11
Lead		8.3E-09	4.6E-06	1.3E-06	1.2E-06	9.1E-08	4.8E-08	3.2E-08	3.1E-08	6.2E-08
Naphthalene	91-20-3	4.4E-13	2.4E-10	6.9E-11	6.1E-11	4.8E-12	2.5E-12	1.7E-12	1.6E-12	3.2E-12

^a LADD_{ing} was calculated from the mean of 35 individual field average concentrations (C_{GI-crumb rubber}) of a chemical detected in artificial gastrointestinal fluids extracts of crumb rubber samples.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

INDIVIDUAL FIELD ASSESSMENT (Table F-448 to Table F-471)

Table F-448. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Athletes 2<6 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	1.4E-09	3.2E-09	0.0E+00	7.8E-09	1.4E-08
Arsenic		9.6E-10	8.2E-09	5.8E-09	7.4E-09	1.5E-08	3.1E-08
Benz[a]anthracene	56-55-3	1.5E-10	2.3E-09	2.5E-09	1.1E-09	7.2E-09	1.0E-08
Benzo[a]pyrene	50-32-8	3.7E-10	1.9E-09	1.6E-09	1.4E-09	5.2E-09	5.6E-09
Benzo[b]fluoranthene	205-99-2	6.5E-10	3.2E-09	2.4E-09	2.1E-09	7.7E-09	9.3E-09
Benzo[k]fluoranthene	207-08-9	8.7E-11	1.0E-09	9.7E-10	6.5E-10	3.0E-09	3.9E-09
Chromium		7.0E-09	3.8E-08	3.7E-08	2.5E-08	8.3E-08	2.1E-07
Chrysene	218-01-9	2.6E-09	1.0E-08	5.5E-09	1.0E-08	1.8E-08	2.8E-08
Cyclopenta[cd]pyrene	27208-37-3	3.9E-10	2.2E-09	2.0E-09	1.9E-09	5.8E-09	1.0E-08
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.4E-10	3.4E-10	0.0E+00	6.2E-10	1.7E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	3.9E-10	6.7E-10	0.0E+00	1.6E-09	2.6E-09
Lead		1.5E-07	7.9E-07	6.9E-07	6.0E-07	2.0E-06	3.6E-06
Naphthalene	91-20-3	0.0E+00	4.2E-11	2.5E-10	0.0E+00	0.0E+00	1.5E-09

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.



° CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.
Values are rounded to two significant figures.

Table F-449. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Athletes 6<11 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	1.2E-09	2.7E-09	0.0E+00	6.5E-09	1.2E-08
Arsenic		8.0E-10	6.8E-09	4.8E-09	6.1E-09	1.2E-08	2.6E-08
Benz[a]anthracene	56-55-3	1.3E-10	1.9E-09	2.1E-09	9.2E-10	6.0E-09	8.4E-09
Benzo[a]pyrene	50-32-8	3.1E-10	1.6E-09	1.3E-09	1.1E-09	4.4E-09	4.6E-09
Benzo[b]fluoranthene	205-99-2	5.4E-10	2.6E-09	2.0E-09	1.8E-09	6.4E-09	7.8E-09
Benzo[k]fluoranthene	207-08-9	7.2E-11	8.6E-10	8.1E-10	5.4E-10	2.5E-09	3.2E-09
Chromium		5.8E-09	3.2E-08	3.1E-08	2.1E-08	6.9E-08	1.7E-07
Chrysene	218-01-9	2.2E-09	8.6E-09	4.6E-09	8.6E-09	1.5E-08	2.4E-08
Cyclopenta[cd]pyrene	27208-37-3	3.2E-10	1.8E-09	1.7E-09	1.6E-09	4.9E-09	8.7E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.1E-10	2.8E-10	0.0E+00	5.2E-10	1.4E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	3.2E-10	5.6E-10	0.0E+00	1.4E-09	2.2E-09
Lead		1.3E-07	6.6E-07	5.7E-07	5.0E-07	1.7E-06	3.0E-06
Naphthalene	91-20-3	0.0E+00	3.5E-11	2.1E-10	0.0E+00	0.0E+00	1.2E-09

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

° CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.
Values are rounded to two significant figures.

Table F-450. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Athletes 11<16 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	9.1E-10	2.1E-09	0.0E+00	5.1E-09	9.1E-09
Arsenic		6.2E-10	5.3E-09	3.7E-09	4.8E-09	9.5E-09	2.0E-08
Benz[a]anthracene	56-55-3	9.7E-11	1.5E-09	1.6E-09	7.1E-10	4.6E-09	6.5E-09
Benzo[a]pyrene	50-32-8	2.4E-10	1.2E-09	1.0E-09	8.7E-10	3.4E-09	3.6E-09
Benzo[b]fluoranthene	205-99-2	4.2E-10	2.1E-09	1.5E-09	1.4E-09	5.0E-09	6.0E-09
Benzo[k]fluoranthene	207-08-9	5.6E-11	6.7E-10	6.3E-10	4.2E-10	1.9E-09	2.5E-09
Chromium		4.5E-09	2.4E-08	2.4E-08	1.6E-08	5.3E-08	1.4E-07
Chrysene	218-01-9	1.7E-09	6.7E-09	3.5E-09	6.6E-09	1.1E-08	1.8E-08



Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Cyclopenta[cd]pyrene	27208-37-3	2.5E-10	1.4E-09	1.3E-09	1.2E-09	3.8E-09	6.7E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	8.8E-11	2.2E-10	0.0E+00	4.0E-10	1.1E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.5E-10	4.4E-10	0.0E+00	1.1E-09	1.7E-09
Lead		1.0E-07	5.1E-07	4.4E-07	3.9E-07	1.3E-06	2.3E-06
Naphthalene	91-20-3	0.0E+00	2.7E-11	1.6E-10	0.0E+00	0.0E+00	9.5E-10

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-451. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Athletes 16<30 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	9.4E-10	2.2E-09	0.0E+00	5.2E-09	9.5E-09
Arsenic		6.5E-10	5.5E-09	3.9E-09	5.0E-09	9.9E-09	2.1E-08
Benz[a]anthracene	56-55-3	1.0E-10	1.6E-09	1.7E-09	7.4E-10	4.8E-09	6.8E-09
Benzo[a]pyrene	50-32-8	2.5E-10	1.3E-09	1.0E-09	9.1E-10	3.5E-09	3.7E-09
Benzo[b]fluoranthene	205-99-2	4.4E-10	2.1E-09	1.6E-09	1.4E-09	5.2E-09	6.3E-09
Benzo[k]fluoranthene	207-08-9	5.8E-11	6.9E-10	6.5E-10	4.4E-10	2.0E-09	2.6E-09
Chromium		4.7E-09	2.5E-08	2.5E-08	1.7E-08	5.5E-08	1.4E-07
Chrysene	218-01-9	1.7E-09	6.9E-09	3.7E-09	6.9E-09	1.2E-08	1.9E-08
Cyclopenta[cd]pyrene	27208-37-3	2.6E-10	1.5E-09	1.4E-09	1.3E-09	3.9E-09	7.0E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	9.2E-11	2.3E-10	0.0E+00	4.2E-10	1.1E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.6E-10	4.5E-10	0.0E+00	1.1E-09	1.8E-09
Lead		1.0E-07	5.3E-07	4.6E-07	4.0E-07	1.3E-06	2.4E-06
Naphthalene	91-20-3	0.0E+00	2.8E-11	1.7E-10	0.0E+00	0.0E+00	9.8E-10

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.



Table F-452. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Athletes 30<40 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	4.5E-10	1.0E-09	0.0E+00	2.5E-09	4.5E-09
Arsenic		3.1E-10	2.6E-09	1.9E-09	2.4E-09	4.7E-09	1.0E-08
Benz[a]anthracene	56-55-3	4.8E-11	7.4E-10	8.1E-10	3.5E-10	2.3E-09	3.2E-09
Benzo[a]pyrene	50-32-8	1.2E-10	6.1E-10	5.0E-10	4.3E-10	1.7E-09	1.8E-09
Benzo[b]fluoranthene	205-99-2	2.1E-10	1.0E-09	7.5E-10	6.8E-10	2.5E-09	3.0E-09
Benzo[k]fluoranthene	207-08-9	2.8E-11	3.3E-10	3.1E-10	2.1E-10	9.7E-10	1.2E-09
Chromium		2.2E-09	1.2E-08	1.2E-08	8.0E-09	2.6E-08	6.7E-08
Chrysene	218-01-9	8.3E-10	3.3E-09	1.7E-09	3.3E-09	5.6E-09	9.0E-09
Cyclopenta[cd]pyrene	27208-37-3	1.2E-10	7.1E-10	6.6E-10	6.1E-10	1.9E-09	3.3E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	4.4E-11	1.1E-10	0.0E+00	2.0E-10	5.5E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.2E-10	2.2E-10	0.0E+00	5.2E-10	8.5E-10
Lead		4.9E-08	2.5E-07	2.2E-07	1.9E-07	6.4E-07	1.2E-06
Naphthalene	91-20-3	0.0E+00	1.3E-11	7.9E-11	0.0E+00	0.0E+00	4.7E-10

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-453. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Athletes 40<50 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	3.8E-10	8.7E-10	0.0E+00	2.1E-09	3.8E-09
Arsenic		2.6E-10	2.2E-09	1.6E-09	2.0E-09	4.0E-09	8.5E-09
Benz[a]anthracene	56-55-3	4.1E-11	6.3E-10	6.9E-10	3.0E-10	1.9E-09	2.7E-09
Benzo[a]pyrene	50-32-8	1.0E-10	5.2E-10	4.2E-10	3.7E-10	1.4E-09	1.5E-09
Benzo[b]fluoranthene	205-99-2	1.8E-10	8.6E-10	6.4E-10	5.8E-10	2.1E-09	2.5E-09
Benzo[k]fluoranthene	207-08-9	2.3E-11	2.8E-10	2.6E-10	1.8E-10	8.2E-10	1.0E-09
Chromium		1.9E-09	1.0E-08	1.0E-08	6.8E-09	2.2E-08	5.7E-08
Chrysene	218-01-9	7.0E-10	2.8E-09	1.5E-09	2.8E-09	4.8E-09	7.7E-09
Cyclopenta[cd]pyrene	27208-37-3	1.1E-10	6.0E-10	5.6E-10	5.2E-10	1.6E-09	2.8E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	3.7E-11	9.3E-11	0.0E+00	1.7E-10	4.6E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.0E-10	1.8E-10	0.0E+00	4.4E-10	7.2E-10



Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Lead		4.2E-08	2.2E-07	1.9E-07	1.6E-07	5.4E-07	9.8E-07
Naphthalene	91-20-3	0.0E+00	1.1E-11	6.7E-11	0.0E+00	0.0E+00	4.0E-10

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-454. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Athletes 50<70 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	7.8E-10	1.8E-09	0.0E+00	4.3E-09	7.8E-09
Arsenic		5.3E-10	4.5E-09	3.2E-09	4.1E-09	8.2E-09	1.7E-08
Benz[a]anthracene	56-55-3	8.3E-11	1.3E-09	1.4E-09	6.1E-10	4.0E-09	5.6E-09
Benzo[a]pyrene	50-32-8	2.1E-10	1.1E-09	8.6E-10	7.5E-10	2.9E-09	3.1E-09
Benzo[b]fluoranthene	205-99-2	3.6E-10	1.8E-09	1.3E-09	1.2E-09	4.3E-09	5.2E-09
Benzo[k]fluoranthene	207-08-9	4.8E-11	5.7E-10	5.4E-10	3.6E-10	1.7E-09	2.1E-09
Chromium		3.9E-09	2.1E-08	2.1E-08	1.4E-08	4.6E-08	1.2E-07
Chrysene	218-01-9	1.4E-09	5.7E-09	3.0E-09	5.7E-09	9.8E-09	1.6E-08
Cyclopenta[cd]pyrene	27208-37-3	2.2E-10	1.2E-09	1.1E-09	1.1E-09	3.2E-09	5.8E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	7.6E-11	1.9E-10	0.0E+00	3.4E-10	9.4E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.1E-10	3.7E-10	0.0E+00	9.1E-10	1.5E-09
Lead		8.5E-08	4.4E-07	3.8E-07	3.3E-07	1.1E-06	2.0E-06
Naphthalene	91-20-3	0.0E+00	2.3E-11	1.4E-10	0.0E+00	0.0E+00	8.1E-10

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-455. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Coaches 16<30 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	2.4E-10	5.4E-10	0.0E+00	1.3E-09	2.4E-09



Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		1.6E-10	1.4E-09	9.8E-10	1.3E-09	2.5E-09	5.3E-09
Benz[a]anthracene	56-55-3	2.6E-11	3.9E-10	4.3E-10	1.9E-10	1.2E-09	1.7E-09
Benzo[a]pyrene	50-32-8	6.3E-11	3.2E-10	2.6E-10	2.3E-10	8.9E-10	9.5E-10
Benzo[b]fluoranthene	205-99-2	1.1E-10	5.4E-10	4.0E-10	3.6E-10	1.3E-09	1.6E-09
Benzo[k]fluoranthene	207-08-9	1.5E-11	1.7E-10	1.6E-10	1.1E-10	5.1E-10	6.6E-10
Chromium		1.2E-09	6.4E-09	6.3E-09	4.2E-09	1.4E-08	3.6E-08
Chrysene	218-01-9	4.4E-10	1.7E-09	9.3E-10	1.7E-09	3.0E-09	4.8E-09
Cyclopenta[cd]pyrene	27208-37-3	6.6E-11	3.7E-10	3.5E-10	3.2E-10	9.9E-10	1.8E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	2.3E-11	5.8E-11	0.0E+00	1.0E-10	2.9E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	6.6E-11	1.1E-10	0.0E+00	2.8E-10	4.5E-10
Lead		2.6E-08	1.3E-07	1.2E-07	1.0E-07	3.4E-07	6.1E-07
Naphthalene	91-20-3	0.0E+00	7.1E-12	4.2E-11	0.0E+00	0.0E+00	2.5E-10

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-456. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Coaches 30<40 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	1.6E-10	3.7E-10	0.0E+00	8.9E-10	1.6E-09
Arsenic		1.1E-10	9.3E-10	6.6E-10	8.4E-10	1.7E-09	3.6E-09
Benz[a]anthracene	56-55-3	1.7E-11	2.6E-10	2.9E-10	1.3E-10	8.2E-10	1.1E-09
Benzo[a]pyrene	50-32-8	4.2E-11	2.2E-10	1.8E-10	1.5E-10	6.0E-10	6.3E-10
Benzo[b]fluoranthene	205-99-2	7.4E-11	3.6E-10	2.7E-10	2.4E-10	8.8E-10	1.1E-09
Benzo[k]fluoranthene	207-08-9	9.8E-12	1.2E-10	1.1E-10	7.4E-11	3.4E-10	4.4E-10
Chromium		8.0E-10	4.3E-09	4.2E-09	2.8E-09	9.4E-09	2.4E-08
Chrysene	218-01-9	2.9E-10	1.2E-09	6.2E-10	1.2E-09	2.0E-09	3.2E-09
Cyclopenta[cd]pyrene	27208-37-3	4.4E-11	2.5E-10	2.3E-10	2.2E-10	6.6E-10	1.2E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.6E-11	3.9E-11	0.0E+00	7.0E-11	1.9E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	4.4E-11	7.7E-11	0.0E+00	1.9E-10	3.0E-10
Lead		1.8E-08	9.0E-08	7.8E-08	6.8E-08	2.3E-07	4.1E-07
Naphthalene	91-20-3	0.0E+00	4.8E-12	2.8E-11	0.0E+00	0.0E+00	1.7E-10

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.



° CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.
Values are rounded to two significant figures.

Table F-457. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Coaches 40<50 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	1.6E-10	3.6E-10	0.0E+00	8.7E-10	1.6E-09
Arsenic		1.1E-10	9.1E-10	6.4E-10	8.2E-10	1.6E-09	3.5E-09
Benz[a]anthracene	56-55-3	1.7E-11	2.6E-10	2.8E-10	1.2E-10	8.0E-10	1.1E-09
Benzo[a]pyrene	50-32-8	4.1E-11	2.1E-10	1.7E-10	1.5E-10	5.8E-10	6.2E-10
Benzo[b]fluoranthene	205-99-2	7.2E-11	3.5E-10	2.6E-10	2.4E-10	8.6E-10	1.0E-09
Benzo[k]fluoranthene	207-08-9	9.6E-12	1.1E-10	1.1E-10	7.3E-11	3.4E-10	4.3E-10
Chromium		7.8E-10	4.2E-09	4.1E-09	2.8E-09	9.2E-09	2.3E-08
Chrysene	218-01-9	2.9E-10	1.1E-09	6.1E-10	1.1E-09	2.0E-09	3.1E-09
Cyclopenta[cd]pyrene	27208-37-3	4.3E-11	2.5E-10	2.3E-10	2.1E-10	6.5E-10	1.2E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.5E-11	3.8E-11	0.0E+00	6.9E-11	1.9E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	4.3E-11	7.5E-11	0.0E+00	1.8E-10	2.9E-10
Lead		1.7E-08	8.8E-08	7.6E-08	6.7E-08	2.2E-07	4.0E-07
Naphthalene	91-20-3	0.0E+00	4.7E-12	2.8E-11	0.0E+00	0.0E+00	1.6E-10

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

° CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.
Values are rounded to two significant figures.

Table F-458. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Coaches 50<70 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	3.1E-10	7.2E-10	0.0E+00	1.7E-09	3.2E-09
Arsenic		2.1E-10	1.8E-09	1.3E-09	1.6E-09	3.3E-09	7.0E-09
Benz[a]anthracene	56-55-3	3.4E-11	5.2E-10	5.7E-10	2.5E-10	1.6E-09	2.2E-09
Benzo[a]pyrene	50-32-8	8.3E-11	4.2E-10	3.5E-10	3.0E-10	1.2E-09	1.2E-09
Benzo[b]fluoranthene	205-99-2	1.4E-10	7.1E-10	5.2E-10	4.8E-10	1.7E-09	2.1E-09
Benzo[k]fluoranthene	207-08-9	1.9E-11	2.3E-10	2.2E-10	1.5E-10	6.7E-10	8.6E-10
Chromium		1.6E-09	8.5E-09	8.3E-09	5.6E-09	1.8E-08	4.7E-08
Chrysene	218-01-9	5.8E-10	2.3E-09	1.2E-09	2.3E-09	3.9E-09	6.3E-09



Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Cyclopenta[cd]pyrene	27208-37-3	8.7E-11	4.9E-10	4.6E-10	4.3E-10	1.3E-09	2.3E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	3.1E-11	7.6E-11	0.0E+00	1.4E-10	3.8E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	8.6E-11	1.5E-10	0.0E+00	3.7E-10	5.9E-10
Lead		3.4E-08	1.8E-07	1.5E-07	1.3E-07	4.5E-07	8.0E-07
Naphthalene	91-20-3	0.0E+00	9.3E-12	5.5E-11	0.0E+00	0.0E+00	3.3E-10

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-459. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Referees 16<30 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	1.0E-10	2.4E-10	0.0E+00	5.8E-10	1.1E-09
Arsenic		7.2E-11	6.1E-10	4.3E-10	5.5E-10	1.1E-09	2.3E-09
Benz[a]anthracene	56-55-3	1.1E-11	1.7E-10	1.9E-10	8.2E-11	5.3E-10	7.5E-10
Benzo[a]pyrene	50-32-8	2.8E-11	1.4E-10	1.2E-10	1.0E-10	3.9E-10	4.2E-10
Benzo[b]fluoranthene	205-99-2	4.8E-11	2.4E-10	1.8E-10	1.6E-10	5.7E-10	6.9E-10
Benzo[k]fluoranthene	207-08-9	6.4E-12	7.7E-11	7.2E-11	4.9E-11	2.2E-10	2.9E-10
Chromium		5.2E-10	2.8E-09	2.8E-09	1.9E-09	6.2E-09	1.6E-08
Chrysene	218-01-9	1.9E-10	7.7E-10	4.1E-10	7.7E-10	1.3E-09	2.1E-09
Cyclopenta[cd]pyrene	27208-37-3	2.9E-11	1.6E-10	1.5E-10	1.4E-10	4.3E-10	7.7E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.0E-11	2.5E-11	0.0E+00	4.6E-11	1.3E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.9E-11	5.0E-11	0.0E+00	1.2E-10	2.0E-10
Lead		1.1E-08	5.9E-08	5.1E-08	4.5E-08	1.5E-07	2.7E-07
Naphthalene	91-20-3	0.0E+00	3.1E-12	1.8E-11	0.0E+00	0.0E+00	1.1E-10

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.



Table F-460. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Referees 30<40 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	7.0E-11	1.6E-10	0.0E+00	3.9E-10	7.1E-10
Arsenic		4.8E-11	4.1E-10	2.9E-10	3.7E-10	7.4E-10	1.6E-09
Benz[a]anthracene	56-55-3	7.5E-12	1.2E-10	1.3E-10	5.5E-11	3.6E-10	5.0E-10
Benzo[a]pyrene	50-32-8	1.9E-11	9.5E-11	7.8E-11	6.7E-11	2.6E-10	2.8E-10
Benzo[b]fluoranthene	205-99-2	3.2E-11	1.6E-10	1.2E-10	1.1E-10	3.8E-10	4.7E-10
Benzo[k]fluoranthene	207-08-9	4.3E-12	5.1E-11	4.9E-11	3.3E-11	1.5E-10	1.9E-10
Chromium		3.5E-10	1.9E-09	1.8E-09	1.2E-09	4.1E-09	1.0E-08
Chrysene	218-01-9	1.3E-10	5.1E-10	2.7E-10	5.1E-10	8.8E-10	1.4E-09
Cyclopenta[cd]pyrene	27208-37-3	1.9E-11	1.1E-10	1.0E-10	9.5E-11	2.9E-10	5.2E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	6.8E-12	1.7E-11	0.0E+00	3.1E-11	8.5E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.9E-11	3.4E-11	0.0E+00	8.2E-11	1.3E-10
Lead		7.7E-09	4.0E-08	3.4E-08	3.0E-08	1.0E-07	1.8E-07
Naphthalene	91-20-3	0.0E+00	2.1E-12	1.2E-11	0.0E+00	0.0E+00	7.3E-11

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-461. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Referees 40<50 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	6.9E-11	1.6E-10	0.0E+00	3.8E-10	6.9E-10
Arsenic		4.7E-11	4.0E-10	2.8E-10	3.6E-10	7.2E-10	1.5E-09
Benz[a]anthracene	56-55-3	7.4E-12	1.1E-10	1.2E-10	5.4E-11	3.5E-10	4.9E-10
Benzo[a]pyrene	50-32-8	1.8E-11	9.3E-11	7.6E-11	6.6E-11	2.6E-10	2.7E-10
Benzo[b]fluoranthene	205-99-2	3.2E-11	1.6E-10	1.1E-10	1.0E-10	3.8E-10	4.6E-10
Benzo[k]fluoranthene	207-08-9	4.2E-12	5.0E-11	4.8E-11	3.2E-11	1.5E-10	1.9E-10
Chromium		3.4E-10	1.9E-09	1.8E-09	1.2E-09	4.0E-09	1.0E-08
Chrysene	218-01-9	1.3E-10	5.0E-10	2.7E-10	5.0E-10	8.6E-10	1.4E-09
Cyclopenta[cd]pyrene	27208-37-3	1.9E-11	1.1E-10	1.0E-10	9.3E-11	2.8E-10	5.1E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	6.7E-12	1.7E-11	0.0E+00	3.0E-11	8.3E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.9E-11	3.3E-11	0.0E+00	8.0E-11	1.3E-10



Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Lead		7.5E-09	3.9E-08	3.4E-08	2.9E-08	9.8E-08	1.8E-07
Naphthalene	91-20-3	0.0E+00	2.0E-12	1.2E-11	0.0E+00	0.0E+00	7.2E-11

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-462. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Referees 50<70 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	1.4E-10	3.1E-10	0.0E+00	7.7E-10	1.4E-09
Arsenic		9.4E-11	8.0E-10	5.7E-10	7.2E-10	1.4E-09	3.1E-09
Benz[a]anthracene	56-55-3	1.5E-11	2.3E-10	2.5E-10	1.1E-10	7.0E-10	9.9E-10
Benzo[a]pyrene	50-32-8	3.7E-11	1.9E-10	1.5E-10	1.3E-10	5.1E-10	5.5E-10
Benzo[b]fluoranthene	205-99-2	6.4E-11	3.1E-10	2.3E-10	2.1E-10	7.6E-10	9.1E-10
Benzo[k]fluoranthene	207-08-9	8.5E-12	1.0E-10	9.5E-11	6.4E-11	3.0E-10	3.8E-10
Chromium		6.9E-10	3.7E-09	3.6E-09	2.5E-09	8.1E-09	2.1E-08
Chrysene	218-01-9	2.5E-10	1.0E-09	5.4E-10	1.0E-09	1.7E-09	2.8E-09
Cyclopenta[cd]pyrene	27208-37-3	3.8E-11	2.2E-10	2.0E-10	1.9E-10	5.7E-10	1.0E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.3E-11	3.3E-11	0.0E+00	6.1E-11	1.7E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	3.8E-11	6.6E-11	0.0E+00	1.6E-10	2.6E-10
Lead		1.5E-08	7.8E-08	6.7E-08	5.9E-08	2.0E-07	3.5E-07
Naphthalene	91-20-3	0.0E+00	4.1E-12	2.4E-11	0.0E+00	0.0E+00	1.4E-10

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-463. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators Third Trimester Fetus<0 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	1.5E-11	3.4E-11	0.0E+00	8.2E-11	1.5E-10



Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		1.0E-11	8.6E-11	6.1E-11	7.7E-11	1.5E-10	3.3E-10
Benz[a]anthracene	56-55-3	1.6E-12	2.4E-11	2.7E-11	1.2E-11	7.5E-11	1.1E-10
Benzo[a]pyrene	50-32-8	3.9E-12	2.0E-11	1.6E-11	1.4E-11	5.5E-11	5.8E-11
Benzo[b]fluoranthene	205-99-2	6.8E-12	3.3E-11	2.5E-11	2.2E-11	8.1E-11	9.8E-11
Benzo[k]fluoranthene	207-08-9	9.1E-13	1.1E-11	1.0E-11	6.9E-12	3.2E-11	4.1E-11
Chromium		7.3E-11	4.0E-10	3.9E-10	2.6E-10	8.7E-10	2.2E-09
Chrysene	218-01-9	2.7E-11	1.1E-10	5.7E-11	1.1E-10	1.8E-10	3.0E-10
Cyclopenta[cd]pyrene	27208-37-3	4.1E-12	2.3E-11	2.1E-11	2.0E-11	6.1E-11	1.1E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.4E-12	3.6E-12	0.0E+00	6.5E-12	1.8E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	4.1E-12	7.1E-12	0.0E+00	1.7E-11	2.8E-11
Lead		1.6E-09	8.3E-09	7.2E-09	6.3E-09	2.1E-08	3.8E-08
Naphthalene	91-20-3	0.0E+00	4.4E-13	2.6E-12	0.0E+00	0.0E+00	1.5E-11

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-464. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 0<2 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	8.2E-09	1.9E-08	0.0E+00	4.6E-08	8.2E-08
Arsenic		5.6E-09	4.8E-08	3.4E-08	4.3E-08	8.6E-08	1.8E-07
Benz[a]anthracene	56-55-3	8.8E-10	1.3E-08	1.5E-08	6.4E-09	4.2E-08	5.9E-08
Benzo[a]pyrene	50-32-8	2.2E-09	1.1E-08	9.1E-09	7.9E-09	3.1E-08	3.3E-08
Benzo[b]fluoranthene	205-99-2	3.8E-09	1.9E-08	1.4E-08	1.2E-08	4.5E-08	5.4E-08
Benzo[k]fluoranthene	207-08-9	5.1E-10	6.0E-09	5.7E-09	3.8E-09	1.8E-08	2.3E-08
Chromium		4.1E-08	2.2E-07	2.2E-07	1.5E-07	4.8E-07	1.2E-06
Chrysene	218-01-9	1.5E-08	6.0E-08	3.2E-08	6.0E-08	1.0E-07	1.6E-07
Cyclopenta[cd]pyrene	27208-37-3	2.3E-09	1.3E-08	1.2E-08	1.1E-08	3.4E-08	6.1E-08
Dibenz[a,h]anthracene	53-70-3	0.0E+00	8.0E-10	2.0E-09	0.0E+00	3.6E-09	9.9E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.3E-09	3.9E-09	0.0E+00	9.6E-09	1.5E-08
Lead		9.0E-07	4.6E-06	4.0E-06	3.5E-06	1.2E-05	2.1E-05
Naphthalene	91-20-3	0.0E+00	2.4E-10	1.4E-09	0.0E+00	0.0E+00	8.5E-09

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.



° CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.
Values are rounded to two significant figures.

Table F-465. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 2<6 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	2.3E-09	5.3E-09	0.0E+00	1.3E-08	2.3E-08
Arsenic		1.6E-09	1.4E-08	9.6E-09	1.2E-08	2.4E-08	5.2E-08
Benz[a]anthracene	56-55-3	2.5E-10	3.8E-09	4.2E-09	1.8E-09	1.2E-08	1.7E-08
Benzo[a]pyrene	50-32-8	6.2E-10	3.1E-09	2.6E-09	2.2E-09	8.7E-09	9.2E-09
Benzo[b]fluoranthene	205-99-2	1.1E-09	5.3E-09	3.9E-09	3.5E-09	1.3E-08	1.5E-08
Benzo[k]fluoranthene	207-08-9	1.4E-10	1.7E-09	1.6E-09	1.1E-09	5.0E-09	6.4E-09
Chromium		1.2E-08	6.3E-08	6.1E-08	4.1E-08	1.4E-07	3.5E-07
Chrysene	218-01-9	4.3E-09	1.7E-08	9.1E-09	1.7E-08	2.9E-08	4.7E-08
Cyclopenta[cd]pyrene	27208-37-3	6.4E-10	3.7E-09	3.4E-09	3.2E-09	9.7E-09	1.7E-08
Dibenz[a,h]anthracene	53-70-3	0.0E+00	2.3E-10	5.7E-10	0.0E+00	1.0E-09	2.8E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	6.4E-10	1.1E-09	0.0E+00	2.7E-09	4.4E-09
Lead		2.6E-07	1.3E-06	1.1E-06	1.0E-06	3.3E-06	6.0E-06
Naphthalene	91-20-3	0.0E+00	6.9E-11	4.1E-10	0.0E+00	0.0E+00	2.4E-09

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

° CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.
Values are rounded to two significant figures.

Table F-466. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 6<11 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	2.0E-09	4.7E-09	0.0E+00	1.1E-08	2.1E-08
Arsenic		1.4E-09	1.2E-08	8.4E-09	1.1E-08	2.1E-08	4.6E-08
Benz[a]anthracene	56-55-3	2.2E-10	3.4E-09	3.7E-09	1.6E-09	1.0E-08	1.5E-08
Benzo[a]pyrene	50-32-8	5.4E-10	2.8E-09	2.3E-09	2.0E-09	7.6E-09	8.1E-09
Benzo[b]fluoranthene	205-99-2	9.4E-10	4.6E-09	3.4E-09	3.1E-09	1.1E-08	1.4E-08
Benzo[k]fluoranthene	207-08-9	1.3E-10	1.5E-09	1.4E-09	9.5E-10	4.4E-09	5.6E-09
Chromium		1.0E-08	5.5E-08	5.4E-08	3.6E-08	1.2E-07	3.1E-07
Chrysene	218-01-9	3.8E-09	1.5E-08	7.9E-09	1.5E-08	2.6E-08	4.1E-08



Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Cyclopenta[cd]pyrene	27208-37-3	5.7E-10	3.2E-09	3.0E-09	2.8E-09	8.5E-09	1.5E-08
Dibenz[a,h]anthracene	53-70-3	0.0E+00	2.0E-10	5.0E-10	0.0E+00	9.0E-10	2.5E-09
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	5.6E-10	9.8E-10	0.0E+00	2.4E-09	3.8E-09
Lead		2.2E-07	1.2E-06	1.0E-06	8.7E-07	2.9E-06	5.2E-06
Naphthalene	91-20-3	0.0E+00	6.1E-11	3.6E-10	0.0E+00	0.0E+00	2.1E-09

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-467. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 11<16 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	1.6E-10	3.7E-10	0.0E+00	8.9E-10	1.6E-09
Arsenic		1.1E-10	9.3E-10	6.6E-10	8.4E-10	1.7E-09	3.6E-09
Benz[a]anthracene	56-55-3	1.7E-11	2.6E-10	2.9E-10	1.3E-10	8.2E-10	1.1E-09
Benzo[a]pyrene	50-32-8	4.2E-11	2.2E-10	1.8E-10	1.5E-10	6.0E-10	6.4E-10
Benzo[b]fluoranthene	205-99-2	7.4E-11	3.6E-10	2.7E-10	2.4E-10	8.8E-10	1.1E-09
Benzo[k]fluoranthene	207-08-9	9.9E-12	1.2E-10	1.1E-10	7.4E-11	3.4E-10	4.4E-10
Chromium		8.0E-10	4.3E-09	4.2E-09	2.8E-09	9.4E-09	2.4E-08
Chrysene	218-01-9	3.0E-10	1.2E-09	6.2E-10	1.2E-09	2.0E-09	3.2E-09
Cyclopenta[cd]pyrene	27208-37-3	4.4E-11	2.5E-10	2.3E-10	2.2E-10	6.6E-10	1.2E-09
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.6E-11	3.9E-11	0.0E+00	7.0E-11	1.9E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	4.4E-11	7.7E-11	0.0E+00	1.9E-10	3.0E-10
Lead		1.8E-08	9.1E-08	7.8E-08	6.9E-08	2.3E-07	4.1E-07
Naphthalene	91-20-3	0.0E+00	4.8E-12	2.8E-11	0.0E+00	0.0E+00	1.7E-10

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.



Table F-468. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Spectators 16<30 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	8.5E-11	1.9E-10	0.0E+00	4.7E-10	8.5E-10
Arsenic		5.8E-11	4.9E-10	3.5E-10	4.5E-10	8.9E-10	1.9E-09
Benz[a]anthracene	56-55-3	9.1E-12	1.4E-10	1.5E-10	6.7E-11	4.3E-10	6.1E-10
Benzo[a]pyrene	50-32-8	2.3E-11	1.1E-10	9.4E-11	8.2E-11	3.2E-10	3.4E-10
Benzo[b]fluoranthene	205-99-2	3.9E-11	1.9E-10	1.4E-10	1.3E-10	4.7E-10	5.6E-10
Benzo[k]fluoranthene	207-08-9	5.2E-12	6.2E-11	5.9E-11	4.0E-11	1.8E-10	2.3E-10
Chromium		4.2E-10	2.3E-09	2.2E-09	1.5E-09	5.0E-09	1.3E-08
Chrysene	218-01-9	1.6E-10	6.2E-10	3.3E-10	6.2E-10	1.1E-09	1.7E-09
Cyclopenta[cd]pyrene	27208-37-3	2.4E-11	1.3E-10	1.2E-10	1.2E-10	3.5E-10	6.3E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	8.3E-12	2.1E-11	0.0E+00	3.7E-11	1.0E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	2.3E-11	4.1E-11	0.0E+00	9.9E-11	1.6E-10
Lead		9.3E-09	4.8E-08	4.1E-08	3.6E-08	1.2E-07	2.2E-07
Naphthalene	91-20-3	0.0E+00	2.5E-12	1.5E-11	0.0E+00	0.0E+00	8.9E-11

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-469. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b**—Combined Gender **Spectators 30<40 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	5.7E-11	1.3E-10	0.0E+00	3.2E-10	5.8E-10
Arsenic		3.9E-11	3.3E-10	2.4E-10	3.0E-10	6.0E-10	1.3E-09
Benz[a]anthracene	56-55-3	6.2E-12	9.4E-11	1.0E-10	4.5E-11	2.9E-10	4.1E-10
Benzo[a]pyrene	50-32-8	1.5E-11	7.8E-11	6.4E-11	5.5E-11	2.1E-10	2.3E-10
Benzo[b]fluoranthene	205-99-2	2.6E-11	1.3E-10	9.6E-11	8.7E-11	3.1E-10	3.8E-10
Benzo[k]fluoranthene	207-08-9	3.5E-12	4.2E-11	4.0E-11	2.7E-11	1.2E-10	1.6E-10
Chromium		2.9E-10	1.5E-09	1.5E-09	1.0E-09	3.4E-09	8.6E-09
Chrysene	218-01-9	1.1E-10	4.2E-10	2.2E-10	4.2E-10	7.2E-10	1.2E-09
Cyclopenta[cd]pyrene	27208-37-3	1.6E-11	9.0E-11	8.4E-11	7.8E-11	2.4E-10	4.2E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	5.6E-12	1.4E-11	0.0E+00	2.5E-11	7.0E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.6E-11	2.8E-11	0.0E+00	6.7E-11	1.1E-10



Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Lead		6.3E-09	3.2E-08	2.8E-08	2.5E-08	8.2E-08	1.5E-07
Naphthalene	91-20-3	0.0E+00	1.7E-12	1.0E-11	0.0E+00	0.0E+00	6.0E-11

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-470. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 40<50 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	5.4E-11	1.2E-10	0.0E+00	3.0E-10	5.5E-10
Arsenic		3.7E-11	3.2E-10	2.2E-10	2.9E-10	5.7E-10	1.2E-09
Benz[a]anthracene	56-55-3	5.8E-12	8.9E-11	9.8E-11	4.3E-11	2.8E-10	3.9E-10
Benzo[a]pyrene	50-32-8	1.4E-11	7.3E-11	6.0E-11	5.2E-11	2.0E-10	2.2E-10
Benzo[b]fluoranthene	205-99-2	2.5E-11	1.2E-10	9.1E-11	8.3E-11	3.0E-10	3.6E-10
Benzo[k]fluoranthene	207-08-9	3.3E-12	4.0E-11	3.8E-11	2.5E-11	1.2E-10	1.5E-10
Chromium		2.7E-10	1.5E-09	1.4E-09	9.7E-10	3.2E-09	8.1E-09
Chrysene	218-01-9	1.0E-10	4.0E-10	2.1E-10	4.0E-10	6.8E-10	1.1E-09
Cyclopenta[cd]pyrene	27208-37-3	1.5E-11	8.5E-11	7.9E-11	7.4E-11	2.3E-10	4.0E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	5.3E-12	1.3E-11	0.0E+00	2.4E-11	6.6E-11
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	1.5E-11	2.6E-11	0.0E+00	6.3E-11	1.0E-10
Lead		6.0E-09	3.1E-08	2.7E-08	2.3E-08	7.7E-08	1.4E-07
Naphthalene	91-20-3	0.0E+00	1.6E-12	9.6E-12	0.0E+00	0.0E+00	5.7E-11

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.

Table F-471. Field-Specific Ingestion Lifetime Average Daily Dose^a for Lifetime Cancer Risk Assessment (LADD_{ing-field}, milligrams per kilogram BW per day) of **Field-Related Carcinogens^b—Combined Gender Spectators 50<70 Years**

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Aniline	62-53-3	0.0E+00	1.1E-10	2.5E-10	0.0E+00	6.1E-10	1.1E-09



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Chemical	CASRN ^c	LADD _{ing-field}					
		Minimum	Mean	Standard Deviation	Median	95th Percentile	Maximum
Arsenic		7.5E-11	6.3E-10	4.5E-10	5.7E-10	1.1E-09	2.4E-09
Benz[a]anthracene	56-55-3	1.2E-11	1.8E-10	2.0E-10	8.6E-11	5.6E-10	7.8E-10
Benzo[a]pyrene	50-32-8	2.9E-11	1.5E-10	1.2E-10	1.0E-10	4.1E-10	4.3E-10
Benzo[b]fluoranthene	205-99-2	5.0E-11	2.5E-10	1.8E-10	1.7E-10	6.0E-10	7.2E-10
Benzo[k]fluoranthene	207-08-9	6.7E-12	8.0E-11	7.5E-11	5.1E-11	2.3E-10	3.0E-10
Chromium		5.4E-10	2.9E-09	2.9E-09	1.9E-09	6.4E-09	1.6E-08
Chrysene	218-01-9	2.0E-10	8.0E-10	4.2E-10	8.0E-10	1.4E-09	2.2E-09
Cyclopenta[cd]pyrene	27208-37-3	3.0E-11	1.7E-10	1.6E-10	1.5E-10	4.5E-10	8.1E-10
Dibenz[a,h]anthracene	53-70-3	0.0E+00	1.1E-11	2.6E-11	0.0E+00	4.8E-11	1.3E-10
Indeno[1,2,3-cd]pyrene	193-39-5	0.0E+00	3.0E-11	5.2E-11	0.0E+00	1.3E-10	2.1E-10
Lead		1.2E-08	6.2E-08	5.3E-08	4.7E-08	1.5E-07	2.8E-07
Naphthalene	91-20-3	0.0E+00	3.2E-12	1.9E-11	0.0E+00	0.0E+00	1.1E-10

^a 35 field-specific LADD_{ing-field} are included in the table.

^b All chemicals were included in the targeted analysis of crumb rubber extracts are designated as Field-Related Chemicals.

^c CASRN: Chemical Abstracts Service Registry Number, CASRN for metals and metalloids are not included as the Study did not speciate the chemicals in the sample analysis.

Values are rounded to two significant figures.



F.7. Draft Report: Determining the Feasibility of Employing Personal Monitoring and Biological Monitoring Methods to Assess Exposures Resulting From Use of Synthetic Turf Fields with Crumb Rubber – by The University of California, Berkeley



Draft Report:

Determining the Feasibility of Employing Personal Monitoring and Biological Monitoring Methods to Assess Exposures Resulting From Use of Synthetic Turf Fields with Crumb Rubber

Principal Investigator: Asa Bradman, PhD, MS
Center for Environmental Research and Children's Health
University of California – Berkeley

Prepared for the California Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Report Authors

Asa Bradman, PhD
Carly Hyland, MS
Rosemary Castorina, PhD

Center for Environmental Research and Children's Health
University of California - Berkeley
1995 University Avenue, Suite 265
Berkeley, CA 94704



Acknowledgements

The authors wish to thank the California Office of Environmental Health Hazard Assessment (OEHHA) for sponsoring this work and especially OEHHA staff members Dr. Patty Wong, Dr. Jocelyn Claude, and Rebecca Belloso. We would also like to thank Drs. Randy Maddalena and Hugo Destailats for their presentation on laboratory measurements of field samples, Scientific Advisory Panel (SAP) members for their input, and the public commenters at the May 25, 2018 SAP meeting. We also thank Arti Kundu for assistance with literature reviews.



Table of Contents

Executive Summary	1
Background.....	1
1. Targeted Analysis	2
a. Volatile Organic Carbons (VOCs).....	3
i. Acetone	6
ii. Benzene	6
iii. Carbon Disulfide	7
iv. Ethylbenzene.....	7
v. Hexane	8
vi. Methylene chloride.....	8
vii. Methyl isobutyl ketone	8
viii. Styrene	9
ix. Toluene	9
x. Xylenes.....	10
xi. Other VOCs.....	10
Summary of VOC Targeted Analysis Recommendations	11
b. Semi-Volatile Organic Carbons (SVOCs)	12
i. Polycyclic Aromatic Hydrocarbons (PAHS).....	15
ii. Phthalates	19
(1) Benzylbutyl phthalate (BBzP).....	20
(2) Dibutyl phthalate (DBP).....	20
(3) Diethyl phthalate (DEP).....	21
(4) Diisononyl phthalate (DINP).....	22
(5) Diisodecyl phthalate (DIDP)	22
(6) Bis(2-ethylhexyl) phthalate (DEHP).....	23
iii. Potential Rubber Curatives, Antioxidants, and Other Chemicals.....	23
(1) Benzothiazole (BZT)	25
(2) Butylated hydroxytoluene (BHT)	25
Summary of SVOC Targeted Analysis Recommendations.....	26
c. Metals.....	27
i. Aluminum	31
ii. Antimony	31
iii. Arsenic	31
iv. Barium	32
v. Cadmium	32
vi. Chromium.....	32
vii. Cobalt.....	33
viii. Copper.....	33
ix. Iron	34
x. Lead	34
xi. Manganese.....	34
xii. Mercury	35
xiii. Nickel.....	35
xiv. Selenium	36



xv.	Zinc	36
	Summary of Metals Targeted Analysis Recommendations.....	37
2.	Metabolomic Analysis to Assess Crumb Rubber Exposure	38
3.	Conclusions	38
	a. Overview	38
	b. Personal Monitoring.....	39
	c. Biological Monitoring	41
4.	Methods and Protocols.....	43
	a. Personal Monitoring.....	43
	b. Biological Monitoring	43
5.	References.....	45
Appendix 1 – US EPA Method TO-15		
Appendix 2 – Radiello Passive Sampler		
Appendix 3 – Silicone Wristband Example Sampling Protocol		
Appendix 4 – Clothing and Skin Patch Example Sampling Protocol		
Appendix 5 – Dermal Wipe Example Sampling Protocol		
Appendix 6 – Breath Example Sampling Protocol		
Appendix 7 – Urine Example Sampling Protocol		
Appendix 8 – Example Participant Recruitment Outline		
Appendix 9 – Example Questionnaire		
Appendix 10 – Example Consent and Assent Forms		
Appendix 11 – Example Field Inspection Documents		
Appendix 12 – Example Biomonitoring Results Return		

For Appendices, see Supplemental Materials Section S4



Tables

Table 1. Summary of studies investigating VOCs on or near synthetic turf fields

Table 2. Summary of available research methods for quantifying exposure to VOCs found in crumb rubber

Table 3. Summary of available research methods for quantifying exposure to SVOCs found in crumb rubber

Table 4. Summary of studies investigating PAHs on or near synthetic turf fields

Table 5. Summary of PAH concentration in environmental samples collected on or near synthetic turf fields

Table 6. Summary of studies investigating phthalates on or near synthetic turf fields and personal monitoring studies (wipes and clothes)

Table 7. Summary of studies investigating potential rubber curatives, antioxidants, and other chemicals on or near synthetic turf fields

Table 8. Summary of studies investigating metals on or near synthetic turf fields

Table 9. Summary of available research methods for quantifying exposure to metals found in crumb rubber



Abbreviations

ATSDR	Agency for Toxic Substances and Disease Registry
CDC	Centers for Disease Control and Prevention
COPC	Contaminants of Potential Concern
CR	Crumb Rubber
LOD	Limit of Detection
NBI	Norwegian Building Institute
NHANES	National Health and Nutrition Examination Survey
PAH	Polycyclic Aromatic Hydrocarbon
STFCR	Synthetic Turf Field with Crumb Rubber
SVOC	Semi-Volatile Organic Carbon
US EPA	United States Environmental Protection Agency
VOC	Volatile Organic Carbon



Summary

The goal of this report is to evaluate the feasibility of studies employing personal and/or biological monitoring to assess chemical exposures resulting from the use of synthetic turf fields containing crumb rubber (STFCRs). We identified 10 Volatile Organic Carbons (VOCs) (i.e. acetone, benzene, carbon disulfide, ethylbenzene, hexane, methylene chloride, methyl isobutyl ketone, styrene, toluene, and m-, p-, and o-xylene), nine Semi-Volatile Organic Carbons (SVOCs) (i.e. polycyclic aromatic hydrocarbons (PAHs), benzylbutyl phthalate, dibutyl phthalate, diethyl phthalate, diisononyl phthalate, diisodecyl phthalate, bis (2-ethylhexyl) phthalate, benzothiazole, and butylated hydroxytoluene), and 15 metals (i.e. aluminum, antimony, arsenic, barium, cadmium, chromium, cobalt, copper, iron, lead, manganese, mercury, nickel, selenium, and zinc) that may be feasible to include in personal monitoring and/or biomonitoring studies related to use of STFCRs. Studies using skin wipe samples or other methods to quantify dermal loading may be especially valuable because exposure can be assessed immediately before and after use of STFCRs, thereby allowing specific attribution of exposure to field use, and providing key information characterizing chemical transfer rates onto skin, information needed to assess both dermal and non-dietary ingestion exposures. Several challenges are anticipated in the development of studies quantifying exposure to chemicals related to the use of STFCRs, particularly for biomonitoring. Notably, many of the chemicals previously identified in environmental or personal samples collected on or near STFCRs have been studied primarily in occupational settings, where levels of exposure may be much higher than those from STFCRs. Currently there are no biomarkers that are unique to STFCRs and humans are routinely exposed to many of the chemicals identified in this report from common sources, including vehicle exhaust, tobacco smoke, food, and personal care products, making it difficult to elucidate the source of exposure. Any future studies of exposures from STFCRs must include careful consideration of control groups.

Background

Hundreds of volatile and semi-volatile chemicals and metals have been identified in crumb rubber. To date, however, few studies have employed personal and/or biological monitoring to assess chemical exposures in athletes and bystanders from synthetic turf fields with crumb rubber (STFCRs). While many of the chemicals present in tires and crumb rubber infill have been studied in occupational settings, there are three primary studies that have or plan to employ personal monitoring and/or biomonitoring to directly examine human exposures from synthetic turf:



1. The 2016 research protocol “Collections Related to Synthetic Turf Fields with Crumb Rubber Infill” from the U.S. Environmental Protection Agency (US EPA) and the Centers for Disease Control and Prevention (CDC) Agency for Toxic Substances and Disease Registry (ATSDR) outlines sampling protocols to collect personal air samples for Volatile Organic Carbons (VOCs) and dermal samples for metals and semi Volatile Organic Carbons (sVOCs) from volunteers engaged in a sports activity on a synthetic turf field with crumb rubber infill.¹ The research protocol also details procedures to collect urine and blood samples from participants before and after the exposure measurement activities; however, the authors indicate the biological samples will be stored in a biorepository indefinitely until researchers are able to 1) determine what chemicals to analyze and 2) potentially develop new analytical methods for chemical analysis.¹
2. Studies by the State of Connecticut in which Ginsberg et al. (2011) and Simcox et al. (2011) collected personal air samples from 3-4 volunteers playing soccer for a 2-hour sampling event at one indoor and four outdoor STFCRs.² Personal air samplers were attached to the belt of players to determine contaminants in the breathing zone of young children and analyzed for 60 VOCs, 120 SVOCs, 7 nitrosamines, and particulate matter less than 10 um range (PM₁₀).² Stationary monitors located 6 inches or 3 feet above the field and upwind samples taken away from the synthetic turf field, usually on a grass field, were used to assess the vertical gradient of contaminants emitted from the field and general background levels of the target analytes.² Investigators identified 22 potentially field-related chemicals of potential concern (COPC) that were elevated at the synthetic turf fields compared to background samples, including 10 VOCs, 11 Polycyclic aromatic hydrocarbons (PAHs), and the SVOC benzothiazole.
3. Van Rooij et al. (2009) tested 1-hydroxypyrene, a metabolite of PAHs, in urine from seven soccer players before and after using STFCRs.³ Of these four volunteers, one had significantly increased hydroxypyrene excretion following study activities on the synthetic turf field, however investigators attributed the increase primarily to dietary factors.

Overall, these current and planned studies demonstrate the potential for human exposure to chemicals from synthetic turf playing fields, including several of toxicological concern. To address existing data gaps, we propose to collect personal (i.e. air and wipe) and biological (urine) samples to test for chemical contaminants or metabolites potentially due to exposures from synthetic turf fields. In this proposal, we identify key categories of chemicals that warrant study and determined the feasibility of employing personal and/or biomonitoring methods to assess exposures among athletes who use synthetic turf fields.

1. Targeted Analysis



Targeted laboratory methods involve measurement of certain, pre-selected chemicals or their breakdown products or metabolites. The chemicals may be selected by reviewing manufacturing materials or environmental or human exposure studies to identify key chemicals of potential toxicological concern. Information about the chemical, such as the retention time and mass spectra, are known.

We propose conducting targeted analysis for chemicals prioritized based on a combination of three factors: 1) chemicals that have been associated with synthetic turf fields and crumb rubber (e.g. chemicals used in the manufacturing of rubber tires); 2) chemicals that have been detected in laboratory and field studies examining potential exposures from STFCRs; and 3) the feasibility of measurements, given potentially low levels and distinguishing exposures from background, non-synthetic turf sources.

Tables 2, 3, and 9 summarize the available and likely feasible methods that have been used to assess exposure to VOCs, sVOCs, and metals. Many compounds have been studied in occupational environments, but these methods may not be feasible at the lower exposure levels expected from synthetic turf. The cells highlighted in green reflect specific chemical and media combinations that we consider feasible to include in personal monitoring and/or biomonitoring studies to detect incremental exposures resulting from use of STFCRs.

In addition to the chemicals highlighted in Tables 2, 3, and 9, we also propose targeting VOCs, SVOCs, and metals for personal and/or biomonitoring that were reported detected in STFCRs during the Fall 2017 Synthetic Turf Exposure Assessment Study field sampling.

a. VOCs

One of the primary challenges anticipated with personal monitoring or biomonitoring of VOCs is determining the amount of exposure attributable to STFCRs. Previous studies have indicated that the use of STFCRs may be associated with elevated health risks;^{1, 2} however, people are exposed to VOCs on a daily basis from many sources. Common activities that contribute to VOC exposure include pumping gas or being near vehicles, living in a home with an attached garage, smoking or being exposed to environmental tobacco, and using gas heating or gas stoves.⁴ Furthermore, while levels of urinary VOCs and VOC metabolites have been measured extensively in



occupational populations, there are data gaps regarding the utility of analyzing personal and biological samples to assess lower exposures in non-occupationally exposed populations exposed for short periods of time.⁵ Most VOCs have relatively short half-lives. For example, the half-life of toluene, xylenes, and phenol is estimated to be <5 hours and it is recommended that collection of samples for biomonitoring occurs within 15-20 minutes following the termination of exposure.⁶

Several laboratory and field studies analyzing tire crumb leachate indicate that athletes using STFCRs may be exposed to a range of VOCs.⁷ Evidence regarding the feasibility of using personal monitoring and/or biomonitoring methods to assess VOC exposure among athletes using STFCRs came primarily from three studies: 1) the study by the Norwegian Building Institute (NBI) in which Dye et al. (2006) analyzed the VOC concentration of air samples collected from three indoor STFCRs; 2) the 2009 investigation in Connecticut in which Ginsberg et al. (2011) compared levels of VOCs, SVOCs, and other contaminants in air samples from four outdoor and one indoor STFCRs to background samples collected at grass fields and upwind from synthetic turf fields;⁸ and 3) the survey in New York in which Vetrano et al. (2009) compared levels of contaminants in stationary air samplers set in the breathing zone of children (i.e. three feet above the ground) at STFCRs with background samples collected at grass fields and upwind from synthetic turf fields⁹ (Table 1). Of these studies, only Ginsberg et al. (2011) employed personal monitoring (personal air sampling) to directly monitor VOC exposure from STFCRs. We did not identify any studies that employed biomonitoring methods to assess exposure to VOCs among athletes using STFCRs.

Table 1. Summary of studies investigating VOCs on or near synthetic turf fields			
Author	Background	VOCs Measured	Results
Dye et al. (2006)	- Study by Norwegian Building Institute - Collected indoor air samples from three indoor artificial turf fields (one with recently laid rubber granulate, one with rubber granulate which had been in use for one year, and one with non-rubber granulate)	Toluene, 4-methyl-2-pentanone, total VOC	VOCs in turf fields with crumb rubber granulate were higher than the field with non-rubber granulate
Ginsberg et al. (2011) ^a	2009 field study in Connecticut - Personal and ambient air samples collected from one indoor and four outdoor synthetic turf fields with crumb rubber; compared levels of contaminants to ambient air	Carbon disulfide, chloromethane, cyclohexane, ethyl benzene, heptane, hexane, methylene chloride, methylisobutyl	Compared to background levels, levels of 10 VOCs were 1.4-12.8 times higher at outdoor fields and 0.9-92.5 times higher at indoor field



	samples collected at one grass field and samples upwind from STFCRs	ketone, toluene, xylenes	
Vetrano et al. (2009)	• 2009 air quality study in New York • Ambient air samples collected under simulated playing conditions at two different outdoor STFCRs; compared levels of contaminants to air samples collected at one grass field and samples upwind from crumb rubber fields	69 VOCs	• Eight VOCs detected in air measurements • Three VOCs (2-butanone, chloroform, and n-hexane) were detected at synthetic turf fields but not in background samples • No consistent pattern to indicate detected VOCs were associated with synthetic turf • Concentrations at turf similar to those found in background samples from grass field and upwind locations
^a The publication by Simcox et al. (2011) cited elsewhere in this report is from the same study			

Table 2 summarizes potential VOC targets and sampling media. We recommend the analysis of personal air samples as the most effective method to quantify inhalation exposure to VOCs resulting from use of synthetic turf fields containing crumb rubber.



Table 2. Summary of available research methods for quantifying exposure to VOCs found in crumb rubber.

Chemical	Personal Monitoring					Biomonitoring				
	Air		Passive badges		Other	Urine		Blood		Other
	Used in Prior Study	Likely Feasible for Turf	Used in Prior Study	Likely Feasible for Turf		Used in Prior Study	Likely Feasible for Turf	Used in Prior Study	Likely Feasible for Turf	
Acetone ⁹⁻¹⁵	Y	Y	Y	N	NA	Y	Y	Y	N	Breath
Aniline ¹⁶⁻²⁰	Y	N	NA	N	NA	Y	N	Y	N	NA
Benzene ^{15, 20}	Y	Y	Y	Y	Wristband	Y	Possibly	Y	Possibly	Breath
Carbon disulfide ^{2, 20-27}	Y	Y	Y	N	NA	Y	N	Y	N	NA
Carbon tetrachloride ^{5, 20, 28}	Y	N	NA	N	NA	Y	N	Y	N	NA
Chloroethane ^{11, 29}	Y	N	NA	N	NA	Y	N	Y	N	Breath
Chloroform ⁹	Y	N	NA	N	NA	Y	N	Y	N	NA
Chloromethane ^{9, 30}	Y	N	NA	N	NA	Y	N	NA	N	NA
Ethylbenzene ^{2, 5, 31}	Y	Possibly	NA	N	Wristband	Y	N	Y	N	NA
Formaldehyde ^{32, 33}	Y	N	NA	N	NA	N	N	N	N	NA
Hexane ^{9, 20, 34}	Y	Y	NA	N	NA	Y	N	Y	N	Breath
Isophorone ^{15, 35-37}	Y	N	Y	N	NA	NA	N	NA	N	NA
Methyl ethyl ketone ^{12, 20, 38, 39}	Y	N	NA	N	NA	Y	N	Y	N	Breath
Methylene chloride ^{2, 9, 40, 41}	Y	Y	NA	N	NA	Y	N	Y	N	Breath
Methyl isobutyl ketone ^{2, 10, 12, 20, 38, 42-45}	Y	Possibly	NA	N	NA	Y	N	Y	N	NA
Styrene ^{5, 13, 16, 31, 34, 46-49}	Y	Y	Y	Y	Wristband	Y	N	Y	N	Breath
Toluene ^{2, 5, 9, 11, 15, 23, 31, 50, 51}	Y	Y	Y	Y	Wristband	Y	Possibly	Y	N	NA



Tetrachloroethylene ^{15, 20, 52}	Y	N	NA	N	NA	Y	N	Y	N	Breath
Trichloroethylene ^{10-12, 15, 20}	Y	N	NA	N	NA	Y	N	Y	N	NA
m-,o-,p-Xylene ^{15, 20, 31, 48}	Y	Y	NA	N	Wristband	Y	Possibly	Y	N	NA
1,1-Dichloroethane ^{15, 28, 42, 53}	Y	N	Y	N	NA	Y	N	Y	N	Breath
1,1,1-Trichloroethane ^{5, 28, 54}	Y	N	NA	N	NA	Y	N	Y	N	Breath
1,1,Trichlorotrifluoroethane ^{5, 28, 55}	Y	N	NA	N	NA	N	N	NA	N	NA
Abbreviations: NA = Data Not Available.										



i. Acetone

Acetone is used as a chemical intermediate and as a solvent in chemical and pharmaceutical applications.⁵⁶ Acetone has been detected at concentrations of 27 and 28 parts per billion (ppb) in two studies evaluating the impact of tire shred fills placed above groundwater tables.^{10, 11}

Personal Monitoring: Instruments such as personal passive dosimeters have been used to measure acetone exposure in the workplace.¹³ In the 2009 air quality survey in New York, investigators detected acetone above STFCRs at concentrations ranging from 9.3-51 $\mu\text{g}/\text{m}^3$.⁹ Levels of acetone in control samples (upwind/background samples and/or samples from a grass field) ranged from non-detectable to 11 $\mu\text{g}/\text{m}^3$.⁹ The concentrations of acetone did not exceed the New York State Department of Environmental Conservation (NYC DEC's) Division of Air Resources (DAR) annual guideline value of 28,000 $\mu\text{g}/\text{m}^3$.⁹

Biological Monitoring: Although acetone exposure has primarily been examined in occupationally exposed populations, researchers in Brazil examined urinary acetone concentrations in 207 non-occupationally exposed volunteers ages 18-70 years.⁵⁷ The mean acetone concentration was 1.12 ± 0.47 mg/L, which is similar to levels found in other countries, ranging from 0.84-3.85 mg/L.⁵⁷ The researchers indicated that headspace/gas chromatography (GC) is a reproducible and reliable method to analyze urinary acetone in non-occupationally exposed populations.⁵⁷

ii. Benzene

Benzene, listed under Proposition 65 as carcinogenic, is used in many industrial processes.⁵⁸ People are routinely exposed to benzene from industrial emissions, motor vehicle exhaust, gas stations, tobacco smoke, and other sources.⁵⁸ Two different field leachate studies reported benzene in tire chip leachate at concentrations of 4 ppb and 1.35 ppb.^{10, 11}

Personal Monitoring: Benzene exposure can be assessed using active personal air samples or passive badges attached to the skin.^{15, 59} We did not identify any studies that used personal monitoring methods to assess benzene exposure among users of STFCRs.



Biomonitoring: Benzene has been measured extensively in urine and blood, however it is uncertain if benzene exposure attributable to the use of STFCRs would be high enough for biomonitoring to be feasible. Various urinary metabolites, including muconic acid, and S-phenylmercapturic acid indicate exposure to benzene. Phenol, which is easy to measure, may be a less reliable indicator of benzene exposure, as it is present in urine from other dietary and environmental sources.^{20, 60} GC or gas chromatography-mass spectroscopy (GC-MS) can be used to directly measure benzene in breath from exhaled air-collection tubes.⁶¹ Headspace/GC has been recommended for the analysis of unmetabolized benzene in blood and urine.⁶¹

iii. Carbon disulfide

Carbon disulfide is used in many industrial processes, including the production of rubber tires.⁶² Carbon disulfide is listed as a chemical causing developmental and reproductive toxicity under Proposition 65.⁶³ A 1991 study by the Scrap Tire Management Council detected carbon disulfide in tire chip leachate, but at levels below the EPAs then-proposed Toxicity Characterization Leaching Procedure (TCLP) regulatory standards.²³

Personal Monitoring: Most studies assessing exposure to carbon disulfide have been conducted in occupational settings. Meuling et al (1990) assessed exposure to carbon disulfide among workers using passive samplers over 8 hour shifts.²⁵ Concentrations in the breathing zone were estimated to be 8.0 mg/m³,²⁵ about 16 times higher than levels among volunteers in the 2009 Connecticut study playing soccer at four outdoor STFCRs (0.47 µg/m³), which in turn were about 3 times higher than concentrations detected while the volunteers played on a natural grass field.²

Biomonitoring: The carbon disulfide metabolite, 2-thioxothiazolidine-4-carboxylic acid (TTCA), has been measured in occupational workers, however some studies have suggested TTCA is not a reliable biomarker in humans with low carbon disulfide exposure.^{21, 24, 27} For example, in a study examining 10 workers with low exposures and 10 people without carbon disulfide exposure, there was no significant difference in TTCA levels among the exposed and unexposed groups.²⁷ Researchers in this study found that TTCA was higher among smokers than non-smokers, however the difference was not significant.²⁷ In a report from Finland where TTCA was measured in 116 people



aged 24-64, investigators found the column-switching method to extract TTCA provided significantly improved sensitivity, even in people with low exposure levels.²²

iv. Ethylbenzene

Ethylbenzene is used in many industries and consumer products, including gasoline, and is also used to make styrene, another chemical commonly detected in studies examining chemical exposure from STFCRs.⁶⁴

Personal Monitoring: In the 2009 study in Connecticut, the maximum concentration of ethylbenzene detected from personal air samples at the outdoor fields and on the grass field was 4.29 $\mu\text{g}/\text{m}^3$ and 2.14 $\mu\text{g}/\text{m}^3$, respectively.^{2, 8}

Biological Monitoring: Ethylbenzene can be measured in urine and blood and has been monitored primarily in occupational populations.^{5, 31}

v. Hexane

N-Hexane is an industrial solvent that was added to the Proposition 65 list in 2017 as a reproductive toxicant.⁶⁵ Occupational groups, including those working in plants manufacturing tires, may be exposed to high levels of n-hexane.⁶⁶ In the air quality study in New York, n-hexane was detected in stationary air samples placed in the breathing zone of children on STFCRs at a maximum concentration of 9 $\mu\text{g}/\text{m}^3$.⁹ Hexane was determined to be unique to synthetic turf fields, as it was not detected in the background grass/upwind sampling locations.⁹

Personal Monitoring: We did not identify any studies related to STFCRs that have used personal monitoring methods to assess hexane exposure. Personal air sampling has been used to monitor hexane exposure in occupational settings.⁶⁷

Biological Monitoring: 2,5-hexanedione, a metabolite of n-hexane, can be measured in urine and blood but may not be detectable in non-occupationally exposed populations.^{68, 69}



vi. Methylene chloride

Methylene chloride, also known as dichloromethane, is a carcinogenic chemical listed on Proposition 65 that is used primarily as a solvent.⁷⁰ In the New York field study, methylene chloride was detected in air samples collected from a synthetic turf and grass fields and one upwind background sample.⁹ However, investigators found that methylene chloride was not detected in the crumb rubber itself and there may have been laboratory contamination.⁹

Personal Monitoring: In the Connecticut investigation, the maximum concentration of methylene chloride detected in the samples from STFCRs was 14.1 $\mu\text{g}/\text{m}^3$, about 13 times the maximum concentration detected at the grass field, suggesting higher exposure from crumb rubber.²

Biomonitoring: Methylene chloride is metabolized into carbon monoxide (CO), which can be measured in blood and breath⁴⁰, however detection of CO in these samples may be confounded by exposure to vehicle exhaust or tobacco smoke.

vii. Methyl isobutyl ketone

Methyl isobutyl ketone is listed on Proposition 65 as a carcinogen and reproductive toxicant.⁷¹ It is used as a solvent and is a component of adhesives used to package, transport, and hold food.⁷¹ Methyl isobutyl ketone has been detected in tire and/or tire crumb rubber samples in field leachate studies at concentrations ranging from trace (<5 ppb) to 69 ppb.^{10, 42-44}

Personal Monitoring: Investigators in the 2009 Connecticut study detected methyl isobutyl ketone in the breathing zone of children at outdoor STFCRs at a maximum concentration of 3.39 $\mu\text{g}/\text{m}^3$.²

Biomonitoring: Methyl isobutyl ketone can be measured in urine and blood. However, it is not likely to be detectable in biological samples from non-occupationally exposed populations.³⁸



viii. Styrene

Styrene is a Proposition 65 listed carcinogen that is used in the production of plastics, rubber, and resins.⁷² People are commonly exposed to styrene from emissions from building materials, consumer products, and tobacco smoke.⁷² In a 2000 study evaluating occupational exposures in a scrap tire factory, styrene was detected in air samples, although levels did not exceed safety standards.¹⁶ Some STFCRs are made from styrene butadiene rubber (SBR) and the chemical was detected above STFCRs in indoor stadiums in 2005 Norwegian study.⁷³

Personal Monitoring: Personal monitoring has been used to assess styrene exposure in occupational settings, including personal passive dosimeters,¹³ monitoring of exhaled and environmental air,⁴⁹ and analysis of carbon and/or cotton cloths fastened to workers.^{34, 49}

Biological Monitoring: Breath may be a useful indicator of styrene exposure.⁴⁹ Unmetabolized styrene can be measured in blood and urine and has been more closely correlated with styrene levels in air and breath than the urinary metabolites mandelic acid (MA) or phenylglyoxylic acid (PGA);⁴⁹ however styrene exposure has primarily been monitored in occupational populations.³¹ It is unclear if current analytical methods could reliably detect styrene in biological samples from non-occupationally exposed populations.

ix. Toluene

Toluene is a development toxicant listed under Proposition 65.⁷⁴ Toluene is used as a solvent and gasoline additive and is a component of car exhaust and tobacco smoke⁷⁴. Toluene has been detected in tire shred leachate from lab leaching studies at levels ranging from 2.6 ppb to 190 ppb.^{11, 23} Dye et al. (2006) found levels of toluene in two indoor fields with crumb rubber ranged from 15.0-85.0 $\mu\text{g}/\text{m}^3$.⁷³ In the 2009 New York study, toluene was detected in samples over STFCRs as well as in background samples; levels did not exceed the NYC DEC's guidance value.⁹

Personal Monitoring: In the 2009 Connecticut study, the highest level of toluene detected in the breathing zone at outdoor synthetic turf fields was 52.7 $\mu\text{g}/\text{m}^3$ and off-gassing was observed.² Toluene exposure has also been quantitatively



assessed in occupational populations using patches attached to the skin and activated charcoal cloth.¹⁵

Biological Monitoring: In a 2003 study in Japan, measurements of toluene in urine was largely attributable to recent bedroom air VOC concentrations, indicating the biomonitoring reflects short term exposures.³¹

x. **Xylenes**

Xylenes, including, m-, p-, and o-xylene isomers, are solvents used in medical technology and industry, including the rubber industry.⁷⁵ In the 2004 investigation by the NBI, investigators analyzed emissions from tire-derived crumb rubber infill samples; m/p-Xylene and o-Xylene were detected at levels ranging from <5 -37 µg /kg.⁷⁶

Personal Monitoring: Xylene exposure can be assessed using diffusive air sampling in the breathing zone⁷⁷ and by attaching personal samplers, such as patches or detection indicators, to participants.¹⁵ In the Connecticut study, the maximum concentration of xylene detected in the breathing zone of participants at the outdoor STFCRs was 14.7 µg/m³.²

Biomonitoring: The parent compound of unmetabolized xylene can be measured in urine and blood. Urinary excretion of methylhippuric acid (MHA) isomers has also been correlated with xylene exposure.⁷⁷

xi. **Other VOCs**

Other VOCs that have been targeted in field studies analyzing chemical leachate from rubber tires and tire crumb include aniline, carbon tetrachloride, chloroethane, chloroform, chloromethane, formaldehyde, isophorone, methyl ethyl ketone, tetrachloroethylene, trichloroethylene, 1,1-dichloroethane, 1,1,1-trichloroethane, and 1,1,2-trichlorotrifluoroethane.⁷⁸ Given current analytical methods and low levels of detection, both in samples collected from STFCRs and among those with occupational exposure, we do not recommend including these VOCs in personal monitoring or biomonitoring studies related to the use of STFCRs.



Summary of VOC Targeted Analysis Recommendations

Given potentially low air levels of VOCs over synthetic turf fields (compared to occupational environments) and current analytical methods, we have identified 10 VOCs (i.e. acetone, benzene, carbon disulfide, ethylbenzene, hexane, methylene chloride, methyl isobutyl ketone, styrene, toluene, and m-, p-, and o-xylene) that may be feasible to analyze using personal monitoring and/or biomonitoring.

Personal Monitoring: Collection of personal air samples may be the most effective method to assess inhalation exposure to VOCs resulting from the use of synthetic turf fields. We propose collecting personal air samples to assess exposure to acetone, benzene, carbon disulfide, ethylbenzene, hexane, methylene chloride, methyl isobutyl ketone, styrene, toluene, and m-, p-, and o-xylene. Analysis of passive patches and badges for benzene, styrene, and toluene and analysis of silicone wristbands⁷⁹ for ethylbenzene, styrene, toluene, and xylenes may also be feasible (Table 2).

Biomonitoring: Biomonitoring for several VOCs is feasible, but distinguishing exposures resulting from use of STFCRs versus background exposures from traffic, building materials, or other sources raises methodological challenges. Candidate VOCs for biomonitoring include acetone, benzene, styrene, toluene, and xylenes in urine samples and acetone, benzene, hexane, methylene chloride, and styrene in breath samples (Table 2).

b. SVOCs

SVOCs are a sub-group of VOCs that have a higher molecular weight and boiling point compared to VOCs.⁸⁰ For this proposal, we focus on three primary groups of SVOCs: PAHs, phthalates, which are known contaminants of tires^{81, 82}, and potential rubber curatives, antioxidants, and other chemicals. A total of 16 PAHs have been listed as toxic pollutants by the US EPA due to their mutagenic and carcinogenic characteristics.⁸³ Phthalates are an integral constituent of many consumer products and building construction materials, tire manufacturing processes, and are commonly used as plasticizers.⁸⁴ Seven phthalates are listed under the Proposition 65 as carcinogens and/or developmental or reproductive toxicants.⁸⁵

Table 3 summarizes potential SVOC targets and sampling media.



Table 3. Summary of available research methods for quantifying exposure to SVOCs found in crumb rubber.								
	Personal Monitoring				Biomonitoring			
	Air		Passive badges on skin and/or clothes		Urine		Blood	
Chemical	Used in Prior Study	Likely Feasible for Turf	Used in Prior Study	Likely Feasible for Turf	Used in Prior Study	Likely Feasible for Turf	Used in Prior Study	Likely Feasible for Turf
PAHs ^{3, 8, 9, 73, 76, 86-89}	Y	Y	Y	N	Y	Y	N	N
Phthalates								
Benzylbutyl phthalate ^{76, 90-95, 96}	Y	Y	Y	Y	Y	Y	Y	N
Dibutyl phthalate ^{73, 76, 93, 97, 98}	Y	Y	Y	N	Y	Y	Y	N
Diethyl phthalate ^{73, 76, 92, 94}	Y	Y	Y	N	Y	Y	Y	N
Diisobutyl phthalate ^{73, 90, 99}	Y	N	Y	N	Y	N	N	N
Diisononyl phthalate ^{76, 100, 101}	Y	Y	N	N	Y	Y	N	N
Dimethyl phthalate ^{73, 90}	Y	N	Y	N	Y	N	N	N
Di-n-octyl phthalate ⁹⁰	Y	N	N	N	Y	N	Y	N
Diisodecyl phthalate ^{76, 102, 103}	Y	Y	Y	N	Y	Y	N	N
Bis(2-ethylhexyl) phthalate ^{73, 92, 95, 104, 105}	Y	Y	Y	Y	Y	Y	Y	N
Others								
2-Mercaptobenzothiazole (MBT) ^{8, 73}	Y	N	N	N	Y	N	NA	N
4-tert-(octyl)-phenol ^{8, 106, 107}	Y	N	N	N	Y	N	Y	N
Benzothiazole ^{2, 108-112}	Y	Y	Y	Y	Y	Possibly	NA	N
Benzothiazolone ⁸	Y	N	NA	N	NA	N	NA	N



Butylated hydroxytoluene ^{8, 113, 114}	Y	Y	Y	Y	Y	Possibly	N	N
Butylbenzene ^{78, 115}	Y	N	NA	N	Y	N	NA	N
Hexadecane ¹¹⁶	NA	N	N	N	N	N	NA	N
2,4-Dimethylphenol ^{7, 117}	Y	N	N	N	Y	N	N	N
Benzoic acid ^{73, 118}	Y	N	NA	N	Y	N	NA	N
Cylcohexane ^{78, 119}	Y	N	N	N	Y	N	NA	N
N-nitrosodiphenylamine ^{8, 120, 121}	Y	N	N	N	Y	N	Y	N
Phenol ^{122, 123}	Y	N	Y	N	Y	N	Y	N
Quinoline ¹⁶	Y	N	NA	N	NA	N	NA	N
Abbreviations: NA = Data Not Available.								



i. Polycyclic Aromatic Hydrocarbons (PAHs):

PAHs are a class of over 100 chemicals that are common constituents of fuels (e.g, coal, oil, and gas) and also combustion products in traffic emissions, smoke (including tobacco), and foods such as smoked or barbequed meats.¹²⁴ Humans can be exposed to PAHs through many pathways, including inhalation of polluted air or cigarette smoke, ingestion of food containing PAHs, and dermal absorption from soil or other materials with PAHs.⁸⁸ The half-life of urinary 1-hydroxypyrene, the most commonly used biomarker for PAH exposure, has been estimated to be as short as 3.9 hours.¹²⁵

We identified six studies that have analyzed environmental samples collected on or near synthetic turf fields for PAHs: 1) a 2018 study in Spain where investigators tested crumb rubber samples collected from 15 soccer fields in Spain;¹²⁶ 2) a 2005 study by the NBI in which Dye et al. (2006) measured VOCs in air samples from three indoor artificial turf fields (one with recently laid rubber granulate, one with rubber granulate which had been in use for one year, and one with non-rubber granulate);⁷³ 3) a study by Marsili et al. (2014) where investigators measured concentration of metals and PAHs in nine crumb rubber samples, including five samples obtained before they were spread on playing surfaces and four samples collected from used fields 1-8 years old;⁸⁶ 4) a study by the NBI in which Plessner & Lund (2004) analyzed PAHs in rubber granule samples from five different distributors of artificial turf systems;⁷⁶ 5) a 2009 investigation in Connecticut in which Simcox et al. (2011) compared levels of VOCs, SVOCs, and other contaminants in air samples from four outdoor and one indoor STFCR to background samples collected at grass fields and upwind from synthetic turf fields;⁸ and 6) an air quality survey in New York in which Vetrano et al. (2009) compared air contaminants in the breathing zone of children above synthetic turf fields with background samples collected on grass fields and upwind from synthetic turf fields⁹ (Table 4).

Table 4. Summary of studies investigating PAHs on or near synthetic turf fields

Author	Background	PAHs Measured	Results
Celeiro et al. (2018)	Crumb rubber samples collected from 15 football fields in Spain and analyzed for PAHs, plasticizers, and antioxidants and vulcanization additives	14 PAHs	- Median concentration of total PAHs in crumb rubber samples = 19.9 ug g⁻¹ (Range: 0.014-18 ug g⁻¹) - Pyrene was the PAH with the greatest median concentration (7 ug g⁻¹)



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Dye et al. (2006)	2005 investigation by Norwegian Building Institute - Air samples collected from three indoor artificial turf fields (one with recently laid rubber granulate, one with rubber granulate which had been in use for one year, and one with non-rubber granulate) analyzed for range of chemicals	38 PAHs in air (gas phase or adhered to particulate matter)	- Total PAHs in air as gas phase and adhered to particulate matter from the two fields with rubber granulate ranged from 174-364 ng/m³ and 6.5-11 ng/m³, respectively - Total PAHs in air as gas phase and adhered to particulate matter from the non-rubber granulate were 121 ng/m³ and 4.9 ng/m³, respectively
Marsili et al. (2014)	- Crumb rubber samples from nine fields in Italy analyzed for heavy metals and PAHs - Five new samples that had not yet been spread on playing fields, including one sample from virgin tires (not recycled tires) - Four samples obtained from fields that had been laid down for 1 to 8 years	14 PAHs	- Load of PAHs in samples installed 3-8 years prior to sampling significantly lower than newer samples - In eight of the samples, the PAHs detected at the highest concentration were benzo(b)fluoranthene or pyrene
Plesser & Lund (2004)	- Study by Norwegian Building Institute - Measured PAH content of crumb rubber samples from five distributors of artificial turf systems	16 PAHs	The total concentration of PAHs in rubber granulates exceeds the Norwegian Pollution Control Authority's normative values for most sensitive land use
Simcox et al. (2011) ^a	2009 field study in Connecticut - Personal and ambient air samples collected from one indoor and four outdoor synthetic turf fields with crumb rubber; compared levels of contaminants to ambient air samples collected at one grass field and samples upwind from crumb rubber fields	- Personal and area sampling for 5 targeted rubber-related SVOCs - Area sampling for PAHs and miscellaneous VOCs	- PAHs generally found at low concentrations (<1 ng/m³) and few found outdoors at higher concentrations than background - For outdoor fields, air concentrations on synthetic turf greater than background levels for 14 miscellaneous SVOCs - Compared to outdoor fields, a larger number of PAHs and miscellaneous SVOCs were detected in indoor fields
Vetrano et al. (2009)	2009 air quality study in New York - Ambient air samples collected under simulated playing conditions at two different outdoor synthetic turf fields; compared levels of contaminants to air samples collected at one	18 SVOCs (17 PAHs and benzothiazole)	None of the 17 PAHs or benzothiazole were detected in any ambient air samples



	grass field and samples upwind from crumb rubber fields		
^a The publication by Ginsberg et al. (2011) cited elsewhere in this report is from the same study			

Table 5 summarizes the concentration of specific PAHs observed in crumb rubber and air samples from the studies listed in Table 4. Vetrano et al. (2009) did not detect PAHs in any ambient air samples.⁹



Table 5. Summary of PAH concentration in environmental samples collected on or near synthetic turf fields

	Dye et al. (2006)		Marsili et al. (2014)		Celeiro et al. (2018)	Plesser & Lund (2004)	Simcox et al. (2011)		
	Range in gas phase air samples (ng/m ³)	Range adhered to particulate matter in air (ng/m ³)	Range in Crumb Rubber Before Use ¹ (ng/g)	Range in Crumb Rubber After Use ² (ng/g)	Range in Crumb Rubber (ng/g)	Range in Crumb Rubber (ng/g)	Outdoor Turf Range (ng/m ³)	Indoor Turf Concentration (ng/m ³)	Background Range (ng/m ³)
1-Methylnaphthalene	16.1-42.5	0.07-0.16	N/A ³	N/A ³	N/A ³	N/A ³	3.7-9.3	ND	4.1-6.9
2,6 Dimethylnaphthalene	N/A ³	N/A ³	N/A ³	N/A ³	N/A ³	N/A ³	ND-7.7	28.7	ND-10
2-Methylnaphthalene	22.3-57.7	0.06-0.13	N/A ³	N/A ³	N/A ³	N/A ³	1.9-4.2	63.4	ND-3.3
Acenaphthene	4.77-14.2	0.02-0.06	352-10,149	405-6,321	21-180	<LOD-320	2.14-3.5	17.4	ND-4.0
Acenaphthylene	6.24-78.1	0.04-0.19	N/A ³	N/A ³	19-46	190-720	ND-6.6	6.8	ND-0.77
Anthracene	0.77-1.86	0.04-0.05	38-283	7.6-182	45-330	<LOD-760	ND-ND	ND	ND-0.02
Benz[a]anthracene	0.05-0.15	0.14-0.56	92-1,613	5.4-389	340-5,700	<LOD-1,900	ND-ND	ND	ND-0.03
Benzo[a]pyrene	N/A ³	N/A ³	120-465	52-663	200-1,900	N/A ³	ND-0.19	ND	ND-0.05
Benzo[b]fluoranthene	<LOD-0.02	0.38-1.15	1,150-15,715	1248.-10,186	1,100-4,000	<LOD-3,900	ND-0.21	ND	ND-0.07
Benzo[e]pyrene	<LOD-0.03	0.39-0.78	N/A ³	N/A ³	N/A ³	120-3,100	ND-0.26	ND	ND-0.06
Benzo[ghi]fluoranthene	0.08-0.17	0.23-0.29	N/A ³	N/A ³	N/A ³	N/A ³	ND-0.08	ND	ND-ND
Bezo[ghi]perylene	N/A ³	N/A ³	396-903	240-544	1,400-6,600	<LOD-5,800	ND-0.14	ND	ND-0.06
Benzo[k]fluoranthene	N/A ³	N/A ³	68-1,203	127-3,616	310-1,400	<LOD-1,500	ND-0.08	ND	ND-0.04
Chrysene	0.10-0.19	0.38-0.76	700-3,422	244-921	200-4,100	<LOD-4,200	ND-0.34	ND	ND-0.04
Dibenz(a,h)anthracene	N/A ³	N/A ³	723-464	109-573	N/A ³	<LOD-440	N/A ³	N/A ³	N/A ³
Fluoranthene	1.66-2.81	0.48-0.67	N/A ³	N/A ³	120-13,000	120-11,000	1.68-6.76	5.6	0.58-4.0
Fluorene	7.86-19.2	0.04-0.10	427-11,025	529-7,145	40-82	<LOD-680	2.21-4.09	53.7	2.4-3.6
Indenol[1,2,3-cd]pyrene	N/A ³	N/A ³	N/A ³	N/A ³	N/A ³	N/A ³	ND-0.05	8.9	ND-0.05
Naphthalene	11.1-56.4	0.014-0.17	360-2,040	223-1,136	14-100	1,000-76,000	5.99-14.57	113	4.5-17



Office of Environmental Health Hazard Assessment
California Environmental Protection Agency

Phenanthrene	14.0-25.0	0.33-0.52	N/A ³	N/A ³	N/A ³	420-5,900	5.07-14.34	32	6.1-13
Pyrene	1.80-3.54	0.64-1.17	N/A ³	N/A ³	1,400-18,000	160-37,000	0.97-6.9	19	0.37-3.2
¹ n=5 crumb rubber samples before they were spread on playing surfaces ² n=4 crumb rubber samples from fields that had been in place for 1-8 years ³ N/A=Not Applicable; PAH not measured in this study									



In addition to studies that have collected samples directly from STFCRs, Llompart et al. (2013) collected 21 crumb rubber mulch samples from nine playgrounds and seven commercial samples of recycled rubber commercial pavers in Spain.⁸² The investigators found that the median level of total PAHs in the playground samples and commercial pavers were 8.42 µg/g and 2,812 µg/g, respectively.⁸²

Many studies that have used personal monitoring or biomonitoring methods to assess exposure to PAHs have done so in occupational settings and have monitored PAHs as a group, rather than by specific compound. In the only studies we identified that used personal or biomonitoring methods to examine PAH exposure from STFCRs, van Rooij & Jongeneelen (2009)³ measured levels of 1-hydroxypyrene among seven volunteers before and after playing on an STFCR and Simcox et al. (2011) compared levels of PAHs in personal air samples collected at STFCRs with background ambient air samples.⁸ We recommend that any studies employing personal monitoring and/or biomonitoring to examine chemical exposure among athletes using STFCRs assess PAH exposure by summing the concentrations of specific PAHs, such as those highlighted in Table 5, or using untargeted analyses to identify the PAHs with the greatest detection frequencies or concentrations.

Personal Monitoring: Various studies have employed personal monitoring methods, such as personal air sampling, to analyze exposure to PAHs, primarily in occupational populations.¹²⁷ For example, Strandberg et al. (2018) used a tape-stripping method to monitor exposure to PAHs among firefighters and office workers.⁸⁷ Studies using personal air sampling to monitor PAH exposure have found that PAH concentrations can be influenced by factors such as season of sampling and traffic intensity in the area of sampling.⁸⁹

Biomonitoring: In the only study to our knowledge to employ biomonitoring methods to assess exposure to chemicals among athletes using STFCRs, van Rooij & Jongeneelen (2009) recruited seven soccer players to take part in a 2.5-hour study on a synthetic turf field.³ Investigators measured urinary 1-hydroxypyrene on the day before, day of, and day after field activities from four volunteers.³ After playing on the field for 2.5 hours, 1-hydroxypyrene was elevated in one of the four participants whose urine was analyzed.³ However investigators noted



that the increased 1-hydroxypyrene excretion in this participant may have been attributable to the consumption of a fried hamburger directly after the sporting activities.³ The investigators concluded that if there is PAH exposure from the use of synthetic turf fields, the level of uptake is within the range of uptake from environmental exposures and/or diet.³ Given the limited available data and the relatively low background population exposures to some PAHs,⁸⁸ additional biomonitoring related to use of STFCRs is warranted.

ii. Phthalates

Phthalates are used as plasticizers in many consumer products, including vehicles, and are present in crumb rubber products.¹²⁸ Humans are exposed to phthalates by inhalation, dermal absorption, ingestion of contaminated food or liquids, and inadvertent consumption of contaminated dust or crumb rubber.^{82, 92, 129} Phthalates are often contaminants in air and dust in residential and school environments.¹²⁹

Phthalates have relatively short biological half-lives of less than 24 hours, indicating that a single spot urine sample may reliably reflect recent exposure.¹³⁰ However, phthalates devolve to several metabolites once they enter the human body, and national biomonitoring data indicate widespread exposure in the U.S. population.¹³¹ Given widespread background exposure, using biomonitoring to assess incremental exposures resulting from use of STFCRs may be challenging.

Few studies have measured phthalates in samples collected from STFCRs (Table 6). We did not identify any studies that have used personal monitoring or biomonitoring methods to directly assess exposure to phthalates among athletes using STFCRs. As take-home exposure through clothing may be particularly relevant to this population, consideration of transfers through clothing on to skin surfaces will be important to assess total exposures.⁹²

Table 6. Summary of studies investigating phthalates on or near synthetic turf fields and personal monitoring studies (wipes and clothes)

Author	Background	Phthalates Measured	Results



Plesser & Lund (2004)	2004 investigation of STFCR by Norwegian Building Institute (NBI) on behalf of the Norwegian Football Association - Phthalate concentrations reported from rubber granulate and artificial turf fiber leachate	8 phthalates: DMP; DEP; DBP; BBzP; DEHP; DOP; DINP; DIDP	- LODs ranged from 0.1-1 µg/L - Significant quantities of phthalates were found in recycled rubber granulates, particularly DINP and DEHP - Levels of DINP and DEHP in rubber granulates ranged from 57-78 mg/kg and 3.9-29 mg/kg, respectively
Dye et al. (2006)	2005 investigation by Norwegian Institute for Air Research (NILU) - Air samples collected from three indoor artificial turf fields (one with recently laid rubber granulate, one with rubber granulate which had been in use for one year, and one with non-rubber granulate) analyzed for a range of chemicals	- Gas phase: DEP, DiBP, DBP - Airborne dust: DMP, DEP, DBP, DEHP, DOP, BBP	Concentrations of total phthalates in airborne dust at the three fields ranged from 37.3 ng/m³ – 134.4 ng/m³
Abbreviations: DMP: dimethylphthalate; DEP: Diethylphthalate; DBP: dibutylphthalate; BBzP: benzylbutylphthalate; DEHP: diethylhexylphthalate; DOP: di n-octylphthalate; DINP: diisononylphthalate; and DIDP: diisodecylphthalate			

Below we summarize information on individual phthalate compounds with relevant personal and biological monitoring data from the studies listed in Table 6.

1) Benzylbutyl phthalate (BBzP)

BBzP has been measured at low levels in two samples of CR leachate from Norwegian STFCRs (Sample 1 = <LOD and Sample 2 = 0.3 ppb).⁷⁶ The reported phthalates LOD ranged from 0.1-1 ppb.

Personal Monitoring: Indoor and personal air samples confirm BBzP exposure in residential populations.⁹¹ In a study by Gong et al. (2016), investigators collected skin wipe samples from 11 adult males (non-athletes) and examined phthalate levels on bare skin (i.e. hand and forehead) and clothing-covered body locations (i.e. arm, back, calf).⁹² Investigators also measured phthalate levels on legs from five adults asked to wear just-washed jeans for 1 day, 5 days, and 10 days.⁹² The detection frequency of BBzP in skin wipe and clothing samples were <35%.⁹² The study authors also



indicated that dermal uptake would be underestimated by a factor of 2-5 if absorption through body locations covered by clothing were neglected. This study validates the use of skin wipes to assess dermal phthalate loading. Additionally, their finding that a significant proportion of total exposure is due to dermal absorption in body parts protected by clothing underscores the need to assess loading on skin after playing on STFCRs over the whole body.^{92, 93}

Biological Monitoring: Monobenzyl phthalate (MBzP) is the primary urinary metabolite of BBP validated as an exposure biomarker.⁹⁰ This metabolite can also be measured in serum and other fluids, but several studies indicate that measurements in urine are more sensitive to exposure.^{94, 95, 96}

2) Dibutyl phthalate (DBP)

DBP has been measured at low levels in two samples of CR leachate from Norwegian STFCRs (Sample 1 = 3.3 ppb and Sample 2 = 2.1 ppb).⁷⁶ The airborne dust concentration (PM_{2.5}) of DBP in the three fields sampled by Dye et al. 2006 ranged from 15.1-ng/m³ – 53.7 ng/m³.⁷³ Vapor phase DBP levels from that study ranged from 0.07 to 0.38 µg/m³, confirming that STFCRS may contribute to DBP inhalation exposures.⁷³ Llompert et al. (2014) indicated that median levels of DBP in samples of crumb rubber mulch and recycled rubber commercial pavers collected in Spain were 0.43 µg/g and 10.1 µg/g, respectively.⁸²

Biological Monitoring: The major metabolite for DBP is monobutyl phthalate (MBP). Koch et al. (2005)⁹⁸ evaluated MBP in German adults and children found that MPB exposures were higher in children (median=161 µg/g creatinine vs. adults=91.8 µg/g creatinine) and suggested that the use of personal care products may have a significant influence on DBP exposure of children.⁹⁸ Blount et al. (2000) tested MBP in U.S. adults. MPB was frequently detected (>75% of subjects) with 95th percentile and maximum MBP levels of 162 µg/g creatinine and 2,760 µg/g creatinine, respectively, confirming widespread background exposure in the U.S. population.⁹⁴



3) Diethyl phthalate (DEP)

Diethyl phthalate (DEP) is used to make plastic more flexible in products ranging from food packaging to toys to automobile parts. The major metabolite for DEP is monoethyl phthalate (MEP).¹³²

The Plesser & Lund (2004) study reported levels of 6.6 ppb and 8.3 ppb of DEP in leached rubber granulate from two samples.⁷⁶ Dye et al. (2006) reported DEP air concentrations that ranged from 2.8 ng/m³ to 27.9 ng/m³.⁷³ The median level of DEP in crumb rubber mulch and commercial rubber paver samples collected in the 2014 study in Spain were 0.37 µg/g and 2.63 µg/g, respectively.⁸²

Personal Monitoring: In a non-athlete population, Gong et al. (2016) found that, compared to bare-skinned subjects, a volunteer wearing clothing had 1/3 the uptake of DEP.⁹² The detection frequency of DEP in skin wipe samples was <35%,⁹² validating the feasibility of skin wipes to assess dermal loading.

Biological Monitoring: Blount et al. (2000) tested MEP in U.S. adults.⁹⁴ MEP was frequently detected (>75% of subjects), with maximum MEP levels of 6,790 µg/g creatinine, confirming widespread background exposure in the U.S. population.⁹⁴

4) Diisononyl phthalate (DINP)

DINP is used as a plasticizer and is listed under Proposition 65 as a carcinogen.¹⁰⁰ California law prohibits the manufacture, sale, or distribution of toys and child care articles intended for use by child under 3 years old if it: 1) contains DINP levels of greater than 0.1%; or 2) if a product can be placed in the child's mouth.¹³³

The Plesser & Lund (2004) study found that levels of DINP in two rubber granulate samples were 57 and 78 mg/kg and that 2.7 ppb and 2.2 ppb of DINP leached from the rubber granulate samples.⁷⁶ Llompарт et al. (2013) reported that the median level of DINP in crumb rubber mulch samples from playgrounds in Spain was 4,179 µg/g.⁸²



The major metabolite for DINP is monoisononyl phthalate (MiNP), which can be further oxidized to secondary metabolites with hydroxyl(MHINP), oxo(MOINP), and carboxyl (MCIOP) as functional groups.¹⁰¹

Biological Monitoring: No biomonitoring has been conducted on populations using STFCRs. The median and 95th percentile of MiNP in the 2009-2010 in the U.S. population was <LOD and 14.6 µ g/L, respectively,¹³¹ indicating widespread exposure.

5) Diisodecyl phthalate (DIDP)

DIDP is used as a plasticizer and is listed under Proposition 65 as a reproductive toxicant.¹⁰² Plessner & Lund (2004) found <1.0 ppb of DIDP in leached rubber granulates.⁷⁶

The major metabolite for DIDP is monoisononyl phthalate (MINP). The oxidative metabolites, including mono(carboxyisooctyl) phthalate (MCIOP), mono(oxoisononyl) phthalate (MOINP), and mono(hydroxyisononyl) phthalate (MHINP) have been measured in the urine of DINP-dosed rats.¹⁰³

Biological Monitoring: No biomonitoring has been conducted on populations using STFCRs. Silva et al. 2006 reported high detection frequencies (>85%) of MCIOP, MHINP, and MOINP in U.S. adults, confirming widespread background exposure in the U.S. population.¹⁰³

6) Bis(2-ethylhexyl) phthalate (DEHP)

DEHP is used to manufacture polyvinyl chloride (PVC). The US EPA has classified DEHP as a probable human carcinogen.¹³⁴ The oxidative metabolites for DEHP are mono-2-ethylhexyl phthalate (mEHP), mono-(2-ethyl-5-hydroxyhexyl) phthalate (mEHHP), mono-(2-ethyl-5-oxohexyl) phthalate (mEOHP), and mono-(2-ethyl-5-carboxypentyl) phthalate (mECP), respectively. DEHP is



ubiquitous in indoor environments, confirming widespread background exposure in the U.S. population.¹⁰⁵

The air concentration of DEHP in the three fields sampled in the Dye et al. (2006) study ranged from 5.5 ng/m³ – 31.3 ng/m³.⁷³ Llompарт et al. (2014) reported that the median level of DEHP in crumb rubber mulch samples from playgrounds and commercial rubber paver samples in Spain were 15.6 µg/g and 184 µg/g, respectively.⁸²

Personal Monitoring: In the study by Gong et al. (2016), DEHP was detected on skin wipe and clothing samples at higher levels than the other five phthalates examined,⁹² validating the feasibility of skin wipes to assess dermal loading.

Biological Monitoring: Studies in Sweden, the United States, and Germany, have detected high levels of DEHP metabolites in urine,⁹⁵ confirming widespread background exposure.

iii. Potential Rubber Curatives, Antioxidants, and Other Chemicals

Table 7 highlights four primary studies that have analyzed environmental samples collected on or near STFCRs for rubber curatives, antioxidants, and other chemicals: 1) a 2018 study in Spain in which investigators analyzed crumb rubber samples collected from 15 football fields in Spain;¹²⁶ 2) a 2005 study by the NBI in which Dye et al. (2006) measured benzothiazole (BZT) in air samples from three indoor artificial turf fields (one with recently laid rubber granulate, one with rubber granulate which had been in use for one year, and one with non-rubber granulate);⁷³ 3) the 2009 investigation in Connecticut in which Ginsberg et al. (2011) compared levels of BZT and butylated hydroxytoluene (BHT) in air samples from four outdoor and one indoor crumb rubber field to background samples collected at grass fields and upwind from synthetic turf fields;⁸ and 4) an air quality survey in New York in which Vetrano et al. (2009) compared BZT in the breathing zone of children above synthetic turf fields with background samples collected on grass fields and upwind from synthetic turf fields.⁹



Table 7. Summary of studies investigating potential rubber curatives, antioxidants, and other chemicals on or near synthetic turf fields

Author	Background	Chemicals Measured	Results
Celeiro et al. (2018)	Crumb rubber samples collected from 15 football fields in Spain and analyzed for PAHs, plasticizers, and antioxidants and vulcanization additives	Benzothiazole, 4- <i>tert</i> -butylphenol, 2-mercaptobenzothiazole, butylated hydroxyanisole, butylated hydroxytoluene	- Benzothiazole, butylated hydroxytoluene, and 4-<i>tert</i>-butylphenol detected in samples - Levels of benzothiazole and butylated hydroxytoluene ranged from 0.90-5.2 µg/g and 0.097-56 µg/g, respectively
Dye et al. (2006)	2005 investigation by Norwegian Building Institute - Air samples collected from three indoor artificial turf fields (one with recently laid rubber granulate, one with rubber granulate which had been in use for one year, and one with non-rubber granulate) analyzed for range of chemicals	Benzothiazole	- Benzothiazole levels on fields with crumb rubber granulate ranged from 4.5-31.7 µg/m³ - Benzothiazole levels on fields with non-rubber granulate ranged from 3.4-3.9 µg/m³
Ginsberg et al. (2011) ^a	2009 field study in Connecticut - Personal and ambient air samples collected from one indoor and four outdoor synthetic turf fields with crumb rubber; compared levels of contaminants to ambient air samples collected at one grass field and samples upwind from crumb rubber fields	Benzothiazole, butylated hydroxytoluene	- Main contributor to SVOC levels detected was benzothiazole - Benzothiazole concentrations were 14-fold higher indoors than outdoors - Butylated hydroxytoluene was not elevated at outdoor fields
Vetrano et al. (2009)	2009 air quality study in New York - Air samples collected under simulated playing conditions at two different outdoor synthetic turf fields; compared levels of contaminants to air samples collected at one grass field and samples upwind from crumb rubber fields	Benzothiazole	Benzothiazole was not detected in any ambient air samples



^aThe publication by Simcox et al. (2011) cited elsewhere in this report is from the same study

Crumb rubber contains many chemicals added during tire manufacturing such as vulcanizing agents, accelerators, activators, antioxidants, retarders, plasticizers, and extenders.⁷⁶ Investigators in the 2009 study in Connecticut found that of the five targeted rubber-related SVOCs measured (i.e. benzothiazole, 2-mercaptobenzothiazole, 4-tert(octyl)phenol, butylated hydroxyanisole, and butylated hydroxytoluene), benzothiazole and butylated hydroxytoluene were the only chemicals detected above background levels at outdoor fields.²⁸

1. Benzothiazole (BZT)

Benzothiazole and benzothiazole derivatives are found in many industrial and consumer products,⁹⁷ including rubber materials, herbicides, fungicides, and food flavors,⁹⁸ and also in clothing.¹⁰³ Several studies have identified benzothiazole in the leachate from rubber tire materials.⁹⁹⁻¹⁰² In a study of crumb rubber samples collected from 15 soccer fields in Spain, benzothiazole was detected in all the playground samples, with a median concentration of 1.9 µg/g.¹²⁶ Llompert et al. (2013) reported that the median level of BZT in crumb rubber mulch samples collected from nine playgrounds and seven samples of recycled rubber commercial pavers in Spain were 6.16 µg/g and 107 µg/g, respectively.⁸² Evidence suggests that benzothiazole may volatilize from crumb rubber, resulting in inhalation exposure among athletes and bystanders¹⁰².

Personal Monitoring: Levels of BZT in personal air samples in the Connecticut study ranged from <81-240 ng/m³ on outdoor STFCRs and 11,000-13,000 ng/m³ on one indoor STFCR, indicating potentially higher exposures related to use of STFCRs.

Biological Monitoring: Several studies report measurements of benzothiazoles and metabolites in urine from populations in Greece, Japan, and the U.S. (3.5 µg/g creatinine up to 11 ng/mL (creatinine-adjusted values not reported)).^{97,98}

2. Butylated hydroxytoluene (BHT)



Butylated hydroxytoluene (BHT) is phenol derivative that is used in a wide range of products in the United States, including as a preservative in food and personal care products.¹¹³ It is also commonly used as an antioxidant in the manufacturing of plastics and rubber products.¹¹⁴ Celeiro et al. (2018) collected crumb rubber samples from 15 soccer fields in Spain and detected BHT in 11 of the samples, with a median level of 1.3 µg/g (range = 0.097-56 µg/g).¹²⁶ In the study in Connecticut, BHT air measurements on outdoor STFCRs were higher than background levels.²⁸ BHT levels in stationary air samplers at the four outdoor fields and one indoor field ranged from <73-160 ng/m³ and 2100-3900 ng/m³, respectively.²⁸

Personal Monitoring: In the 2009 study in Connecticut, BHT was detected in personal air samples at one of the outdoor fields and the indoor field.²⁸ The concentration of BHT in personal air samples at these fields ranged from <81-97 ng/m³ and 1300-1800 ng/m³, respectively.²⁸

Biomonitoring: BHT can be measured in urine using GC-MS, but has primarily been monitored in populations that are occupationally exposed.¹¹⁴ Given use of this material in food, is a lower priority for biomonitoring examining exposures from STFCRs.

Summary of SVOC Targeted Analysis Recommendations

We have identified 9 SVOCs (i.e. PAHs, benzylbutyl phthalate, dibutyl phthalate, diethyl phthalate, diisononyl phthalate, diisodecyl phthalate, bis (2-ethylhexyl) phthalate, benzothiazole, and butylated hydroxytoluene) that may be feasible to target for studies using personal and/or biomonitoring to assess human exposures resulting from use of STFCRs.

Personal Monitoring: We recommend the analysis of dermal wipe samples collected from the hand, forehead, forearm, and leg of soccer players to assess SVOC exposures and transfer coefficients resulting from use of synthetic turf fields. We also propose using targeted and untargeted analyses to monitor exposure to SVOCs in clothing and skin patch samples. Because players can be exposed to SVOCs from other environmental sources and it may be difficult to attribute exposures to the use of STFCRs, we recommend comparing levels of chemicals in dermal wipe samples collected before and after study activities taking place on a synthetic turf field.



Biological Monitoring: Biomonitoring for several SVOCs, including PAHs, is feasible. However, it may be difficult to attribute exposures to the use of STFCRs, as humans are exposed to SVOCs from multiple sources on a daily basis. For future biomonitoring assessing PAH exposure, we recommend measuring 1-hydroxypyrene in urine as a marker of total PAH exposure.

c. Metals

Potential exposure to heavy metals has been a primary concern surrounding the use of STFCRs. Several studies where turf fibers, crumb rubber, and particulate matter (PM_{2.5}) were tested for metals have largely concluded that there is a low risk of exposure to metals resulting from use of STFCRs.^{9, 76, 78} For example, investigators from a 2009 study in Italy examined the metal content of CR samples collected from 32 playgrounds and found that median concentrations of all but one metal (zinc) were well below acceptable thresholds according to Italian regulatory standards.¹³⁵ People are routinely exposed to metals through a variety of sources, including food, drinking water, tobacco smoke, and personal care products,¹³⁶⁻¹³⁸ presenting challenges in attributing metal exposure among participants involved in a personal monitoring and/or biomonitoring study to the use of STFCRs.

Metals can be measured ICP/MS^{139, 140} in a variety of biological samples, including blood, urine, hair, saliva, and feces; however, some studies have not observed high correlations between external trace metal exposure assessed via personal monitors and concentrations detected in biological samples¹⁴¹. Additionally, hair samples may not be a useful medium to assess recent exposures from STFCRs, as metal concentrations in hair would reflect exposures from the past 6-12 months and cannot be attributed to a particular source.^{142, 143}

The majority of information reviewed in this section regarding the feasibility of assessing exposure to metals using personal monitoring and/or biomonitoring methods is derived from four studies: 1) a 2009 study in Italy where investigators analyzed the metal content of rubber granules collected from 32 different playgrounds in Italy¹³⁵ 2) the 2010 report produced by the California Department of Resources Recycling and Recovery (Cal Recycle) where investigators monitored levels of PM_{2.5} and metals in air samples collected above three STFCRs in the San Francisco Bay Area during active field use and in ambient air samples collected upwind from the fields;⁷⁸ 3) the study by Marsili et al. (2014) in which investigators measured the concentration of



metals and PAHs in nine crumb rubber samples, including five samples that were obtained before they were spread on playing surfaces, and four samples collected from fields that had been in place for 1-8 years;⁸⁶ and 4) a 2004 investigation by the NBI that analyzed the metal content of rubber granulate samples collected from five different distributors of artificial turf systems and two types of artificial turf fiber⁷⁶ (Table 8). Additionally, in its 2016 research protocol, the US EPA proposed using dermal wipes to assess exposure to metals among athletes using synthetic turf fields.

Table 8 summarizes the studies that have monitored metals in samples collected from STFRs, or have used personal monitoring methods to assess exposure to metals that may be applicable to future studies of athletes using synthetic turf fields. We did not identify any studies that have previously employed personal monitoring and/or biological monitoring methods to assess exposure to metals in studies related to STFCRs.

Table 8. Summary of studies investigating metals on or near synthetic turf fields			
Author	Background	Metals Measured	Results
Bocca et al. (2009)	Crumb rubber samples from 32 playgrounds in Italy analyzed for metals by microwave digestion	Aluminum, arsenic, barium, beryllium, cadmium, cobalt, chromium, copper, mercury, iron, lithium, magnesium, manganese, molybdenum, nickel, lead, rubidium, antimony, selenium, tin, strontium, thallium, vanadium, tungsten, zinc	- Highest leaching observed for zinc and magnesium (median = 2300 µg /l and 2500 µg /l, respectively) - Median leaching from tires <Limit of Quantification (LoQ) for beryllium, mercury, selenium, tin, thallium, and tungsten
California Department of Resources Recycling and Recovery (2010)	- Air samples collected on two separate days from above three synthetic turf crumb rubber fields in San Francisco Bay Area during active field use - Upwind air samples collected	Arsenic, cadmium, chromium, lead, manganese, mercury, nickel, zinc, sodium, calcium, potassium, sulfur, chlorine	- Synthetic turf fields sampled were not a significant source of PM_{2.5} containing these elements 32 elements analyzed were belows the LOD and/or laboratory minimum detectable limit
Marsili et al. (2014)	Crumb rubber samples from nine fields in Italy analyzed for heavy metals and PAHs	Cadmium, lead, chromium, nickel, copper, zinc, iron	Cadmium exceeded Italian National Amateur League limit (2.0 mg/kg) in 3 samples (range = 2.05-2.68 mg/kg)



	• 5 new samples that had not yet been spread on playing fields, including one sample from virgin tires (not recycled tires) • 4 samples obtained from fields that had been laid down for 1 to 8 years		• Zinc exceeded Italian National Amateur League limit (150 mg/kg) in all 9 samples (range = 3723-132102 mg/kg) • Lead, chromium, nickel, copper, iron below limit in all samples
Plesser & Lund (2004) ^a	• 2003 investigation by the Norwegian Building Institute (NBI) analyzed the metal content of two types of artificial turf fiber and rubber granule samples from five different distributors of artificial turf systems	Arsenic, lead, cadmium, copper, chromium, mercury, nickel, zinc	• Copper and zinc were detected in the artificial turf fiber samples • Artificial turf fiber samples from both companies exceeded the Norwegian Pollution Control Authority's normative value for zinc • Lead, cadmium, copper, and zinc detected in rubber granules but did not exceed the Norwegian Pollution Control's normative values
^a The publication by Dye et al. (2006) cited elsewhere in this report is also from the Norwegian Building Institute			

Table 9 summarizes the available research methods for quantifying exposure to metals found in crumb rubber based on our literature search. The majority of the studies that have assessed exposure to metals have taken place in occupational settings and levels observed in these studies may not be comparable to levels expected from STFCRs.



Table 9. Summary of available research methods for quantifying exposure to metals found in crumb rubber.

	Personal Monitoring				Biomonitoring			
	Air		Dermal wipes/pads		Urine		Blood	
Chemical	Used in Prior Study	Likely Feasible for Turf	Used in Prior Study	Likely Feasible for Turf	Used in Prior Study	Likely Feasible for Turf	Used in Prior Study	Likely Feasible for Turf
Aluminum ^{135, 144-146}	Y	N	Y	Y	Y	N	Y	N
Antimony ¹⁴⁷⁻¹⁴⁹	Y	N	Y	Y	Y	N	Y	N
Arsenic ^{9, 76, 131, 135, 142, 150, 151 152-154}	Y	N	Y	Y	Y	N	Y	N
Barium ^{9, 131, 135, 147, 155}	Y	N	Y	Y	Y	N	Y	N
Cadmium ^{9, 76, 135, 150, 152-154, 156}	Y	N	Y	Y	Y	N	Y	N
Chromium ^{9, 76, 135, 147, 150, 152, 153, 156-158}	Y	N	Y	Y	Y	N	Y	N
Cobalt ^{131, 135, 152, 154, 159}	Y	N	Y	Y	Y	N	Y	N
Copper ^{9, 76, 135, 147, 152, 154, 157}	Y	N	Y	Y	Y	N	Y	N
Iron ^{9, 135}	Y	N	Y	Y	Y	N	Y	N
Lead ^{9, 76, 131, 150, 152-154, 156, 160}	Y	N	Y	Y	Y	N	Y	N
Manganese ^{9, 135, 150, 152, 153}	Y	N	Y	Y	Y	N	Y	N
Mercury ^{76, 131, 135, 147, 150}	Y	N	Y	Y	Y	N	Y	N
Nickel ^{9, 76, 135, 147, 150, 152, 153, 157}	Y	N	Y	Y	Y	N	Y	N
Selenium ^{95, 141, 147}	Y	N	Y	Y	Y	N	Y	N
Zinc ^{9, 76, 135, 152, 153, 156}	Y	N	Y	Y	Y	N	Y	N



i. Aluminum

People are routinely exposed to aluminum through food, drinking water, consumer products, pharmaceuticals, and the environment.¹⁶¹ Aluminum is listed in the California Public Health Goals under risk categories for neurotoxicity and immunotoxicity.¹⁶¹ Bocca et al. (2009) found that the median level of aluminum in playground crumb rubber samples was 755 mg/kg.¹³⁵ Two lab leachate studies measured aluminum in tire chip leachate at concentrations of 550 ppb and 790 ppb.^{144, 145}

Personal Monitoring: Personal air samples have been used to monitor aluminum exposure in non-occupationally exposed individuals.¹⁵³

Biological Monitoring: Aluminum can be measured in urine and blood.¹⁴⁶

ii. Antimony

Antimony is used in cosmetics and medicines and is often alloyed with other metals.¹⁶²

Personal Monitoring: Personal air samplers can be employed to monitor antimony exposure, primarily in occupational settings.¹⁶³

Biomonitoring: Antimony can be measured in blood and urine, however levels are expected to be low in non-occupationally exposed populations.¹⁴⁹ Hair can also be measured for antimony levels, however there are several limitations that limit the utility of hair analysis in the examination of exposures from the use of synthetic turf fields.¹⁴⁸ For example, metal concentrations in hair would reflect exposures from the past 6-12 months and cannot be attributed to a particular source.^{142, 143}

iii. Arsenic

Arsenic is listed under Proposition 65 as a carcinogen.¹⁶⁴ Data from the 2010 California Department of Resources Recycling and Recovery (CalRecycle)



study indicate that arsenic was not detected in any of the on-field air samples.⁷⁸ Plesser & Lund (2004) did not detect arsenic in synthetic turf fibers and crumb rubber granules collected from five different suppliers (detection limit = 0.03 mg/kg).⁷⁶ The median concentration of arsenic in CR samples analyzed by Bocca et al. (2009) was 0.24 mg/kg,¹³⁵ which is lower than background soil levels.¹⁶⁵

Personal Monitoring: Personal air samples have been used to measure arsenic exposure in non-occupationally exposed populations.^{152, 153}

Biomonitoring: According to the ATSDR, measurement of arsenic in urine is the most reliable means of detecting recent arsenic exposure.¹⁴² Data from the U.S. population indicated that As exposure are generally low.¹³¹

iv. Barium

Barium salts are used in a variety of commercial applications, such as in the manufacture of paints, soap, paper, rubber, ceramics, and glass.¹⁶⁶ The median level of barium detected in CR tested by Bocca et al. (2009) was 22 mg/kg.¹³⁵

Personal Sampling: Personal air and dermal wipe samples have been used to assess exposure to barium, primarily in occupational settings.^{167, 168}

Biomonitoring: Barium can be monitored in blood and urine, but levels are expected to be very low in non-occupationally exposed populations.¹³¹

v. Cadmium

Cadmium (Cd) listed under Proposition 65 chemical as a carcinogen and reproductive toxicant.¹⁶⁹ People who work with Cd have the highest exposures, but the general public can also be exposed from tobacco smoke, jewelry, glassware, and some paints.¹⁶⁹ Low levels of Cd have been detected in on-field air, turf fiber, and crumb rubber samples. Investigations by CalRecycle and the NBI did not detect Cd in on-field air samples or in turf fibers and CR granules collected from artificial turf suppliers.^{76, 170} Additionally, the New York study reported similar Cd levels on and off fields.⁹ Bocca et al. (2009) reported a median Cd level in CR granules from playgrounds of 6.2 mg/kg.¹³⁵ Marsili et al.



(2014) reported cadmium levels in new and field-collected CR of 0.47-2.68 mg/kg and 0.47-2.38 mg/kg, respectively.⁸⁶

Personal Monitoring: Personal air samples have been used to measure Cd exposure in non-occupationally exposed populations.^{152, 153}

Biomonitoring: Cadmium can be measured in blood and urine. The median urinary and blood cadmium levels in the 2013-2014 in the U.S. population were 0.04 µg /L and 0.1 µg /L, respectively.¹³¹

vi. Chromium

Chromium (Cr) has two common oxidation states: Cr⁺³ is an essential nutrient, whereas Cr⁺⁶ (hexavalent chromium) is listed under Proposition 65 as a carcinogen and developmental toxicant.¹⁷¹ In the New York study, background and on-field air levels of Cr were similar.⁹ In California, on-field Cr air levels were below detection limits,⁷⁸ and were also below the LOD in turf fiber samples collected in the study in Norway (Plessner & Lund 2004).⁷⁶ The median level of chromium detected in CR granules in the study by Bocca et al. (2009) was 6.2 mg/kg.¹³⁵ Cr levels in new versus field-collect CR samples were 2.84-17.5 mg/kg and 1.9-5.4 mg/kg, respectively.⁸⁶

Personal Monitoring: Exposure to chromium has been monitored in occupational settings using dermal wipe sampling.¹⁵⁸

Biomonitoring: Cr can be analyzed in urine samples using graphite furnace atomic absorption spectroscopy.¹⁵⁷ Cr can also be measured in blood. U.S. population-based studies suggest most Cr⁺⁶ exposures are occupational.¹⁵⁷

vii. Cobalt

Cobalt (Co) and Co compounds are used to make superalloys and in pigment manufacture. Cobalt is an essential nutrient in humans and some forms are likely carcinogens. Occupational exposure occurs primarily among workers in the hard metal industry.¹⁷² The median level of Co in rubber granules collected from the playgrounds in Italy was 15 mg/kg.¹³⁵



Personal Monitoring: Co can be measured in personal air samples and dermal wipe samples and has primarily been monitored in occupational settings.^{154, 158, 159}

Biomonitoring: Co can be measured in biological samples including blood and urine. Also like Cr, U.S. population-based studies suggest most Co exposures are occupational.^{157, 173}

viii. Copper

Copper is used extensively in industry and household products and can be alloyed with zinc to form brass.¹⁷⁴ The median levels of copper in CR granules from Italian playgrounds was 12 mg/kg.¹³⁵ Plessner & Lund (2004) reported that concentrations of in crumb rubber granules and turf fiber ranged from 20-70 mg/kg and 59-68 mg/kg, respectively.⁷⁶ In the study by Marsili et al. (2014), the concentration of copper in the five new crumb rubber samples ranged from 9.50-84.49 mg/kg and the concentration in four samples collected from fields ranged from 5.49-65.11 mg/kg.⁸⁶

Personal Monitoring: Copper can be measured in personal air samples.¹⁵⁴ Personal monitoring methods have not been used to examine copper exposure from STFCRs.

Biomonitoring: Copper can be analyzed in urine samples using ICP-MS.¹⁵⁷

ix. Iron

Iron (Fe) is an essential nutrient in humans that has been detected at high levels in occupational populations, such as welders.¹⁷⁵ Median Fe levels detected in CR playground samples in Bocca et al. (2009) were 305 mg/kg.¹³⁵ Fe levels in new versus field-collect CR samples were 129.12-7,256.00 mg/kg and 262.20-1,577.50 mg/kg, respectively.⁸⁶

Personal Monitoring: Iron can be measured in personal air samples¹⁷⁶ but has not been tested in SFTCR studies.



Biomonitoring: Iron can be measured in biological samples such as blood and urine using ICP/MS.¹⁷⁷

x. Lead

Lead (Pb) is listed as a probable human carcinogen and reproductive and developmental toxicant under Proposition 65.¹⁷⁸ Lead is a widespread environmental contaminant and has been used in piping, paints, cable coverings, bullets, radiation shielding materials, and as a gasoline additive.¹⁷⁸ Lead concentrations in on-field air samples from STFCRs in California were below the LOD.⁷⁸ Lead concentrations were also below the LOD in turf fiber samples collected by Plesser & Lund (2004), however the CR Pb concentrations ranged in this study ranged from 8-20 mg/kg.⁷⁶ Additionally, the median level of Pb in the crumb rubber granules analyzed in the study by Bocca et al. (2009) was 22 mg/kg.¹³⁵ Pb levels in new versus field-collect CR samples were 11-34 and 11-39 mg/kg, respectively.⁸⁶ These levels are similar to background soil lead levels.¹⁶⁵

Personal Monitoring: Personal air samples have been used to monitor Pb exposure in non-occupationally exposed populations¹⁵³, however personal monitoring methods have not been used to examine Pb exposure from STFCRs.

Biological Monitoring: Lead can be measured in blood using ICP-MS or atomic absorption spectroscopy. Most U.S. residents have low-level exposures to Pb.¹³¹

xi. Manganese

Manganese (Mn) and manganese compounds are used in a variety of products, including in the manufacturing of steel and in batteries, ceramics, textile printing, fertilizers, and food additives and supplements.¹⁷⁹ Manganese is an essential nutrient but neurotoxic at high levels. Manganese concentrations in on-field air samples from STFCRs in California were below the LOD.⁷⁸ Median manganese levels detected in CR playground samples in Bocca et al. (2009) were 5.2 mg/kg.¹³⁵



Personal Monitoring: Personal air samples have been used to monitor manganese exposure in non-occupationally exposed populations¹⁵³, however personal monitoring methods have not been used to examine Mn exposure resulting from use of STFCRs.

Biomonitoring: Manganese can be measured in blood, urine, and deciduous teeth.¹³¹ Mn is homeostatically regulated and short-term exposure biomarkers using body fluids may not reflect overall exposure trends.¹⁸⁰

xii. Mercury

Mercury (Hg) and mercury compounds are listed under Proposition 65 as developmental toxicants.¹⁸¹ People are commonly exposed to mercury from consumption of fish, and mercury has also been used in fluorescent lights, glass thermometers, some skin creams, and dietary supplements and traditional medicine.¹⁸¹ Various studies have detected mercury below the LOD or at very low concentrations in samples collected on or near STFCRs. Mercury concentrations in on-field air samples from STFCRs in California were below the LOD.⁷⁸ Mercury concentrations were also below the LOD in turf fiber samples collected by Plessner & Lund (2004), however the CR mercury concentrations ranged in this study from 0.03-0.04 mg/kg.⁷⁶ Additionally, the median level of Hg in the crumb rubber granules analyzed in the study by Bocca et al. (2009) was 0.07 mg/kg.¹³⁵

Personal Monitoring: Exposure to mercury, primarily in occupational settings, can be analyzed via personal wipe samples.¹⁸²

Biomonitoring: Mercury can be measured in blood and urine. The median concentration of mercury in urine from the 2013-2014 NHANES was 0.13 µg /L.¹³¹ The median concentration of total mercury in blood from the 2013-2014 NHANES was 0.99 µg /L.¹³¹

xiii. Nickel

Nickel (Ni) and nickel compounds are listed on Proposition 65 due to evidence of carcinogenicity and are under consideration to be listed for reproductive toxicity.⁸⁵ Nickel has been detected below the LOD or at low concentrations in



studies evaluating air, turf fiber, and CR samples collected from synthetic turf fields. Nickel concentrations in on-field air samples from STFCRs in California and turf fiber samples analyzed by Plesser and Lund (2004) were below the LOD.^{76, 78} The median Ni concentration in samples of crumb rubber granules in Bocca et al. (2009) was 2.0 mg/kg.¹³⁵ Nickel levels in new versus field-collect CR samples were 4.1-26 and 3.9-5.8 mg/kg, respectively.⁸⁶ These levels are similar to background soil Ni levels.¹⁶⁵

Personal Monitoring: Nickel can be measured in personal air and dermal wipe samples using ICP/MS.^{153, 154, 158}

Biomonitoring: Nickel can be measured in blood and urine.¹⁵⁷

xiv. Selenium

Selenium (Se) is used in rubber, pharmaceuticals, fungicides and insecticides and is a component of tobacco smoke.¹⁸³ Selenium sulfide is listed as a carcinogen under Proposition 65.⁸⁵ The median level of selenium detected in the crumb rubber samples in Bocca et al. (2009) was <0.3 mg/kg.¹³⁵ These levels are similar to background soil Se levels.¹⁶⁵

Personal Monitoring: Personal air samples can be used to monitor exposure to selenium, primarily in occupational settings.¹⁴¹

Biomonitoring: Selenium can be measured in urine and blood and evidence suggests occupational exposures account for a majority of total selenium exposure.^{141 131}

xv. Zinc

Zinc and zinc compounds, including zinc oxide, zinc chloride, zinc sulfate, and zinc sulfide, are used in a variety of applications, including in nutritional supplements, herbicides, drinking water systems, metal plating, and vulcanization acceleration for rubber.¹⁸⁴ While zinc concentrations in on-field air samples from STFCRs in California were below the LOD,⁷⁸ Plesser & Lund (2004) found that levels in turf fibers and rubber granules exceeded the Norwegian Pollution Control Authority's normative values for zinc.⁷⁶



Additionally, Bocca et al. (2009) found the median level of zinc in rubber granules collected from playgrounds was 10,339 mg/kg.¹³⁵ Zinc levels in new versus field-collect CR samples were 3,474-13,202 mg/kg and 4,168-6,006 mg/kg, respectively.⁸⁶

Personal Monitoring: Exposure to zinc can be monitored using methods such as the analysis of personal air samples and skin wipe samples.¹⁵³

Biomonitoring: Zinc can be measured in a variety of biological samples, including blood, urine, feces, hair, saliva, and nails¹³⁶, however no studies have used biological monitoring methods to assess exposure to zinc resulting from use of STFCRs.

Summary of Metals Target Analysis Recommendations

There are various factors that may limit the feasibility of employing diverse personal monitoring and biomonitoring methods to assess exposure to metals from the use of STFCRs. Data from NHANES indicate widespread low-level exposure to metals in the general U.S. population,¹³¹ and various studies in occupational settings have indicated that occupational exposures account for the majority of total metal exposure. Additionally, the studies reviewed here that have analyzed the metal content of crumb rubber and air samples collected on or near STFCRs indicate that exposures to various metals from STFCRs may be relatively low. For example, arsenic, cadmium, chromium, lead, manganese, mercury, and nickel were each below the LOD in air, turf fiber, or CR samples analyzed in at least one of these four studies. While some metals, such as zinc, were detected at higher concentrations in environmental samples, background exposure to metals from other sources, such as food, personal care products, local soils, and tobacco smoke^{136-138, 165} pose concerns regarding the ability to elucidate the source of exposure among athletes who use STFCRs.

Personal Monitoring: Following the US EPA's proposal, we recommend the use of dermal wipe samples to analyze exposure to aluminum, antimony, arsenic, barium, cadmium, chromium, cobalt, copper, iron, lead, manganese, mercury, nickel, selenium, and zinc. We also recommend monitoring exposure to metals in using clothing and skin wipe or patch samples.

Biological Monitoring: Metals can be measured in a variety of biological samples, however evidence suggests that for many metals, levels of exposure from STFCRs are



low enough that biomonitoring may not be an effective method to assess exposure. We do not recommend biomonitoring for metals at this time.



2. Metabolomic analysis to assess crumb rubber exposure

Environmental metabolomics is the comprehensive analysis of small molecules in biofluids that can elucidate mechanisms of toxicity due to xenobiotic exposures.¹⁸⁵⁻¹⁸⁸ Exposure of an organism to an external stressor can result in changes to gene expression and protein production, both of which are subjected to a variety of homeostatic controls and feedback mechanisms. These changes are amplified at the level of the metabolome, potentially making metabolomics a sensitive indicator of the external stress.^{189, 190} Studying the products of metabolism comprehensively allows for the full characterization of biochemical changes in metabolites that occur during xenobiotic metabolism, and can therefore contribute to understanding the impact of environmental chemical exposure on human health.¹⁸⁵⁻¹⁸⁸

Recent studies have reported urinary metabolomics data measured in populations environmentally exposed to phthalates, one chemical found in CR,^{191, 192} confirming that urine metabolomics is a powerful tool to inform understanding of toxicological mechanisms of environmental stressors. For example, a metabolomics study of Chinese men exposed to phthalates indicated increased oxidative stress and fatty acid oxidation and decreased prostaglandin metabolism consistent with inflammation and metabolic disorders related to phthalate exposure.¹⁹² In California, Zhou et al. also found evidence that phthalate exposure was associated with enhanced lipid biogenesis, inflammation and altered nucleic acid metabolism.¹⁹¹ To date, no studies have examined metabolome patterns within and between populations using synthetic turf fields containing crumb rubber.

We propose initial pilot studies to investigate the effects of crumb rubber chemical exposure on the urinary and plasma metabolomes of soccer players in California. This information can be used to identify novel exposure biomarkers in urine and determine whether metabolomic profiles change with synthetic turf field use, providing insight on potential toxicological impacts of CR-related chemical exposures.

3. Conclusions

a. Overview

The US EPA has proposed collecting the following samples from study participants after an on-field sports activity: 1) three dermal wipe samples from the hand, forearm, and leg (either the calf or thigh, depending on which had more exposed skin during the sports activity) from one side of the body



for metal analysis; 2) three dermal wipe samples from the hand, forearm, and leg on the other side of the body for sVOC analysis; and 3) personal air samples from a passive sampler attached to study participants before, during, and after engaging in an on-fields sports activity for VOC analysis.¹ The US EPA has also proposed collecting urine and blood biological specimens from participants pre and post-exposure use of STFCRs, which will be stored in a biorepository indefinitely until researchers may identify which chemicals to target and potentially develop new analytical techniques to analyze the chemicals.

Based on the US EPA protocol, evidence from studies assessing exposure from crumb rubber presented in this report, and methods employed in studies examining exposures other than crumb rubber (i.e. phthalates), we conclude that it is feasible to employ personal monitoring techniques to assess chemical exposures among athletes using STFCRs, including use of 1) personal air samples, patches, and/or wristbands worn by participants for VOCs; and 2) dermal wipe samples and wristbands worn by participants for sVOCs; and 3) dermal wipe samples for metals. Importantly, studies assessing dermal loading and characterizing transfer rates of chemicals from STFCRs onto skin will provide critical information needed to assess dermal and non-dietary ingestion exposures.

Biomonitoring can provide useful exposure assessment information that can't be derived from questionnaires and personal monitoring studies. However, in studies related to use of STFCRs, the benefits of biomonitoring need to be weighed against potential challenges. While many of the chemicals that have been identified in rubber tires and tire crumb have been studied in occupational populations, evidence suggests that the methods used to analyze these chemicals and their metabolites may not be sensitive or specific enough to assess exposures at background levels in non-occupationally exposed populations.^{24, 27} Furthermore, humans are exposed to these chemicals from numerous every-day activities and the chemicals identified in this report do not provide unique CR biomarkers. Thus, it may be difficult to attribute chemical or metabolite concentrations in biofluids directly to the use of STFCRs, rather than other sources, such as eating grilled foods or pumping gas. Because of these anticipated challenges, we recommend that any biomonitoring studies target chemicals that are expected to be elevated among those who use STFCRs and are relatively easy to measure in biological samples, including PAHs, phthalates, and possibly metals such as zinc.



b. Personal Monitoring

VOCs:

Based on the studies reviewed, we recommend that any personal monitoring studies investigating chemical exposures resulting from use of STFCRs prioritize acetone, benzene, carbon disulfide, ethylbenzene, hexane, methylene chloride, methyl isobutyl ketone, styrene, toluene, and m-, p-, and o-xylene in air samples. Analysis of passive patches worn by participants has been conducted for benzene, styrene and toluene in occupational settings and may be feasible in studies assessing exposures from synthetic turf fields (Table 2).^{15, 34, 49, 59} Additionally, silicone wristbands have been used as passive samplers to monitor a wide array of environmental exposures and are a non-invasive method that could be employed to characterize exposure to ethylbenzene, styrene, toluene, and xylenes (Table 2).⁷⁹

sVOCs:

The US EPA has proposed collecting three dermal wipe samples from participants following an on-field sports activity from the hand, forearm, and leg of one the right side of the body.¹ In addition to these three samples, we also propose collection of a dermal wipe samples from the forehead of participants. We recommend collecting dermal wipe samples before and after on-field study activities. We also recommend analyzing patches attached to the skin and clothes of participants. Patches could be attached directly on the shin on the left side of the body and on the sock over the shin on the right side of the body. We recommend using targeted analyses to monitor exposure to SVOCs in dermal wipe, skin, and clothing samples, including chemicals identified by untargeted studies conducted by the Lawrence Berkeley National Laboratory. Additionally, silicone wristbands could be used to monitor exposure to PAHs while playing on STFCRs.⁷⁹ We propose using these personal monitoring methods to assess exposure to PAHs, benzylbutyl phthalate, dibutyl phthalate, diethyl phthalate, diisononyl phthalate, diisodecyl phthalate, bis (2-ethylhexyl) phthalate, benzothiazole, and butylated hydroxytoluene (Table 3).

Metals:



Consistent with the 2016 US EPA research proposal, we conclude that the use of dermal wipes may be the most feasible method to monitor exposure to metals from the use of synthetic turf fields. We propose collecting dermal wipe samples from the forehead, leg, forearm, and hand from the left side of the body before and after on-field study activities. We also recommend analyzing patches attached to the skin and clothes of participants. Patches could be attached directly on the shin on the left side of the body and on the sock over the shin on the right side of the body. We recommend assessing dermal wipe samples and skin/clothes patches for all metals listed in Table 9 (i.e. aluminum, antimony, arsenic, barium, cadmium, chromium, cobalt, copper, iron, lead, manganese, mercury, nickel, selenium, and zinc). Zinc in particular should be a focus of field studies. Although unlikely to be toxicologically significant, zinc is abundant in crumb rubber and easy to measure, and field studies will provide reliable information on transfer rates and dermal loading that can be applied to other chemicals in crumb rubber.

Summary

In summary, we recommend analysis of the following samples for personal monitoring:

- Personal air samples collected during on-field study activities to be analyzed for VOCs
- Wristbands worn during on-field study activities to be monitored for VOCs (ethylbenzene, styrene, toluene, and xylenes) and sVOCs (PAHs)
- Dermal wipe samples taken before and after on-field study activities to be analyzed for sVOCs and metals using targeted and untargeted analyses:
 - Dermal wipe samples from the forehead, leg, forearm, and hand on the right side of the body analyzed for sVOCs
 - Dermal wipe samples from the forehead, leg, forearm, and hand on the left side of the body analyzed for metals
- Passive patch dosimeters attached directly to the skin over the shin on the left side of the body for VOCs, sVOCs, and metals
- Passive patch dosimeters attached on the sock over the shin on the right side of the body for VOCs, sVOCs, and metals

Using data from the dermal wipe samples, clothing and skin patches, and



concentrations of chemicals from the field sampling during the exposure assessment study, we propose calculating transfer coefficients for particular metals and/or organic compounds, such as zinc, which is easy to measure and has been detected at high levels on or near synthetic turf fields. These transfer coefficients could be applied to other metals and organic compounds to better understand transfer from the source (CR) to the receptors (players). For example, in a 2009 intervention study with strawberry harvesters in Monterey County, we calculated malathion transfer coefficients using hand rinse malathion levels and dislodgeable foliar residue (DFR) data.¹⁹³

Because people are commonly exposed to many of the chemicals that have been detected on or near STFCRs, we recommend careful consideration of a comparison group. Samples could be collected before and after participants play on a synthetic turf field. If feasible, it would be preferable to collect samples from participants before and after field activities on both natural grass and STFCRs.

c. Biological Monitoring

VOCs:

Various investigations, primarily in occupational settings, have analyzed VOCs in breath, blood, and urine. Because humans are exposed to these VOCs from many sources, however, the literature suggests current analytical methods may not be sensitive enough to detect incremental exposures resulting from use of STFCRs. Based on the information we reviewed, we propose targeting the analysis of acetone, benzene, styrene, toluene, and m-, p-, and o-xylene in urine and acetone, benzene, hexane, methylene chloride, and styrene in breath samples (Table 2).

The analysis of breath samples may present a non-invasive technique to assess VOC exposures among users of synthetic turf fields. Over 500 VOCs can be detected in human breath,¹⁹⁴ however there is no single standard procedure for breath sampling.¹⁹⁵ In a review of the US EPA's single breath canister (SBC) method to test for exhaled VOC biomarkers, investigators indicated that when applied properly, this method can demonstrate recent VOC exposure and is ideal for assessing exposures to complex mixtures.¹⁹⁶



SVOCs:

Data gaps exist regarding the best biomarker to monitor exposure to SVOCs among athletes who use STFCRs. We recommend measuring 1-hydroxypyrene in urine as a marker of total PAH exposure (Table 3). If feasible, we recommend comparing levels of urinary 1-hydroxypyrene among those use synthetic turf fields to a control group of individuals who do not use synthetic turf fields, or among the same group of individuals before and after playing on a synthetic turf field. Using the Geometric Mean (GM=150 ng/L) and Geometric Standard Deviation (GSD=300 ng/L) of 1-hydroxypyrene levels among children in the United States from 2011-2012 NHANES data, a study employing biological monitoring to assess 1-hydroxypyrene levels among athletes using STFCRs would have 80% power to detect a difference of 120 ng/L with 50 participants and 175 ng/L with 25 participants.

Metals:

Metals can easily be monitored in biological samples such as blood and urine, however humans are exposed to metals through numerous sources, presenting challenges in attributing exposures to the use of STFCRs. At this time we do not recommend biological monitoring for metals.

Summary

In summary, we recommend analysis of the following samples for biomonitoring:

- Breath for VOCs
- Urine for SVOCs

Biomonitoring is a valuable exposure assessment tool, however there are challenges anticipated in studies assessing chemical exposures resulting from use of STFCRs. Notably, there are no unique biomarkers that would indicate exposure to chemicals found solely from STFCRs and people are exposed to many of the chemicals in crumb rubber from numerous common sources, including air pollution, tobacco smoke, and food. Thus, it may be difficult to attribute chemical exposures specifically to the use of STFCRs, rather than other sources, and we recommend that any biomonitoring studies include an appropriate control group. For example, urine or blood samples of



people who do and do not play on STFCRs could be compared, or urine and blood samples could be collected from the same group of participants 1) after on-field study activities taking place on natural grass fields, and 2) after on-field study activities taking place on STFCRs.

4. **Methods and Protocols**

a. **Personal Monitoring**

VOCs:

We propose using the US EPA method TO-15 to analyze personal air samples for VOCs and any appropriate modifications identified by LBNL. Information on method TO-15 can be found in Appendix 1.

The US EPA has proposed attaching a passive sampler (Radiello™ passive diffusive body containing Carbopack X sorbent, SigmaAldrich) to participant's shirt or jersey during a sports activity on a synthetic turf field with crumb rubber infill.¹ These samplers can be used to analyze exposure to over 70 VOCs, including BTEX compounds (benzene, toluene, ethylbenzene, o-xylene, m-xylene, and p-xylene).¹⁹⁷ See Appendix 2 for information on the Radiello™ sampler from the manufacturer.

See Appendix 3 for an example sampling protocol to administer silicone wristbands to study participants to analyze VOC exposure during on-field activities.

See Appendix 4 for an example sampling protocol to collect clothing and skin patch samples.

sVOCs:

See Appendix 3 for an example sampling protocol to administer silicone wristbands to study participants to analyze sVOC exposure during on-field activities.

See Appendix 4 for an example sampling protocol to collect clothing and skin



patch samples.

The US EPA research protocol proposed collecting wipe samples for sVOC analysis using a wetted (1:1 water:isopropanol) cotton wipe material that has been evenly wetted with 3mL of 1:1 deionized water:isopropanol using a transfer pipette. (M.G. Chemicals, Cleanroom Twill, 10.2 X 10.2 cm, cotton). See Appendix 6 for an example sampling protocol to collect dermal wipe samples for sVOC analysis, adapted from the US EPA research protocol.

Metals:

See Appendix 4 for an example sampling protocol to collect clothing and skin patch samples.

The US EPA research protocol proposed collecting wipe samples for metal analysis using a wet (water) wipe (Environmental Express, Ghost Wipe No. 4210) conforming to American Society for Testing and Materials 9ASTM E1792 requirements (ASTM-03, 2016a). See Appendix 5 for an example sampling protocol to collect dermal wipe samples for metal analysis, adapted from the US EPA research protocol.¹

b. Biological Monitoring

VOCs:

In a review of the US EPA's single breath canister (SBC) method to test for exhaled VOC biomarkers, investigators indicated that when used properly, this method reflects recent VOC exposure and is ideal for assessing exposures to complex mixtures.¹⁹⁶ See Appendix 6 for an example sampling protocol to collect breath samples for VOC analysis.

SVOCs:

See Appendix 7 for an example sampling protocol to collect urine samples for sVOC analysis, adapted from the basic urine collection protocol in the US EPA research protocol.¹



References:

1. USEPA, CDC, and ATSDR, *Research Protocol: Collections Related to Synthetic Turf Fields with Crumb Rubber Infill*. 2016, U.S. Environmental Protection Agency, Centers for Disease Control and Prevention, Agency for Toxic Substances and Disease Registry.
2. Ginsberg, G., et al., *Human health risk assessment of synthetic turf fields based upon investigation of five fields in Connecticut*. J Toxicol Environ Health A, 2011. **74**(17): p. 1150-74.
3. van Rooij, J.G. and F.J. Jongeneelen, *Hydroxypyrene in urine of football players after playing on artificial sports field with tire crumb infill*. Int Arch Occup Environ Health, 2010. **83**(1): p. 105-10.
4. Su, F.-C., B. Mukherjee, and S. Batterman, *Determinants of personal, indoor and outdoor VOC concentrations: An analysis of the RIOPA data*. Environmental research, 2013. **126**: p. 192-203.
5. Blount, B.C., et al., *Quantification of 31 volatile organic compounds in whole blood using solid-phase microextraction and gas chromatography-mass spectrometry*. Journal of Chromatography B, 2006. **832**(2): p. 292-301.
6. Heinrich-Ramm, R., et al., *Biological Monitoring for Exposure to Volatile Organic Compounds (VOCs) (IUPAC Recommendations 2000)*. Pure and Applied Chemistry, 2000. **72**(3): p. 385-436.
7. California Office of Environmental Health Hazard Assessment (OEHHA), *Evaluation of Health Effects of Recycled Waste Tires in Playground and Track Products*. 2007.
8. Simcox, N.J., et al., *Synthetic turf field investigation in Connecticut*. J Toxicol Environ Health A, 2011. **74**(17): p. 1133-49.
9. Vetrano, K. and G. Ritter, *Air quality survey of synthetic turf fields containing crumb rubber infill*. 2009, New York City Department of Health and Mental Hygiene: New York, NY.
10. Humphrey, D.N. and L.E. Katz, *Field study of water quality effects of tire shreds placed below the water table*. 2001, Proceedings of the Conference on Beneficial Use of Recycled Materials in Transportation Applications, Air and Waste Management Association: Pittsburgh, PA.
11. Exponent, *Tire shred leachate study: Chemical composition and aquatic toxicity for above- and below-water-table applications*. 2003, Exponent: Oakland, CA.
12. Gunter, M., et al., *The environmental sustainability of scrap tire chip in environmental and civil engineering applications: a laboratory investigation*, in *Environmental Geotechnics Report*. 1999, Department of Civil and Environmental Engineering. University of Wisconsin: Madison, Wisconsin.



13. Prieto, M.J., D. Marhuenda, and A. Cardona, *Analysis of styrene and its metabolites in blood and urine of workers exposed to both styrene and acetone*. J Anal Toxicol, 2002. **26**(1): p. 23-8.
14. Fujino, A., et al., *Biological monitoring of workers exposed to acetone in acetate fibre plants*. British Journal of Industrial Medicine, 1992. **49**(9): p. 654-657.
15. Creta, M., et al., *A Method to Quantitatively Assess Dermal Exposure to Volatile Organic Compounds*. Ann Work Expo Health, 2017. **61**(8): p. 975-985.
16. Chien, Y.-C., et al., *Assessment of occupational health hazards in scrap-tire shredding facilities*. Science of The Total Environment, 2003. **309**(1): p. 35-46.
17. Chen, C.F., et al., *Automated spectrophotometric assay for urine p-aminophenol by an oxidative coupling reaction*. Ann Clin Lab Sci, 2004. **34**(3): p. 336-40.
18. Dierkes, G., et al., *N-Acetyl-4-aminophenol (paracetamol), N-acetyl-2-aminophenol and acetanilide in urine samples from the general population, individuals exposed to aniline and paracetamol users*. Int J Hyg Environ Health, 2014. **217**(4-5): p. 592-9.
19. Ward, E.M., et al., *Monitoring of Aromatic Amine Exposures in Workers at a Chemical Plant With a Known Bladder Cancer Excess*. JNCI: Journal of the National Cancer Institute, 1996. **88**(15): p. 1046-1053.
20. Occupational Safety and Health Administration (OSHA), *Surface Contaminants, Skin Exposure, Biological Monitoring and Other Analyses*. 2014, United States Department of Labor.
21. Kivisto, H., *TTCA measurements in biomonitoring of low-level exposure to carbon disulphide*. Int Arch Occup Environ Health, 2000. **73**(4): p. 263-9.
22. Riihimaki, V., et al., *Assessment of exposure to carbon disulfide in viscose production workers from urinary 2-thiothiazolidine-4-carboxylic acid determinations*. Am J Ind Med, 1992. **22**(1): p. 85-97.
23. Scrap Tire Management Council, *The RMA TCLP Assessment Project: Radian Report. Educational seminar on scrap tire management*. 1991: Washington D.C.
24. Chang, H.Y., et al., *Biological monitoring of carbon disulphide: kinetics of urinary 2-thiothiazolidine-4-carboxylic acid (TTCA) in exposed workers*. Toxicol Ind Health, 2002. **18**(1): p. 1-14.
25. Meuling, W.J., C.L. Bragt Pc Fau - Braun, and C.L. Braun, *Biological monitoring of carbon disulfide*. 1990(0271-3586).
26. Krstev, S., et al., *Environmental and biological monitoring in carbon disulfide exposure assessment*. Med Lav, 1993. **84**(6): p. 473-81.
27. Lovreglio, P., et al., *Validity of urinary 2-thiothiazolidine-4-carboxylic acid (TTCA) as biomarker of exposure to very low concentrations of carbon disulphide: preliminary results*. G Ital Med Lav Ergon, 2008. **30**(1): p. 55-60.



28. Simcox, N.J., A. Bracker, and J. Meyer, *Artificial Turf Field Investigation in Connecticut Final Report* 2010, University of Connecticut Health Center, Section of Occupational and Environmental Medicine.
29. Agency for Toxic Substances and Disease Registry (ATSDR), *Toxicological Profile for Chloroethane*. 1998, U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.
30. Agency for Toxic Substances and Disease Registry (ATSDR), *Toxicological profile for chloromethane*. 1998, U.S. Department of Health and Human Services: Atlanta, GA.
31. Wang, B.L., et al., *Unmetabolized VOCs in urine as biomarkers of low level exposure in indoor environments*. J Occup Health, 2007. **49**(2): p. 104-10.
32. Moretto, R. *Environmental and Health Assessment of the Use of Elastomer Granulates (Virgin and from Used Tyres) as Filling in Third-Generation Artificial Pitches*. 2007; Available from: https://www.aliapur.fr/pdf/1_environmental_evaluation_0.pdf
33. Agency for Toxic Substances and Disease Registry (ATSDR). *Public Health Statement Formaldehyde*. 2008; Available from: <https://www.atsdr.cdc.gov/ToxProfiles/tp111-c1-b.pdf>.
34. Eriksson, K. and L. Wiklund, *Dermal Exposure to Styrene in the Fibreglass Reinforced Plastics Industry*. The Annals of Occupational Hygiene, 2004. **48**(3): p. 203-208.
35. Transportation, M.D.o., *A comparative study of the toxicity of chipped tires and wood chips leachate*. Report # 95-10-6161. 1995.
36. Agency for Toxic Substances and Disease Registry (ATSDR), *Toxicological Profile for Isophorone*. 1989, Agency for Toxic Substances and Disease Registry (ATSDR) U.S. Public Health Service, U.S. Environmental Protection Agency (EPA).
37. Samimi, B., *Exposure to isophorone and other organic solvents in a screen printing plant*. Am Ind Hyg Assoc J, 1982. **43**(1): p. 43-8.
38. Kawai, T., et al., *Methyl isobutyl ketone and methyl ethyl ketone in urine as biological markers of occupational exposure to these solvents at low levels*. Int Arch Occup Environ Health, 2003. **76**(1): p. 17-23.
39. Toxicology Data Network. *Methyl ethyl ketone*. [cited 2018; Available from: <https://toxnet.nlm.nih.gov/cgi-bin/sis/search/a?dbs+hsdb:@term+@DOCNO+99>.
40. Hoffer, E., et al., *Monitoring of occupational exposure to methylene chloride: sampling protocol and stability of urine samples*. J Anal Toxicol, 2005. **29**(8): p. 794-8.
41. Health & Safety Laboratory. *Guidance Sheet for Dichloromethane (DCM) Monitored by analysis of carbon monoxide (CO) in breath*. Biological Monitoring



- Guidance Values [cited 2018; Available from:
[https://www.hsl.gov.uk/media/1188353/dichloromethane \(dcm\) in breath.pdf](https://www.hsl.gov.uk/media/1188353/dichloromethane_(dcm)_in_breath.pdf).
42. Humphrey, D.N.a.K., L.E., *Five-year study of the water quality effects of tire shreds placed above the water table*. 2000, Transportation Research Board, 79th Annual Meeting, January 9-13, 2000, Paper No. 00-0892.: Washington, D.C.
 43. Miller, W.L.a.C., P.A., *A study of waste tire leachability in potential disposal and usage environments*. . 1993, College of Engineering, University of Florida, Department of Environmental Engineering Sciences.
 44. Edil, T.B., Park, J.K. and Kim, J.Y. J. , *Effectiveness of scrap tire chips as sorptive drainage material*. . Environ. Engineer, 2003. **130**(7).
 45. Granvil, C.P., M. Sharkawi, and G.L. Plaa, *Metabolic fate of methyl n-butyl ketone, methyl isobutyl ketone and their metabolites in mice*. Toxicology Letters, 1994. **70**(3): p. 263-267.
 46. Chang, F.-H., et al., *Emission characteristics of VOCs from athletic tracks*. Journal of Hazardous Materials, 1999. **70**(1): p. 1-20.
 47. Highsmith, R., Thomas, K.W., Williams R.W. , *A Scoping-Level Field Monitoring Study of Synthetic Turf Fields and Playgrounds*. 2009, U.S. Environmental Protection Agency: Washington, DC.
 48. Janasik, B., et al., *Unmetabolized VOCs in urine as biomarkers of low level occupational exposure*. Int J Occup Med Environ Health, 2010. **23**(1): p. 21-6.
 49. Ong, C.N., et al., *Biological monitoring of exposure to low concentrations of styrene*. Am J Ind Med, 1994. **25**(5): p. 719-30.
 50. Tatlisoz, N., Edil, T., Benson, C., Park, J. and Kim, J., *Review of environmental suitability of scrap tires*. *Environmental Geotechnics Report No: 96-7*. 1996, Department of Civil and Environmental Engineering, University of Wisconsin, Madison.
 51. Florida Community College at Jacksonville Nassau Center, *Study of the suitability of ground rubber tire as a parking lot surface, prepared for: State of Florida Department of Environmental Protection*. 1999.
 52. McKernan, L.T., et al., *Biological exposure assessment to tetrachloroethylene for workers in the dry cleaning industry*. Environmental Health, 2008. **7**: p. 12-12.
 53. Delaware Health and Social Services Division of Public Health. *1,1-Dichloroethane*. 2013; Available from:
<http://www.dhss.delaware.gov/dph/files/dichlethane11faq.pdf>.
 54. Agency for Toxic Substances and Disease Registry (ATSDR). *Public Health Statement 1,1,1-Trichloroethane*. 2006; Available from:
<https://www.atsdr.cdc.gov/ToxProfiles/tp70-c1-b.pdf>.
 55. Santa Cruz Biotechnology. *1,1,2-Trichloro-1,2,2-trifluoroethane*. [cited 2018; Available from: <http://datasheets.scbt.com/sc-251541.pdf>.



56. U.S. Environmental Protection Agency, *Toxicological Review of Acetone* (CAS No. 67-64-1) *In Support of Summary Information on the Integrated Risk Information System (IRIS)*. 2003: Washington, DC.
57. Oliveira, D., Maria E., *A simple and rapid method for urinary acetone analysis by headspace/gas chromatography*. Química Nova, 2007. **30**: p. 1362-1364.
58. California Office of Environmental Health Hazard Assessment (OEHHA). *Benzene Proposition 65 Warnings* 2016 [cited 2018; Available from: https://www.p65warnings.ca.gov/sites/default/files/downloads/factsheets/benzene_fact_sheet.pdf].
59. Weisel, C.P., *Benzene exposure: An overview of monitoring methods and their findings*. Chemico-biological interactions, 2010. **184**(0): p. 58-66.
60. Agency for Toxic Substances and Disease Registry (ATSDR). *Toxicological Profile for Benzene*. Available from: <https://www.atsdr.cdc.gov/ToxProfiles/tp3.pdf>.
61. Ong, C.N. and B.L. Lee, *Determination of benzene and its metabolites: application in biological monitoring of environmental and occupational exposure to benzene*. J Chromatogr B Biomed Appl, 1994. **660**(1): p. 1-22.
62. Agency for Toxic Substances and Disease Registry (ATSDR), *Toxicological profile for carbon disulfide*. 1996, U.S. Department of Health and Human Services Public Health Service: Atlanta, GA.
63. California Office of Environmental Health Hazard Assessment (OEHHA). *Carbon disulfide*. [cited 2018; Available from: <https://oehha.ca.gov/chemicals/carbon-disulfide>].
64. Agency for Toxic Substances and Disease Registry (ATSDR). *Public Health Statement Ethylbenzene*. 2010; Available from: <https://www.atsdr.cdc.gov/ToxProfiles/tp110-c1-b.pdf>.
65. California Office of Environmental Health Hazard Assessment (OEHHA). *Consideration of n-Hexane for Listing under Proposition 65 as Known to Cause Reproductive Toxicity*. 2017 [cited 2018; Available from: <https://oehha.ca.gov/media/downloads/proposition-65/chemicals/n-hexanehid090117.pdf>].
66. Agency for Toxic Substances and Disease Registry (ATSDR). *Public Health Statement for n-Hexane*. 1999 [cited 2018; Available from: <https://www.atsdr.cdc.gov/ToxProfiles/tp113-c1-b.pdf>].
67. Periago, J.F., et al., *Correlation between concentrations of n-hexane and toluene in exhaled and environmental air in an occupationally exposed population*. J Appl Toxicol, 1994. **14**(1): p. 63-7.
68. Agency for Toxic Substances and Disease Registry. *Toxicological Profile for n-Hexane*. 1999; Available from: <https://www.atsdr.cdc.gov/toxprofiles/tp113.pdf>.



69. Cardona, A., et al., *Biological monitoring of occupational exposure to n-hexane by measurement of urinary 2,5-hexanedione*. Int Arch Occup Environ Health, 1993. **65**(1): p. 71-4.
70. California Office of Environmental Health Hazard Assessment (OEHHA). *Methylene Chloride (Dichloromethane)*. [cited 2018; Available from: <https://oehha.ca.gov/chemicals/methylene-chloride-dichloromethane>.
71. California Office of Environmental Health Hazard Assessment (OEHHA). *Methyl Isobutyl Ketone*. [cited 2018; Available from: <https://oehha.ca.gov/chemicals/methyl-isobutyl-ketone>.
72. U.S. Environmental Protection Agency. *Styrene*. 2000; Available from: <https://www.epa.gov/sites/production/files/2016-09/documents/styrene.pdf>.
73. Dye, C., A. Bjerke, and S. Schmidbauer, *Measurement of air pollution in indoor artificial turf halls*. 2006, Norwegian Pollution Control Authority/Norwegian Institute for Air Research.
74. California Office of Environmental Health Hazard Assessment (OEHHA). *Toluene*. [cited 2018; Available from: <https://oehha.ca.gov/chemicals/toluene>.
75. Kandyala, R., S.P.C. Raghavendra, and S.T. Rajasekharan, *Xylene: An overview of its health hazards and preventive measures*. Journal of Oral and Maxillofacial Pathology : JOMFP, 2010. **14**(1): p. 1-5.
76. Plesser, T.S. and O.J. Lund, *Potential health and environmental effects linked to artificial turf systems-final report*. 2004, Norwegian Building Institute: Trondheim, Norway.
77. Inoue, O., et al., *Excretion of methylhippuric acids in urine of workers exposed to a xylene mixture: comparison among three xylene isomers and toluene*. Int Arch Occup Environ Health, 1993. **64**(7): p. 533-9.
78. California Department of Resources Recycling and Recovery, *Safety Study of Artificial Turf Containing Crumb Rubber Infill Made From Recycled Tires: Measurements of Chemicals and Particulates in the Air, Bacteria in the Turf, and Skin Abrasions Caused by Contact with the Surface*. 2010, California Office of Environmental Health Hazard Assessment (OEHHA).
79. Oregon State University. *Methods*. College of Agricultural Sciences. Food Safety and Environmental Stewardship Program; Available from: <http://fses.oregonstate.edu/methods>.
80. U.S. Environmental Protection Agency. *Technical Overview of Volatile Organic Compounds*. Available from: <https://www.epa.gov/indoor-air-quality-iaq/technical-overview-volatile-organic-compounds>.
81. Pavilonis, B.T., et al., *Bio-accessibility and Risk of Exposure to Metals and SVOCs in Artificial Turf Field Fill Materials and Fibers*. Risk analysis : an official publication of the Society for Risk Analysis, 2014. **34**(1): p. 44-55.



82. Llompарт, M., et al., *Hazardous organic chemicals in rubber recycled tire playgrounds and pavers*. Chemosphere, 2013. **90**(2): p. 423-431.
83. Perelo, L.W., *Review: In situ and bioremediation of organic pollutants in aquatic sediments*. J Hazard Mater, 2010. **177**(1-3): p. 81-9.
84. ChemRisk, I., *Review of the Human Health & Ecological Safety of Exposure to Recycled Tire Rubber found at Playgrounds and Synthetic Turf Fields*. 2008: Pittsburgh, PA.
85. California Office of Environmental Health Hazard Assessment (OEHHA). *Chemicals Known to the State to Cause Cancer or Reproductive Toxicity*. 2018; Available from: <https://oehha.ca.gov/media/downloads/proposition-65/p65list052518.pdf>.
86. Marsili, L.C., D; Bianchi, N; Maltese, S; Bianchi, M; Fossi, MC., *Release of Polycyclic Aromatic Hydrocarbons and Heavy Metals from Rubber Crumb in Synthetic Turf Fields: Preliminary Hazard Assessment for Athletes*. Journal of Environmental and Analytical Toxicology, 2014. **5**(2).
87. Strandberg, B., et al., *An improved method for determining dermal exposure to polycyclic aromatic hydrocarbons*. Chemosphere, 2018. **198**: p. 274-280.
88. Li, Z., et al., *Concentration and profile of 22 urinary polycyclic aromatic hydrocarbon metabolites in the US population*. Environmental Research, 2008. **107**(3): p. 320-331.
89. Perico, A., et al., *Assessment of Exposure to Polycyclic Aromatic Hydrocarbons in Police in Florence, Italy, through Personal Air Sampling and Biological Monitoring of the Urinary Metabolite 1-Hydroxypyrene*. Archives of Environmental Health: An International Journal, 2001. **56**(6): p. 506-512.
90. Genuis, S.J., et al., *Human Elimination of Phthalate Compounds: Blood, Urine, and Sweat (BUS) Study*. The Scientific World Journal, 2012. **2012**: p. 615068.
91. Adibi, J.J., et al., *Characterization of phthalate exposure among pregnant women assessed by repeat air and urine samples*. Environ Health Perspect, 2008. **116**(4): p. 467-73.
92. Gong, M., C.J. Weschler, and Y. Zhang, *Impact of Clothing on Dermal Exposure to Phthalates: Observations and Insights from Sampling Both Skin and Clothing*. Environmental Science & Technology, 2016. **50**(8): p. 4350-4357.
93. Gong, M., Y. Zhang, and C.J. Weschler, *Measurement of phthalates in skin wipes: estimating exposure from dermal absorption*. Environ Sci Technol, 2014. **48**(13): p. 7428-35.
94. Blount, B.C., et al., *Levels of seven urinary phthalate metabolites in a human reference population*. Environmental Health Perspectives, 2000. **108**(10): p. 979-982.



95. Högberg, J., et al., *Phthalate Diesters and Their Metabolites in Human Breast Milk, Blood or Serum, and Urine as Biomarkers of Exposure in Vulnerable Populations*. Environmental Health Perspectives, 2008. **116**(3): p. 334-339.
96. Ferguson, K.K., et al., *Urinary phthalate metabolites and biomarkers of oxidative stress in pregnant women: a repeated measures analysis*. Environ Health Perspect, 2015. **123**(3): p. 210-6.
97. Otake, T., J. Yoshinaga, and Y. Yanagisawa, *Exposure to phthalate esters from indoor environment*. J Expo Anal Environ Epidemiol, 2004. **14**(7): p. 524-8.
98. Koch, H.M., et al., *Exposure of nursery school children and their parents and teachers to di-n-butylphthalate and butylbenzylphthalate*. Int Arch Occup Environ Health, 2005. **78**(3): p. 223-9.
99. Johns, L.E., et al., *Exposure assessment issues in epidemiology studies of phthalates*. Environ Int, 2015. **85**: p. 27-39.
100. California Office of Environmental Health Hazard Assessment (OEHHA), *Evidence of the Carcinogenicity of Diisononyl Phthalate (DINP)*. 2013.
101. Lin, S., et al., *Phthalate exposure in pregnant women and their children in central Taiwan*. Chemosphere, 2011. **82**(7): p. 947-55.
102. California Office of Environmental Health Hazard Assessment (OEHHA). *Proposition 65 Maximum Allowable Dose Level (MADL) for Reproductive Toxicity for Di-isodecyl Phthalate (DIDP)*. 2010; Available from: <https://oehha.ca.gov/media/downloads/cnr/didpmadlfinalrisk042310.pdf>.
103. Silva, M.J., et al., *Oxidative metabolites of diisononyl phthalate as biomarkers for human exposure assessment*. Environ Health Perspect, 2006. **114**(8): p. 1158-61.
104. Agency for Toxic Substances and Disease Registry (ATSDR). *Toxicological Profile for di(2-ethylhexyl) phthalate*. 2002; Available from: <https://www.atsdr.cdc.gov/toxprofiles/tp9.pdf>.
105. Rudel, R.A., et al., *Phthalates, Alkylphenols, Pesticides, Polybrominated Diphenyl Ethers, and Other Endocrine-Disrupting Compounds in Indoor Air and Dust*. Environmental Science & Technology, 2003. **37**(20): p. 4543-4553.
106. Inoue, K., et al., *Measurement of 4-nonylphenol and 4-tert-octylphenol in human urine by column-switching liquid chromatography–mass spectrometry*. Analytica Chimica Acta, 2003. **486**(1): p. 41-50.
107. Hamelin, G., et al., *Determination of p-tert-octylphenol in blood and tissues by gas chromatography coupled with mass spectrometry*. J Anal Toxicol, 2008. **32**(4): p. 303-7.
108. He, G., B. Zhao, and M.S. Denison, *Identification of Benzothiazole Derivatives and Polycyclic Aromatic Hydrocarbons as Aryl Hydrocarbon Receptor Agonists Present in Tire Extracts*. Environmental Toxicology and Chemistry / Setac, 2011. **30**(8): p. 1915-1925.



109. Asimakopoulos, A.G., et al., *Benzotriazoles and benzothiazoles in human urine from several countries: a perspective on occurrence, biotransformation, and human exposure*. Environ Int, 2013. **59**: p. 274-81.
110. Asimakopoulos, A.G., et al., *Determination of benzotriazoles and benzothiazoles in human urine by liquid chromatography-tandem mass spectrometry*. Anal Chem, 2013. **85**(1): p. 441-8.
111. Reddy, C.M. and J.G. Quinn, *Environmental Chemistry of Benzothiazoles Derived from Rubber*. Environmental Science & Technology, 1997. **31**(10): p. 2847-2853.
112. Avagyan, R., et al., *Determination of benzothiazole and benzotriazole derivatives in tire and clothing textile samples by high performance liquid chromatography–electrospray ionization tandem mass spectrometry*. Journal of Chromatography A, 2013. **1307**: p. 119-125.
113. Lanigan, R.S. and T.A. Yamarik, *Final report on the safety assessment of BHT(1)*. Int J Toxicol, 2002. **21 Suppl 2**: p. 19-94.
114. *Butylated hydroxytoluene (BHT) [Biomonitoring Methods, 2010]*, in *The MAK-Collection for Occupational Health and Safety*.
115. Han-Na, S., et al., *Simultaneous determination of volatile organic compounds with a wide range of polarities in urine by headspace solid-phase microextraction coupled to gas chromatography/mass spectrometry*. Rapid Communications in Mass Spectrometry, 2017. **31**(7): p. 613-622.
116. Mattina, M., Isleyen, M., Berger, W., Ozdemir, S., *Examination of Crumb Rubber Produced from Recycled Tires*. The Connecticut Agricultural Experiment Station. Department of Analytical Chemistry: New Haven, Connecticut
117. *Phenols and Aromatic Alcohols (Phenol; 2- and 4-Methylphenol; DL-1- and 2-Phenylethanol; 3-Methylbenzyl alcohol; 2-Ethylphenol; 2,4-, 2,3- and 3,4-Dimethylphenol) [Biomonitoring Methods, 1988]*, in *The MAK-Collection for Occupational Health and Safety*.
118. Aylward, L.L., et al., *Screening-level Biomonitoring Equivalents for tiered interpretation of urinary 3-phenoxybenzoic acid (3-PBA) in a risk assessment context*. Regul Toxicol Pharmacol, 2018. **92**: p. 29-38.
119. Perico, A., et al., *Biological monitoring of occupational exposure to cyclohexane by urinary 1,2- and 1,4-cyclohexanediol determination*. Int Arch Occup Environ Health, 1999. **72**(2): p. 115-20.
120. Spiegelhalder, B. and R. Preussmann, *In vivo nitrosation of amidopyrine in humans: use of 'ethanol effect' for biological monitoring of N-nitrosodimethylamine in urine*. Carcinogenesis, 1985. **6**(4): p. 545-8.
121. Agency for Toxic Substances and Disease Registry (ATSDR). *Toxicological Profile for N-Nitrosodiphenylamine*. Available from: <https://www.atsdr.cdc.gov/toxprofiles/tp16.pdf>.



122. Agency for Toxic Substances and Disease Registry (ATSDR). *Toxicological Profile for Phenol*. 2008; Available from: <https://www.atsdr.cdc.gov/toxprofiles/tp115.pdf>.
123. Hotz, P., et al., *Biological monitoring of vehicle mechanics and other workers exposed to low concentrations of benzene*. Int Arch Occup Environ Health, 1997. **70**(1): p. 29-40.
124. Agency for Toxic Substances and Disease Registry (ATSDR). *Public Health Statement: Polycyclic Aromatic Hydrocarbons (PAHs)*. 1995; Available from: <https://www.atsdr.cdc.gov/ToxProfiles/tp69-c1-b.pdf>.
125. Li, Z., et al., *Excretion profiles and half-lives of ten urinary polycyclic aromatic hydrocarbon metabolites after dietary exposure*. Chem Res Toxicol, 2012. **25**(7): p. 1452-61.
126. Celeiro, M., T. Dagnac, and M. Llompарт, *Determination of priority and other hazardous substances in football fields of synthetic turf by gas chromatography-mass spectrometry: A health and environmental concern*. Chemosphere, 2018. **195**: p. 201-211.
127. Strandberg, B., et al., *Evaluation of polyurethane foam passive air sampler (PUF) as a tool for occupational PAH measurements*. Chemosphere, 2018. **190**: p. 35-42.
128. Canepari, S., et al., *Release of particles, organic compounds, and metals from crumb rubber used in synthetic turf under chemical and physical stress*. Environ Sci Pollut Res Int, 2018. **25**(2): p. 1448-1459.
129. Gaspar, F.W., et al., *Phthalate exposure and risk assessment in California child care facilities*. Environ Sci Technol, 2014. **48**(13): p. 7593-601.
130. Watkins, D.J., et al., *Variability and Predictors of Urinary Concentrations of Phthalate Metabolites during Early Childhood*. Environmental Science & Technology, 2014. **48**(15): p. 8881-8890.
131. Centers for Disease Control and Prevention, *Fourth National Report on Human Exposure to Environmental Chemicals, Updated Tables, January 2017*. 2009, U.S. Department of Health and Human Services
132. Parlett, L.E., A.M. Calafat, and S.H. Swan, *Women's exposure to phthalates in relation to use of personal care products*. Journal of exposure science & environmental epidemiology, 2013. **23**(2): p. 197-206.
133. California Office of Environmental Health Hazard Assessment (OEHHA). *Diisononyl Phthalate (DINP)*. Available from: <https://www.p65warnings.ca.gov/fact-sheets/diisononyl-phthalate-dinp>.
134. U.S. Environmental Protection Agency. *Bis(2-ethylhexyl) phthalate (DEHP)*. [cited 2018; Available from: <https://www.epa.gov/sites/production/files/2016-09/documents/bis-2-ethylhexyl-phthalate.pdf>.



135. Bocca, B., et al., *Metals contained and leached from rubber granulates used in synthetic turf areas*. Science of The Total Environment, 2009. **407**(7): p. 2183-2190.
136. Lewis, R.C., et al., *Urinary metal concentrations among mothers and children in a Mexico City birth cohort study*. International Journal of Hygiene and Environmental Health, 2018.
137. Mahaffey, K.R., et al., *Heavy metal exposure from foods*. Environmental Health Perspectives, 1975. **12**: p. 63-69.
138. Caruso, R.V., et al., *Toxic Metal Concentrations in Cigarettes Obtained from U.S. Smokers in 2009: Results from the International Tobacco Control (ITC) United States Survey Cohort*. International Journal of Environmental Research and Public Health, 2014. **11**(1): p. 202-217.
139. Nisse, C., et al., *Blood and urinary levels of metals and metalloids in the general adult population of Northern France: The IMEPOGE study, 2008–2010*. International Journal of Hygiene and Environmental Health, 2017. **220**(2, Part B): p. 341-363.
140. Pino, A., et al., *Human biomonitoring data analysis for metals in an Italian adolescents cohort: An exposome approach*. Environmental Research, 2017. **159**: p. 344-354.
141. Goen, T., et al., *Biological monitoring of exposure and effects in workers employed in a selenium-processing plant*. Int Arch Occup Environ Health, 2015. **88**(5): p. 623-30.
142. Agency for Toxic Substances and Disease Registry (ATSDR). *Public Health Statement Arsenic*. 2007; Available from: <https://www.atsdr.cdc.gov/ToxProfiles/tp2-c1-b.pdf>.
143. Agency for Toxic Substances and Disease Registry (ATSDR). *Analysis of hair samples: how to hair sampling results relate to environmental exposures?* [cited 2018; Available from: https://www.atsdr.cdc.gov/HAC/hair_analysis/03-0330HairSampleTesting-Scientific.pdf.
144. Sengupta, S. and H.J. Miller, *Preliminary investigation of tire shreds for use in residential subsurface leaching field systems*. 1999, Chelsea Center for Recycling and Economic Development, University of Massachusetts, Lowell.
145. O'Shaughnessy, V. and V.K. Garga, *Tire-reinforced earthfill. Part 3: Environmental assessment*. Canadian Geotechnical Journal, 2000. **37**(1): p. 117-131.
146. Riihimaki, V. and A. Aitio, *Occupational exposure to aluminum and its biomonitoring in perspective*. Crit Rev Toxicol, 2012. **42**(10): p. 827-53.
147. Mangold, J. and R. Simon. *Synthetic Turf Field Study*. 2016; Available from: http://www.landtekgroup.com/wp-content/uploads/2016/02/2016_Field-Turf-New-York-Seminar_Berkeley_Final_Public.pdf.



148. Agency for Toxic Substances and Disease Registry (ATSDR). *Public Health Statement for Antimony and Compounds*. [cited 2018; Available from: <https://www.atsdr.cdc.gov/toxprofiles/tp23-c1.pdf>.
149. Filella, M., N. Belzile, and Y.-W. Chen, *Human Exposure to Antimony. IV. Contents in Human Blood*. Critical Reviews in Environmental Science and Technology, 2013. **43**(19): p. 2071-2105.
150. California Office of Environmental Health Hazard Assessment (OEHHA), *Chemicals and particulates in the air above the new generation of artificial turf playing fields, and artificial turf as a risk factor for infection by methicillin-resistant Staphylococcus aureus (MRSA). Literature review and data gap identification*. 2009, California Environmental Protection Agency.
151. Centers for Disease Control and Prevention. *Arsenic Factsheet*. National Biomonitoring Program [cited 2018; Available from: https://www.cdc.gov/biomonitoring/Arsenic_FactSheet.html.
152. Cao, Z., et al., *Preliminary assessment on exposure of four typical populations to potentially toxic metals by means of skin wipes under the influence of haze pollution*. Science of The Total Environment, 2018. **613-614**: p. 886-893.
153. Rasmussen, P.E., et al., *Monitoring personal, indoor, and outdoor exposures to metals in airborne particulate matter: Risk of contamination during sampling, handling and analysis*. Atmospheric Environment, 2007. **41**(28): p. 5897-5907.
154. Occupational Safety and Health Administration (OSHA). *Arsenic, Cadmium, Cobalt, Copper, Lead, and Nickel (Open Vessel Microwave Digestion/ICP-MS Analysis)* [cited 2018; Available from: <https://www.osha.gov/dts/sltc/methods/mdt/mdt1006/1006.pdf>.
155. Lukasik-Glebocka, M., et al., *Barium determination in gastric contents, blood and urine by inductively coupled plasma mass spectrometry in the case of oral barium chloride poisoning*. J Anal Toxicol, 2014. **38**(6): p. 380-2.
156. Zhang, J., et al., *Hazardous chemicals in synthetic turf materials and their bioaccessibility in digestive fluids*. Journal Of Exposure Science And Environmental Epidemiology, 2008. **18**: p. 600.
157. Komaromy-Hiller, G., et al., *Comparison of representative ranges based on U.S. patient population and literature reference intervals for urinary trace elements*. Clinica Chimica Acta, 2000. **296**(1): p. 71-90.
158. Julander, A., et al., *Skin deposition of nickel, cobalt, and chromium in production of gas turbines and space propulsion components*. Ann Occup Hyg, 2010. **54**(3): p. 340-50.
159. Maria, K., et al., *Biological monitoring of dermal and air exposure to cobalt at a Swedish hard metal production plant: does dermal exposure contribute to uptake?* Contact Dermatitis, 2017. **77**(4): p. 201-207.



160. Yantasee, W., et al., *Electrochemical Sensors for the Detection of Lead and Other Toxic Heavy Metals: The Next Generation of Personal Exposure Biomonitorers*. Environmental Health Perspectives, 2007. **115**(12): p. 1683-1690.
161. California Office of Environmental Health Hazard Assessment (OEHHA), *Public Health Goal for Aluminum in Drinking Water*. 2001, California Protection Agency.
162. California Office of Environmental Health Hazard Assessment (OEHHA), *Public Health Goal for Antimony in Drinking Water*. 2016, California Environmental Protection Agency.
163. Kentner, M., et al., *External and internal antimony exposure in starter battery production*. Int Arch Occup Environ Health, 1995. **67**(2): p. 119-23.
164. California Office of Environmental Health Hazard Assessment (OEHHA). *Arsenic*. [cited 2018; Available from: <https://oehha.ca.gov/chemicals/arsenic>.
165. Kearny Foundation of Soil Science Division of Agriculture and Natural Resources University of California, *Background Concentrations of Trace and Major Elements in California Soils* 1996.
166. California Office of Environmental Health Hazard Assessment (OEHHA), *Public Health Goals for Chemicals in Drinking Water - Barium*. 2003.
167. New Jersey Department of Health and Senior Services. *Hazardous Substance Fact Sheet: Barium Oxide*. 2002 [cited 2018; Available from: <http://nj.gov/health/eoh/rtkweb/documents/fs/0187.pdf>.
168. Almaguer, D., et al., *Control Technology and Exposure Assessment for Electronic Recycling Operations File No: EPHB 326-12a*. 2008, U.S. Department of Health and Human Services, Centers for Disease Control and Prevention: Cincinnati, Ohio.
169. California Office of Environmental Health Hazard Assessment (OEHHA). *Cadmium Factsheet*. [cited 2018; Available from: <https://oehha.ca.gov/media/downloads/faqs/cadmiumfacts.pdf>.
170. State of California-Office of Environmental Health Hazard Assessment. *Evaluation of Health Effects of Recycled Waste Tires in Playground and Track Products* 2017; Available from: <http://www.calrecycle.ca.gov/Publications/Documents/1206/62206013.pdf>.
171. California Office of Environmental Health Hazard Assessment (OEHHA). *Chromium-hexavalent*. [cited 2018; Available from: <https://oehha.ca.gov/chemicals/chromium-hexavalent>.
172. U.S. Environmental Protection Agency. *Cobalt Compounds*. [cited 2018; Available from: <https://www.epa.gov/sites/production/files/2016-09/documents/cobalt-compounds.pdf>.
173. Agency for Toxic Substances and Disease Registry (ATSDR). *Toxicological Profile for Cobalt*. 2004; Available from: <https://www.atsdr.cdc.gov/toxprofiles/tp33.pdf>.



174. California Office of Environmental Health Hazard Assessment (OEHHA), *Copper in Drinking Water California Public Health Goal*. 2008.
175. Li, G.J., et al., *Occupational Exposure to Welding Fume among Welders: Alterations of Manganese, Iron, Zinc, Copper, and Lead in Body Fluids and the Oxidative Stress Status*. Journal of occupational and environmental medicine / American College of Occupational and Environmental Medicine, 2004. **46**(3): p. 241-248.
176. Chillrud, S.N., et al., *Steel dust in the New York City subway system as a source of manganese, chromium, and iron exposures for transit workers*. J Urban Health, 2005. **82**(1): p. 33-42.
177. Saravanabhavan, G., et al., *Human biomonitoring reference values for metals and trace elements in blood and urine derived from the Canadian Health Measures Survey 2007–2013*. International Journal of Hygiene and Environmental Health, 2017. **220**(2, Part A): p. 189-200.
178. California Office of Environmental Health Hazard Assessment (OEHHA), *Lead in Drinking Water California Public Health Goal*. 2009.
179. California Office of Environmental Health Hazard Assessment (OEHHA). *OEHHA Acute, 8-hour and Chronic Reference Exposure Level (REL) Summary*. [cited 2018; Available from: <https://oehha.ca.gov/media/downloads/crn/appendixd1final.pdf>].
180. Gunier, R.B., et al., *Biomarkers of manganese exposure in pregnant women and children living in an agricultural community in California*. Environ Sci Technol, 2014. **48**(24): p. 14695-702.
181. California Office of Environmental Health Hazard Assessment (OEHHA). *Mercury and Mercury Compounds Proposition 65 Warnings 2016* [cited 2018; Available from: <https://oehha.ca.gov/media/downloads/proposition-65/chemicals/mercurymethylmercurycompfactsheet.pdf>].
182. Beaucham, C., D. Ceballos, and B. King, *Lessons learned from surface wipe sampling for lead in three workplaces*. Journal of Occupational and Environmental Hygiene, 2017. **14**(8): p. 609-617.
183. California Office of Environmental Health Hazard Assessment (OEHHA). *Selenium*. Available from: <https://oehha.ca.gov/chemicals/selenium>.
184. U.S. Environmental Protection Agency, *Toxicological Review of Zinc and Zinc Compounds (CAS No 7440-66-6) In Support of Summary Information on the Integrated Risk Information System (IRIS)*. 2005: Washington, D.C.
185. Johnson, C.H., et al., *Xenobiotic metabolomics: major impact on the metabolome*. Annu Rev Pharmacol Toxicol, 2012. **52**: p. 37-56.
186. Lankadurai, B.P., E.G. Nagato, and M.J. Simpson, *Environmental metabolomics: an emerging approach to study organism responses to environmental stressors*. Environmental Reviews, 2013. **21**(3): p. 180-205.



187. Rappaport, S.M. and M.T. Smith, *Epidemiology. Environment and disease risks*. Science, 2010. **330**(6003): p. 460-1.
188. Wild, C.P., *The exposome: from concept to utility*. International Journal of Epidemiology, 2012. **41**(1): p. 24-32.
189. Nicholson, J.K., J.C. Lindon, and E. Holmes, *'Metabonomics': understanding the metabolic responses of living systems to pathophysiological stimuli via multivariate statistical analysis of biological NMR spectroscopic data*. Xenobiotica, 1999. **29**(11): p. 1181-9.
190. van Ravenzwaay, B., et al., *The use of metabolomics for the discovery of new biomarkers of effect*. Toxicol Lett, 2007. **172**(1-2): p. 21-8.
191. Zhou, M., Ford B., Lee, D., Huen, K., Tran, V., Bradman, A., Gunier, R., Eskenazi, B., Nomura, D., Holland, N., *Metabolomic Markers of Phthalate Exposure in Plasma and Urine of Pregnant Women*. Environmental Health, 2018(Submitted to a Special Issue "Exposure to Endocrine-Disrupting Chemicals and Cardiometabolic Disease: A Developmental Origins Approach" in Frontiers in Public Health).
192. Zhang, J., et al., *Low-Level Environmental Phthalate Exposure Associates with Urine Metabolome Alteration in a Chinese Male Cohort*. Environ Sci Technol, 2016. **50**(11): p. 5953-60.
193. Bradman, A., et al., *Community-Based Intervention to Reduce Pesticide Exposure to Farmworkers and Potential Take-Home Exposure to their Families*. Journal of exposure science & environmental epidemiology, 2009. **19**(1): p. 79-89.
194. O'Hara, M.E., et al., *Development of a protocol to measure volatile organic compounds in human breath: a comparison of rebreathing and on-line single exhalations using proton transfer reaction mass spectrometry*. Physiol Meas, 2008. **29**(3): p. 309-30.
195. Miekisch, W., J.K. Schubert, and G.F.E. Noeldge-Schomburg, *Diagnostic potential of breath analysis—focus on volatile organic compounds*. Clinica Chimica Acta, 2004. **347**(1): p. 25-39.
196. Lindstrom, A.B. and J.D. Pleil, *A review of the USEPA's single breath canister (SBC) method for exhaled volatile organic biomarkers*. Biomarkers, 2002. **7**(3): p. 189-208.
197. Supelco. *Raidello Diffusive Sampling System*. Available from: [https://www.sigmaaldrich.com/content/dam/sigma-aldrich/docs/Sigma/General Information/radiello bro 2nd ed 2007.pdf](https://www.sigmaaldrich.com/content/dam/sigma-aldrich/docs/Sigma/General%20Information/radiello_bro_2nd_ed_2007.pdf).