



# Health Advisory and Guidelines for Eating Fish from Big Bear Lake (San Bernardino County)

January 2018



Fish, Ecotoxicology, and Water Section  
Pesticide and Environmental Toxicology Branch  
Office of Environmental Health Hazard Assessment  
California Environmental Protection Agency

# LIST OF CONTRIBUTORS

## ***Office of Environmental Health Hazard Assessment***

### ***Authors***

Lori Chumney, M.S.  
Huyen Tran Pham, M.P.H.

### ***Primary Reviewers***

Susan A. Klasing, Ph.D., Section Chief  
Wesley Smith, Ph.D.  
Shannon R. Murphy, Ph.D.

### ***Final Reviewers***

David Ting, Ph.D., Branch Chief  
David Siegel, Ph.D., Assistant to the Deputy Director  
Allan Hirsch, Chief Deputy Director

### ***Director***

Lauren Zeise, Ph.D.

## ACKNOWLEDGMENTS

Developing fish consumption advisories depends on sampling and analysis of fish. The Office of Environmental Health Hazard Assessment (OEHHA) acknowledges the contribution of information from the following entities: the State Water Resources Control Board (SWRCB), the California Department of Fish and Wildlife (CDFW) and its analytical resources, the Moss Landing Marine Laboratories and the Water Pollution Control Laboratory. Data were obtained from the California Environmental Data Exchange Network (<http://www.ceden.us/AdvancedQueryTool>). The map was created using ArcMap (10.3) from Environmental Systems Resource Institute (ESRI, Redlands, California).

### ***For further information, contact:***

Pesticide and Environmental Toxicology Branch  
Office of Environmental Health Hazard Assessment  
California Environmental Protection Agency

1515 Clay Street, 16<sup>th</sup> Floor  
Oakland, California 94612  
Telephone: (510) 622-3170  
Email address: [fish@oehha.ca.gov](mailto:fish@oehha.ca.gov)

1001 I Street, P.O. Box 4010  
Sacramento, CA 95812-4010  
Telephone: (916) 324-7572

Big Bear Lake Fish Advisory

## LIST OF ACRONYMS AND ABBREVIATIONS

ATL	Advisory Tissue Level
CDFW	California Department of Fish and Wildlife
DDT(s)	dichlorodiphenyltrichloroethane (DDT) and its metabolites dichlorodiphenyldichloroethane (DDD) and dichlorodiphenyldichloroethylene (DDE)
DHA	docosahexaenoic acid
EPA	eicosapentaenoic acid
FDA	Food and Drug Administration
Hg	mercury
MDL	method detection limit
MLML	Moss Landing Marine Laboratories
mm	millimeters
OEHHA	Office of Environmental Health Hazard Assessment
PBDEs	polybrominated diphenyl ethers
PCBs	polychlorinated biphenyls
ppb	parts per billion
RL	reporting limit
RWB8	Regional Water Board 8 (Santa Ana Region)
Se	selenium
SWAMP	Surface Water Ambient Monitoring Program
TSMF	Toxic Substances Monitoring Program
USDA	United States Department of Agriculture

USDHHS                    United States Department of Health and Human Services

US EPA                    United States Environmental Protection Agency

## PREFACE

The Office of Environmental Health Hazard Assessment (OEHHA), a department in the California Environmental Protection Agency, is responsible for evaluating potential public health risks from chemical contamination of sport fish. This includes issuing fish consumption advisories, when appropriate, for the State of California. OEHHA's authorities to conduct these activities are based on mandates in the:

- California Health and Safety Code
  - Section 59009, to protect public health
  - Section 59011, to advise local health authorities
- California Water Code
  - Section 13177.5, to issue health advisories

The health advisories are published in the California Department of Fish and Wildlife Sport Fishing Regulations in the section on public health advisories.

This report presents guidelines for eating fish from Big Bear Lake in San Bernardino County. The report provides background information and a technical description of how the guidelines were developed. The resulting advice is summarized in the pictorial after the Table of Contents and List of Figures and Tables.

## TABLE OF CONTENTS

A GUIDE TO EATING FISH FROM BIG BEAR LAKE .....	6
INTRODUCTION .....	7
<i>Location</i> .....	7
<i>Approach Used</i> .....	8
CHEMICALS OF POTENTIAL CONCERN.....	8
DATA SOURCES.....	10
<i>Toxic Substances Monitoring Program (TSMP)</i> .....	10
<i>Surface Water Ambient Monitoring Program (SWAMP): Contaminants in Fish From California Lakes and Reservoirs, 2007-2008</i> .....	10
<i>Regional Water Quality Control Board, Santa Ana Fish Contamination Study (RWB8)</i> .....	10
FISH SAMPLED FROM BIG BEAR LAKE.....	11
CHEMICAL CONCENTRATIONS .....	12
<i>Mercury</i> .....	12
<i>PCBs, PBDEs, and Pesticides</i> .....	12
<i>Selenium</i> .....	13
DEVELOPMENT OF GUIDELINES FOR EATING FISH FROM BIG BEAR LAKE .....	15
CONSUMPTION ADVICE FOR FISH FROM BIG BEAR LAKE .....	17
<i>Black Bass Species (Largemouth and Smallmouth)</i> .....	17
<i>Common Carp</i> .....	17
<i>Crappie</i> .....	18
<i>Rainbow Trout</i> .....	18
RECOMMENDED MAXIMUM NUMBER OF SERVINGS.....	19
REFERENCES .....	20
Big Bear Lake Fish Advisory .....	4

APPENDIX I. Advisory Tissue Levels.....23

## LIST OF FIGURES AND TABLES

Figure 1. Location of Big Bear Lake ..... 7


Table 1. Fish Samples Evaluated for the Big Bear Lake Advisory ..... 11

Table 2. Mercury Concentrations in Fish from Big Bear Lake ..... 14

Table 3. PCB Concentrations in Fish from Big Bear Lake..... 14

Table 4. Recommended Maximum Number of Servings per Week for Fish from Big Bear Lake..... 19

Advisory Tissue Levels for Selected Analytes..... 23



**Women**  
(18-45 Years)

**Children**  
(1-17 Years)

**5** TOTAL SERVINGS A WEEK


OR

**5** TOTAL SERVINGS A WEEK

OR

**1** TOTAL SERVING A WEEK

**Serving Size**  
A serving of fish is about the size and thickness of your hand. Give children smaller servings.



**Women**  
(46+ Years)

**Men**  
(18+ Years)



**7** TOTAL SERVINGS A WEEK

OR

**5** TOTAL SERVINGS A WEEK

OR

**2** TOTAL SERVINGS A WEEK


**For Adults**  **For Children** 

## A GUIDE TO EATING FISH


*from*  
**BIG BEAR LAKE**

(SAN BERNARDINO COUNTY)


**Eat the Good Fish**  
Eating fish that are low in chemicals may provide health benefits to children and adults.



**Avoid the Bad Fish**  
Eating fish with higher levels of chemicals like mercury or PCBs may cause health problems in children and adults.




**Choose the Right Fish**  
Chemicals may be more harmful to unborn babies and children.




**Crappie**

---




**Rainbow Trout**  
♥ *high in omega-3s*


---



**Common Carp**




**Black Bass Species**  
♥ *high in omega-3s*




**California Office of Environmental Health Hazard Assessment**

web [www.oehha.ca.gov/fish](http://www.oehha.ca.gov/fish)  
email [fish@oehha.ca.gov](mailto:fish@oehha.ca.gov)  
phone (916) 324-7572

**Eat only the skinless fillet**



**Eat only the meat**



Some chemicals are higher in the skin, fat, and guts.




## INTRODUCTION

This report presents a guideline for eating fish from Big Bear Lake (Figure 1), in San Bernardino County, CA. Big Bear Lake is located about 25 miles northeast of the city of San Bernardino, and can be accessed by state highways 38 and 18.

### LOCATION

Big Bear Lake is a snow-fed reservoir in the San Bernardino National Forest. It is a popular high-elevation lake that provides 22 miles of shoreline and stores 73,730 acre feet of water. The Big Bear Municipal Water District manages the lake with the primary goal of stabilizing water levels for recreational, economic, and environmental purposes.<sup>1</sup>

FIGURE 1. LOCATION OF BIG BEAR LAKE



<sup>1</sup> Information regarding Big Bear Lake was obtained from the Big Bear Municipal Water District. Online at: <http://www.bbmwd.com/current-lake-status/>

### *APPROACH USED*

The Office of Environmental Health Hazard Assessment (OEHHA) used the results from three monitoring studies described in this report to develop the Big Bear Lake Advisory. OEHHA uses the following general process in developing consumption advice for sport fish:

- 1) Evaluation of all fish contaminant data available from a water body and selection of appropriate data that meet data quality criteria and sampling plan guidelines.
- 2) Determination of fish species for which adequate data are available to issue fish consumption advice.
- 3) Calculation of an appropriate measure of central tendency (often a weighted arithmetic mean<sup>2</sup>) and other descriptive statistics of the contaminant data, as appropriate, for a chemical of potential concern for the selected fish species.
- 4) Comparison of the chemical concentrations with the OEHHA Advisory Tissue Levels (ATLs) for each chemical of potential concern.
- 5) Development of final advice based on a thorough review of the data and best professional judgment relating to the benefits and risks of consuming a particular fish species.

The ATLs (discussed further in a subsequent section of this report) are chemical levels in fish tissue that are considered acceptable, based on chemical toxicity, for a range of consumption rates. Development of the ATLs also included consideration of health benefits associated with including fish in the diet (OEHHA, 2008). The ATLs should not be interpreted as static “bright lines”, but one component of a complex process of data evaluation and interpretation used by OEHHA in the assessment and communication of the benefits and risks of consuming sport fish.

## CHEMICALS OF POTENTIAL CONCERN

Certain chemicals are considered to be of potential concern for people who eat fish because of their toxicity and their ability to accumulate in fish tissue. The majority of fish consumption advisories in California are issued because of mercury (Hg), followed by polychlorinated biphenyls (PCBs) and, in a few cases, selenium (Se) or some legacy pesticides (pesticides that are no longer used but remain in the environment).

Mercury is a natural element found in some rock and soil. Human activities, such as burning coal and the historic use of mercury to mine gold, also add mercury to the environment. If mercury enters waterways, it can be converted to a more toxic form

---

<sup>2</sup> Means are an arithmetic average of individual values and/or a weighted average of composites. A weighted average of composites is calculated by multiplying the chemical concentration in each composite by the number of fish in that composite for each species. Products are then summed and divided by the total number of fish in all composites for that species, combined.

known as methylmercury – which can pass into and build up in fish. High levels of methylmercury can harm the brain, especially in fetuses and children.

PCBs are industrial chemicals previously used in electrical transformers, plastics, and lubricating oils, often as flame retardants or electrical insulators. Their use was banned in the 1970s, but they persist in the environment because they do not break down easily and can accumulate in fish. Depending on the exposure level, PCBs may cause cancer or other health effects, including neurotoxicity, in humans.

Selenium is a naturally occurring metalloid and at low doses is an essential nutrient for many important human health processes, including thyroid regulation and vitamin C metabolism. Higher doses cause selenium toxicity, which can include symptoms ranging from hair loss and gastrointestinal distress to dizziness and tremors.

Chlordanes, dichlorodiphenyltrichloroethane (DDT), dieldrin, and toxaphene are pesticides that were banned from use in 1973 (DDT), the late 1980s (chlordanes and dieldrin) and 1990 (toxaphene), but are still found in some fish in certain water bodies in California. Depending on the exposure level, these chemicals may cause cancer or adverse effects on the nervous system.

Polybrominated diphenyl ethers (PBDEs) are a class of flame retardants historically used in a variety of consumer products including furniture, textiles, automotive parts, and electronics. The use of PBDEs in new products was largely phased out by 2013 but, due to their wide usage and persistence in the environment, they are still being detected in fish tissues. PBDEs may affect hormone levels or learning and behavior in children.

Detailed discussion of the toxicity of these chemicals and references are presented in “Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Chlordane, DDTs, dieldrin, methylmercury, PCBs, selenium, and toxaphene” (OEHHA, 2008) and “Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Polybrominated Diphenyl Ethers (PBDEs)” (OEHHA, 2011).

All fish species or specie groups collected from Big Bear Lake and used in advisory development were analyzed for mercury (as a measure of methylmercury) as well as the legacy pesticides chlordanes, dieldrin, DDTs (DDT and its metabolites dichlorodiphenyldichloroethane [DDD] and dichlorodiphenyldichloroethylene [DDE]), PCBs, and selenium. Common Carp and Largemouth Bass were additionally analyzed for the legacy pesticide toxaphene, and Common Carp were further analyzed for PBDEs. Among the chemicals analyzed in fish tissue samples from Big Bear Lake, only mercury and PCB levels were sufficiently high to impact consumption advice. For this reason, levels of other contaminants are not shown in this report.

## DATA SOURCES

The guidelines for eating fish from Big Bear Lake are based on the chemicals detected in the fish collected for the three monitoring studies described below. These studies met OEHHA's data quality criteria, including adequate documentation of sample collection, fish preparation method (e.g., skinning or filleting), chemical analyses, quality assurance, and sufficiently low detection limits. "Sample," as used in this report, refers to an individual fish or a composite of multiple fish for which contaminant data was reported. "Sampling" or "sampled" refers to the act of collecting fish for chemical analysis.

### *TOXIC SUBSTANCES MONITORING PROGRAM (TSMP)*

The TSMP (1976-2003) was a state water quality-monitoring program managed by the State Water Resources Control Board (SWRCB, 2007 and 2013). Its objective was to provide statewide information on the occurrence of toxic substances by monitoring water bodies with known or suspected water quality impairment. Staff from the California Department of Fish and Wildlife (CDFW), then known as the California Department of Fish and Game, collected Common Carp, Largemouth Bass, and Smallmouth Bass from Big Bear Lake between 1984 and 2001, as part of the program. Common Carp and Largemouth Bass were analyzed for chlordanes, DDTs, dieldrin, mercury, selenium, and toxaphene. Smallmouth Bass were analyzed for mercury.

### *SURFACE WATER AMBIENT MONITORING PROGRAM (SWAMP): CONTAMINANTS IN FISH FROM CALIFORNIA LAKES AND RESERVOIRS, 2007- 2008*

SWAMP, operated by the SWRCB in cooperation with Santa Ana Regional Water Quality Control Board Region 8 (RWB8) staff, monitors water quality in California's surface waters. This study was part of a SWAMP statewide sampling effort to survey contaminants in sport fish found in California lakes and reservoirs (SWRCB, 2010). The program collected Common Carp from Big Bear Lake in 2007, and analyzed fish tissue for chlordanes, DDTs, dieldrin, mercury, PBDEs, PCBs, and selenium.

### *REGIONAL WATER QUALITY CONTROL BOARD, SANTA ANA FISH CONTAMINATION STUDY (RWB8)*

The RWB8 coordinates ongoing sampling efforts to monitor contaminant levels, including mercury, in sport fish caught from lakes and reservoirs within the region. RWB8 collected Common Carp, Crappie, Largemouth Bass, and Rainbow Trout from Big Bear Lake in 2004-2005 and analyzed fish tissue for chlordanes, DDTs, dieldrin, mercury, PCBs, and selenium.

## FISH SAMPLED FROM BIG BEAR LAKE

The fish sampling data used in this advisory were retrieved from the California Environmental Data Exchange Network (CEDEN), the State's repository for environmental data. Samples were excluded when the fish were not legal size to take or did not meet OEHHA's criteria for minimum "edible" size based on species size at maturity, and professional judgment (as described in OEHHA, 2005). A summary of all fish species evaluated for this advisory is shown in Table 1, including the name of the species, number of samples collected, total number of fish, project name, year sampled, and contaminants analyzed.

TABLE 1. FISH SAMPLES EVALUATED FOR THE BIG BEAR LAKE ADVISORY

Common Name	Scientific Name	Number of Samples	Total Number of Fish	Project	Year Collected	Contaminants Analyzed <sup>b</sup>
Common Carp	<i>Cyprinus carpio</i>	3	18	TSMP <sup>a</sup>	2000-2001	Chlordanes, DDTs, Dieldrin, Hg, Se, Toxaphene
		1	4	TSMP <sup>a</sup>	1984	Hg
		1	6	TSMP <sup>a</sup>	1988	Hg, Se
		3	15	SWAMP	2007	Chlordanes, DDTs, Dieldrin, Hg, PBDEs, PCBs
		1	15	SWAMP	2007	Se
		6	23	RWB8	2004-2005	Chlordanes, DDTs, Dieldrin, Hg, PCBs, Se
Crappie	<i>Pomoxis spp.</i>	3	9	RWB8	2005	Chlordanes, DDTs, Dieldrin, Hg, PCBs, Se
Largemouth Bass	<i>Micropterus salmoides</i>	1	6	TSMP <sup>a</sup>	2001	Chlordanes, DDTs, Dieldrin, Hg, Se, Toxaphene
		1	6	TSMP <sup>a</sup>	1984	Hg
		5	20	RWB8	2004-2005	Chlordanes, DDTs, Dieldrin, Hg, PCBs, Se
Rainbow Trout	<i>Oncorhynchus mykiss</i>	5	15	RWB8	2004	Chlordanes, DDTs, Dieldrin, Hg, PCBs, Se
Smallmouth Bass	<i>Micropterus dolomieu</i>	1	5	TSMP <sup>a</sup>	1989	Hg

<sup>a</sup>Study report did not specify whether skin was removed from fillets prior to tissue analysis.

<sup>b</sup>Organic data (chlordanes, DDTs, dieldrin, PCBs or toxaphene) generated prior to 2000 were excluded from the analysis because more recent data are considered more reliable because of improved analytical methods.

## CHEMICAL CONCENTRATIONS

As shown in Table 1, samples were analyzed for one or more of the following: total mercury, selenium, chlordanes, DDTs, dieldrin, PBDEs, PCBs (54-55 congeners<sup>3</sup>), and toxaphene. All fish samples were prepared as skinless fillets, except for the TSMP study where the fillet preparation method for Common Carp, Largemouth Bass, and Smallmouth Bass was not recorded. Samples were analyzed as individual fish or composites.

Composites were prepared from equal amounts of tissue from several similarly sized individual fish of a species. Ideally, for composite samples, the total length of the smallest fish in a composite sample is at least 75% of the length of the largest fish in the sample (US EPA, 2000a). This information is not available for samples collected in the TSMP program; however, OEHHA assumes that the data complies with the 75% rule. All other composite samples from Big Bear Lake met this requirement.

For this advisory, OEHHA used the weighted (by the number of individual fish) arithmetic mean (average) of the chemical concentrations (in wet weight) for each fish species to estimate average human exposure.

### *MERCURY*

Samples were analyzed for total mercury, as either individual fish or composite samples, using a direct mercury analyzer (DMA) at the CDFW Moss Landing Marine Laboratories (MLML). The DMA method utilizes thermal decomposition and atomic absorption. OEHHA assumed all mercury detected was methylmercury, which is the most common form found in fish and is also the more toxic form (Bloom, 1992). Table 2 shows the averages and ranges for total length<sup>4</sup>, as well as mercury concentrations in each fish species. The DMA method detection limit (MDL)<sup>5</sup> and the reporting limit (RL)<sup>6</sup> for total mercury were reported at 9 or 12 and 12 or 26 parts per billion (ppb), respectively. Although mercury was detected at commonly found concentrations in the TSMP study, the MDL and RL for mercury were not reported.

### *PCBs, PBDEs, AND PESTICIDES*

Some composite samples were analyzed for PCBs, PBDEs, and the legacy pesticides (chlordanes, DDTs, dieldrin, and toxaphene). Pesticides, PBDEs and PCBs were analyzed by gas chromatography at the CDFW Water Pollution Control Laboratory. For

---

<sup>3</sup> Congeners are related compounds with similar chemical forms. Of the 209 possible PCB congeners, 54-55 are generally reported.

<sup>4</sup> Total length is the maximum length of the fish, measured from the tip of the closed mouth to the tip of the pinched tail fin.

<sup>5</sup> The MDL is the lowest quantity of a chemical that can be distinguished (as greater than zero) in a sample.

<sup>6</sup> The RL is the lowest quantity of a chemical that can be accurately quantified in a sample.

chlordanes, DDTs, PCBs, and PBDEs, each of the concentrations presented was the sum of the detected parent compound, congeners, or metabolites, where applicable. Since the MDLs or RLs were relatively low ( $\leq 5$  ppb), individual congeners or metabolites with concentrations reported as non-detects were assumed to be zero. This is a standard method of handling non-detect values for PCBs and other chemicals with multiple congeners or metabolites in a given sample when detection levels are adequate (US EPA, 2000a). Table 3 shows the averages and ranges for total length, as well as PCB concentrations in each fish species. Concentrations of chlordanes, dieldrin, DDTs, PBDEs, and toxaphene were not sufficiently high to alter consumption advice and are not shown.

#### *SELENIUM*

Common Carp, Crappie, Largemouth Bass, and Rainbow Trout were analyzed for selenium using Inductively Coupled Plasma-Mass Spectrometry with the MDL and RL reported at 40 and 120 ppb, respectively. The selenium concentration was not sufficiently high to alter consumption advice and is not shown.

TABLE 2. MERCURY CONCENTRATIONS IN FISH FROM BIG BEAR LAKE

Species from Big Bear Lake	Number of Samples	Total Number of Fish	Mean* Total Length (mm)	Range of Total Lengths** (mm)	Mercury (ppb)	
					Mean*	Range**
Common Carp	14	66	468	331 - 638	151	32 - 248
Crappie	3	9	203	179 - 239	41	38 - 44
Rainbow Trout	5	15	390	305 - 490	16	13 - 21
Black Bass species***	8	37	401	353 - 466	344	218 - 600
Largemouth Bass	7	32	408	370 - 466	354	218 - 600
Smallmouth Bass	1	5	353	n/a	280	n/a

\*Means are an arithmetic average of individual values and/or a weighted average of composites.

\*\*Range of individuals and/or range of the composites.

\*\*\* Largemouth and Smallmouth bass were combined for "Black Bass species" for the purpose of developing consumption advice.

n/a = not applicable due to a single sample.

TABLE 3. PCB CONCENTRATIONS IN FISH FROM BIG BEAR LAKE

Species from Big Bear Lake	Number of Samples	Total Number of Fish	Mean* Total Length (mm)	Range of Total Lengths** (mm)	PCB (ppb)	
					Mean*	Range**
Common Carp	9	38	498	331 - 638	38	5 - 104
Crappie	3	9	203	179 - 239	3	2 - 4
Largemouth Bass	5	20	427	380 - 466	13	8 - 30
Rainbow Trout	5	15	390	305 - 490	12	9 - 17

\*Means are an arithmetic average of individual values and/or a weighted average of composites.

\*\*Range of individuals and/or range of the composites.



## DEVELOPMENT OF GUIDELINES FOR EATING FISH FROM BIG BEAR LAKE

The OEHHA fish advisory process considers the health benefits of fish consumption as well as the risk from exposure to the chemical contaminants found in fish. Benefits are included in the advisory process because there is considerable evidence and scientific consensus that fish should be part of a healthy, well-balanced diet. Fish contain many nutrients that are important for general health and, in particular, help promote optimal growth and development of babies and young children, and may reduce the incidence of heart disease in adults (FDA/US EPA, 2017; American Heart Association, 2014; OEHHA, 2008; Institute of Medicine, 2007; Kris-Etherton et al., 2002). Fish are a significant source of the specific omega-3 fatty acids, docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA), thought to be associated with these beneficial health effects (USDA/USDHHS, 2015; Weaver et al., 2008).

The 2015-2020 U.S. Dietary Guidelines recommend that 1) the general population “consume eight or more ounces per week (less for young children)” of a variety of seafood<sup>7</sup> “for the total package of nutrients that seafood provides, including its EPA and DHA content” and 2) “women who are pregnant or breastfeeding should consume at least eight and up to twelve ounces of a variety of seafood per week from choices that are lower in methylmercury” (USDA/USDHHS, 2015). The particular fish that people eat is an important factor in determining the net beneficial effects of fish consumption. For example, studies have shown that children of mothers who ate low-mercury fish during pregnancy scored better on cognitive tests compared to children of mothers who did not eat fish or ate high-mercury fish (Oken et al., 2005 and 2008). Accordingly, because of the high mercury content of certain fish species, the US Food and Drug Administration (FDA) and the US Environmental Protection Agency (US EPA) recommend that women who are pregnant (or might become pregnant) or breastfeeding, and young children avoid consuming shark, swordfish, tilefish (Gulf of Mexico), bigeye tuna, marlin, orange roughy, and king mackerel (FDA/US EPA, 2017).

In order to address the potential health concerns associated with exposure to contaminants in sport fish, OEHHA has established ATLS for chemicals that are known to accumulate in the edible tissues of fish. ATLS consider both the toxicity of the chemical and potential benefits of eating fish. OEHHA uses the ATLS to determine the maximum number of servings per week that consumers can eat, for each species and at each location, to limit their exposure to these contaminants. Consumers can use OEHHA’s guidance when choosing which fish and how much to eat as part of an overall healthy diet.

---

<sup>7</sup> “Marine animals that live in the sea and in freshwater lakes and rivers. Seafood includes fish, such as salmon, tuna, trout, and tilapia, and shellfish, such as shrimp, crab, and oysters” (USDHHS/USDA, 2015).

There are two sets of ATLS for methylmercury in fish because of the age-related toxicity of this chemical (OEHHA, 2008). The fetus and children are more sensitive to the toxic effects of methylmercury. Thus, the ATLS for the sensitive population, including women who might become pregnant (typically 18 to 45 years of age) and children 1-17 years, are lower than those for women 46 years and older, and men 18 years and older. The lower ATL values for the sensitive population provide additional protection to allow for normal growth and development of the brain and nervous system of unborn babies and children. Detailed discussion about the toxicity of common fish contaminants and health benefits of fish consumption, as well as derivation of the ATLS, are provided in “Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Chlordane, DDTs, dieldrin, methylmercury, PCBs, selenium, and toxaphene” (OEHHA, 2008) and “Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Polybrominated Diphenyl Ethers (PBDEs)” (OEHHA, 2011). A list of the ATLS used in this report is presented in Appendix I.

For each fish species in this advisory, OEHHA compared the mean mercury and PCB concentrations detected in the fillet to the corresponding ATLS to establish the maximum number of servings per week that could be consumed (see Appendix I).

The consumption advice for a fish species is initially based on the chemical with the lowest allowable number of servings per week. Because some chemicals, such as mercury and PCBs, are known to have similar adverse effects, additivity of toxicity is assumed in such cases and may be assessed using multiple chemical exposure methodology (US EPA, 1989 and 2000b). If two or more chemicals with similar adverse effects are present in fish tissue at levels above the corresponding ATL values for daily consumption, multiple chemical exposure methodology is employed. This may result in advising the sensitive population to consume fewer meals per week than would be the case for the presence of one chemical alone, in a similar concentration. The potential effect of multiple chemical exposures (mercury and PCBs) was assessed in Common Carp, black bass species, and Rainbow Trout, and did not affect advice. Advice for all species in this advisory was based solely on mercury or PCB concentrations.

OEHHA recommends that individuals strive to meet the US Dietary Guidelines seafood consumption recommendations, while also adhering to federal and OEHHA recommendations to limit the consumption of fish with higher contaminant levels. The advice discussed in the following section represents the maximum recommended number of servings per week for different fish species. People should eat no more than the recommended number of servings for each fish species or species group. OEHHA’s consumption advice for a particular fish species can be extended to other closely related fish species<sup>8</sup> known to accumulate similar levels of contaminants.

---

<sup>8</sup> Fish species within the same genus are most closely related, and Family is the next level of relationship.

Consumption advice should not be combined. That is, if a person chooses to eat a fish from the “one-serving-a-week” category, then they should not eat any other fish from any source (including commercial) until the next week. If a person chooses to eat a fish from the “two-servings-per-week” category, they can combine fish species from that category, or eat one fish from that category and one from a category that recommends more than two-servings-per-week (if available), for a total of two servings in that week. Then they should not eat any other fish from any source (including commercial) until the following week.

## CONSUMPTION ADVICE FOR FISH FROM BIG BEAR LAKE

OEHHA’s advisory protocol requires at least nine fish of a species to be collected from a water body before an advisory can be developed for the primary contaminant of concern. This is to ensure the sample dataset is representative of the fish species population in the water body. In some cases, an exception is made for species that are commonly caught and consumed from a given water body but where available data may be limited. For Big Bear Lake, the sample size criterion was met for the following species: Black Bass species, Common Carp, Crappie, and Rainbow Trout. There were not sufficient data to evaluate other species that may be found in this water body.

### *BLACK BASS SPECIES (LARGEMOUTH AND SMALLMOUTH)*

The mean combined mercury concentration in black bass species from Big Bear Lake was 344 ppb. The mercury levels in individual black bass species were 354 ppb (Largemouth Bass) and 280 ppb (Smallmouth Bass), respectively. The mean PCB concentration in Largemouth Bass was 13 ppb. PCB concentration was not analyzed in Smallmouth Bass. Based on the concentration of mercury in these species, OEHHA recommends a maximum of one serving a week of black bass species for the sensitive population (women 18 to 45 years and children 1 to 17 years), and a maximum of two servings a week for the general population (women 46 years and older, and men 18 years and older).

OEHHA evaluated mercury concentrations in black bass species in many water bodies in California and has found a similar range of mercury concentrations when two or more of these species were caught from the same water body. Therefore, OEHHA extends the consumption advice for Largemouth Bass and Smallmouth Bass to other black bass species that may be found in Big Bear Lake.

### *COMMON CARP*

The mean mercury and PCB concentrations in Common Carp from Big Bear Lake were 151 and 38 ppb, respectively. OEHHA recommends a maximum of one serving a week of Common Carp for the sensitive population (women 18 to 45 years and children 1 to 17 years) based on mercury, and a maximum of two servings a week for the general population (women 46 years and older, and men 18 years and older), based on PCBs.

*CRAPPIE*

The mean mercury and PCB concentrations in Crappie from Big Bear Lake were 41 and 3 ppb, respectively. Based on the concentration of mercury in these crappie, OEHHA recommends a maximum of five servings a week of Crappie for the sensitive population (women 18 to 45 years and children 1 to 17 years) and a maximum of seven servings a week for the general population (women 46 years and older, and men 18 years and older).

*RAINBOW TROUT*

The mean mercury and PCB concentrations in Rainbow Trout from Big Bear Lake were 16 and 12 ppb, respectively. Based on the concentration of PCBs in these Rainbow Trout, OEHHA recommends a maximum of five meals per week for the sensitive (women 18 to 45 years and children 1 to 17 years) and general populations (women 46 years and older, and men 18 years and older).

**RECOMMENDED MAXIMUM NUMBER OF SERVINGS**

The recommended maximum numbers of servings per week for fish from Big Bear Lake are shown in Table 4.

TABLE 4. RECOMMENDED MAXIMUM NUMBER OF SERVINGS PER WEEK FOR FISH FROM BIG BEAR LAKE

<b>Fish Species</b>	<b>Women 18–45 years and Children 1–17 years</b>	<b>Women 46 years and older and Men 18 years and older</b>
Black Bass species	1	2
Common Carp	1	2
Crappie	5	7
Rainbow Trout	5	5

## REFERENCES

American Heart Association. 2014. Fish and Omega-3 Fatty Acids. Online at: <http://www.americanheart.org/presenter.jhtml?identifier=4632>.

Bloom, N.S. 1992. On the chemical form of mercury in edible fish and marine invertebrate tissue. *Can. J. Fish. Aquat. Sci.* 49(5):1010-1017.

FDA/USEPA. 2017. Eating Fish: What pregnant women and parents should know. Advice by FDA and USEPA/January, 2017. Online at: <http://www.fda.gov/downloads/Food/FoodborneIllnessContaminants/Metals/UCM537120.pdf>.

Institute of Medicine. 2007. Seafood choices, balancing benefits and risks. Committee on Nutrient Relationships in Seafood: Selections to Balance Benefits and Risks. Institute of Medicine, Food and Nutrition Board. The National Academies Press, Washington, D.C.

Kris-Etherton, P.M., W.S. Harris, and L.J. Appel. 2002. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circ.* 106:2747-2757.

OEHHA. 2005. General Protocol for Sport Fish Sampling and Analysis. Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Sacramento, California. Online at: <http://oehha.ca.gov/media/downloads/fish/document/fishsamplingprotocol2005.pdf>.

OEHHA. 2008. Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Chlordane, DDTs, dieldrin, methylmercury, PCBs, selenium, and toxaphene. Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Sacramento, California. Online at: <http://oehha.ca.gov/media/downloads/fish/report/atlmhgandothers2008c.pdf>.

OEHHA. 2011. Development of Fish Contaminant Goals and Advisory Tissue Levels for Common Contaminants in California Sport Fish: Polybrominated diphenyl ethers (PBDEs). Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Sacramento, California. Online at: <http://oehha.ca.gov/media/downloads/fish/report/pbdes052311.pdf>.

Oken, E., R.O. Wright, K.P. Kleinman, D. Bellinger, C.J. Amarasiriwardena, H. Hu, J.W. Rich-Edwards, and M.W. Gillman. 2005. Maternal fish consumption, hair mercury, and infant cognition in a U.S. cohort. *Environ. Health Perspect.* 113(10):1376-1380.

Oken, E., J.S. Radesky, R.O. Wright, D. Bellinger, C.J. Amarasiriwardena, K.P. Kleinman, H. Hu, J.W. Rich-Edwards, and M.W. Gillman. 2008. Maternal fish intake

during pregnancy, blood mercury levels, and infant cognition at age 3 years in a U.S. cohort. *Am. J. Epidemiol.* 167(10):1171-1181.

SARWQCB. 2012. Fact Sheet Region 8: Santa Ana Regional Water Board. State Water Resources Control Board. Online at: [https://www.waterboards.ca.gov/water\\_issues/programs/swamp/docs/factsheets/rb8\\_cw101.pdf](https://www.waterboards.ca.gov/water_issues/programs/swamp/docs/factsheets/rb8_cw101.pdf)

SWRCB. 2007. Bioaccumulation of Pollutants in California Waters: A Review of Historic Data and Assessment of Impacts on Fishing and Aquatic Life. State Water Resources Control Board, California Environmental Protection Agency, Sacramento, California. Online at: [http://www.waterboards.ca.gov/water\\_issues/programs/swamp/docs/bop/cw117\\_swrcb\\_report.pdf](http://www.waterboards.ca.gov/water_issues/programs/swamp/docs/bop/cw117_swrcb_report.pdf).

SWRCB. 2010. Contaminants in Fish from California Lakes and Reservoirs, 2007-2008: Summary Report on a Two-Year Screening Survey. State Water Resources Control Board, California Environmental Protection Agency, Sacramento, California. Online at: [http://www.waterboards.ca.gov/water\\_issues/programs/swamp/docs/lakes\\_study/lake\\_survey\\_yr2\\_no\\_app.pdf](http://www.waterboards.ca.gov/water_issues/programs/swamp/docs/lakes_study/lake_survey_yr2_no_app.pdf).

SWRCB. 2013. State Mussel Watch (SMW) Program/Toxic Substances Monitoring (TSM) Program. State Water Resources Control Board, California Environmental Protection Agency, Sacramento, California. Online at: [http://www.waterboards.ca.gov/water\\_issues/programs/swamp/mussel\\_watch.shtml](http://www.waterboards.ca.gov/water_issues/programs/swamp/mussel_watch.shtml).

USDA/USDHHS. 2015. 2015-2020 Dietary Guidelines for Americans. 8<sup>th</sup> Edition. U.S. Government Printing Office, Washington, D.C. December. Online at: <http://health.gov/dietaryguidelines/2015/guidelines/>.

US EPA. 1989. Risk Assessment Guidance for Superfund Volume I: Human Health Evaluation Manual (Part A) Interim Final. EPA/5401-89/002, December 1989. Office of Emergency and Remedial Response, U.S. Environmental Protection Agency, Washington, D.C. Online at: <https://rais.onl.gov/documents/HHEMA.pdf>.

US EPA. 2000a. Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories: Volume 1. Fish Sampling and Analysis. 3<sup>rd</sup> Ed. EPA 823-B00-007. Office of Water, U.S. Environmental Protection Agency, Washington, D.C.

US EPA. 2000b. Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories: Volume 2. Risk Assessment and Fish Consumption Limits, 3<sup>rd</sup> Edition. EPA 823-B-00-007. Office of Water, U.S. Environmental Protection Agency, Washington, D.C.

Weaver, K.L., P. Ivester, J.A. Chilton, M.D. Wilson, P. Pandey, and F.H. Chilton. 2008. The content of favorable and unfavorable polyunsaturated fatty acids found in commonly eaten fish. *J. American Dietetic Assoc.* 108:1178-1185.



## APPENDIX I. ADVISORY TISSUE LEVELS

Advisory Tissue Levels (ATLs) guide the development of advice for people eating sport fish. ATLs are levels of contaminants found in fish that correspond to the maximum numbers of recommended fish servings. OEHHA uses ATLs to provide advice to prevent consumers from being exposed to:

- More than the average daily reference dose<sup>9</sup> for chemicals not known to cause cancer, such as methylmercury, or
- For cancer-causing chemicals, a risk level greater than one additional cancer case in a population of 10,000 people consuming fish at the given consumption rate over a lifetime. This cancer endpoint is the maximum acceptable risk level recommended by the US EPA (2000b) for fish advisories.

For each chemical, ATLs were determined for both cancer and non-cancer risk, if appropriate, for one to seven eight-ounce servings per week. The most health-protective ATLs for each chemical, selected from either cancer or non-cancer based risk, are shown in the table below for zero to seven servings per week. When the guidelines for eating fish from Big Bear Lake are followed, exposure to chemicals in fish from Big Bear Lake would be at or below the average daily reference dose or the cancer risk probability of one in 10,000.

ADVISORY TISSUE LEVELS FOR SELECTED ANALYTES

Contaminant	Consumption Frequency Categories (8-ounce servings/week) <sup>a</sup> and ATLs (in ppb)							
	7	6	5	4	3	2	1	0
Chlordanes	≤ 80	>80-90	>90-110	>110-140	>140-190	>190-280	>280-560	>560
DDTs	≤ 220	>220-260	>260-310	>310-390	>390-520	>520-1,000	>1,000-2,100	>2,100
Dieldrin	≤ 7	>7-8	>8-9	>9-11	>11-15	>15-23	>23-46	>46
MeHg (Women 18-45 and children 1-17)	≤ 31	>31-36	>36-44	>44-55	>55-70	>70-150	>150-440	>440
MeHg (Women > 45 and men)	≤ 94	>94-109	>109-130	>130-160	>160-220	>220-440	>440-1,310	>1,310
PBDEs	≤ 45	>45-52	>52-63	>63-78	>78-100	>100-210	>210-630	>630
PCBs	≤ 9	>9-10	>10-13	>13-16	>16-21	>21-42	>42-120	>120
Selenium	≤ 1000	>1,000-1200	>1,200-1,400	>1,400-1,800	>1,800-2,500	>2,500-4,900	>4,900-15,000	>15,000
Toxaphene	≤ 87	>87-100	>100-120	>120-150	>150-200	>200-300	>300-610	>610

<sup>a</sup> Serving sizes (prior to cooking, wet weight) are based on an average 160-pound person. Individuals weighing less than 160 pounds should eat proportionately smaller amounts.

<sup>9</sup> The reference dose is an estimate of the maximum daily exposure to a chemical likely to be without significant risk of harmful health effects during a lifetime.