

Office of Environmental Health Hazard Assessment



Linda S. Adams
Secretary for Environmental Protection

Joan E. Denton, Ph.D., Director
Headquarters • 1001 I Street • Sacramento, California 95814
Mailing Address: P.O. Box 4010 • Sacramento, California 95812-4010
Oakland Office • Mailing Address: 1515 Clay Street, 16th Floor • Oakland, California 94612



Arnold Schwarzenegger
Governor

MEMORANDUM

TO: Allan Hirsch, Deputy Director
Office of Environmental Health Hazard Assessment

VIA: George V. Alexeeff, Ph.D., D.A.B.T.
Deputy Director for Scientific Affairs

VIA: Anna M. Fan, Ph.D., Chief
Pesticide and Environmental Toxicology Branch

Robert A. Howd, Ph.D., Chief
Water Toxicology Section
Pesticide and Environmental Toxicology Branch

FROM: Hristo Hristov, MD, Ph.D., Staff Toxicologist
Applied Risk Assessment Section
Integrated Risk Assessment Branch

DATE: August 13, 2009

SUBJECT: UPDATE OF PHG – 1,2-DICHLOROBENZENE

Under the Calderon-Sher California Safe Drinking Water Act of 1996, the Office of Environmental Health Hazard Assessment (OEHHA) develops public health goals (PHGs) for regulated chemicals in drinking water and reviews and updates the risk assessments every five years (Health and Safety Code Section 116365(e)(1)). This memorandum represents an update of the literature review and evaluation of the existing PHG for 1,2-dichlorobenzene (OEHHA, 1997). Our re-evaluation supports the previous PHG derivation in 1997, and no new data would justify a significant change to the document.

Summary of Review

OEHHA developed the PHG of 0.6 mg/L (0.6 ppm) for 1,2-dichlorobenzene (also known as *ortho*-dichlorobenzene) and published it in December 1997. 1,2-Dichlorobenzene is used as a solvent in the manufacture of toluene diisocyanate, waxes, rubbers, oils, asphalts, and dyes; as a

California Environmental Protection Agency

The energy challenge facing California is real. Every Californian needs to take immediate action to reduce energy consumption.

degreaser for metals, leather, and wood; as a deodorizing agent for garbage and sewage treatment (OEHHA, 1997); and in the production of chemicals and herbicides (NTP, 1985).

No adequate cancer or genotoxicity studies were identified in the scientific literature. Accordingly, the International Agency for Research on Cancer (IARC) (1982) and U.S. Environmental Protection Agency (U.S. EPA) (1992) defined 1,2-dichlorobenzene as not classifiable as to human carcinogenicity. The 1997 PHG was based on a no-observed-adverse-effect level (NOAEL) of 125 mg/kg-day (the critical effect being hepatotoxicity in rats) estimated in a 13-week subchronic experiment carried out as part of the dose-range finding for the chronic study (NTP, 1985). The PHG is the same as the U.S. EPA's Maximum Contaminant Level Goal (MCLG) and the California and federal Maximum Contaminant Level (MCL) of 0.6 mg/L (U.S. EPA, 1995).

In the most recent toxicity review for 1,2-dichlorobenzene, the Agency for Toxic Substances and Disease Registry (ATSDR) (2006) calculated an oral chronic minimal risk level (MRL) of 0.3 mg/kg-day using the results of the chronic component of the National Toxicology Program (NTP) (1985) study and the benchmark approach. In this NTP study there was a significant increase in renal tubular regeneration in male B6C3F₁ mice only at the highest dose (controls, 8/48; 60 mg/kg-day, 12/50; 120 mg/kg-day, 17/49). ATSDR estimated the benchmark dose associated with 10 percent extra risk (BMDL₁₀) at 30.74 mg/kg-day based on the kidney lesions after exposure for 103 weeks (ATSDR, 2006). This is a lower value than the NOAEL chosen for the 1997 PHG of 125 mg/kg-day, from a subchronic study in F-344 rats that was part of the same NTP (1985) studies.

In the 1997 PHG, OEHHA had rejected the mouse kidney endpoint for risk assessment because concurrent experiments conducted by NTP showed incidences as high for this effect among other control animal populations as for those in the high-dose group exposed to 1,2-dichlorobenzene. In addition, single acute doses of 1,2-dichlorobenzene up to 600 mg/kg caused liver but not kidney damage in mice in the study of Ban *et al.* (1998). Thus, the significance or robustness of the renal endpoint remains unclear.

Other recent studies on 1,2-dichlorobenzene toxicity have addressed the mechanism of hepatotoxicity. Studies of the biochemical correlates of strain-dependent hepatotoxicity in rats (Gunawardhana and Sipes, 1991; Kulkarni *et al.*, 1999; Younis *et al.*, 2000, 2003) have helped explain the critical effects of 1,2-dichlorobenzene in hepatocytes. In addition, Hissink *et al.* (1997) developed a physiologically based pharmacokinetic (PBPK) model for 1,2-dichlorobenzene that they suggested would be useful as "a quantitative tool for evaluating human risk for two different toxicity scenarios, namely covalent binding of reactive metabolites and depletion of GSH" with regard to liver toxicity. The cross-species extrapolation was based on their earlier study comparing 1,2-dichlorobenzene metabolism in rat and human liver microsomes (Hissink *et al.*, 1996), suggesting that humans should be

much less sensitive. However, in our opinion, the uncertainties regarding hepatotoxicity mechanisms as well as in the parameters of human liver metabolism are not yet well-enough established to utilize these data in our risk assessment.

The ATSDR divided its BMDL₁₀ by an uncertainty factor of 100 (10 for interspecies extrapolation, and 10 for human variability) to derive the MRL. Our previous risk estimate used an uncertainty factor of 1,000, with the extra 10-fold related to the fact that our calculation was based on a subchronic study rather than chronic exposure. In addition, our current preference is to use drinking water consumption rates based on the 95th percentile consumption estimates calculated by U.S. EPA (2004) rather than the earlier default rate of 2 L/day.

If we were to calculate a health-protective concentration using the ATSDR BMDL₁₀ approach and a higher drinking water consumption rate, the value would be about double the current PHG of 0.6 mg/L. Keeping the same liver toxicity endpoint and using the higher drinking water consumption rate would decrease the value by about a third. However, neither change seems critical, since this chemical has relatively low-toxicity and is rarely found at significant levels in drinking water.

In conclusion, we have considered several new studies, the revised interpretation of the NTP (1985) mouse data provided by ATSDR (2006), and our new approach to drinking water consumption rates. We conclude that incorporating these new data and interpretations does not provide an adequate rationale for revision of our earlier PHG. Therefore a complete update and revision of the PHG document is considered to be unnecessary.

References

ATSDR (2006). Toxicological Profile for Dichlorobenzenes, Agency for Toxic Substances and Disease Registry, August 2006.

Ban M, Hettich D, Goutet M, Binet S (1998). Serum-borne factor(s) of 1,1-dichloroethylene and 1,2-dichlorobenzene-treated mice inhibited in vitro antibody forming cell response and natural killer cell activity. *Toxicol Lett* 94(2):93-101.

Gunawardhana L, Sipes IG (1991). Dichlorobenzene hepatotoxicity strain differences and structure activity relationships. *Adv Exp Med Biol* 283:731-4.

Hissink AM, Oudshoorn MJ, Van Ommen B, Haenen GR, Van Bladeren PJ (1996). Differences in cytochrome P450-mediated biotransformation of 1,2-dichlorobenzene by rat and man: implications for human risk assessment. *Chem Res Toxicol* 9(8):1249-56.

Hissink AM, Van Ommen B, Krüse J, Van Bladeren PJ (1997). A physiologically based pharmacokinetic (PB-PK) model for 1,2-dichlorobenzene linked to two possible parameters of toxicity. *Toxicol Appl Pharmacol* 145(2):301-10.

IARC (1982). Ortho- and para-dichlorobenzenes. International Agency for Research on Cancer. *IARC Monogr Eval Carcinog Risks Hum* 29:213-38.

Kulkarni SG, Harris AJ, Casciano DA, Mehendale HM (1999). Differential protooncogene expression in Sprague Dawley and Fischer 344 rats during 1,2-dichlorobenzene-induced hepatocellular regeneration. *Toxicology* 139(1-2):119-27.

NTP (1985). Toxicology and carcinogenesis studies of 1,2-dichlorobenzene (o-dichlorobenzene) in F344/N rats and B6C3F1 mice (gavage studies). National Toxicology Program, Technical Report Series 255.

OEHHA (1997). Public Health Goal for 1,2-Dichlorobenzene in Drinking Water. Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, December 1997.

U.S. EPA (1992). 1,2-Dichlorobenzene, carcinogenicity assessment for lifetime exposure. Integrated Risk Information System. U.S. Environmental Protection Agency, Washington, DC.

U.S. EPA (1995). National primary drinking water regulations: o-dichlorobenzene. Office of Water, U.S. Environmental Protection Agency, Washington, DC. EPA 811-F-95-004 e-C.

U.S. EPA (2004). Estimated Per Capita Water Ingestion and Body Weight in the United States - An Update. U.S. Environmental Protection Agency, Washington, DC. EPA-822-R-00-001.

Younis HS, Hoglen NC, Kuester RK, Gunawardhana L, Sipes IG (2000). 1,2-Dichlorobenzene-mediated hepatocellular oxidative stress in Fischer-344 and Sprague-Dawley rats. *Toxicol Appl Pharmacol* 163(2):141-8.

Younis HS, Parrish AR, Sipes IG (2003). The role of hepatocellular oxidative stress in Kupffer cell activation during 1,2-dichlorobenzene-induced hepatotoxicity. *Toxicol Sci* 76(1):201-11.