

PUBLIC NOTICE
Initiation of Risk Assessments for Chemicals in Drinking Water
July 2010

A. Requirements

The Calderon-Sher California Safe Drinking Water Act of 1996 requires the Office of Environmental Health Hazard Assessment (OEHHA) to post notices on its web site of water contaminants for which it is initiating development of public health goals (PHGs) for the chemicals in drinking water. The law also describes the intent and general context of the PHGs. PHGs are concentrations of chemicals in drinking water that are not anticipated to produce adverse health effects following long-term exposures. OEHHA is required to consider potential adverse effects on members of subgroups that comprise a meaningful proportion of the population, including but not limited to infants, children, pregnant women, the elderly, and individuals with a history of serious illness. The public health goals are non-regulatory in nature but are to be used as the health basis to update the state's primary drinking water standards (maximum contaminant levels, or MCLs) established by the California Department of Public Health (DPH) for chemicals subject to regulation.

The act requires PHGs to be developed for the approximately 90 chemicals for which state or federal MCLs are provided, and review and update the risk assessments that form the basis for the PHGs as appropriate at least every five years. Other chemicals may be added to the list by legislative or interdepartmental request. Opportunities for public comment and peer review are provided.

B. Implementation

OEHHA has published 85 PHGs as of July 2010. Two MCLs, for gross alpha and gross beta radionuclides, represent screening levels for contaminants rather than specific regulatory standards; for these, OEHHA has provided risk assessments and guidance memoranda. In addition, re-evaluations of the original PHG have been completed for 18 chemicals. OEHHA concluded that no new information was available on some of these that would require significant changes to the PHG document. The re-reviews of several other chemicals required more extensive efforts, for which complete updated documents have been published. The supporting documents are available at <http://www.oehha.ca.gov/water/phg/index.html>.

PHG review documents for all the other chemicals that have state MCLs are currently in preparation, as well as updates of the PHG for several chemicals. Drafts for public comment on eight of these chemicals (antimony, benzo(a)pyrene, hexavalent chromium, methoxychlor, selenium, styrene, TCDD (dioxin) and trihalomethanes) are currently posted on the OEHHA web site. Draft PHG documents for several more chemicals should be released this year. At the initial posting, a 45-day public comment period is provided, followed by a public

workshop. All comments received are considered in the preparation of a second draft, which is then posted for a 30-day public comment period. After consideration of comments, the final PHG is then published on the OEHHA web site for public reference and use by DPH in developing California MCLs.

Evaluation is now being initiated for two more chemicals for which PHGs were developed earlier (see Section D). These are now being re-reviewed as part of the ongoing PHG update process. Information relevant to the development of PHGs is requested on each of these chemicals.

C. PHGs to be released for public review

Draft documents for updated PHGs for three chemicals (alachlor, fluoride, and perchlorate) are nearing completion, and second postings for methoxychlor, selenium, and trihalomethanes are also being finalized.

Risk assessments are currently in progress for several other chemicals for which initiation of review has previously been announced. Workload considerations and in a few cases, complicated and difficult toxicological interpretations, do not allow us to project completion schedules for these draft updates:

- Atrazine/Simazine
- Cyanide
- Diethylhexylphthalate
- Dibromochloropropane (DBCP)
- 1,4-Dichlorobenzene
- 1,1-Dichloroethylene
- 1,2-Dichloropropane
- Endothall
- Haloacetic acids
- Hexachlorocyclopentadiene
- Nitrate/nitrite
- Picloram
- 1,2,4-Trichlorobenzene
- Trichlorofluoromethane (Freon 11)
- Trichlorotrifluoroethane (Freon 113)

D. Initiation of risk assessments

Risk assessment is being initiated for the following chemicals for which PHGs have already been released, but are being updated after prioritization on the basis of availability of new data, concern about environmental exposures, and potential significance as drinking water contaminants:

- Ethylbenzene
- Xylene

A brief description of these chemicals is provided below. This announcement solicits the submission of pertinent information on these contaminants that could assist our office in updating the risk assessment and deriving a revised PHG.

Information submitted to OEHHA in response to this request should not be proprietary in nature, because all information submitted is a matter of public record. Information should be submitted by September 1, 2010 to:

Michael Baes
PHG Project
Pesticide and Environmental Toxicology Branch
Office of Environmental Health Hazard Assessment
1515 Clay St., 16th floor
Oakland, California 94612

All data submitted will be considered in the development of the PHG for these chemicals. If substantive revisions to the original PHG documents are required, the draft documents will be available for discussion in a public workshop and public comment will be solicited as described above in Section B. The final risk assessments will be utilized by DPH for potential revisions to the MCLs for the chemical in drinking water, as described in more detail on the DPH Web site at <http://www.cdph.ca.gov/certlic/drinkingwater/Pages/MCLsandPHGs.aspx>.

E. Descriptions of chemicals or substances for review initiation:

ETHYLBENZENE

Ethylbenzene (CAS Registry Number: 100-41-4) is a volatile aromatic hydrocarbon derived from petroleum. Ethylbenzene is a natural constituent of crude petroleum and is a standard component of gasoline and diesel fuels. It is also used as a chemical intermediate, primarily in the production of styrene. Ethylbenzene may contaminate drinking water from both natural (oil seeps) and human sources (petroleum spills, waste sites, gasoline storage tank leaks, etc.). Ethylbenzene enters the atmosphere from emissions from industrial facilities and other localized sources, and from evaporation of gasoline in refueling and other activities.

Ethylbenzene was classified as Group 2B (possibly carcinogenic to humans) by the International Agency for Research on Cancer in 2000. The classification of ethylbenzene as a carcinogen was based upon studies by the National Toxicology Program (NTP, 1999) which found clear evidence of a carcinogenic effect in male rats, and some evidence of a carcinogenic effect in female rats and male and female mice. These data were judged appropriate for a satisfactory cancer risk assessment. Ethylbenzene was listed under Proposition 65 as a chemical known to the State of California to cause cancer in 2004. Ethylbenzene

was identified as a Toxic Air Contaminant (TAC) by OEHHA (2007) as a result of its listing as a U.S. Hazardous Air Pollutant.

OEHHA adopted a unit risk value for ethylbenzene of $2.5 \times 10^{-6} (\mu\text{g}/\text{m}^3)^{-1}$ and an inhalation cancer potency value of $0.0087 (\text{mg}/\text{kg}\cdot\text{d})^{-1}$ (OEHHA, 2007). These values are based on the incidence of kidney cancer (renal tubule adenoma or carcinoma) in male rats. They are used in the OEHHA Air Toxics Hot Spots program for estimating the cancer risk associated with inhalation exposures to ethylbenzene. An oral cancer potency value of $0.011 (\text{mg}/\text{kg}\cdot\text{d})^{-1}$ has also been adopted. The support document that describes the derivation of these values has undergone public and peer review and was approved by the Scientific Review Panel for Toxic Air Contaminants. For assessment of non-cancer effects, the OEHHA Air Toxics Program also adopted a Chronic Reference Exposure Level for ethylbenzene of 0.4 ppm in air in 2000, based upon renal, pituitary, and liver effects in both rats and mice (NTP, 1999). Other effects such as hearing loss are also observed (Cappaert *et al.*, 1999; Fechter *et al.*, 2007).

A Proposition 65 “no significant risk level” (NSRL) for ethylbenzene was developed by OEHHA in 2008. A NSRL is defined in regulation as the daily intake level posing a 10^{-5} lifetime risk of cancer. The inhalation and oral human cancer potency estimates for ethylbenzene were estimated as 0.013 and 0.017 $(\text{mg}/\text{kg}\cdot\text{day})^{-1}$, respectively, resulting in corresponding NSRLs of 54 and 41 $\mu\text{g}/\text{day}$ for the two exposure routes (OEHHA, 2008).

Soil screening levels for chemicals are also developed by OEHHA for risk assessments of properties for school sites. A draft has been released on ethylbenzene for this purpose (OEHHA, 2009), based on carcinogenicity and an inhalation cancer potency value of $0.0087 (\text{mg}/\text{kg}\cdot\text{d})^{-1}$ from the OEHHA (2007) evaluation.

The update of the risk assessment for the ethylbenzene PHG will include an evaluation of the studies on which these recent OEHHA risk assessments were based, as well as a broader review of the relevant literature since publication of the original PHG in 1997. A few additional studies, noted below, have been published since the recent OEHHA reviews. This evaluation will be used to estimate appropriate health-protective levels for ethylbenzene in drinking water, with an adequate margin of safety, for all exposed populations.

Selected References

- Cappaert NL, Klis SF, Muijser H, de Groot JC, Kulig BM, Smoorenburg GF (1999). The ototoxic effects of ethyl benzene in rats. *Hear Res* 137(1-2):91-102.
- Chan PC, Hasemani JK, Mahleri J, Aranyi C (1998). Tumor induction in F344/N rats and B6C3F1 mice following inhalation exposure to ethylbenzene. *Toxicol Lett* 99(1):23-32.
- Chen CS, Hseu YC, Liang SH, Kuo JY, Chen SC (2008). Assessment of genotoxicity of methyl-tert-butyl ether, benzene, toluene, ethylbenzene, and xylene to human lymphocytes using comet assay. *J Hazard Mater* 153:351-6.

Cruzan G, Bus J, Banton M, Gingell R, Carlson G (2009). Mouse specific lung tumors from CYP2F2-mediated cytotoxic metabolism: an endpoint/toxic response where data from multiple chemicals converge to support a mode of action. *Regul Toxicol Pharmacol* 55(2):205-18.

De Celis R, Feria-Velasco A, González-Unzaga M, Torres-Calleja J, Pedrón-Nuevo N (2000). Semen quality of workers occupationally exposed to hydrocarbons. *Fertil Steril* 73(2):221-8.

Dennison JE, Bigelow PL, Mumtaz MM, Andersen ME, Dobrev ID, Yang RS (2005). Evaluation of potential toxicity from co-exposure to three CNS depressants (toluene, ethylbenzene, and xylene) under resting and working conditions using PBPK modeling. *J Occup Environ Hyg* 2(3):127-35.

Faber WD, Roberts LS, Stump DG, Beck M, Kirkpatrick D, Regan KS, Tort M, Moran E, Banton M (2007). Inhalation developmental neurotoxicity study of ethylbenzene in Crl-CD rats. *Birth Defects Res B Dev Reprod Toxicol* 80:34-48.

Faber WD, Roberts LS, Stump DG, Tardif R, Krishnan K, Tort M, Dimond S, Dutton D, Moran E, Lawrence W (2006). Two generation reproduction study of ethylbenzene by inhalation in Crl-CD rats. *Birth Defects Res B Dev Reprod Toxicol* 77(1):10-21.

Fechter LD, Gearhart C, Fulton S, Campbell J, Fisher J, Na K, Cocker D, Nelson-Miller A, Moon P, Pouyatos B (2007). Promotion of noise-induced cochlear injury by toluene and ethylbenzene in the rat. *Toxicol Sci* 98:542-51.

Gagnaire F, Langlais C (2005). Relative ototoxicity of 21 aromatic solvents. *Arch Toxicol* 79(6):346-54.

Gagnaire F, Langlais C, Grossmann S, Wild P (2007). Ototoxicity in rats exposed to ethylbenzene and to two technical xylene vapours for 13 weeks. *Arch Toxicol* 81(2):127-43.

Hard GC (2002). Significance of the renal effects of ethyl benzene in rodents for assessing human carcinogenic risk. *Toxicol Sci* 69(1):30-41.

Henderson L, Brusick D, Ratpan F, Veenstra G (2007). A review of the genotoxicity of ethylbenzene. *Mutat Res* 635(2-3):81-9.

López E, Schuhmacher M, Domingo JL (2008) Human health risks of petroleum-contaminated groundwater. *Environ Sci Pollut Res Int* 15:278-88.

Mellert W, Deckardt K, Kaufmann W, van Ravenzwaay B (2007). Ethylbenzene: 4- and 13-week rat oral toxicity. *Arch Toxicol* 81(5):361-70.

NTP (1999). Toxicology and Carcinogenesis Studies of Ethylbenzene (CAS No. 100-41-4) in F344/N Rats and B6C3F1 Mice (Inhalation Studies). National Toxicology Program. Tech Rep Series No. 466. NIH Publication No. 99-3956. U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health, 1999.

OEHHA (2007). Notice of Adoption of Unit Risk Value for Ethylbenzene. Attachment: Long-Term Health Effects of Exposure to Ethylbenzene. Air

Toxicology and Epidemiology Branch, Office of Environmental Health Hazard Assessment, California Environmental Protection Agency. Accessed at: http://www.oehha.ca.gov/air/hot_spots/ebenz111407.html.

OEHHA (2008). No Significant Risk Levels (NASRLs) for the Proposition 65 Carcinogen Ethylbenzene. Reproductive and Cancer Hazard Assessment Branch, Office of Environmental Health Hazard Assessment, California Environmental Protection Agency. Accessed at: www.oehha.ca.gov/prop65/law/pdf_zip/EthylbenzeneNSRL032808.pdf - 2008-03-27

OEHHA (2009). California Human Health Screening Levels for Ethylbenzene. Draft, Nov 2009. Integrated Risk Assessment Branch, Office of Environmental Health Hazard Assessment, California Environmental Protection Agency. Accessed at: www.oehha.ca.gov/risk/pdf/120409Ethylbenzene.pdf.

Saghir SA, Rick DL, McClymont EL, Zhang F, Bartels MJ, Bus JS (2009). Mechanism of ethylbenzene-induced mouse-specific lung tumor: metabolism of ethylbenzene by rat, mouse, and human liver and lung microsomes. *Toxicol Sci* 107(2):352-66.

Saghir SA, Zhang F, Rick DL, Kan L, Bus JS, Bartels MJ (2010). In vitro metabolism and covalent binding of ethylbenzene to microsomal protein as a possible mechanism of ethylbenzene-induced mouse lung tumorigenesis. *Regul Toxicol Pharmacol* 57(2-3):129-35.

Saillenfait AM, Gallissot F, Morel G, Bonnet P (2003). Developmental toxicities of ethylbenzene, ortho-, meta-, para-xylene and technical xylene in rats following inhalation exposure. *Food Chem Toxicol* 41(3):415-29.

Saillenfait AM, Gallissot F, Sabaté JP, Bourges-Abella N, Cadot R, Morel G, Lambert AM (2006). Developmental toxicity of combined ethylbenzene and methylethylketone administered by inhalation to rats. *Food Chem Toxicol* 44(8):1287-98.

Saillenfait AM, Gallissot F, Sabaté JP, Bourges-Abella N, Muller S (2007). Developmental toxic effects of ethylbenzene or toluene alone and in combination with butyl acetate in rats after inhalation exposure. *J Appl Toxicol* 27(1):32-42.

Stott WT, Johnson KA, Bahnemann R, Day SJ, McGuirk RJ (2003). Evaluation of potential modes of action of inhaled ethylbenzene in rats and mice. *Toxicol Sci* 71(1):53-66.

Symanski E, Stock TH, Tee PG, Chan W (2009). Demographic, residential, and behavioral determinants of elevated exposures to benzene, toluene, ethylbenzene, and xylenes among the U.S. population: results from 1999-2000 NHANES. *J Toxicol Environ Health A* 72(14):915-24.

U.S. EPA (2002). Non-Confidential Inventory Update Reporting Production Volume Information. Toxic Substances Control Act (TSCA) Inventory. U.S. Environmental Protection Agency, Washington, DC. Accessed at: <http://www.epa.gov/oppt/iur/tools/data/2002-vol.htm>.

Vyskocil A, Leroux T, Truchon G, Lemay F, Gendron M, Gagnon F, El Majidi N, Viau C (2008). Ethyl benzene should be considered ototoxic at occupationally relevant exposure concentrations. *Toxicol Ind Health* 24(4):241-6.

Zhang M, Wang Y, Wang Q, Yang J, Yang D, Liu J, Li J (2010). Involvement of mitochondria-mediated apoptosis in ethylbenzene-induced renal toxicity in rat. *Toxicol Sci* 115(1):295-303.

XYLENE

Xylene is an organic solvent consisting of a benzene ring with two attached methyl groups. The methyl groups can be in different relative positions on the six-carbon benzene ring, resulting in three slightly different forms, or isomers. Xylene is commonly utilized as a mixture of the three isomers, called ortho-, meta- and para-xylene (o-, m- and p-xylene), which have similar properties and are thus isolated together from their mixed hydrocarbon sources. Xylene is found in small proportions in petroleum and coal tar and also can be formed by the catalytic processes used in a petroleum refinery. Technical grade xylene consists of the three isomers in the proportions of 45 to 55% m-xylene and about 20% of each of the o- and p-isomers, with the remainder being ethylbenzene.

Xylene is one of the most widely used chemicals, with a United States production of several billion pounds per year. It is used in paint thinners, varnishes, as a rubber solvent in the tire industry, in printing and paper manufacturing, as a solvent in the plastic industry and as a component of gasoline and fuel oils. The individual isomers are used in the manufacture of plastics. This widespread use results in great opportunities for spills, leaks and process losses (such as in the drying of paints), as well as potential occupational and residential exposures.

Xylene is volatile and can be rapidly absorbed following both inhalation and oral exposure. Once absorbed, it is relatively rapidly metabolized in the body, mostly in the liver. Its lipophilicity results in uptake into fat and the brain. Exposure to the vapors can cause irritation to eyes, nose, and throat, and sedation at high doses. Xylene does not appear to be carcinogenic in animals or humans. The 1997 OEHA risk assessment for the PHG estimates a health-protective level based on behavioral effects with chronic human exposures to xylene.

In animal studies, toxic effects have not been remarkable. The U.S. EPA reference dose (RfD) of 0.2 mg/kg-day (with a 1000-fold uncertainty factor) is based on decreased body weight and increased mortality in a chronic study in rats, with a No Observed Adverse Effect Level (NOAEL) of 250 mg/kg-day (U.S. EPA, 2003). Xylene exposure can cause ototoxicity (hearing loss) with relatively high exposures (Crofton *et al.*, 1994; Maguin *et al.*, 2006; Gagnaire *et al.*, 2007). A large number of new studies have been published since the earlier PHG review; a selection of these is listed below.

Selected References

- Adams JC, Dills RL, Morgan MS, Kalman DA, Pierce CH (2005). A physiologically based toxicokinetic model of inhalation exposure to xylenes in Caucasian men. *Regul Toxicol Pharmacol* 43(2):203-14.
- Chatterjee A, Babu RJ, Ahaghotu E, Singh M (2005). The effect of occlusive and unocclusive exposure to xylene and benzene on skin irritation and molecular responses in hairless rats. *Arch Toxicol* 79(5):294-301.
- Chen CS, Hseu YC, Liang SH, Kuo JY, Chen SC (2008). Assessment of genotoxicity of methyl-tert-butyl ether, benzene, toluene, ethylbenzene, and xylene to human lymphocytes using comet assay. *J Hazard Mater* 153:351-6.
- Crofton KM, Lassiter TL, Rebert CS. Solvent-induced ototoxicity in rats: an atypical selective mid-frequency hearing deficit. *Hear Res.* 1994 Oct;80(1):25-30.
- Cruzan G, Bus J, Banton M, Gingell R, Carlson G (2009). Mouse specific lung tumors from CYP2F2-mediated cytotoxic metabolism: an endpoint/toxic response where data from multiple chemicals converge to support a mode of action. *Regul Toxicol Pharmacol* 55(2):205-18.
- d'Azevedo PA, Tannhauser M, Tannhauser SL, Barros HM (1996). Hematological alterations in rats from xylene and benzene. *Vet Hum Toxicol* 38(5):340-4.
- De Celis R, Feria-Velasco A, González-Unzaga M, Torres-Calleja J, Pedrón-Nuevo N (2000). Semen quality of workers occupationally exposed to hydrocarbons. *Fertil Steril* 73(2):221-8.
- Dennison JE, Bigelow PL, Mumtaz MM, Andersen ME, Dobrev ID, Yang RS (2005). Evaluation of potential toxicity from co-exposure to three CNS depressants (toluene, ethylbenzene, and xylene) under resting and working conditions using PBPK modeling. *J Occup Environ Hyg* 2(3):127-35.
- Draper TH, Bamiou DE (2009). Auditory neuropathy in a patient exposed to xylene: case report. *J Laryngol Otol* 123(4):462-5.
- Edelfors S, Hass U, Ravn-Jonsen A, Lund SP (1996). The effect of ageing and in vitro exposure to xylene and KCl on $[Ca^{2+}]_i$ in synaptosomes from rats exposed prenatally to xylene. *Pharmacol Toxicol* 78(6):409-12.
- Fay M, Eisenmann C, Diwan S, de Rosa C (1998). ATSDR evaluation of health effects of chemicals. V. Xylenes: health effects, toxicokinetics, human exposure, and environmental fate. *Toxicol Ind Health* 14(5):571-781.
- Foy JW, Schatz RA (2004). Inhibition of rat respiratory-tract cytochrome P-450 activity after acute low-level m-xylene inhalation: role in 1-nitronaphthalene toxicity. *Inhal Toxicol* 16(3):125-32.

- Foy JW, Silverman DM, Schatz RA (1996). Low-level m-xylene inhalation alters pulmonary and hepatic cytochrome P-450 activity in the rat. *J Toxicol Environ Health* 47(2):135-44.
- Fuente A, McPherson B (2006). Organic solvents and hearing loss: The challenge for audiology. *Int J Audiol* 45(7):367-81.
- Gagnaire F, Langlais C (2005). Relative ototoxicity of 21 aromatic solvents. *Arch Toxicol* 79(6):346-54.
- Gagnaire F, Langlais C, Grossmann S, Wild P (2007). Ototoxicity in rats exposed to ethylbenzene and to two technical xylene vapours for 13 weeks. *Arch Toxicol* 81(2):127-43.
- Gagnaire F, Marignac B, Blachère V, Grossmann S, Langlais C (2007). The role of toxicokinetics in xylene-induced ototoxicity in the rat and guinea pig. *Toxicology* 231(2-3):147-58.
- Gralewicz S, Wiaderna D (2001). Behavioral effects following subacute inhalation exposure to m-xylene or trimethylbenzene in the rat: a comparative study. *Neurotoxicology* 22(1):79-89.
- Hass U, Lund SP, Simonsen L (1997). Long-lasting neurobehavioral effects of prenatal exposure to xylene in rats. *Neurotoxicology* 18(2):547-51.
- IARC (1999). Xylenes. *IARC Monogr Eval Carcinog Risks Hum* 71 Pt 3:1189-208.
- Kükner A, Canpolat L, Ozan E, Gökçimen A, Ozan S, Doğrul M (1997-1998). The effect of xylene inhalation on the rat liver. *Acta Physiol Hung* 85(3):231-41.
- Kum C, Sekkin S, Kiral F, Akar F (2007). Effects of xylene and formaldehyde inhalations on renal oxidative stress and some serum biochemical parameters in rats. *Toxicol Ind Health* 23(2):115-20.
- López E, Schuhmacher M, Domingo JL (2008) Human health risks of petroleum-contaminated groundwater. *Environ Sci Pollut Res Int* 15:278-88.
- Maguin K, Lataye R, Campo P, Cossec B, Burgart M, Waniusiow D (2006). Ototoxicity of the three xylene isomers in the rat. *Neurotoxicol Teratol* 28(6):648-56.
- Mao IF, Chang FK, Chen ML (2007). Delayed and competitively inhibited excretion of urinary hippuric acid in field workers coexposed to toluene, ethyl benzene, and xylene. *Arch Environ Contam Toxicol* 53(4):678-83.
- Miller MJ, Edwards JW (1999). Possible preferential metabolism of xylene isomers following occupational exposure to mixed xylenes. *Int Arch Occup Environ Health* 72(2):89-97.
- Morel G, Bonnet P, Cossec B, Morel S, Cour C, Lambert AM, Roure MB, Brondeau MT (1998). The role of glutathione and cysteine conjugates in the nephrotoxicity of o-xylene in rats. *Arch Toxicol* 72(9):553-8.

Park SH, Schatz RA (1999). Effect of low-level short-term o-xylene inhalation on benzo[a]pyrene (BaP) metabolism and BaP-DNA adduct formation in rat liver and lung microsomes. *J Toxicol Environ Health A* 58(5):299-312.

Rabinowitz PM, Galusha D, Slade MD, Dixon-Ernst C, O'Neill A, Fiellin M, Cullen MR (2008). Organic solvent exposure and hearing loss in a cohort of aluminium workers. *Occup Environ Med* 65(4):230-5.

Revilla AS, Pestana CR, Pardo-Andreu GL, Santos AC, Uyemura SA, Gonzales ME, Curti C (2007). Potential toxicity of toluene and xylene evoked by mitochondrial uncoupling. *Toxicol In Vitro* 21(5):782-8.

Saillenfait AM, Gallissot F, Morel G, Bonnet P (2003). Developmental toxicities of ethylbenzene, ortho-, meta-, para-xylene and technical xylene in rats following inhalation exposure. *Food Chem Toxicol* 41(3):415-29.

Sandikci M, Seyrek K, Aksit H, Kose H (2009). Inhalation of formaldehyde and xylene induces apoptotic cell death in the lung tissue. *Toxicol Ind Health* 25(7):455-61.

Symanski E, Stock TH, Tee PG, Chan W (2009). Demographic, residential, and behavioral determinants of elevated exposures to benzene, toluene, ethylbenzene, and xylenes among the U.S. population: results from 1999-2000 NHANES. *J Toxicol Environ Health A* 72(14):915-24.

U.S. EPA (2003). Toxicological Review of Xylenes (CAS No. 1330-20-7), In Support of Summary Information on the Integrated Risk Information System (IRIS). U.S. Environmental Protection Agency, Washington, D.C. January 2003. EPA 635/R-03/001.

Vaidyanathan A, Foy JW, Schatz R (2003). Inhibition of rat respiratory-tract cytochrome P-450 isozymes following inhalation of m-xylene: possible role of metabolites. *J Toxicol Environ Health A* 66(12):1133-43.

Vineis P, Miligi L, Costantini AS (2007). Exposure to solvents and risk of non-Hodgkin lymphoma: clues on putative mechanisms. *Cancer Epidemiol Biomarkers Prev* 16(3):381-4.